

theory model and with the Peaslee excited-intermediate-state model shows that: (i) The predictions of the Fermi theory, in its present formulation, do not agree at this energy with the experimental results. In particular, it predicts a ratio σ_3/σ_1 which is at least 7 times smaller than the experimental one. (ii) The Peaslee model agrees, at least from a qualitative point of view, with experimental results. There is also an indication of the possible existence of an intermediate excited state which decays with $200 \leq Q \leq 250$ Mev, although

the statistics do not allow a positive statement concerning the existence of such an intermediate state.

VII. ACKNOWLEDGMENTS

We wish to thank Dr. M. Kaplon for his advice and encouragement throughout the course of this experiment and Mrs. V. Miller and Mrs. J. Milks for the assistance in scanning the plates. We are also indebted to the Brookhaven Cosmotron staff for their assistance in obtaining the exposure.

Pais-Piccioni Experiment*

K. M. CASE

H. M. Randall Laboratory of Physics, University of Michigan, Ann Arbor, Michigan

(Received April 30, 1956)

Two experiments to check the Gellmann-Pais "particle mixture" suggestion are analyzed, by using a detailed, if crude, phenomenological description. A cloud-chamber experiment of the type proposed by Pais and Piccioni is compared to an experiment using a liquid-xenon bubble chamber. It is found that for likely values of the parameters involved, both experiments are considerably more difficult than envisaged by Pais and Piccioni. For intensity reasons, the bubble-chamber experiment seems preferable. However, if accidentally the relevant cross sections have rather special values, experiments of this type would be particularly easy.

I. INTRODUCTION

RECENTLY Pais and Piccioni¹ have proposed a cloud-chamber experiment to verify the Gellmann-Pais² suggestion that the θ^0 should be considered as a "particle mixture." In view of the availability of a xenon-filled bubble chamber,³ it is of some interest to compare the feasibility of a bubble-chamber experiment with the Pais-Piccioni experiment.

Below, a set of phenomenological equations describing a crude model of these experiments is given. Solutions for the two types of experiments are obtained. From these it follows that for likely values of the parameters involved: (a) both experiments are considerably more difficult than anticipated by Pais and Piccioni; (b) bubble-chamber experiments are somewhat more feasible; (c) if the $\bar{\theta}^0$ absorption cross section and the θ_1^0 , θ_2^0 mass difference are miraculously just right, experiments of this sort will be especially straightforward.

II. PHENOMENOLOGICAL DESCRIPTION

Our model is the following: The wave functions describing θ^0 and $\bar{\theta}^0$ particles are to be linear combinations with prescribed phases of functions describing θ_1^0

and θ_2^0 particles. Specifically, we have

$$\Psi(\theta^0) = (\Psi_1 + i\Psi_2)/\sqrt{2}, \tag{1a}$$

$$\Psi(\bar{\theta}^0) = (\Psi_1 - i\Psi_2)/\sqrt{2}. \tag{1b}$$

The θ_1^0 is to undergo the familiar two- π decay with lifetime $\tau_1 \sim 1.5 \times 10^{-10}$ second. The θ_2^0 has a completely different set of decay modes with a lifetime $\tau_2 \gg \tau_1$. (To obtain detailed numerical results, we will idealize this and put $\tau_2 = \infty$.) In passing through matter, the $\bar{\theta}^0$ can be absorbed while the θ^0 cannot. (For simplicity we follow reference 1 and omit consideration of all other processes.) This absorption will be described by an effective lifetime τ . Clearly,

$$1/\tau = N\rho v\sigma_a, \tag{2}$$

where v is the velocity of the $\bar{\theta}^0$, ρ the density of absorber material, N the number of absorbing particles per gram, and σ_a the absorption cross section per particle.

Let the wave function describing the state at time t after a θ^0 is produced be

$$\Psi(t) = \alpha_1(t)\Psi_1 + \alpha_2(t)\Psi_2. \tag{3}$$

We have

$$\frac{d\Psi(t)}{dt} = \left[\frac{d\Psi(t)}{dt} \right]_{ph} + \left[\frac{d\Psi(t)}{dt} \right]_a + \left[\frac{d\Psi(t)}{dt} \right]_s. \tag{4}$$

Here $[d\Psi(t)/dt]_{ph}$ describes the phase change due to

* Supported in part by the Office of Naval Research, U. S. Navy Department.

¹ A. Pais and O. Piccioni, Phys. Rev. **100**, 1487 (1955).

² M. Gell-Mann and A. Pais, Phys. Rev. **97**, 1387 (1955).

³ Brown, Glaser, and Perl, Phys. Rev. **102**, 586 (1956).

the kinetic energies. Thus, as in reference 1, we have

$$\left[\frac{d\Psi}{dt} \right]_{\text{ph}} = i\omega_1\alpha_1(t)\Psi_1 - i\omega_2\alpha_2(t)\Psi_2, \quad (5)$$

where

$$\omega_i = [c^2p^2 + m_i^2c^4]^{1/2}/\hbar \quad (i=1, 2). \quad (6)$$

The expression $[d\Psi(t)/dt]_d$ is due to the θ^0 decay. Again, as in reference 1, we can write it as

$$\left[\frac{d\Psi}{dt} \right]_d = -\frac{1}{2\tau_1}\alpha_1\Psi_1 - \frac{1}{2\tau_2}\alpha_2\Psi_2. \quad (7)$$

The expression $[d\Psi(t)/dt]_a$ is due to the absorption of θ^0 's. Phenomenologically it is

$$\left[\frac{d\Psi(t)}{dt} \right]_a = -\frac{1}{2\tau} [\text{Component of } \theta^0\text{'s in } \Psi(t)]. \quad (8)$$

Using Eqs. (1) and (3), we obtain

$$\Psi(t) = \frac{1}{\sqrt{2}}[\alpha_1(t) - i\alpha_2(t)]\Psi(\theta^0) + \frac{[\alpha_1(t) + \alpha_2(t)]}{\sqrt{2}}\Psi(\bar{\theta}^0). \quad (9)$$

Hence,

$$\left[\frac{d\Psi(t)}{dt} \right]_a = -\frac{1}{2\tau} \frac{[\alpha_1(t) + i\alpha_2(t)]}{\sqrt{2}}\Psi(\bar{\theta}^0) \quad (10a)$$

$$= -\frac{1}{4\tau}[\alpha_1(t) + i\alpha_2(t)]\Psi_1 + \frac{i}{4\tau}[\alpha_1(t) + i\alpha_2(t)]\Psi_2. \quad (10b)$$

Since

$$\frac{d\Psi(t)}{dt} = \frac{d\alpha_1}{dt}\Psi_1 + \frac{d\alpha_2}{dt}\Psi_2, \quad (11)$$

we obtain, on inserting (5), (7), and (10) into (4) and equating coefficients of Ψ_1 and Ψ_2 ,

$$\frac{d\alpha_1}{dt} = -\left(\lambda_1^0 + \frac{1}{4\tau}\right)\alpha_1 - \frac{i}{4\tau}\alpha_2, \quad (12)$$

$$\frac{d\alpha_2}{dt} = \frac{i}{4\tau}\alpha_1 - \left(\lambda_2^0 + \frac{1}{4\tau}\right)\alpha_2, \quad (13)$$

where

$$\lambda_i^0 = i\omega_i + \frac{1}{2\tau_i} \quad (i=1, 2).$$

III. SOLUTIONS

For initial values $\alpha_1(0)$ and $\alpha_2(0)$, the solution of (12) is

$$\alpha(t) = \begin{pmatrix} \alpha_1(t) \\ \alpha_2(t) \end{pmatrix} = \frac{\alpha_1(0) - R\alpha_2(0)}{1 - R^2} e^{-\lambda_1 t} \begin{pmatrix} 1 \\ R \end{pmatrix} + \frac{\alpha_2(0) - R\alpha_1(0)}{1 - R^2} e^{-\lambda_2 t} \begin{pmatrix} R \\ 1 \end{pmatrix}, \quad (14)$$

where

$$\left. \begin{matrix} \lambda_1 \\ \lambda_2 \end{matrix} \right\} = \frac{\lambda_1^0 + \lambda_2^0}{2} + \frac{1}{4\tau} + \left\{ \pm \frac{1}{2} [(\lambda_1^0 - \lambda_2^0)^2 + 1/4\tau^2]^{1/2} \right\}, \quad (15)$$

and

$$R = -(i/2\tau) \{ (\lambda_1^0 - \lambda_2^0) + [(\lambda_1^0 - \lambda_2^0)^2 + 1/4\tau^2]^{1/2} \}^{-1}. \quad (16)$$

The probability of the two- π decay at time t is

$$p_{2\pi}(t) = |\alpha_1(t)|^2/\tau_1. \quad (17)$$

Hence, the ratio of the intensity of decays at time t after production to those at time zero is

$$\frac{I}{I_0} = \frac{\left| \frac{[\alpha_1(0) - R\alpha_2(0)]}{1 - R^2} e^{-\lambda_1 t} + \frac{[\alpha_2(0) - R\alpha_1(0)]}{1 - R^2} R e^{-\lambda_2 t} \right|^2}{|\alpha_1(0)|^2}. \quad (18)$$

IV. VALUES OF THE PARAMETERS

Measuring time in units of τ_1 , we see that the relative intensities given by (18) depend on the two parameters $\beta = \tau_1/\tau$ and $\gamma = \tau_1\Delta\omega$. Here $\Delta\omega = \omega_1 - \omega_2$ arises from any difference in the masses of θ_1^0 and θ_2^0 . Since such differences presumably would originate from the same interactions which occasion the difference in lifetimes, it seems reasonable that $\gamma \sim 1$. To see the effects of differing values of γ , we have computed the results to be expected for $\gamma = 0, 1$, and 2π .

While the decay lifetime τ_1 is known to be $\sim 1.5 \times 10^{-10}$ seconds, comparably accurate information for the $\bar{\theta}^0$ absorption lifetime τ does not seem to be available at present.⁴

To obtain some idea of τ , let us assume each nucleon can absorb $\bar{\theta}^0$'s with a cross section σ_a . Then

$$\tau = \frac{1}{N_0 \rho v \sigma_a}, \quad (19)$$

where N_0 is Avogadro's number and ρ is the absorber density. Thus,

$$\beta = \frac{\tau_1}{\tau} \sim \left(\frac{\rho}{2.3} \right) \left(\frac{v}{c} \right) \left(\frac{\sigma_a}{10^{-27}} \right) \times 6.2 \times 10^{-3}. \quad (20)$$

Assuming $v \sim c$ and the absorption cross section per nucleon as 1 millibarn, we have

$$\beta(\text{Xe}) \approx 6.2 \times 10^{-3}, \quad (21a)$$

$$\beta(\text{Pb}) \approx 3.0 \times 10^{-2}. \quad (21b)$$

⁴ The evidence of D. M. Fournet and M. Widgoff [Phys. Rev. 102, 929 (1956)] and D. M. Ritson (Proceedings of the Sixth Annual Rochester Conference) seems to indicate the θ^- cross section is approximately geometrical and primarily due to absorption. If the $\bar{\theta}^0$ does, indeed, form an isotopic doublet, σ_a should be almost geometrical.

With either substance we are certainly in a situation of little absorption.

To obtain an extreme upper limit for β , let us assume $\sigma_a \sim 5 \times 10^{-26}$ cm² (i.e., roughly geometrical per nucleon). Then

$$\beta(\text{Xe}) \approx 0.3, \quad (22a)$$

$$\beta(\text{Pb}) \approx 1.5. \quad (22b)$$

From (21) and (22), one can conclude that with either xenon or lead we do not have a situation in which β is very large. Upper limits in the two cases are of the order of $\frac{1}{3}$ and 1, respectively, with somewhat lower values being likely.

V. CLOUD-CHAMBER EXPERIMENT

By this we mean an experiment of the type proposed by Pais and Piccioni. A beam of θ^0 's produced in a lead plate are to decay in a cloud chamber until all θ_1^0 's are removed. The remainder then passes through a lead plate which removes the $\bar{\theta}^0$ component. Since, effectively, θ_1^0 's are produced by removing $\bar{\theta}^0$'s, the familiar two- π decay can be expected to reappear below the second plate. As indicated in reference 1, the intensity of decays below the second plate may be expected to be of the order of $\frac{1}{4}$ of that below the first. However, implicit in the derivation of this result is the assumption that $\tau(\text{Pb}) \ll \tau_1$ (i.e., $\beta \gg 1$). In the other limit of $\beta \ll 1$, qualitatively different results will occur. If $\tau(\text{Pb}) \gg \tau_1$, most θ_1^0 's arising from θ^0 absorption will decay within the lead. The number of decays occurring below the second plate will be very small.

To analyze the situation in detail, we can use Eqs. (12) and (18). On production in the first plate, we have $\alpha_2(0) = i\alpha_1(0)$. After passing through a region with $\tau \sim \infty$ for a time T long compared to τ_1 , we will have

$$\alpha_1(T) = 0, \quad (23)$$

and

$$\begin{aligned} \alpha_2(T) &= \alpha_2(0) \\ &= i\alpha_1(0). \end{aligned} \quad (24)$$

Using these as the initial values when the beam enters a second plate, the ratio of two- π decays occurring at a time t later to those occurring just below the first plate is

$$\begin{aligned} \frac{I(t)}{I(0)} &= \frac{|R|^2}{|1-R^2|^2} \left| \frac{e^{-\lambda_2 t} - e^{-\lambda_1 t}}{1-R^2} \right|^2 = \frac{2|R|^2}{|1-R^2|^2} e^{-(1+\beta)t/2\tau_1} \\ &\quad \times [\cosh(\delta_1 t/\tau_1) - \cos(\delta_2 t/\tau_1)], \end{aligned} \quad (25)$$

where

$$\delta_1 + i\delta_2 = \frac{1}{2} [(1+2i\gamma)^2 + \beta^2]^{\frac{1}{2}}, \quad (26)$$

and

$$R = -\frac{i\beta}{[(1+2i\gamma) + [(1+2i\gamma)^2 + \beta^2]^{\frac{1}{2}}]}. \quad (27)$$

Let us first consider some limiting cases.

(1) $\beta \gg 1$.—Here $\delta_1 \sim \beta$, $\delta_2 \sim 0$, $R \sim -i$, and hence,

$$\frac{I(t)}{I(0)} \approx \frac{1}{2} e^{-(1+\beta)t/2\tau_1} \left[\cosh\left(\frac{\beta t}{2\tau_1}\right) - 1 \right]. \quad (28)$$

The maximum number of θ^0 decays occurring below the plate is clearly obtained by choosing the thickness of the plate so that the time of traversal t_t satisfies

$$\tau \ll t_t \ll \tau_1. \quad (29)$$

In this case we have

$$\frac{I(t_t)}{I(0)} = \frac{1}{4}, \quad (30)$$

(which is just the Pais-Piccioni prediction).

(2) $\beta \ll 1$.—In this case, we have

$$\delta_1 \sim \frac{1}{2}, \quad \delta_2 = \gamma, \quad R = -\frac{i\beta}{2(1+2i\gamma)}. \quad (31)$$

$$\frac{I(t)}{I(0)} \approx \frac{\beta^2}{2|1+2i\gamma|^2} e^{-t/2\tau_1} [\cosh(t/2\tau_1) - \cos(\gamma t/\tau_1)]. \quad (32)$$

Now the maximum number of decays occurring below the plate is of the order of magnitude of the coefficient in (32), i.e.,

$$\left[\frac{I(t)}{I(0)} \right]_{\max} \approx \frac{\beta^2}{2|1+2i\gamma|^2}. \quad (33)$$

Thus, for $\beta = 0.1$ and choosing the most favorable case of $\gamma = 0$, we have

$$\left[\frac{I(t)}{I(0)} \right]_{\max} \approx 0.005, \quad (34)$$

as compared with the value 0.25 of (30).

(3) In Fig. 1 the maximum ratio of the number of decays occurring below the second plate to that below the first is shown as a function of β for $\gamma = 0, 1, 2\pi$. (Maximum means having chosen the thickness of the second plate to make this ratio greatest.) It is striking how slowly the value 0.25 is reached as a function of β . For $\gamma = 2\pi$ and $\beta = 1.5$ [which we have seen is a probable upper limit for $\beta(\text{Pb})$], we have

$$[I/I_0]_{\max} = 0.007.$$

VI. BUBBLE-CHAMBER EXPERIMENT

It should be possible to produce θ^0 's directly within a liquid-xenon bubble chamber. Since one would be able to observe θ^0 's moving in all directions from the point of production, a considerable increase in the number of θ^0 's observed would result as compared with the number observed if one produced them in a lead plate above a cloud chamber. The experiment we have in mind consists of measuring the distance of the θ^0 decay from the




FIG. 1. Cloud-chamber experiment—maximum relative intensity as a function of θ^0 absorption.

point of production. From a knowledge of the velocity of the θ^0 the time of decay is known. Equation (18) [with $\alpha_2(0) = i\alpha_1(0)$] shows that the relative number of decays at time t after production is given by

$$\frac{I(t)}{I(0)} = \frac{e^{-t/2\tau_1}}{[(1-R_1^2+R_2^2)]^2+4R_1^2R_2^2} \{ \},$$

$$\{ \} = [(1+R_2)^2+R_1^2]e^{-(\delta_1+\beta/2)t/\tau_1} + [(R_1^2+R_2^2)][(1-R_2)^2+R_1^2]e^{-(\beta/2-\delta_1)t/\tau_1} + e^{-\beta t/2\tau_1} \{ -2[R_1^2(2-R_2) + R_2(1-R_2^2)] \cos(\delta_2 t/\tau_1) - 2R_1(1-R_1^2-R_2^2-2R_2) \sin(\delta_2 t/\tau_1) \}. \quad (35)$$

Here $R = R_1 + iR_2$ is given by (27) and δ_1 and δ_2 are given by (26). Now, of course, β refers to the xenon.

(1) $\beta \gg 1$.—Equation (35) becomes

$$I(t)/I(0) \cong e^{-t/2\tau_1}. \quad (36)$$

In this limit the θ^0 's would behave as if their lifetime in xenon was double that found in cloud-chamber studies. This would certainly be easy to observe. Unfortunately, the estimates of Sec. IV indicate that we are far from this limit.

(2) $\beta \ll 1$.—In this case,

$$\frac{I(t)}{I(0)} \cong e^{-t/\tau_1} + \frac{\beta^2}{4(1+4\gamma^2)} + \frac{\beta e^{-t/2\tau_1}}{1+4\gamma^2} [\cos(\gamma t/\tau_1) + 2\gamma \sin(\gamma t/\tau_1)]. \quad (37)$$

Thus, only when the number of decays has dropped

quite low will there be any deviations from what one could expect for a particle with lifetime τ_1 .

(3) In Figs. 2, 3, and 4, decay curves for various values of β are given, corresponding to $\gamma = 0, 1$, and 2π , respectively.




FIG. 2. Bubble-chamber experiment—relative decay probability as a function of time. ($\gamma = 0$)

From Fig. 2 ($\gamma=0$), we see that for β small ($\beta \lesssim 1$) significant deviations from $\beta=0$ (no θ^0 absorption) occur only when the number of decays is quite small. Thus, for $\beta=1/3$ there are twice as many decays as for $\beta=0$ only when the latter is approximately 1/70 of its initial value.

Figure 3 ($\gamma=1$) leads to similar conclusions, except for one salient feature. For $\beta \sim 1/3$ there is a tremendous decrease in the number of two- π decays expected in the region where, for no absorption of θ^0 's, we would expect 1% of the decays to occur. While the observation of this effect would be difficult, it is hard to imagine that a decrease by a factor of 100 could not be seen. It should be pointed out that this phenomenon only occurs for a




FIG. 3. Bubble-chamber experiment—relative decay probability as a function of time. ($\gamma=1$.)

region of values $\gamma \sim 1$, $\beta \sim 1/3$. However, the discussion in Sec. IV by no means rules out this possibility.

The curves of Fig. 4 for $\gamma=2\pi$ are quite remarkable. Increasing β from zero to infinity does not change the curves smoothly from the straight line describing the decay e^{-t/τ_1} to that for $e^{-t/2\tau_1}$. As β increases from zero, the decay curves develop characteristic oscillations [due to the sine and cosine terms in (35)]. Since these oscillations probably would be averaged out in an experiment, we have also given curves for this average. These oscillations are most pronounced for $\beta=1$. The decay occurs increasingly rapidly with increasing β until $\beta \cong 12.5$. In this region the decay curves look as if the θ^0




FIG. 4. Bubble-chamber experiment—relative decay probability as a function of time. ($\gamma=2\pi$.)

lifetime is considerably shorter than 1.5×10^{-10} sec. Increasing β further the decay curves swing back up, becoming essentially straight. Two phenomena should be observed:

(a) Extremely large values of β are required before the curve for $\beta=\infty$ is even approximately realized. Thus, $\beta=200$ is clearly far from this limit.

(b) The decay curves for $12.5 \lesssim \beta \lesssim 60$ coincide with the averaged curves for $12.5 > \beta > 0$. For example, the curve for $\beta=40$ falls, within the accuracy with which one can read the figure, exactly on the averaged curve for $\beta=1$. Thus, if one did see an effect of the kind indicated—namely, an apparent increased decay rate—one could only conclude that there are two possible absorption lifetimes τ compatible with the results.

VII. CONCLUSION

We have seen that, in general, for reasonable values of the parameters the “mixture property” of Gell-Mann and Pais will affect only a small proportion of the θ^0 's—in either a bubble chamber or a cloud chamber. Under such circumstances the possibility of observing more θ^0 decays makes the bubble chamber preferable for experiments of this type. If accidentally, $\tau \sim 1$, $\beta \sim 1/3$ for xenon, the bubble chamber experiment would be especially easy.

It is a pleasure to thank Professor D. A. Glaser and Professor G. E. Uhlenbeck for helpful discussions.