

Elastic Scattering of Deuterons by He^{4†}

J. L. GAMMEL, B. J. HILL,* AND R. M. THALER‡
Los Alamos Scientific Laboratory, Los Alamos, New Mexico

(Received February 5, 1960)

A model of the $d+\text{He}^4$ interaction is developed and compared to the data on the ground state of Li⁶ and the $d+\text{He}^4$ elastic scattering data up to 4.5 Mev (deuteron laboratory energy). New phase-shift analyses of the 8- and 10.3-Mev elastic scattering data are made, and quantities relevant to the production or analysis of beams of polarized deuterons are calculated.

I. INTRODUCTION

A MODEL of the $d+\text{He}^4$ interaction is developed. The model is in agreement with the ground-state properties of Li⁶ as deduced by Foldy¹ from the (γ,d) reaction in Li⁶. Phase shifts calculated from this model are in agreement with phase shifts found in Galonsky and McEllistrem's² analysis of Galonsky et al.'s³ experimental data on elastic $d+\text{He}^4$ scattering in the energy range up to 4.5 Mev (laboratory deuteron energy). Phase shifts calculated from the model make it possible to extend the phase-shift analysis to the experimental data of Allred et al. at 10.3 Mev, and Burge et al. at 8 Mev.⁴ From the phase shifts for 8 and 10.3 Mev, we have calculated the quantities relevant to experiments designed to produce or analyze beams of polarized deuterons.

II. THE MODEL

We reduce the problem to a three-body problem by treating He⁴ as a "fundamental particle" or "lump," and imagining the nucleons in the deuteron to interact with He⁴ "lump" via the nucleon-He⁴ optical model potential⁵ and with each other via the nucleon-nucleon potential. We solve this three-body problem only approximately by applying a no distortion approximation; that is, we imagine the deuteron is not distorted in the course of the $d+\text{He}^4$ interaction.

Thus we neglect the possibility of deuteron breakup in the interaction. We hope to estimate the effects of the breakup in future work (in progress with Mr. John H. Christy). Our calculation of the deuteron spin polarization quantities is thus suspect at 8 and 10.3 Mev but may serve as a guide in the planning of experiments.

In order to formulate the model mathematically, we

wrote the deuteron wave function in the form

$$\begin{aligned}\varphi(\mathbf{r}) &= [u(r) + (1/\sqrt{8})w(r)S_{12}]\chi_{1^m}, \\ S_{12} &= (3\boldsymbol{\sigma}_n \cdot \mathbf{r}\boldsymbol{\sigma}_p \cdot \mathbf{r})/r^2 - 1,\end{aligned}\quad (1)$$

where \mathbf{r} is $\mathbf{r}_p - \mathbf{r}_n$, χ_{1^m} is a triplet spin function, and $\boldsymbol{\sigma}_n$ and $\boldsymbol{\sigma}_p$ are spin operators for neutron and proton, respectively. The total wave function is taken to be

$$\Psi = \varphi(\mathbf{r})F(\mathbf{q}), \quad (2)$$

where \mathbf{q} is the vector from the He⁴ lump to the center of gravity of the deuteron

$$\mathbf{q} = \frac{1}{2}(\mathbf{r}_p + \mathbf{r}_n) - \mathbf{r}_{\text{He}}. \quad (3)$$

Substituting Ψ into the Schrödinger equation, we obtain the no distortion approximation by multiplying by $u(r) + w(r)S_{12}$ and integrating over \mathbf{r} . In this way we obtain an equation for $F(\mathbf{q})\chi_{1^m}$, in which $V(\mathbf{q})$ appears as a potential, where

$$\begin{aligned}V(\mathbf{q}) &= \int d\mathbf{r} [u + (1/\sqrt{8})wS_{12}] [2V_c + V_{LS}(\boldsymbol{\sigma}_n + \boldsymbol{\sigma}_p) \\ &\quad \cdot (\mathbf{q} + \frac{1}{2}\mathbf{r}) \times (\frac{1}{2}\boldsymbol{\nabla}_q + \boldsymbol{\nabla}_r)] [u + (1/\sqrt{8})wS_{12}].\end{aligned}\quad (4)$$

In Eq. (4), V_c and V_{LS} (the nucleon He⁴ optical model potentials) have $|\mathbf{q} + \frac{1}{2}\mathbf{r}|$ as argument. Terms with $\mathbf{q} - \frac{1}{2}\mathbf{r}$, which may also appear with $\frac{1}{2}\boldsymbol{\nabla}_q - \boldsymbol{\nabla}_r$, have been transformed by the substitution $\mathbf{r} \rightarrow -\mathbf{r}$ in deriving Eq. (4).

One may wonder about the question of conservation of total angular momentum. The $V(\mathbf{q})$ given by Eq. (4) is an operator; we have not yet made a partial wave expansion of $F(\mathbf{q})\chi_{1^m}$.

We find by performing the azimuthal integrations in Eq. (4) (referring \mathbf{r} to \mathbf{q} as polar axis)

$$\begin{aligned}V(\mathbf{q}) &= V_c(q) + V_{LS}(q)(\boldsymbol{\sigma}_n + \boldsymbol{\sigma}_p) \cdot \mathbf{q} \\ &\quad \times \boldsymbol{\nabla}_q + V_T(q)[(\boldsymbol{\sigma}_n \cdot \mathbf{q}\boldsymbol{\sigma}_p \cdot \mathbf{q})/q^2],\end{aligned}\quad (5)$$

where⁶

$$\begin{aligned}V_c(q) &= 2 \int d\mathbf{r} u'(r) V_c(|\mathbf{q} + \frac{1}{2}\mathbf{r}|) u'(r) \\ &\quad + 2 \int d\mathbf{r} u'(r) V_c(|\mathbf{q} - \frac{1}{2}\mathbf{r}|) u'(r),\end{aligned}$$

⁶ We use the symbols V_c and V_{LS} in two different ways, but there is no reason to confuse them.

† This work done under the auspices of the U. S. Atomic Energy Commission.

* Now at the Southwestern College, Weatherford, Oklahoma.

‡ Now at the Case Institute of Technology, Cleveland, Ohio.

¹ L. L. Foldy (to be published).

² A. Galonsky and M. T. McEllistrem, Phys. Rev. **98**, 590 (1955).

³ A. Galonsky, R. A. Douglas, W. Haerberli, M. T. McEllistrem, and H. T. Richards, Phys. Rev. **98**, 586 (1955).

⁴ J. C. Allred, D. K. Froman, A. M. Hudson, and L. Rosen, Phys. Rev. **82**, 786 (1951). E. J. Burge, H. B. Burrows, and W. M. Gibson, Proc. Roy. Soc. (London) **A210**, 534 (1952).

⁵ J. L. Gammel and R. M. Thaler, Phys. Rev. **109**, 2041 (1958).




FIG. 1. (a) The central term in the $d+\text{He}^4$ interaction. The solid curve is calculated from the model; the dashed curve is an equivalent square well potential adjusted to fit certain data. (b) The tensor and spin orbit terms in the $d+\text{He}^4$ interaction. The solid curves are calculated from the model; the dashed curves are equivalent square well potentials adjusted to fit certain data.

$$V_{LS}(q) = \frac{1}{2} \int d\mathbf{r} u'(r) V_{LS}(|\mathbf{q} + \frac{1}{2}\mathbf{r}|) u'(r) + \frac{1}{4} \int d\mathbf{r} u'(r) V_{LS}(|\mathbf{q} + \frac{1}{2}\mathbf{r}|) u'(r) \cos(\mathbf{q}, \mathbf{r}) - \frac{1}{2} \int d\mathbf{r} u'(r) V_{LS}(\mathbf{q} + \frac{1}{2}\mathbf{r}) w'(r) \sin^2(\mathbf{q}, \mathbf{r}), \quad (6)$$

$$V_T(q) = 4 \int d\mathbf{r} u'(r) V_C(|\mathbf{q} + \frac{1}{2}\mathbf{r}|) w'(r) P_2(\mathbf{q}, \mathbf{r}).$$

In Eq. (6), we have used the substitutions

$$\begin{aligned} u' &= u - w/\sqrt{8}, \\ w' &= 3w/\sqrt{8}. \end{aligned} \quad (7)$$

Also, (\mathbf{q}, \mathbf{r}) means the angle between \mathbf{q} and \mathbf{r} . We have

neglected integrals involving w'^2 (because the percent of D state in the deuteron is small.)

In calculations, we have used for the deuteron wave function⁷

$$\begin{aligned} u(r) &= N \cos\delta [1 - \exp(-\beta x)] \exp(-x), \\ w(r) &= N \sin\delta [1 - \exp(-\gamma x)] \exp(-x) \\ &\quad \{1 + 3[1 - \exp(-\gamma x)]/x \\ &\quad + 3[1 - \exp(-\gamma x)]^2/x^2\}, \quad (8) \\ x &= \alpha r, \end{aligned}$$

$N = 0.875041$, $\gamma = 2.0170$, $\cos\delta = 0.99947$,
 $\beta = 4.7533$, $\sin\delta = 0.03356$, $\alpha = 0.23181750 \text{ f}^{-1}$,
 and for the nucleon He^4 optical model potential⁵

$$V_C(r) = V_C \left[1 + \left(\frac{r}{D} - 1 \right) \exp\left(\frac{r-R}{D} \right) \right], \quad (9)$$

$$V_{LS}(r) = V_{LS} \exp(r-R/D) \left[\quad \right]^2,$$

where (these values are rough average values from reference 5)

$$\begin{aligned} V_C &= 57.6 \text{ Mev}, \\ V_{LS} &= 16.75 \text{ Mev}, \\ R &= 1.775 \text{ f}, \\ D &= 0.8875 \text{ f}. \end{aligned}$$

Graphs of $V_C(q)$, $V_{LS}(q)$, and $V_T(q)$ so calculated are shown in Fig. 1.

III. QUALITATIVE FEATURES OF THE MODEL

The nucleon He^4 optical model predicts a bound state of Li^5 (or He^5) if the exclusion principle is not taken




FIG. 2. ${}^3S_1 + {}^3D_1$ phase shifts calculated from the model as a function of energy.

⁷ L. Hulthén and L. T. Hedin, Kgl. Norske Videnskab. Selskabs, Forh. 31, No. 3 (1958).

into account. When the exclusion principle is taken into account, the bound state disappears because the amplitude of the properly antisymmetrized wave function formed from the unantisymmetrized wave function corresponding to this bound state vanishes. Essentially, one is trying to put three protons (or neutrons) in the 1S shell in antisymmetrizing the wave function. These spurious bound states arise in other problems⁸ and are well understood.

We must anticipate that spurious bound states will arise from the deep central potential shown in Fig. 1. In all, two S_1 bound states and one P_0 , one P_1 , and one P_2 bound states (the first and last with admixtures of D_1 and F_2 states, respectively) are found to exist with the potentials shown in Fig. 1.

The lowest S state is deep (more than 10 Mev of binding), as are the three P states (more than 5 Mev of binding). It is known from the mass defects of d , He⁴, and Li⁶ that the binding energy of Li⁶ relative to sepa-

TABLE I. Phase shifts versus energy calculated from the square well potentials shown in Fig. 1.^a The φ_l are Coulomb phase shifts $\varphi_0=0$, $\varphi_l=\varphi_{l-1}+2 \arctan(\eta/l)$.

E (Mev)	$\delta(P_0)+\frac{1}{2}\varphi_1$	$\delta(P_1)+\frac{1}{2}\varphi_1$	$\delta(P_2)+\frac{1}{2}\varphi_1$	$\delta(D_2)+\frac{1}{2}\varphi_2$	$\delta(D_3)+\frac{1}{2}\varphi_2$
	(radians)				
1	0.3692	0.3749	0.3828	0.6428	0.8114
2	0.1377	0.1530	0.1757	0.4969	0.3531
3	-0.0417	-0.0189	0.0165	0.5509	0.2237
4	-0.1947	-0.1663	-0.1206	1.1065	0.1168
5	-0.3290	-0.2964	-0.2424	-0.9075	0.0226
6	-0.4489	-0.4131	-0.3520	-0.5949	-0.0632
7	-0.5572	-0.5187	-0.4515	-0.5368	-0.1429
8	-0.6558	-0.6150	-0.5422	-0.5435	-0.2178
9	-0.7461	-0.7033	-0.6252	-0.5734	-0.2887
10	-0.8291	-0.7844	-0.7012	-0.6133	-0.3562
11	-0.9057	-0.8592	-0.7710	-0.6576	-0.4207
14	-1.1029	-1.0509	-0.9468	-0.7980	-0.6532

^a The square well potentials in Fig. 1 would give the D phase shifts shown; however, the potential actually used in this calculation is $V_c=22.100$ Mev, $V_{ls}=1.3667$ Mev, $V_\pi=0$, so that the P phase shifts should be slightly different. However, the main point is that the P phase shifts are negative and not much split; their precise magnitude is not important.

rated d and He⁴ is 1.52 Mev, and the binding energy of the second S state agrees closely with this value.

Without further study, we accept the lowest S level and the three P levels as spurious in the sense that if we knew how to antisymmetrize the corresponding wave functions, we would find the antisymmetrized wave function to vanish. We accept the $2S$ level as the ground state of Li⁶.

From a study of the γd reaction in Li⁶, Foldy¹ has concluded that $\langle r^2 \rangle$ in Li⁶ is given approximately by the size of a deuteron, and that $\langle q^2 \rangle$ is much larger than $\langle r^2 \rangle$. We get automatically that $\langle r^2 \rangle$ is given by the size of a deuteron in our model, of course. $\langle q^2 \rangle$ will turn out to be large for two reasons: first, it will be related to the binding energy of d and He⁴ relative to Li⁶ (1.52 Mev) which is less than the deuteron binding energy (2.22 Mev). The fact that the bound state is a $2S$ level means that there is a node in the wave function, which further

⁸ P. Swan, Proc. Roy. Soc. (London) A228, 10 (1955).

TABLE II. Phase-shift analysis for 8 and 10.3 Mev.^a

E	8 Mev		10.3 Mev	
	Starting point	End point	Starting point	End point
$\delta(^3S_1)$	0.9840	0.9489	0.7120	0.7739
$\delta(^3D_1)+\frac{1}{2}\varphi_2$	1.5411	1.7764	2.0516	2.2595
2ϵ	0.3990	0.3990	+0.2500	+0.2500
$\delta(^3D_2)+\frac{1}{2}\varphi_2$	-0.5435	-1.266	-0.6133	-0.6263
$\delta(^3D_3)+\frac{1}{2}\varphi_2$	-0.2178	-0.2965	-0.3562	-0.1760
$\delta(^3P_0)+\frac{1}{2}\varphi_1$	-0.6558	-0.1566	-0.8291	-0.1411
$\delta(^3P_1)+\frac{1}{2}\varphi_1$	-0.6150	-0.1566	-0.7844	-0.1212
$\delta(^3P_2)+\frac{1}{2}\varphi_1$	-0.5422	-0.1566	-0.7012	-0.1627

^a For 8 Mev, the Coulomb phase shifts are $\varphi_1/2=0.1566$ radians, $\varphi_2/2=0.2354$ radian, and for 10.5 Mev, $\varphi_1/2=0.1383$ radian, $\varphi_2/2=0.2077$ radian.

increases $\langle q^2 \rangle$ over what one might expect. We find

$$\begin{aligned} \langle r^2 \rangle &= 11 \text{ f}^2 \text{ calculated} & \langle r^2 \rangle &= 12 \text{ f}^2 \text{ Foldy,} \\ \langle q^2 \rangle &= 15 \text{ f}^2, & \langle q^2 \rangle &= 22 \text{ f}^2, \end{aligned} \quad (11)$$

whereas, as Foldy points out

$$\begin{aligned} \langle r^2 \rangle &= 41.80 \text{ f}^2 \text{ shell model,} \\ \langle q^2 \rangle &= 10.45 \text{ f}^2. \end{aligned} \quad (12)$$

It has to be concluded that the d -He⁴ model of Li⁶ is superior to the shell model in this respect. We now calculate the location of the virtual (unbound) D levels in order to show that the d -He⁴ model also works for these levels. We treat the virtual D levels by calculating scattering phase shifts,⁹ of course.

We anticipate the following qualitative features. The S phase shift starts at 2π at zero energy and decreases with energy; that is, (when the 2π is subtracted from it) the S phase shift will look something like the S phase shift for hard sphere scattering for some radius hard sphere. This behavior agrees with the result of Galonsky and McEllistrem.² The P phase shifts start at π at zero energy and decrease with energy. Because the tensor and spin orbit terms shown in Fig. 1 are weak, compared

TABLE III. Quality of fit to differential cross section at 8 Mev.

θ (c.m.) (degrees)	σ_{exp} (barns)	σ_{calc} (barns)
30	0.3300	0.3520
40	0.1760	0.1838
50	0.0920	0.0820
60	0.0395	0.0385
70	0.0415	0.0395
80	0.0660	0.0623
90	0.0830	0.0837
100	0.0950	0.0900
110	0.0860	0.0822
120	0.0730	0.0734
130	0.0750	0.0809
140	0.1070	0.1153
150	0.1860	0.1747
160	0.2400	0.2431

⁹ The literature on the theory of d -He⁴ scattering (definition and calculation of phase shifts and connection between phase shifts and scattering amplitudes) is reviewed in the Appendix.

TABLE IV. Quality of fit to differential cross section at 10.3 Mev.

θ (c.m.) (degrees)	σ_{exp} (barns)	σ_{calc} (barns)	θ (c.m.) (degrees)	σ_{exp} (barns)	σ_{calc} (barns)
18.4	0.452	0.5445	100.6	0.0679	0.0713
22.0	0.356	0.3797	101.2	0.0709	0.0710
23.0	0.326	0.3479	103.6	0.0668	0.0694
29.4	0.186	0.2093	109.2	0.0616	0.0629
34.0	0.143	0.1460	110.6	0.0587	0.0608
36.6	0.111	0.1176	112.4	0.0576	0.0580
40.8	0.0761	0.0803	114.6	0.0519	0.0542
43.8	0.0571	0.0593	119.8	0.04447	0.0453
48.2	0.0369	0.0361	120.6	0.0418	0.0440
51.0	0.0223	0.0256	124.6	0.0378	0.0381
55.4	0.0105	0.0155	127.2	0.0362	0.0351
58.4	0.00975	0.0125	129.3	0.0370	0.0332
62.5	0.0118	0.0129	133.6	0.0354	0.0312
65.4	0.0171	0.0158	134.4	0.0371	0.0312
69.4	0.0217	0.0225	137.8	0.0400	0.0320
72.2	0.02535	0.0285	140.6	0.0436	0.0340
76.1	0.0343	0.0377	141.7	0.0473	0.0351
78.8	0.04115	0.0444	145.3	0.04655	0.0401
82.6	0.0481	0.0532	147.4	0.05365	0.0437
85.2	0.0548	0.0586	149.0	0.05845	0.0468
85.4	0.05785	0.0590	152.3	0.0706	0.0540
91.6	0.0629	0.0684	154.1	0.07275	0.0582
95.2	0.0694	0.0712	155.4	0.07425	0.0613
97.7	0.0691	0.0718	157.4	0.0824	0.0662

to the central term, there will not be much J splitting unless there is a resonance. Calculation shows there is no resonance and (as expected) little J splitting. This behavior also agrees with the results of Galonsky and McEllistrem, who found small, negative unsplit P phase shifts.

By inspection of Fig. 1, and from a knowledge of the interval rules¹⁰ for the spin orbit and tensor terms in Eq. (5), it can be seen that the order of the D levels as given by Galonsky and McEllistrem is the same as the order given by the $d+\text{He}^4$ model.

Only detailed computation can confirm whether or not the magnitude of the S phase shift is correct, whether or not the D phase shifts pass through 90° at the right energy and with the right level width, and whether or not the tensor coupling of the $^3S_1+^3D_1$ state is negligible

TABLE V. The function t vs θ .

θ (c.m.) (degrees)	4.5 Mev	8 Mev	10.3 Mev
20	0.2155	...	-0.005
30	0.3327	-0.1044	-0.045
40	0.1938	-0.2256	-0.115
50	-0.0899	-0.4663	-0.285
60	-0.3618	-0.6060	-0.240
70	-0.2698	-0.0285	0.150
80	0.1778	0.3170	0.220
90	0.4155	0.3712	0.235
100	0.2675	0.3192	0.250
110	-0.0790	0.1777	0.255
120	-0.2920	-0.0621	0.230
130	-0.1749	-0.2526	0.080
140	0.1300	-0.2323	-0.195
150	0.4247	-0.1285	-0.305
160	0.6339	-0.0465	-0.310
170	0.7527	...	-0.310

¹⁰ See the Appendix for the interval rules.

as Galonsky and McEllistrem assumed. We may summarize our calculations very shortly by saying simply that everything calculated from the $d+\text{He}^4$ model does, in fact, agree with the results of Galonsky and McEllistrem, and that the $d+\text{He}^4$ model makes it possible to extrapolate their work to 8 and 10.3 Mev, at which energies the tensor coupling is important.

IV. SUMMARY OF CALCULATIONS BASED ON THE MODEL

The $^3S_1+^3D_1$ phase shifts, including the coupling constant, are shown as a function of energy in Fig. 2. The coupling parameter 2ϵ is negligible for $E_d < 4.5$ Mev, and the 3S_1 and 3D_1 phase shifts are in reasonable (if not excellent) agreement with the phase shifts of Galonsky and McEllistrem.

Rather than calculating the 3D_2 and 3D_3 phase shifts directly from the model, we have adjusted a "phenomenological" square well potential central, tensor,

TABLE VI. The function u vs θ .

θ (c.m.) (degrees)	4.5 Mev	8 Mev	10.3 Mev
20	0.2449	...	0.23
30	0.4961	0.6615	0.395
40	0.7173	0.9326	0.595
50	0.9201	1.093	0.61
60	0.9944	0.382	-0.40
70	0.7324	-0.740	-0.76
80	0.2773	-0.514	-0.31
90	0	-0.245	0
100	-0.1056	0.433	0.285
110	-0.1726	0.810	0.57
120	-0.2509	0.884	0.875
130	-0.3019	0.423	0.98
140	-0.2934	-0.093	0.52
150	-0.2403	-0.273	0.095
160	-0.1660	-0.243	...
170	-0.0841

and spin orbit terms to fit the following data: the $2S$ level must have the right binding energy (1.52 Mev), the 3D_3 and 3D_2 phase shifts must pass through 90° at the energies given by Galonsky and McEllistrem, and the width of the 3D_3 resonance must agree with the width given by Galonsky and McEllistrem. This last condition determines the range of the square well (we assume that all three terms have the same range) and the first three conditions determine the depth of each term.¹¹ This square well potential is compared with the model potential in Fig. 1. The agreement is reasonable. P and 3D_2 and 3D_3 phase shifts calculated from the square well potential are given in Table I as a function of energy.

We have used phase shifts taken from Table I and Fig. 2 as a starting point for a phase-shift analysis of

¹¹ This is not really so since as indicated in Fig. 1 we have used different central terms in the S and D states. Thus the square well potential in Fig. 1 is "a" potential (and not "the" potential) which fits the D_2 and D_3 phase shifts of Galonsky and McEllistrem (see also remarks in footnote to Table I).

the 8- and 10.3-Mev data. The phase shifts which we find to give the best fit to the data are similar to the phase shifts used as a starting point as shown in Table II; however, the magnitude of the P phase shift is much smaller than expected from Table I, and also smaller than expected from a graphical extrapolation of Galonsky and McEllistrem's P phase shift. The quality of the fit to the data is shown in Tables III and IV; the rms deviation is a little over twice the errors assigned by the experimentalists in the case of 10.3 Mev.

The quantities relevant to experiments designed to produce or analyze beams of polarized deuterons are the expectation values of four operators.¹² These four quantities are denoted differently by various authors.

Wolfenstein's notation ¹³	=	Stapp's notation	
$\langle T_{20} \rangle$	=	$t/\sqrt{2}$	
$\langle T_{11} \rangle$	=	$-u/\sqrt{3}$	(13)
$\langle T_{21} \rangle$	=	$-v/\sqrt{3}$	
$\langle T_{22} \rangle$	=	$-w/2\sqrt{3}$	

TABLE VII. The function v vs θ .

θ (c.m.) (degrees)	4.5 Mev	8 Mev	10.3 Mev
20	0.1139	...	0
30	0.3292	0.0424	0
40	0.5421	0.0910	0
50	0.7426	0.1650	0
60	0.8433	0.1913	-0.06
70	0.6664	0.0043	-0.11
80	0.2916	-0.1044	-0.115
90	0	-0.1421	-0.12
100	-0.2155	-0.1762	-0.125
110	-0.4485	-0.2270	-0.115
120	-0.6624	-0.2794	-0.07
130	-0.7369	-0.2647	0.08
140	-0.6596	-0.1812	0.24
150	-0.5074	-0.1060	0.225
160	-0.3363	-0.0577	0.13
170	-0.1664

The quantities t , u , v , w are given in Tables V-VIII as a function of angle for the two energies and also 4.5 Mev. They have been calculated at lower energies (in the vicinity of the 3D_3 resonance, namely 1.1 Mev) by Pondrom¹⁴ and also at 2.5 and 3.5 Mev by Phillips.¹⁵ We have checked these calculations.

V. SUMMARY

The model we have given for the d -He⁴ interaction may be of interest for d -heavy nucleus interaction.

It is our hope that the analysis of the 8- and 10.3-Mev data has led to correct estimates of the quantities required to plan experiments with polarized deuteron beams.

¹² See the Appendix for an outline of the theory.

¹³ See W. Lakin, Phys. Rev. **98**, 139 (1955).

¹⁴ Lee G. Pondrom, Phys. Rev. Letters **2**, 386 (1959).

¹⁵ R. J. N. Phillips, Phys. Rev. Letters **3**, 101 (1959).

TABLE VIII. The function w vs θ .

θ (c.m.) (degrees)	4.5 Mev	8 Mev	10.3 Mev
20	0.0885	...	0.045
30	0.3802	0.2434	0.163
40	0.8268	0.5699	0.44
50	1.3568	1.115	0.98
60	1.5953	1.226	0.8
70	0.8091	-0.3131	-0.60
80	-0.6021	-1.029	-0.73
90	-1.2460	-1.047	-0.65
100	-0.8842	-0.8240	-0.52
110	0.0782	-0.3703	-0.315
120	0.9736	0.3351	0.055
130	1.2392	0.8482	0.62
140	0.9820	0.7421	0.85
150	0.5860	0.4186	0.52
160	0.2615	0.1743	0.23
170	0.0648

APPENDIX

The partial wave expansion of $F(\mathbf{q})\chi_1^m$ is exactly the same as it is for the triplet state in nucleon nucleon scattering. The interval rule for the term $(\sigma_n + \sigma_p) \cdot \mathbf{q} \times \nabla_q$ is the same as the interval rule for the spin orbit term; namely

$$\frac{1}{2}[J(J+1) - L(L+1) - S(S+1)],$$

and since $\sigma_n \cdot \mathbf{q} \sigma_p \cdot \mathbf{q}/q^2$ is $(S_{12}+1)/3$, its properties when operating on states of definite J and L may be read from a table of the properties of S_{12} (see, for example, Ashkin and Wu¹⁶).

We have used the so called nuclear bar phase shifts in parametrizing the scattering matrix. The formulas for the elements of the scattering matrix in terms of these phase shifts are given by Stapp.¹⁷ Galonsky and McEllistrem introduce 5 scattering amplitudes A , B , C , D , E and give formulas for these in terms of the elements of the scattering matrix (see their Appendix I, final equation). Stapp, on the other hand, uses only 4 scattering amplitudes a , b , c , d . [See his Eq. (22), p. 75.] Therefore Galonsky and McEllistrem's amplitudes must be connected; we find

$$E \sin^2\theta/\sqrt{2} = A - B + \cos\theta(C - D),$$

and

$$b = -\frac{1}{2}(C + D) \sin\theta i,$$

$$d = -(C + D)/2,$$

$$a = \frac{1}{2}(2A + B),$$

$$c = -2A + 2B - \frac{3}{2}\cos\theta + \frac{3}{2}D \cos\theta.$$

This last formula is, indirectly [via Galonsky and McEllistrem's last equation in their Appendix I and Stapp's Eq. (A.20)], an expression for Stapp's amplitudes in terms of the bar phase shifts.

For the calculation of quantities relating to deuteron spin polarization phenomena, we use Stapp's equations on his p. 76 ff.

¹⁶ J. Ashkin and T.-Y. Wu, Phys. Rev. **73**, 982 (1948), Eq. (26).

¹⁷ H. P. Stapp, thesis, University of California Radiation Laboratory Report UCRL-3098 (unpublished), p. 107, Eq. (A.20).