

a volume of dimensions about 1 in.  $\times$   $\frac{1}{2}$  in.  $\times$  20 in. The mean angular spread from this source combined with the original beam spread is shown by the resolution lines in Fig. 3.

### APPENDIX III

Several minor effects in the experiment were estimated and found to be small. Nuclear absorption in

the helium reduced the beam by only 0.1%, but for the large scattering angles ( $44^\circ$  and  $90^\circ$  c.m.) where the protons traversed air rather than He the value was 0.7% to 0.9%. The root mean square scattering angle for the He in the bag was  $0.02^\circ$ , for air in the case of the larger angles  $0.1^\circ$ , for the 0.006 in. of aluminum in the container  $0.043^\circ$  and for styrofoam  $0.024^\circ$  for a 1-cm thick layer.

## Mu-Mesonic Molecules. I. Three-Body Problem\*

STANLEY COHEN,<sup>†</sup> DAVID L. JUDD, AND ROBERT J. RIDDELL, JR.<sup>‡</sup>  
Lawrence Radiation Laboratory, University of California, Berkeley, California

(Received July 27, 1959)

An approximate method is developed for treating a generalized hydrogen-molecule ion in which two heavy particles have positive unit charges and one light particle has a negative unit charge. The expansion parameter of this approximation is the ratio of the light to the heavy mass. In first order, the method requires finding a solution to a pair of ordinary, second-order differential equations, which are coupled unless the masses of the heavy particles are equal. Explicit expressions for the coefficients in these equations are derived. The asymptotic forms of these coefficients for large nuclear separations give to first order the reduced-mass corrections to the binding energy of the light particle on either of the two heavy particles. The usual scattering theory is extended to obtain formulas for the various possible cross sections associated with this system. An iterative, variational technique for obtaining eigenvalues and eigenfunctions for bound states of the system is presented.

### I. INTRODUCTION

THE experimental observation of  $\mu^-$ -meson-induced fusion in a hydrogen bubble chamber<sup>1</sup> has led to an increased interest in the three-body system consisting of a light negatively charged particle in the presence of two heavier positively charged nuclei. This system, the generalized hydrogen molecular ion, has been treated in the past by the approximation of Born and Oppenheimer.<sup>2</sup> In this approximation the expansion parameter is the fourth root of the ratio of the mass of the light particle to that of the heavier particles. For electronic molecules this quantity is small ( $\sim 1/7$ ) and the approximation is sufficiently accurate to be useful in many calculations. For  $\mu$ -mesonic molecules, however, the corresponding value is nearly one ( $\sim \frac{2}{3}$ ), and the approximation is open to question.

In this paper we develop a method based on a variational approximation to the wave function of this three-body system. Although this method has the same starting point as the Born-Oppenheimer approximation

—namely, the solution for the motion of the light particle with the heavy ones held fixed—it leads to an expansion parameter that is the ratio of the masses themselves. In the present approximate treatment first-order terms in this parameter have been included, while second-order ones are ignored. When the masses of the two nuclei are not equal, it is essential that the first-order terms be included, because they lead to the distinctive features of the unequal-mass case. Thus, for example, the difference in binding energy of the light particle on one or the other of the two nuclei is contained in these terms; clearly, if the positions of the nuclei are fixed, their mass differences can play no role. In this unequal-mass case, it will be shown that the wave function of the system is obtained from the solution of a pair of coupled, ordinary, second-order differential equations in which the coupling terms come from the first-order corrections. On the other hand, if the masses of the two nuclei are equal, the pair of equations is uncoupled and the first-order terms serve only to improve the accuracy of the calculation. The development of the equations for the wave functions is given in Sec. II.

In Sec. III, the scattering states for these systems are treated. By use of the asymptotic behavior of the system of equations, explicit expressions for the elastic and exchange cross sections are derived. For unequal nuclear masses one obtains different expressions depending on whether the total energy is less than or

\* This work was performed under the auspices of the U. S. Atomic Energy Commission.

<sup>†</sup> Present address: Argonne National Laboratory, Lemont, Illinois.

<sup>‡</sup> On leave of absence to U. S. Atomic Energy Commission, Washington, D. C.

<sup>1</sup> Alvarez, Bradner, Crawford, Falk-Vairant, Good, Gow, Rosenfeld, Solmitz, Stevenson, Ticho, and Tripp, *Phys. Rev.* **105**, 1127 (1957).

<sup>2</sup> M. Born and J. R. Oppenheimer, *Z. Physik* **46**, 814 (1928); *Z. Physik* **50**, 347 (1928).

greater than the binding energy of the light particle on the lighter nucleus. Finally, in Sec. IV a variational procedure for the determination of the eigenvalues and eigenfunctions of the bound states of the system is given. This method involves an iteration scheme that converges rapidly to the desired eigensolutions.

In a subsequent paper,<sup>3</sup> the techniques that have been developed in this paper will be applied to the problem of muon-catalyzed fusion. It will be seen that close agreement with the experimental results is obtained.

## II. THREE-BODY WAVE FUNCTION

### A. General Equations

In this section we treat the Schrödinger equation for the generalized hydrogen molecular ion consisting of two positively charged nuclei and a light negatively charged particle referred to as a meson. All particles are assumed to have unit electronic charge.

In the development which follows, a convenient choice for a coordinate system is one in which the center-of-mass motion, the relative motion of the two nuclei, and the motion of the meson relative to the center of mass of the two nuclei are separated. If  $\mathbf{r}_1$ ,  $\mathbf{r}_2$ , and  $\mathbf{r}_\mu$  are the position vectors of nucleus 1, nucleus 2, and the meson, respectively, and  $m_1$ ,  $m_2$ , and  $m_\mu$  their masses, then the position of the center of mass,  $\mathbf{r}_c$ , the internuclear separation,  $\mathbf{r}_n$ , and the position of the meson relative to the center of mass of the two nuclei,  $\mathbf{R}_\mu$ , are

$$\mathbf{r}_c = \rho_1 \mathbf{r}_1 + \rho_2 \mathbf{r}_2 + \rho_\mu \mathbf{r}_\mu,$$

$$\mathbf{r}_n = \mathbf{r}_1 - \mathbf{r}_2,$$

$$\mathbf{R}_\mu = \mathbf{r}_\mu - f_1 \mathbf{r}_1 - f_2 \mathbf{r}_2,$$

and

where

$$\rho_i = m_i/M_t = m_i/(m_1 + m_2 + m_\mu) \quad (\text{for } i = 1, 2, \mu),$$

and

$$f_i = m_i/(m_1 + m_2) \quad (\text{for } i = 1, 2).$$

The wave function,  $\Psi$ , for the three-body system satisfies the Schrödinger equation,

$$\left[ -\frac{\hbar^2}{2} \left( \frac{1}{M_t} \nabla_c^2 + \frac{1}{M_n} \nabla_n^2 + \frac{1}{M_\mu} \nabla_\mu^2 \right) + V \right] \Psi = W \Psi,$$

where  $W$  is the energy and  $M_n$  and  $M_\mu$  are the appropriate reduced masses, i.e.,

$$M_n = m_1 m_2 / (m_1 + m_2) \quad \text{and} \quad M_\mu = m_\mu (m_1 + m_2) / M_t.$$

The subscripts on the Laplacians refer to derivatives with respect to the appropriate coordinates.

If all particles have unit charge,  $e$ , the potential  $V$  can be written as

$$V(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_\mu) = \frac{e^2}{r_n} \frac{e^2}{r_{\mu 1}} \frac{e^2}{r_{\mu 2}} = \frac{e^2}{r_n} \frac{e^2}{|\mathbf{R}_\mu - f_2 \mathbf{r}_n|} \frac{e^2}{|\mathbf{R}_\mu + f_1 \mathbf{r}_n|},$$

where  $r_{\mu 1}$  and  $r_{\mu 2}$  are the distances between the meson and nuclei 1 and 2 respectively;  $r_n$  is the magnitude of  $\mathbf{r}_n$ .

The dependence of the wave function on the center-of-mass motion is removed by the usual substitution

$$\Psi(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_\mu) = \exp(i\mathbf{P}_c \cdot \mathbf{r}_c / \hbar) \psi(\mathbf{R}_\mu, \mathbf{r}_n),$$

where  $P_c$  is the momentum associated with the motion of the center of mass. The resultant wave function  $\psi(\mathbf{R}_\mu, \mathbf{r}_n)$  can then be expanded in terms of a complete set of functions,  $\psi_i$ , which are functions of the variable  $\mathbf{R}_\mu$  and may contain the variable  $\mathbf{r}_n$  as an independent parameter. Thus we may write

$$\psi(\mathbf{R}_\mu, \mathbf{r}_n) = \sum_i \psi_i(\mathbf{R}_\mu, \mathbf{r}_n) \chi_i(\mathbf{r}_n).$$

Here the functions  $\chi_i(\mathbf{r}_n)$  are to be determined and are dependent on the choice for the  $\psi_i(\mathbf{R}_\mu, \mathbf{r}_n)$ .

It is convenient to choose for the  $\psi_i$  the complete set of solutions for the wave functions of the meson in the Coulomb potential of the fixed nuclei of unit charge.<sup>4</sup> With such a choice the adiabatic effects of the presence of the meson on the motion of the two nuclei can be replaced by an effective potential. In this case the  $\chi_i(\mathbf{r}_n)$  represent to lowest order the wave functions describing the motion of the two nuclei.

The  $\psi_i$  are therefore the solutions to the equations

$$\mathcal{H}_\mu \psi_i(\mathbf{R}_\mu, \mathbf{r}_n) = W_i(r_n) \psi_i(\mathbf{R}_\mu, \mathbf{r}_n),$$

where

$$\mathcal{H}_\mu = -\frac{\hbar^2}{2M_\mu} \nabla_\mu^2 - \frac{e^2}{r_{\mu 1}} - \frac{e^2}{r_{\mu 2}},$$

and  $W_i(r_n)$  is the energy associated with this system as a function of the parameter  $r_n$ .

If we insert the expansion for  $\psi(\mathbf{R}_\mu, \mathbf{r}_n)$  into the Schrödinger equation, multiply by  $\psi_j$ , and integrate over all values of  $\mathbf{R}_\mu$ , we obtain a set of equations

$$-\frac{\hbar^2}{2M_n} \int \psi_j(\mathbf{R}_\mu, \mathbf{r}_n) \nabla_n^2 \sum_i \chi_i(\mathbf{r}_n) \psi_i(\mathbf{R}_\mu, \mathbf{r}_n) d^3 R_\mu + \left[ W_j(r_n) + \frac{e^2}{r_n} \right] \chi_j(\mathbf{r}_n) = W \chi_j(\mathbf{r}_n),$$

where  $W$  is now the energy of the three-body system in its center of mass. When the indicated differentiations

<sup>3</sup> S. Cohen, D. L. Judd, and R. J. Riddell, Phys. Rev. **117**, 384 (1960).

<sup>4</sup> The ground state and some of the excited states for this system have been studied, e.g., by E. A. Hylleraas, Z. Physik **71**, 739 (1931); Edward Teller, Z. Physik **61**, 458 (1930); and Bates, Ledsham, and Stewart, Phil. Trans. Roy. Soc. (London) **A246**, 215 (1953-4).

are carried out, the first term in this equation may be rewritten as

$$\begin{aligned} & \int \psi_j(\mathbf{R}_\mu, \mathbf{r}_n) \sum_i \nabla_n^2 \chi_i(\mathbf{r}_n) \psi_i(\mathbf{R}_\mu, \mathbf{r}_n) d^3 R_\mu \\ &= \nabla_n^2 \chi_j + 2 \sum_i \nabla_n \chi_i \cdot \int \psi_j(\mathbf{R}_\mu, \mathbf{r}_n) \nabla_n \psi_i(\mathbf{R}_\mu, \mathbf{r}_n) d^3 R_\mu \\ & \quad + \sum_i \chi_i \cdot \int \psi_j(\mathbf{R}_\mu, \mathbf{r}_n) \nabla_n^2 \psi_i(\mathbf{R}_\mu, \mathbf{r}_n) d^3 R_\mu. \end{aligned}$$

Here the last term can be made more symmetric by an integration by parts:

$$\begin{aligned} \int \psi_j \nabla_n^2 \psi_i d^3 R_\mu &= \nabla_n \cdot \int \psi_j \nabla_n \psi_i d^3 R_\mu \\ & \quad - \int (\nabla_n \psi_j) \cdot (\nabla_n \psi_i) d^3 R_\mu. \end{aligned}$$

Finally, if we define

$$\mathbf{f}_{ij} \equiv \int \psi_i \nabla_n \psi_j d^3 R_\mu = -\mathbf{f}_{ji},$$

and

$$g_{ij} \equiv \int (\nabla_n \psi_i) \cdot (\nabla_n \psi_j) d^3 R_\mu = g_{ji},$$

the set of Schrödinger equations becomes

$$\begin{aligned} -\frac{\hbar^2}{2M_n} \{ \nabla_n^2 \chi_j(\mathbf{r}_n) + \sum_i [2\mathbf{f}_{ji} \cdot \nabla_n \chi_i + (\nabla_n \cdot \mathbf{f}_{ji}) \chi_i - g_{ji} \chi_i] \} \\ + \left( W_j(\mathbf{r}_n) + \frac{e^2}{r_n} \right) \chi_j(\mathbf{r}_n) = W \chi_j(\mathbf{r}_n). \end{aligned}$$

For convenience, we introduce the dimensionless parameters

$$\mathbf{x} = \mathbf{r}_n / a_\mu, \quad v = W / W_\mu,$$

where  $a_\mu$  is the Bohr radius for the mesonic atom having a reduced mesonic mass  $M_\mu$ , [ $a_\mu = \hbar^2 / (M_\mu e^2)$ ], and  $W_\mu$  is the corresponding mesonic Rydberg, [ $W_\mu = e^4 M_\mu / (2\hbar^2)$ ]. All distances are measured in units of  $a_\mu$  and all energies in units of  $W_\mu$ . The dimensionless Schrödinger equations are then

$$\begin{aligned} -\frac{M_\mu}{M_n} \{ \nabla_n^2 \chi_j(\mathbf{r}_n) + \sum_i [2\mathbf{f}_{ji} \cdot \nabla_n \chi_i + (\nabla_n \cdot \mathbf{f}_{ji}) \chi_i - g_{ji} \chi_i] \} \\ + \left[ W_j(\mathbf{r}_n) + \frac{2}{r_n} \right] \chi_j(\mathbf{r}_n) = W \chi_j(\mathbf{r}_n), \end{aligned}$$

where the definitions of the symbols have been altered to refer to the dimensionless variables. We may write these equations as

$$\left\{ \nabla_n^2 + \frac{M_n}{M_\mu} \left[ W - W_j(\mathbf{r}_n) - \frac{2}{r_n} \right] \right\} \chi_j(\mathbf{r}_n) = -\sum_i \Theta_{ji} \chi_i(\mathbf{r}_n), \quad (1)$$

where

$$\Theta_{ji} = 2\mathbf{f}_{ji} \cdot \nabla_n + (\nabla_n \cdot \mathbf{f}_{ji}) - g_{ji}.$$

In the lowest Born-Oppenheimer approximation to the solution to these equations the dynamic correction terms  $\Theta_{ji}$  are assumed to be zero and only the  $\chi_j$  corresponding to the lowest  $W_j$  is retained. Furthermore, the effective potential  $W_j(\mathbf{r}_n) + 2/r_n$  is expanded about its minimum value in a power series in the displacement of  $\mathbf{r}_n$  from its value at this minimum. While such approximations are reasonable for the treatment of the electronic molecular ions, the larger mass of the meson present in the mesonic molecular ions makes these approximations less reliable.

An alternative approach developed here includes the lowest-order dynamic corrections and makes use of exact solutions to a simplified set of Schrödinger equations. In this treatment it is necessary to separate the cases for identical and distinguishable nuclei. If the nuclei are identical, the wave functions must be either symmetric, +, or antisymmetric, -, with respect to an interchange of the two nuclei, and the two types are not coupled in the set of equations; i.e.,  $\mathbf{f}_{ij}$  and  $g_{ij}$  are zero if  $i$  and  $j$  correspond to states of opposite symmetries. If the nuclei are not identical these terms do not vanish, and furthermore, since there is a degeneracy between the unperturbed symmetric and antisymmetric energies for large nuclear separation (corresponding to the equality of binding energy of the meson on either of the two fixed charge centers), it is necessary to include states of both symmetries in the wave function.

In our treatment we restrict ourselves to treating only the states corresponding to the lowest values of  $W_i(\mathbf{r}_n)$ , designated by a zero subscript, for either of the two possible symmetries. Therefore for distinguishable (unequal-mass) nuclei our wave function is of the form

$$\psi = \psi_0^+ \chi_0^+ + \psi_0^- \chi_0^-,$$

while for the equal-mass case it is

$$\text{either } \psi = \psi_0^+ \chi_0^+ \quad (\text{symmetric case}),$$

$$\text{or } \psi = \psi_0^- \chi_0^- \quad (\text{antisymmetric case}).$$

The errors introduced by the omission of the higher excited mesonic states cannot be accurately determined. We may, however, estimate these errors to some extent by use of a simple perturbation expansion. If we consider the states of lowest  $W_i$  to be the dominant ones and treat  $M_\mu / M_n = \epsilon$  as an expansion parameter, then for the equal-mass case we may write

$$\chi_0 = \chi_0^{(1)} + \epsilon \chi_0^{(2)} \dots,$$

$$\chi_i = \epsilon \chi_i^{(1)} + \epsilon^2 \chi_i^{(2)} \dots, \quad \text{for } i \neq 0,$$

where  $\chi_0$  is the state corresponding to the lowest  $W_i$ . Inserting these expressions in the coupled equations and considering only those terms of the lowest power

in  $\epsilon$ , we obtain

$$\chi_i^{(1)} = (W - W_i - 2r_n)^{-1} \epsilon \mathcal{O}_{i0} \chi_0^{(1)};$$

hence, to lowest order in  $\epsilon$ ,

$$\chi_i = (W - W_i - 2r_n)^{-1} \epsilon \mathcal{O}_{i0} \chi_0.$$

If we consider the effects of the  $i$ th excited state on the equation for  $\chi_0$ , we note that they enter the equation only in second order in  $\epsilon$ . Thus the omission of the excited states introduces errors of order  $\epsilon^2$  in the calculation of the energy of bound states. Similarly, for free states at energies such that mesonic excitations are energetically impossible even for large nuclear separations, only  $\chi_0$  is necessary to determine cross sections. The errors in these cross sections are also of order  $\epsilon^2$ . In addition, for the treatment of bound states and scattering states of low energy, the denominator in the above expression is in general large due to the large separation of the excited states of the meson in the molecular ions which we shall treat.

In the unequal-mass case, a small perturbation can cause large changes in the wave function. It is therefore necessary to treat the two states of lowest  $W_i$  corresponding to opposite symmetries,  $\chi_0^+$  and  $\chi_0^-$ , together. The effects of the remaining higher states will, as before, introduce second-order corrections to the binding energies and cross sections.

### B. Solution for $\psi_i$

Although it is possible in principle to obtain exact numerical solutions to the mesonic problem with two fixed centers,<sup>5</sup> the ultimate accuracy of our approximations has been shown to be limited. We therefore felt justified in using approximate variational solutions for this part of the problem.

For the symmetric solution  $\psi_0^+$ , we assumed a variational solution of the form<sup>6</sup>

$$\psi_0^+ = A_+ \cosh(q_+ r_n \eta / 2) e^{-p_+ r_n \xi / 2},$$

where  $A_+$  is the normalization constant and  $p_+$  and  $q_+$  are variational parameters which minimize the expectation value of  $\mathcal{H}_\mu$  for a given value of  $r_n$ . The variables  $\xi$  and  $\eta$  are the usual confocal elliptic coordinates  $\xi = (r_{\mu 1} + r_{\mu 2}) / r_n$  and  $\eta = (r_{\mu 1} - r_{\mu 2}) / r_n$ .

A similar function was chosen for the antisymmetric solutions, i.e.,

$$\psi_0^- = A_- \{\xi\} \sinh(q_- r_n \eta / 2) e^{-p_- r_n \xi / 2}.$$

Here the  $\{\xi\}$  indicates that a factor of  $\xi$  was included in the expression if this led to a lower expectation value. Specifically, for values of  $r_n$  less than  $r_c$  ( $r_c \sim 1.70$ ) this factor was included; for values of  $r_n$  larger than this it was omitted. For small values of  $r_n$  this additional

factor is essential in order for the solution to approach the hydrogenlike  $2p$  function as  $r_n$  approaches zero. In the neighborhood of  $r_c$  the two sets of solutions were smoothly joined.

The desired expectation values can be expressed as the sum of two terms,

$$W_i = \int \psi_i \mathcal{H}_\mu \psi_i dt_\mu = \langle T_\mu \rangle + \langle V_\mu \rangle,$$

where

$$\langle T_\mu \rangle = \int \psi_i T_\mu \psi_i dt_\mu = \int \nabla_\mu \psi_i \cdot \nabla_\mu \psi_i dt_\mu,$$

and

$$\langle V_\mu \rangle = \int \psi_i V_\mu \psi_i dt_\mu = \int \psi_i \left[ -2 \left( \frac{1}{r_{\mu 1}} + \frac{1}{r_{\mu 2}} \right) \right] \psi_i dt_\mu.$$

For confocal elliptic coordinates the volume element is

$$dt_\mu = \frac{1}{8} r_n^3 (\xi^2 - \eta^2) d\xi d\eta d\phi,$$

where the limits on the variables are

$$1 \leq \xi < \infty, \quad -1 \leq \eta \leq 1, \quad \text{and} \quad 0 \leq \phi < 2\pi.$$

The integrands in the above expressions may also be expressed in these coordinates, i.e.,

$$V_\mu = -\frac{8}{r_n} \frac{\xi}{\xi^2 - \eta^2},$$

and

$$\nabla_\mu \psi_i \cdot \nabla_\mu \psi_i = \frac{4}{r_n^2} \left[ (\xi^2 - 1) \left( \frac{d\psi_i}{d\xi} \right)^2 + (1 - \eta^2) \left( \frac{d\psi_i}{d\eta} \right)^2 \right].$$

The integrals that occur in these and other expressions in this paper can be conveniently expressed in terms of the definite integrals

$$E_n = E_n(P) = \int_1^\infty e^{-P\xi} \xi^n d\xi,$$

$$C_{2n} = C_{2n}(Q) = \int_{-1}^1 \eta^{2n} \cosh^2(Q\eta/2) d\eta,$$

$$C_{2n}' = C_{2n}'(Q) = \int_{-1}^1 \eta^{2n} \sinh^2(Q\eta/2) d\eta,$$

and

$$C_{2n+1} = C_{2n+1}' = \int_{-1}^1 \eta^{2n} \sinh(Q\eta/2) \cosh(Q\eta/2) d\eta,$$

where  $P = pr_n$  and  $Q = qr_n$ .

For the symmetric solution the explicit results are

$$\langle V_\mu \rangle_+ = -2\pi r_n^2 A_+^2 E_1 C_0,$$

$$\langle T_\mu \rangle_+ = \frac{1}{4} \pi r_n A_+^2 [P_+^2 (E_2 - E_0) C_0 + Q_+^2 (C_0' - C_2') E_0].$$

The normalization constant  $A_+$  is determined by the relationship

$$\frac{1}{4} \pi A_+^2 r_n^3 [E_2 C_0 - E_0 C_2] = 1.$$

<sup>5</sup> E. A. Hylleraas, reference 4.

<sup>6</sup> This form was used in treating the hydrogen molecular ion by V. Guillemin, Jr., and C. Zener, Proc. Natl. Acad. Sci. U. S. 15, 314 (1929).




FIG. 1. Static molecular-ion potentials for the lowest symmetric ( $W_+$ ) and antisymmetric ( $W_-$ ) mesonic states. Here  $\blacksquare$  indicates values obtained by Teller,<sup>4</sup> and  $\bullet$  the exact values of Hylleraas.<sup>4</sup>

Similar expressions can be obtained for the antisymmetric solutions.

The minimization of the expectation values of  $\mathcal{H}_\mu$  for values of  $r_n$  between 0 and 20 in intervals of 0.05 was carried out with the aid of an IBM 650 digital computer. The expression for  $W_i$  was minimized to an accuracy of eight figures; however, because of the extremal properties of  $W_i$ , errors due to rounding made the determination of  $P$  and  $Q$  less accurate. The results of these calculations are given graphically in Figs. 1 and 2. For comparison the results of some previous calculations of the values of  $W_i$  are included in Fig. 1. For the symmetric case with our approximate solutions we obtain  $W_i = -1.20489$  for a value of  $r_n$  of 2.00. This is to be compared to the exact value of  $-1.20527$  obtained by Hylleraas.<sup>5</sup> Similar agreement is found for the other values calculated by him. In view of the other




FIG. 2. Mesonic wave-function parameters,  $p_\pm(r) = P_\pm(r)/r$  and  $q_\pm(r) = Q_\pm(r)/r$ , which minimize the static Hamiltonian. For the antisymmetric states, different curves are presented for  $r < r_c$ ,  $r > r_c$  (see text).

approximations made in these calculations we felt that this close agreement indicated a satisfactory solution for the mesonic part of the wave function.

By making use of the analytic forms for the mesonic wave function it was possible to analytically evaluate the first-order dynamic correction terms,  $f_{ij}$  and  $g_{ij}$ , discussed in the preceding section. The specific analytic forms for these terms are given in the Appendix.

For the equal-mass case only "diagonal" correction terms occur, because the states of different symmetries are not coupled. As a consequence of the relationship  $f_{ij} = -f_{ji}$  it follows that  $f_{ii} \equiv 0$ . Thus the only first-order correction for the equal-mass case can be considered as a correction to the potential. This term is of the form

$$g_{ii} = \int \nabla_n \psi_i \cdot \nabla_n \psi_i dt_\mu,$$




FIG. 3. First-order dynamic corrections to the molecular-ion potentials for the equal-mass case.

where  $i$  denotes either  $+$  or  $-$ . This correction has been computed numerically by using the parameters obtained from the variational calculation. The results for both the symmetric and antisymmetric states in the equal-mass case are shown in Fig. 3. There has been a certain amount of controversy concerning these corrections; our results for the symmetric case are in general agreement with those of Dalgarno and McCarroll.<sup>7</sup>

For the unequal-mass case in which the ratio of the two masses is 1:2, similar diagonal correction terms were computed. In addition, the off-diagonal terms were obtained for this case. The results for these calculations are shown in Figs. 4 and 5.

<sup>7</sup> A. Dalgarno and R. McCarroll, Proc. Roy. Soc. (London) A237, 383 (1956). The results of T. Y. Wu, J. Chem. Phys. 24, 444 (1956), and T. Y. Wu and A. B. Bhatia, J. Chem. Phys. 24, 48 (1956) are in disagreement with ours.

**C. Behavior for Small  $r_n$**

The study of the behavior of the mesonic solutions for small values of  $r_n$  is of considerable interest, both for the general understanding of the three-body problem and for the development of solutions to the differential equations for  $\chi_i(r_n)$ .

The behavior of the parameters  $p$  and  $q$  in this limit can be obtained by expressing the energy  $W_i$  in powers of the parameters  $r_n$ ,  $P$ , and  $Q$ . For the symmetric case to lowest order in  $P_+$  and  $Q_+$  we have

$$-W_+ r_n^2 \cong -P_+^2 + 4P_+ r_n - \frac{P_+^2 Q_+^4}{240} - (P_+^2 - 4P_+ r_n) \frac{P_+^2 Q_+^2}{90}.$$

Minimizing this expression for  $W_+$  with respect to the




FIG. 4. First-order dynamic correction terms to the molecular-ion potentials for the unequal-mass case ( $m_1/m_2=2$ ).

parameters  $P_+$  and  $Q_+$ , we find for the limit as  $r_n \rightarrow 0$ ,

$$P_+ \rightarrow 2r_n, \quad p_+ \rightarrow 2, \\ Q_+ \rightarrow (4/\sqrt{3})r_n, \quad q_+ \rightarrow 4/\sqrt{3},$$

and

$$W_+ \rightarrow -4.$$

These results are consistent with the hydrogenlike  $1s$  solution which would be expected in this limiting case. We note that the energy is relatively insensitive to the parameter  $q_+$  in this region, occurring in terms of order  $r_n^4$ .

In a similar manner we may obtain limiting values for the parameters  $p_-$  and  $q_-$  for the antisymmetric solution. In this case we have

$$-W_- r_n^2 = (-P_-^2 + 2r_n P_-) \left( 1 + \frac{P_-^2 Q_-^2}{2100} \right) - \frac{P_-^2 Q_-^2}{2520},$$




FIG. 5. First-order dynamic coupling terms between the lowest symmetric and antisymmetric molecular-ion states for the unequal-mass case ( $m_1/m_2=2$ ).

from which it follows, for the limit as  $r_n \rightarrow 0$ ,

$$P_- \rightarrow r_n, \quad p_- \rightarrow 1, \\ Q_- \rightarrow (\sqrt{3}/5)r_n, \quad q_- \rightarrow \sqrt{3}/5,$$

and

$$W_- \rightarrow -1,$$

which indicates that our solution  $\psi_-$  approaches a hydrogenlike  $2p$  solution with  $m_z=0$ , where the  $z$  axis is aligned in the direction of  $\mathbf{r}_n$ .

The asymptotic forms for  $g_{++}$  and  $g_{--}$  can be readily obtained from the complete expressions given in the Appendix. It is found that while  $g_{++}$  tends to zero in the limit of vanishing  $r_n$ ,  $g_{--}$  is divergent, having a leading term of the form  $2/r_n^2$ . This asymptotic behavior is in fact necessary for a consistent set of solutions to the three-body system for a state in which the total angular momentum is zero. We have already seen that  $\psi_-$  approaches a  $p$  state as  $r_n$  tends to zero, hence for the total angular momentum of the system to be preserved the two nucleons must be in a relative  $p$  state. This angular dependence must be carried entirely by  $\chi_-(\mathbf{r}_n)$ , because  $\psi_-$  (or  $\psi_+$ ) is a function only of the parameters  $r_\mu, r_n$ , and  $\mathbf{r}_\mu \cdot \mathbf{r}_n$  and hence is invariant with respect to rotations of the entire system. For  $s$  states the radial wave functions,  $\chi_-$ , associated with  $\psi_-$  satisfy an equation which in the limit of small  $r_n$  is of the form

$$\frac{1}{r_n^2} \frac{d}{dr_n} \left( r_n^2 \frac{d\chi_-}{dr_n} \right) + \frac{2}{r_n^2} \chi_- + \text{terms of order } \frac{1}{r_n} = 0,$$

where the singularity in  $g_{--}$  provides the term necessary to correct the form for the  $\chi_-$  equation to agree with that of the usual  $p$ -state equation.

For the case in which the total angular momentum of the system is one, the situation is somewhat less clear. In this case, if the meson is in a  $p$  state it is necessary only that the nuclei be in relative  $s$  or  $d$  states. With our choice of approximate wave functions we have in fact chosen a linear combination of these states such that the potential for small  $r_n$  is  $4/r_n^2$ . A similar situation arises for states of higher total angular momentum, so that for small values of  $r_n$  the wave functions for antisymmetric meson states are not accurately described. For symmetric states no such ambiguities appear. This difficulty for small values of  $r_n$  is associated with the degeneracy of the various  $2p$  states that occur for  $r_n=0$ , and is therefore unimportant for larger values of  $r_n$ . To treat the inner region correctly would require the introduction of the two other  $2p$  states and their associated  $\chi_i$ 's. We expect that such a treatment would, however, make small corrections to the wave functions at large distances and would be significant only for small values of  $r_n$ . For the scattering states the energies of interest to us are such that the contributions for other than  $s$  states are negligible. For the bound states, only the unequal-mass cases involve  $\psi_-$ , and the effect of this term is small except for large values of  $r_n$ . We have therefore felt justified in omitting these additional complications in our treatment.

The asymptotic behavior of  $f_{+-}$  and  $g_{+-}$  in this limit is also of interest. From the expressions in the Appendix we obtain

$$\lim_{r_n \rightarrow 0} \mathbf{f}_{+-} = 4\pi A_+ A_- (f_2 - f_1) \frac{p_+ q_-}{(p_+ + p_-)^4} \mathbf{e}_{r_n} \\ = \frac{16\sqrt{2}}{81} (f_2 - f_1) \mathbf{e}_{r_n},$$

and

$$\lim_{r_n \rightarrow 0} g_{+-} = -2f_{+-}/r_n,$$

where  $f_i = m_i/(m_1 + m_2)$ , and  $\mathbf{e}_{r_n}$  is a unit vector in the direction of  $\mathbf{r}_n$  and  $f_{+-} = \mathbf{f}_{+-} \cdot \mathbf{e}_{r_n}$ . The term  $g_{+-}$  is therefore seen to be divergent in this limit. As we shall show, the particular form of this divergence is crucial for the satisfactory solution of the differential equations.

The radial equations for a state of total angular momentum  $l$  can be obtained from Eqs. (1) by the usual substitution of  $\phi_i/r_n$  for  $\chi_i$ . These equations are

$$\frac{d^2 \phi_+}{dr_n^2} + \frac{1}{\epsilon} [W - V_+(r_n)] \phi_+ \\ = -2f_{+-} \frac{d\phi_-}{dr_n} - \frac{df_{+-}}{dr_n} \phi_- + g_{+-} \phi_-, \quad (2a)$$

and

$$\frac{d^2 \phi_-}{dr_n^2} + \frac{1}{\epsilon} [W - V_-(r_n)] \phi_- \\ = 2f_{+-} \frac{d\phi_+}{dr_n} + \frac{df_{+-}}{dr_n} \phi_+ + g_{+-} \phi_+, \quad (2b)$$

where

$$V_+(r_n) = W_+ + 2r_n^{-1} + \epsilon g_{++} + \epsilon l(l+1)r_n^{-2}, \\ \text{and} \\ V_-(r_n) = W_- + 2r_n^{-1} + \epsilon g_{--} + \epsilon l(l+1)r_n^{-2}.$$

In order to obtain the behavior of the solutions to these equations for small values of  $r_n$  it is convenient to express the solutions in a power series in  $r_n$ ; i.e., we assume

$$\phi_+(r_n) = \sum_{t=0}^{\infty} a_t r_n^{t+K},$$

and

$$\phi_-(r_n) = \sum_{t=0}^{\infty} b_t r_n^{t+K}.$$

In addition it is necessary to expand the various other functions which appear in the equations in power series, thus

$$f_{+-} = \sum_{t=0}^{\infty} F_t r_n^t,$$

$$g_{+-} = \sum_{t=-1}^{\infty} g_t r_n^t,$$

$$V_+ = \epsilon l(l+1)r_n^{-2} + \sum_{t=-1}^{\infty} v_t r_n^t = \sum_{t=-2}^{\infty} v_t r_n^t,$$

and

$$V_- = \epsilon [l(l+1) + 2] r_n^{-2} + \sum_{t=-1}^{\infty} \mu_t r_n^t = \sum_{t=-2}^{\infty} \mu_t r_n^t.$$

Inserting these expressions into Eqs. (2a) and (2b) and equating terms with equal powers of  $r_n$ , we obtain the recursion relationships

$$[(t+K)(t+K-1) - l(l+1)]a_t \\ - \frac{1}{\epsilon} \sum_{t'=0}^{t-1} v_{t'-1} a_{t-t'-1} + \frac{W}{\epsilon} a_{t-2} \\ = \sum_{t'=0}^{t-1} [g_{t'-1} - (2t+2K-t'-2)F_{t'}] b_{t-t'-1},$$

and

$$[(t+K)(t+K-1) - 2 - l(l+1)]b_t \\ - \frac{1}{\epsilon} \sum_{t'=0}^{t-1} \mu_{t'-1} b_{t-t'-1} + \frac{W}{\epsilon} b_{t-2} \\ = \sum_{t'=0}^{t-1} [g_{t'-1} + (2t+2K-t'-2)F_{t'}] b_{t-t'-1}.$$

From these equations we obtain the pair of indicial equations

$$[K(K-1) - l(l+1)]a_0 = 0,$$

and

$$[K(K-1) - 2 - l(l+1)]b_0 = 0.$$

Thus if  $a_0$  is not zero  $K$  is either  $-l$  or  $l+1$ , while if  $b_0$  is not zero  $K$  is equal to  $\frac{1}{2} \pm [9/4 + l(l+1)]^{1/2}$ . In both these cases the solutions with the minus sign do not satisfy the conditions of integrability and may be discarded. For  $l \neq 0$ , these two cases constitute the two possible solutions to the equations. For  $l=0$ , on the other hand, these two values for  $K$  differ by an integer, and therefore further investigation is necessary to determine whether or not two regular and independent solutions to the equations exist. It is clear that two such solutions must exist, since we must be able to describe states in which the meson is associated with either of the two nuclei for large separations of the nuclei. If we examine the recursion equation which determines the values of  $b_1$  in the case where  $K=1$ , i.e., for  $a_0 \neq 0$ , we find

$$[(1+K)K - l(l+1) - 2]b_1 = (g_{-1} + 2b_0)a_0,$$

or

$$0 \cdot b_1 = (g_{-1} + 2b_0)a_0;$$

this would lead to an inconsistency unless the multiplier of  $a_0$  were zero.<sup>8</sup> This is, in fact, the condition which we have shown to be true from the asymptotic behavior of the functions  $f_{+-}$  and  $g_{+-}$ . This being true, the value of  $b_1$  is undetermined. Thus the constant  $b_1$ , which is arbitrary in the solution with  $K=1$ , represents the fact that one can add an arbitrary amount of the solution with  $K=2$  to the solution and still retain a valid power-series expansion for small values of  $r_n$ .

#### D. Unitarity Current

It is of interest that one can obtain an invariant relationship between the various solutions for the system of equations describing the motion of the nuclei. If we consider two sets of solutions  $\chi_{\pm}^{(1)}$  and  $\chi_{\pm}^{(2)}$  to these equations, with eigenvalues  $W^1$  and  $W^2$ , respectively, then the following equation can be constructed:

$$\begin{aligned} & \int \left\{ \chi_{+}^{(2)*} \left[ \left( \nabla_n^2 - \frac{1}{\epsilon} V_+ \right) \chi_{+}^{(1)} + \Theta_{+-} \chi_{-}^{(1)} \right] \right. \\ & \quad + \chi_{-}^{(2)*} \left[ \left( \nabla_n^2 - \frac{1}{\epsilon} V_- \right) \chi_{-}^{(1)} + \Theta_{-+} \chi_{+}^{(1)} \right] \\ & \quad - \chi_{+}^{(1)} \left[ \left( \nabla_n^2 - \frac{1}{\epsilon} V_+ \right) \chi_{+}^{(2)} + \Theta_{+-} \chi_{-}^{(2)} \right]^* \\ & \quad \left. - \chi_{-}^{(1)} \left[ \left( \nabla_n^2 - \frac{1}{\epsilon} V_- \right) \chi_{-}^{(2)} + \Theta_{-+} \chi_{+}^{(2)} \right]^* \right\} d\tau_n \\ & = -\frac{1}{\epsilon} \left\{ W^1 \int (\chi_{+}^{(2)*} \chi_{+}^{(1)} + \chi_{-}^{(2)*} \chi_{-}^{(1)}) d\tau_n \right. \\ & \quad \left. - W^2 \int (\chi_{+}^{(1)} \chi_{+}^{(2)*} + \chi_{-}^{(1)} \chi_{-}^{(2)*}) d\tau_n \right\}. \end{aligned}$$

<sup>8</sup> If such a conflict exists, the second solution will be of a logarithmic type.

If  $W^1$  is equal to  $W^2$  then the right-hand side of the equation is zero and we find

$$\int \nabla \cdot \mathbf{j}_{12} d\tau_n = 0,$$

where

$$\mathbf{j}_{12} = [\chi_{+}^{(2)*} \nabla_n \chi_{+}^{(1)} - \chi_{+}^{(1)} \nabla_n \chi_{+}^{(2)*} + \chi_{-}^{(2)*} \nabla_n \chi_{-}^{(1)} - \chi_{-}^{(1)} \nabla_n \chi_{-}^{(2)*} + 2f_{+-} (\chi_{+}^{(2)*} \chi_{-}^{(1)} - \chi_{-}^{(2)*} \chi_{+}^{(1)})].$$

We shall call  $\mathbf{j}_{12}$  the "unitarity current." (For  $\chi^{(1)} \equiv \chi^{(2)}$  this reduces to the usual expression for the probability current.) If  $\chi^{(1)}$  and  $\chi^{(2)}$  have the same angular dependence, then we may write

$$\begin{aligned} r_n^2 \mathbf{j}_{12} \cdot \mathbf{e}_{r_n} = r_n^2 j_{12} = & \chi_{+}^{(2)*} \frac{d\chi_{+}^{(1)}}{dr_n} - \chi_{+}^{(1)} \frac{d\chi_{+}^{(2)*}}{dr_n} \\ & - \chi_{-}^{(2)*} \frac{d\chi_{-}^{(1)}}{dr_n} - \chi_{-}^{(1)} \frac{d\chi_{-}^{(2)*}}{dr_n} \\ & + 2f_{+-} (\chi_{+}^{(2)*} \chi_{-}^{(1)} - \chi_{+}^{(1)} \chi_{-}^{(2)*}) = \text{constant}. \end{aligned}$$

Furthermore, if these solutions are regular solutions to the differential equations, their contribution to this quantity is zero because  $r_n^2 j_{12}$  vanishes at the origin. Irregular solutions, on the other hand, contribute a finite amount to this expression. It follows that if  $r_n^2 j_{12}$  is not zero when evaluated for any value of  $r_n$ , some irregular solutions must be present.

### III. SCATTERING WAVE FUNCTIONS

#### A. Asymptotic Behavior for Large Values of $r_n$

In the limit of large values of  $r_n$  the parameters  $P$  and  $Q$  which describe the mesonic wave functions also become large. In this limit the binding energies may be expressed approximately by

$$W_+ r_n^2 = W_- r_n^2 = (P+Q-4r_n)PQ/(P+Q).$$

The values of  $P$  and  $Q$  which minimize this expression for the energies are  $P=Q=r_n$ , for which  $W_+ = W_- = -1$ . This expresses qualitatively the fact that the symmetric and antisymmetric solutions for fixed nuclei can be formed from the solutions in which the meson is centered on either of the two nuclei. These eigenvalues, however, are not exactly the binding energies for such a separated system, because the units in which the eigenvalues are measured use the reduced mass for the meson with respect to the sum of the nuclear masses. The necessary corrections to the energies in this limit are contained in the asymptotic behavior of coupling terms  $g_{ii}$ . In addition, for the unequal-mass case it is necessary to obtain the splitting in the energies corresponding to the fact that the meson is more tightly bound on the heavier of the nuclei. The removal of this degeneracy in energy for this limit is contained in the off-diagonal term  $g_{+-}$ . As will be shown, both these



corrections lead to expressions accurate to first order in the parameter  $\epsilon$ .

The asymptotic values of  $g_{++}$  and  $g_{--}$  come entirely from the terms  $I_{\xi\xi}$  and  $I_{\eta\eta}$  (see Appendix) because the derivatives which occur in the remaining terms vanish in this limit. We find

$$\lim_{r_n \rightarrow \infty} g_{++} = \lim_{r_n \rightarrow \infty} g_{--} = \frac{1}{2}(f_1^2 + f_2^2).$$

For the equal-mass case we have  $f_1 = f_2 = \frac{1}{2}$ ; hence

$$\lim_{r_n \rightarrow \infty} g_{++} = \lim_{r_n \rightarrow \infty} g_{--} = \frac{1}{4},$$

and the effective potentials are corrected to give

$$\lim_{r_n \rightarrow \infty} V_+ = \lim_{r_n \rightarrow \infty} V_- = -1 + \frac{1}{4}\epsilon = -1 + m_\mu/2m_1.$$

To first order in  $\epsilon$  this is identical to  $M_\mu^*/M_\mu$ , where  $M_\mu^*$  is the reduced mass of the meson with respect to one of the nuclei. For the unequal-mass case let us consider a system which consists of a proton (nucleon 1), a deuteron (nucleon 2), and a meson. In this case we have  $f_1 = \frac{1}{3}$  and  $f_2 = \frac{2}{3}$ , and hence

$$\lim_{r_n \rightarrow \infty} g_{++} = \lim_{r_n \rightarrow \infty} g_{--} = 5/18.$$

The only term for  $g_{+-}$  which does not vanish in this limit is the term  $g_{\xi\eta}$ , and this leads to the result

$$\lim_{r_n \rightarrow \infty} g_{+-} = -(f_2 - f_1)/2 = -\frac{1}{6}.$$

To interpret these results let us consider the asymptotic form for the radial differential equations,

$$\frac{d^2\phi_+}{dr_n^2} + \frac{1}{\epsilon} \left( W + 1 - \frac{5}{18}\epsilon \right) \phi_+ = -\frac{1}{6}\phi_-,$$

and

$$\frac{d^2\phi_-}{dr_n^2} + \frac{1}{\epsilon} \left( W + 1 - \frac{5}{18}\epsilon \right) \phi_- = -\frac{1}{6}\phi_+.$$

From these equations we obtain a new set of equations,

$$\frac{d^2\phi_p}{dr_n^2} + \frac{1}{\epsilon} \left[ W + 1 - \epsilon \left( \frac{5}{18} + \frac{1}{6} \right) \right] \phi_p = 0, \quad (3a)$$

and

$$\frac{d^2\phi_d}{dr_n^2} + \frac{1}{\epsilon} \left[ W + 1 - \epsilon \left( \frac{5}{18} - \frac{1}{6} \right) \right] \phi_d = 0, \quad (3b)$$

where

$$\phi_p = (\phi_+ - \phi_-)/\sqrt{2} \quad \text{and} \quad \phi_d = (\phi_+ + \phi_-)/\sqrt{2}.$$

This particular choice for  $\phi_p$  and  $\phi_d$  is such that asymptotically  $\phi_p$  corresponds to a total wave function in which the meson is centered on nucleon 1, while  $\phi_d$  is

the corresponding case for nucleon 2. The binding energy of the meson is that value of  $W$  for which the kinetic energy of the relative nuclear motion is zero ( $d^2\phi/dr_n^2 = 0$ ). Thus we find, for the binding energies,

$$W_p = -1 + 4\epsilon/9,$$

and

$$W_d = -1 + \epsilon/9.$$

These expressions are the correct binding energies to first order in the parameter  $\epsilon$ .<sup>9</sup>

## B. Scattering Cross Sections

In the treatment which follows, we restrict ourselves to the consideration of the scattering states of zero total angular momentum; the extension to states of higher angular momentum, however, is straightforward. For the investigation of scattering phenomena, as in the usual treatments, we need study only the asymptotic behavior of the wave functions for the separated system.

For the unequal-mass case there are four asymptotic functions to consider, i.e.,  $\chi_+$  and  $\chi_-$  for each of the two solutions regular at the origin. In addition, as we have shown in the preceding section, the degeneracy of the binding energies of the meson on the two separated nuclei has been removed by the dynamic correction terms. It is therefore necessary to distinguish between the case in which the energy of the system lies between these two binding energies and the case in which it is larger than either of them. These two energy ranges correspond to different physical situations. In the former the only scattering states allowed are those in which the meson is bound to the heavier nucleus for large separations. In the latter the meson may be bound to either of the nuclei; exchange processes are also possible in this case.

It is once again convenient to use wave functions that asymptotically describe the meson centered on one of the two nuclei. We therefore define

$$\phi_p^i = (\phi_+^i - \phi_-^i)/\sqrt{2},$$

and

$$\phi_d^i = (\phi_+^i + \phi_-^i)/\sqrt{2}.$$

Here  $\phi_p$  is the radial wave function whose form corresponds to the meson bound on nucleus 1, the lighter nucleus, with an energy  $W_p$ ;  $\phi_d$  and  $W_d$  are defined in a similar manner for the heavier nucleus. (If we consider the system of a proton, a deuteron, and a meson, the expressions of the preceding section define  $W_p$  and  $W_d$ .) The superscripts  $i$  ( $i = 1, 2$ ) refer to either of the two regular solutions to the radial differential equations.

<sup>9</sup> By generalizing the functions  $\psi_+$  and  $\psi_-$  to the form  $\psi = \alpha\psi_+ + \beta\psi_-$  and including in the Hamiltonian those terms in  $g_{++}$ ,  $g_{--}$ , and  $g_{+-}$  that are independent of derivatives of  $p$ ,  $q$ , etc., one could obtain the exact binding energy as  $r_n \rightarrow \infty$ , without changing the form of the differential equation for the  $\chi$ 's. Because physical processes would still involve unknown terms of order  $\epsilon^2$ , it was felt that the additional labor involved in such a treatment was not justified.

If the energy,  $W$ , is such that  $W_d \leq W \leq W_p$ , then we can write the asymptotic behavior for  $\phi_p$  and  $\phi_d$  in the form

$$\phi_p = a_p^i e^{\alpha r_n} + b_p^i e^{-\alpha r_n}$$

and

$$\phi_d = a_d^i \sin(k_d r_n + \delta_d^i),$$

where  $\alpha = [(W_p - W)/\epsilon]^{\frac{1}{2}}$  and  $k_d = [(W - W_d)/\epsilon]^{\frac{1}{2}}$ . The parameters  $a_p^i$ ,  $b_p^i$ ,  $a_d^i$ , and  $\delta_d^i$  are determined by the value of  $\phi_p^i$ ,  $d\phi_p^i/dr_n$ ,  $\phi_d^i$ , and  $d\phi_d^i/dr_n$  evaluated for some large value of  $r_n$ . In order to completely specify the wave function, it is necessary to determine that linear combination of the two solutions  $i=1$  and  $i=2$  for which no increasing exponential remains in the asymptotic expression for  $\phi_p$ . We may also normalize these solutions to the incident part of the plane wave. If this is done we find, for the corresponding wave functions  $\chi_p$  and  $\chi_d$  in this asymptotic limit, the forms<sup>10</sup>

$$\chi_p \propto e^{-\alpha r_n}/r_n,$$

and

$$\chi_d = e^{ik_d z} + \left( \frac{M-1}{2ik_d} \right) \frac{e^{ik_d r_n}}{r_n},$$

where the  $z$  axis is in the direction of the incident-particle beam. The quantity  $M$  is defined as

$$M = \frac{a_p^1 a_d^2 \exp(i\delta_d^2) - a_p^2 a_d^1 \exp(i\delta_d^1)}{a_p^1 a_d^2 \exp(-i\delta_d^2) - a_p^2 a_d^1 \exp(-i\delta_d^1)}.$$

If we replace  $M$  by the quantity  $e^{2i\theta_d}$ , then  $\theta_d$  may be considered the phase shift for this scattering, and the scattering cross section can be written in the usual form,

$$\sigma = (4\pi/k_d^2) \sin^2 \theta_d.$$

If the energy is larger than  $W_p$ , then both solutions have asymptotic sinusoidal behavior, i.e.,

$$\phi_p^i = a_p^i \sin(k_p r_n + \delta_p^i),$$

and

$$\phi_d^i = a_d^i \sin(k_d r_n + \delta_d^i),$$

where  $k_p = [(W - W_p)/\epsilon]^{\frac{1}{2}}$ . There are now two possible physical states corresponding to incident states in which the meson is centered on either of the two nuclei. In this case, however, we may separately choose those linear combinations of the two sets of solutions that correspond to no incident part for either the  $\chi_p$  or the

<sup>10</sup> It might be pointed out that this asymptotic form has a defect in that the meson current is zero. This is consistent with the assumption that the meson velocity about the nuclei is large compared with the nuclear velocities, and therefore, in the region where the particles interact strongly, such additional velocities represent a small correction. A correct asymptotic form for the plane-wave part would be

$$\exp\{i[kZ_\mu + k'Z_1 - (k+k')Z_2]\} f(\mathbf{r}_\mu - \mathbf{r}_1),$$

where  $k/m_1 = k'/m_\mu$  and  $f$  is the wave function for the meson about nucleus 1. Our wave function thus neglects  $k' = (m_\mu/m_1)k$ .

$\chi_d$ . In either case we may normalize the solutions to the incident wave.

For the scattering of the  $p$ -mesonic system from a  $d$  nucleus, we obtain the asymptotic forms

$$\chi_p = e^{ik_p z} + h_{pp} e^{ik_p r_n}/r_n,$$

$$\chi_d = h_{pd} e^{ik_d r_n}/r_n,$$

where

$$h_{pp} = \{[a_p^2 a_d^1 \exp(i(\delta_p^2 - \delta_d^1)) - a_p^1 a_d^2 \exp(i(\delta_p^1 - \delta_d^2))]\} D^{-1} - 1\} / 2ik_p,$$

$$h_{pd} = a_d^1 a_d^2 \sin(\delta_d^1 - \delta_d^2) / Dk_p,$$

and

$$D = a_p^2 a_d^1 \exp[-i(\delta_p^2 + \delta_d^1)] - a_p^1 a_d^2 \exp[-i(\delta_p^1 + \delta_d^2)].$$

Here  $h_{pp}$  gives the amplitude for normal scattering and  $h_{pd}$  the corresponding amplitude for exchange scattering in which the meson is captured by the heavier nucleus. The cross sections for these cases are

$$\sigma_{pp} = 4\pi |h_{pp}|^2,$$

and for the exchange process,

$$\sigma_{pd} = 4\pi |h_{pd}|^2 k_d/k_p,$$

where the factor  $k_d/k_p$  is necessary to correct for the change in velocity of the incident and outgoing particles in this inelastic collision.

In a similar fashion the scattering of the  $d$ -mesonic system from a  $p$  nucleus can be obtained. In this case we find

$$h_{dd} = \{[a_p^2 a_d^1 \exp(i(\delta_d^1 - \delta_p^2)) - a_p^1 a_d^2 \exp(i(\delta_d^2 - \delta_p^1))]\} D^{-1} - 1\} / 2ik_d,$$

and

$$h_{dp} = a_p^1 a_p^2 \sin(\delta_p^1 - \delta_p^2) / Dk_d.$$

The corresponding cross sections are

$$\sigma_{dd} = 4\pi |h_{dd}|^2,$$

and

$$\sigma_{dp} = 4\pi |h_{dp}|^2 k_p/k_d.$$

From the conservation of unitarity current one can show  $h_{pd} = h_{dp}$  and that the two exchange cross sections are simply related.

For the equal-mass case the phase shifts for the symmetric and antisymmetric scattering states may be independently evaluated by use of the asymptotic forms

$$\chi_i = a_i \sin(kr_n + \delta_i)/r_n, \quad (\text{for } i = +, -)$$

where  $k = [(W - 1 + \frac{1}{4}\epsilon)/\epsilon]^{\frac{1}{2}}$ . In this case, however, a further complication is introduced because the nuclei are generally identical particles. In such cases the total

wave functions for the system must be properly symmetrized. As before, it is convenient to introduce combinations of wave functions which describe the states corresponding to the meson centered on each of the two nuclei for large separations. For the case in which the meson is centered on nucleus 1, the mesonic wave function is

$$\psi_1 = (\psi_+ - \psi_-)/\sqrt{2},$$

with a corresponding nucleonic wave function

$$\chi_1 = (\chi_+ - \chi_-)/\sqrt{2}.$$

The wave functions for the other case are

$$\psi_2 = (\psi_+ + \psi_-)/\sqrt{2},$$

and

$$\chi_2 = (\chi_+ + \chi_-)/\sqrt{2}.$$

The total wave function is then of the form

$$\Psi = \chi_1\psi_1 + \chi_2\psi_2.$$

Before symmetrization, the solution which corresponds to an incident system in which the meson is associated with nucleus 1 or with nucleus 2 has the asymptotic forms

$$\chi_1 = e^{ikz} + h_{11}r_n^{-1}e^{ikr_n},$$

$$\chi_2 = h_{12}r_n^{-1}e^{ikr_n},$$

and

$$\chi_1 = h_{21}r_n^{-1}e^{ikr_n},$$

$$\chi_2 = e^{-ikz} + h_{22}r_n^{-1}e^{ikr_n},$$

respectively. Using the asymptotic forms for  $\chi_+$  and  $\chi_-$ , we find

$$h_{11} = h_{22} = (4ik)^{-1}[e^{2i\delta_+} + e^{2i\delta_-} - 2],$$

and

$$h_{12} = h_{21} = (4ik)^{-1}[e^{2i\delta_+} - e^{2i\delta_-}].$$

For a system consisting of a meson and two spin- $\frac{1}{2}$  particles such as protons, the total wave function must be symmetric for singlet states (*s*) and antisymmetric for triplet states (*t*). Thus for states of zero total angular momentum the total wave functions are<sup>10</sup>

$$\Psi_s = e^{ikz}\psi_1 + e^{-ikz}\psi_2 + (h_{11} + h_{12})(e^{ikr_n}/r_n)(\psi_1 + \psi_2),$$

$$\Psi_t = e^{ikz}\psi_1 - e^{-ikz}\psi_2 + (h_{11} - h_{12})(e^{ikr_n}/r_n)(\psi_1 - \psi_2).$$

From this wave function we obtain for the cross sections<sup>11</sup>

$$\sigma_{\text{singlet}} = 4\pi |h_{11} + h_{12}|^2 = 4\pi k^{-2} \sin^2\delta_+,$$

$$\sigma_{\text{triplet}} = 4\pi |h_{11} - h_{12}|^2 = 4\pi k^{-2} \sin^2\delta_-.$$

<sup>11</sup> It is perhaps of interest that this result is the same as that which would have been obtained for nonidentical particles. The presence of the meson on one of the two particles provides some basis for distinguishing between them. The result follows directly from second quantization of the system.

The total cross section is

$$\sigma_{pp} = 4\pi k^{-2} [\frac{1}{4} \sin^2\delta_+ + \frac{3}{4} \sin^2\delta_-].$$

Similarly we find, for the case in which the two nuclei are deuterons,

$$\sigma_{dd} = 4\pi k^{-2} [\frac{2}{3} \sin^2\delta_+ + \frac{1}{3} \sin^2\delta_-].$$

#### IV. BOUND STATES

This section is devoted to a discussion of an iterative scheme by which it is possible to obtain the bound-state eigenvalues and eigenfunctions for the system of differential equations describing the nuclear motion. In this development it is assumed that the integration of the differential equations can be carried out by either exact or numerical methods.

For the unequal-mass case, the Hamiltonian for the nuclear system can be written as

$$H = \int_0^\infty \left[ \left( \frac{d\phi_+}{dr_n} \right)^2 + \left( \frac{d\phi_-}{dr_n} \right)^2 + \frac{V_+}{\epsilon} \phi_+^2 + \frac{V_-}{\epsilon} \phi_-^2 + \frac{l(l+1)}{r_n^2} (\phi_+^2 + \phi_-^2) - 2g_{+-} \phi_+ \phi_- + 2f_{+-} \left( \phi_+ \frac{d\phi_-}{dr_n} - \phi_- \frac{d\phi_+}{dr_n} \right) \right] dr_n.$$

If we consider variations of  $\phi_+$ , where  $\phi_+$  is assumed to be continuous with a piecewise continuous first derivative, subject to the constraint that  $\int_0^\infty (\phi_+^2 + \phi_-^2) dr_n = 1$ , then the variation of  $H + \lambda N$  is

$$\delta(H + \lambda N) = \int_0^\infty 2 \left[ -\frac{d^2\phi_+}{dr_n^2} + \left( \frac{V_+}{\epsilon} + \frac{l(l+1)}{r_n^2} \right) \phi_+ - g_{+-} \phi_- + 2f_{+-} \frac{d\phi_-}{dr_n} + \phi_- \frac{df_{+-}}{dr_n} \right] \delta\phi_+ dr_n + \lambda \int_0^\infty 2\phi_+ \delta\phi_+ dr_n + \sum_i \delta\phi_+(r_i) \Delta_i \left( \frac{d\phi_+}{dr_n} \right).$$

Here  $r_i$  indicates the values of  $r_n$  at the points of discontinuity of  $d\phi_+/dr_n$ , and  $\Delta_i(d\phi_+/dr_n)$  are the changes in  $d\phi_+/dr_n$  between  $r_i - |\nu|$  and  $r_i + |\nu|$  in the limit of vanishing  $\nu$ . If the parameter  $\lambda$  is  $-W/\epsilon$ , then the conditions for an extremum in  $H$  are that the radial differential equation for  $\phi_+$  be satisfied and  $d\phi_+/dr_n$  be continuous. In a similar manner the variations of  $H$  with respect to variations of  $\phi_-$  lead to the radial equations for  $\phi_-$ . The value of  $H$  obtained for this extremum is in fact  $W/\epsilon$ .

If we now use trial wave functions  $\phi_+^{(0)}$  and  $\phi_-^{(0)}$  that: (a) satisfy the differential equations for an energy  $W^0$ , (b) are continuous for all values of  $r_n$ , and (c) have continuous first derivatives except at one point  $r_0$ , then

the true eigenvalue  $W^T$  may be expressed as

$$\begin{aligned}
 W^T &= \int_0^\infty (\phi_+^{(0)2} + \phi_-^{(0)2}) dr_n \\
 &= \int_0^\infty \phi_+^{(0)} \left\{ -\frac{d^2\phi_+^{(0)}}{dr_n^2} + \frac{V_+}{\epsilon} \phi_+^{(0)} + \frac{l(l+1)}{r_n^2} \phi_+^{(0)} \right. \\
 &\quad \left. - g_{+-} \phi_-^{(0)} + 2f_{+-} \frac{d\phi_-^{(0)}}{dr_n} + \phi_-^{(0)} \frac{df_{+-}}{dr_n} \right\} dr_n \\
 &\quad + \int_0^\infty \phi_-^{(0)} \left\{ -\frac{d^2\phi_-^{(0)}}{dr_n^2} + \frac{V_-}{\epsilon} \phi_-^{(0)} + \frac{l(l+1)}{dr_n^2} \phi_-^{(0)} \right. \\
 &\quad \left. - g_{+-} \phi_+^{(0)} - 2f_{+-} \frac{d\phi_+^{(0)}}{dr_n} - \phi_+^{(0)} \frac{df_{+-}}{dr_n} \right\} dr_n \\
 &\quad - \left\{ \phi_+^{(0)}(r_0) \Delta_0 \left( \frac{d\phi_+^{(0)}}{dr_n} \right) + \phi_-^{(0)}(r_0) \Delta_0 \left( \frac{d\phi_-^{(0)}}{dr_n} \right) \right\}.
 \end{aligned}$$

As a result of condition (a) and the extremal nature of  $H$  for this solution, this may be rewritten as

$$\begin{aligned}
 W^T &= W^0 \\
 &= \frac{[\phi_+^{(0)}(r_0) \Delta_0 (d\phi_+^{(0)}/dr_n) + \phi_-^{(0)}(r_0) \Delta_0 (d\phi_-^{(0)}/dr_n)]}{\int_0^\infty \{\phi_+^{(0)2} + \phi_-^{(0)2}\} dr_n} \\
 &\quad + \mathcal{O}(\xi^2),
 \end{aligned}$$

where  $\xi$  is a parameter of smallness that indicates the deviation between the trial and true solutions.

The integration for the bound states can be divided into two regions, with  $r_m$  as the common boundary. At this point four independent quantities can be specified, namely,  $\phi_+^{(0)}$ ,  $d\phi_+^{(0)}/dr_n$ ,  $\phi_-^{(0)}$ , and  $d\phi_-^{(0)}/dr_n$ . By choosing appropriate linear combinations of the two regular inner solutions and the two bounded outer solutions, three of the four quantities can be made continuous at  $r_m$ .<sup>12</sup> For the correct eigenvalue the fourth quantity will also be continuous. For a trial eigenvalue in general one of the four quantities will not be continuous, however. If, for example, we allow this discontinuity to occur in  $d\phi_+^{(0)}/dr_n$ , then a better approximation to the correct eigenvalue,  $W^n$ , is expressed in terms of the trial eigenvalue and the value of the function at  $r_m$  as

$$W^n = W^0 + \frac{\phi_+^{(0)} [(d\phi_+^{(0)}/dr_n)_{r-|r|} - (d\phi_+^{(0)}/dr_n)_{r+|r|}]}{\int_0^\infty (\theta_+^{(0)2} + \phi_-^{(0)2}) dr_n}$$

<sup>12</sup> If, as was done for the bound states, the integration is divided into two regions for free states with  $W_D \leq W < W_P$ , three bounded outer solutions exist (two sinusoidal ones associated with  $\psi_D$  and a decreasing-exponential one associated with  $\psi_H$ ). This gives the necessary freedom to make all four quantities continuous at  $r_m$ .

By using this new improved eigenvalue to obtain a new trial solution and thus iterate the solution, we can converge upon the correct eigenvalue and consequently the correct eigenfunction.

This variational procedure is not restricted to the ground state, but may be applied to the higher excited states as well. The determination of a specific state merely requires the specification of a boundary condition on the number of nodes allowed in the solution.

For the equal-mass case the development given above is equally applicable. In this case the  $\phi_+$  and  $\phi_-$  equations are not coupled. Bound states, however, occur only for the  $\phi_+$  solutions. It therefore suffices to impose on the development given above the added constraint  $\phi_- \equiv 0$ .

### APPENDIX

In this Appendix we obtain explicit analytic expressions for the first-order dynamic corrections for the case in which the mesonic wave functions are assumed to be those used in the text, i.e.,

$$\psi_+ = A_+ \cosh(q_+ r_n \eta / 2) e^{-p_+ r_n \xi / 2},$$

and

$$\psi_- = A_- \{\xi\} \sinh(q_- r_n \eta / 2) e^{-p_- r_n \xi / 2}.$$

The dynamic correction terms are of the form

$$g_{ij} = g_{ji} = \int \nabla_n \psi_i \cdot \nabla_n \psi_j d\tau_\mu,$$

and

$$\mathbf{f}_{ij} = -\mathbf{f}_{ji} = \int \psi_i \nabla_n \psi_j d\tau_\mu,$$

where, from the previous development, the indicated differentiations,  $\nabla_n$ , must be carried out with the mesonic variable  $\mathbf{R}_\mu$  fixed.

For the symmetric case, we may write

$$\begin{aligned}
 \frac{\nabla_n \psi_+}{\psi_+} &= \mathbf{e}_{r_n} \left\{ \frac{1}{A_+} \frac{dA_+}{dr_n} + \frac{r_n \eta}{2} \frac{dq_+}{dr_n} \tanh \frac{q_+ r_n \eta}{2} - \frac{r_n \xi}{2} \frac{dp_+}{dr_n} \right\} \\
 &\quad + \frac{q_+}{2} \nabla_n(r_n \eta) \tanh \frac{q_+ r_n \eta}{2} - \frac{p_+}{2} \nabla_n(r_n \xi),
 \end{aligned}$$

where  $\mathbf{e}_{r_n}$  is a unit vector in the direction of  $\mathbf{r}_n$ . For reference we designate these five terms by the subscripts  $A$ ,  $q$ ,  $p$ ,  $\eta$ , and  $\xi$ , respectively. Similarly for the anti-symmetric cases for  $r_n > r_c$ , we have

$$\begin{aligned}
 \frac{\nabla_n \psi_-}{\psi_-} &= \mathbf{e}_{r_n} \left\{ \frac{1}{A_-} \frac{dA_-}{dr_n} + \frac{r_n \eta}{2} \frac{dq_-}{dr_n} \coth \frac{q_- r_n \eta}{2} - \frac{r_n \xi}{2} \frac{dp_-}{dr_n} \right\} \\
 &\quad + \frac{q_-}{2} \nabla_n(r_n \eta) \coth \frac{q_- r_n \eta}{2} - \frac{p_-}{2} \nabla_n(r_n \xi),
 \end{aligned}$$

while for  $r_n < r_c$ , we write

$$\frac{\nabla_n \psi_-}{\psi_-} = \mathbf{e}_{r_n} \left[ \left( \frac{1}{A_-} \frac{dA_-}{dr_n} - \frac{1}{r_n} \right) + \frac{r_n \eta}{2} \frac{dq_-}{dr_n} \coth \frac{q_- r_n \eta}{2} - \frac{r_n \xi}{2} \frac{d\dot{p}_-}{dr_n} \right] + \frac{q_-}{2} \nabla_n(r_n \eta) \coth \frac{q_- r_n \eta}{2} - \left( \frac{\dot{p}_-}{2} - \frac{1}{r_n \xi} \right) \nabla_n(r_n \xi).$$

In order to obtain the expressions for  $g_{ij}$  and  $f_{ij}$  the following identities are useful:

$$\begin{aligned} \nabla_n(r_n \xi) \cdot \nabla_n(r_n \xi) &= f_1^2 + f_2^2 - 2f_1 f_2 \frac{\xi^2 + \eta^2 - 2}{\xi^2 - \eta^2}, \\ \nabla_n(r_n \eta) \cdot \nabla_n(r_n \eta) &= f_1^2 + f_2^2 + 2f_1 f_2 \frac{\xi^2 + \eta^2 - 2}{\xi^2 - \eta^2}, \\ \nabla_n(r_n \xi) \cdot \nabla_n(r_n \eta) &= f_2 - f_1, \\ \nabla_n(r_n \xi) \cdot \mathbf{e}_{r_n} &= \frac{1}{\xi^2 - \eta^2} [\xi(1 - \eta^2) + (f_2 - f_1)\eta(\xi^2 - 1)], \\ \nabla_n(r_n \eta) \cdot \mathbf{e}_{r_n} &= \frac{1}{\xi^2 - \eta^2} [(f_2 - f_1)\xi(1 - \eta^2) + \eta(\xi^2 - 1)]. \end{aligned}$$

Using the above expressions, we may evaluate each of the terms that occur.

For  $g_{++}$  we obtain the sum of the following terms:

$$\begin{aligned} I_{A,A^+} &= \left( \frac{1}{A_+} \frac{dA_+}{dr_n} \right)^2, \\ I_{p,p^+} &= \frac{1}{4} N r_n^2 (d\dot{p}_+/dr_n)^2 (E_4 C_0 - E_2 C_2), \\ I_{q,q^+} &= \frac{1}{4} N r_n^2 (dq_+/dr_n)^2 (E_2 C_2' - E_0 C_4'), \\ I_{pq^+} &= -\frac{1}{2} N r_n^2 (d\dot{p}_+/dr_n) (dq_+/dr_n) (E_3 C_1 - E_1 C_3), \\ I_{p,A^+} &= -N \left( \frac{1}{A_+} \frac{dA_+}{dr_n} \right) r_n \left( \frac{d\dot{p}_+}{dr_n} \right) (E_3 C_0 - E_1 C_2), \\ I_{q,A^+} &= N \left( \frac{1}{A_+} \frac{dA_+}{dr_n} \right) r_n \left( \frac{dq_+}{dr_n} \right) (E_2 C_1 - E_0 C_3), \\ I_{\eta,\eta^+} &= \frac{1}{4} N q_+^2 \{ (f_1^2 + f_2^2) [E_2 C_0' - E_0 C_2'] \\ &\quad + 2f_1 f_2 [E_2 C_0' + E_0 C_2' - 2E_0 C_0'] \}, \\ I_{\xi,\eta^+} &= 0, \\ I_{\eta,p^+} &= -N r_n q_+ C_1 (E_3 - E_1) / 2, \\ I_{\eta,q^+} &= N r_n q_+ C_2' (E_2 - E_0) / 2, \\ I_{\xi,A^+} &= -N \left( \frac{1}{A_+} \frac{dA_+}{dr_n} \right) \dot{p}_+ E_1 (C_0 - C_2), \end{aligned}$$

$$\begin{aligned} I_{\eta,A^+} &= N \left( \frac{1}{A_+} \frac{dA_+}{dr_n} \right) q_+ C_1 (E_2 - E_0), \\ I_{\xi,\xi^+} &= \frac{1}{4} \dot{p}_+^2 \{ f_1^2 + f_2^2 - 2f_1 f_2 N [E_2 C_0 + E_0 C_2 - 2E_0 C_0] \}, \\ I_{\xi,q^+} &= -N r_n \dot{p}_+ E_1 (C_1 - C_3) / 2, \\ I_{\xi,p^+} &= N r_n \dot{p}_+ E_2 (C_0 - C_2) / 2, \end{aligned}$$

where  $N = (E_2 C_0 - E_0 C_2)^{-1}$ . The subscripts indicate the pairs of terms involved.

A similar set of terms is obtained for  $g_{--}$ . For  $r_n > r_0$  these differ from the above not only in that  $\dot{p}_+$  and  $q_+$  are replaced by  $\dot{p}_-$  and  $q_-$ , but also by the interchange of  $C_{2n}$  and  $C_{2n}'$ . For  $r_n < r_0$  several additional changes occur:

- (a)  $\frac{1}{A_-} \frac{dA_-}{dr_n}$  is replaced by  $\left( \frac{1}{A_-} \frac{dA_-}{dr_n} - \frac{1}{r_n} \right)$ .
- (b)  $E_n$  is replaced by  $E_{n+2}$ .
- (c) in the last two terms  $\dot{p}_+ E_n$  is replaced by  $\dot{p}_- E_{n+2} - (2/r_n) E_{n+1}$ .
- (d)  $I_{\xi,\xi^-}$  has the form

$$\begin{aligned} I_{\xi,\xi^-} &= \frac{1}{4} \dot{p}_-^2 \{ f_1^2 + f_2^2 - 2f_1 f_2 N [E_4 C_0' + E_2 C_2' - 2E_2 C_0'] \} \\ &\quad + \frac{\dot{p}_-}{2r_n} f_1 f_2 N [E_3 C_0' + E_1 C_2' - 2E_1 C_0'] \\ &\quad - \frac{1}{r_n^2} f_1 f_2 N [E_2 C_0' + E_0 C_2' - 2E_0 C_0']. \end{aligned}$$

In the calculation for  $f_{+-}$ , only those terms containing either  $\nabla_n(r_n \xi)$  or  $\nabla_n(r_n \eta)$  give a result different from zero because the others lead to integrands odd in the variable  $\eta$ . Thus we find

$$\begin{aligned} f_{+-} &= \frac{1}{16} \pi r_n^3 A_+ A_- (f_2 - f_1) \{ \dot{p}_+ (E_2 - E_0) [C_1(\Sigma) + C_1(\Delta)] \\ &\quad - q_+ E_1 [C_0(\Sigma) - C_0(\Delta) - C_2(\Sigma) + C_2(\Delta)] \}, \end{aligned}$$

where  $\Sigma = (q_+ + q_-)/2$ , and  $\Delta = (q_+ - q_-)/2$ . The argument of the terms  $E_n$  is  $(\dot{p}_+ + \dot{p}_-)$ . This expression is valid for  $r \geq r_c$ ; for  $r < r_c$ ,  $E_n$  becomes  $E_{n+1}$ .

Finally, we obtain the expression for  $g_{+-}$  as  $(f_2 - f_1) A_+ A_- \pi r_n^3 / 4$  times the sum of the terms

$$\begin{aligned} g_{\xi,\eta} &= -\frac{1}{4} \dot{p}_+ q_- \{ E_2 [C_0(\Sigma) + C_0(\Delta)] \\ &\quad - E_0 [C_2(\Sigma) + C_2(\Delta)] \}, \\ g_{\eta,\xi} &= -\frac{1}{4} q_+ \dot{p}_- \{ E_2 [C_0(\Sigma) - C_0(\Delta)] \\ &\quad - E_0 [C_2(\Sigma) - C_2(\Delta)] \}, \\ g_{A,\xi} &= -\frac{1}{A_+} \frac{dA_+}{dr_n} \frac{\dot{p}_-}{2} (E_2 - E_0) [C_1(\Sigma) - C_1(\Delta)], \\ g_{\xi,A} &= -\frac{1}{A_-} \frac{dA_-}{dr_n} \frac{\dot{p}_+}{2} (E_2 - E_0) [C_1(\Sigma) - C_1(\Delta)], \end{aligned}$$

$$g_{p,\xi} = + \frac{r_n \dot{p}_- \dot{d} p_+}{4 \dot{d} r_n} (E_3 - E_1) [C_1(\Sigma) - C_1(\Delta)],$$

$$g_{p,\eta} = - \frac{r_n q_- \dot{d} p_+}{4 \dot{d} r_n} E_1 [C_0(\Sigma) + C_0(\Delta) - C_2(\Sigma) - C_2(\Delta)],$$

$$g_{\xi,p} = \frac{r_n \dot{p}_+ \dot{d} p_-}{4 \dot{d} r_n} (E_3 - E_1) [C_1(\Sigma) - C_1(\Delta)],$$

$$g_{\eta,p} = - \frac{r_n q_+ \dot{d} p_-}{4 \dot{d} r_n} E_2 [C_0(\Sigma) - C_0(\Delta) - C_2(\Sigma) + C_2(\Delta)],$$

$$g_{q,\xi} = - \frac{r_n \dot{p}_- \dot{d} q_+}{4 \dot{d} r_n} (E_2 - E_0) [C_2(\Sigma) - C_2(\Delta)],$$

$$g_{q,\eta} = \frac{r_n q_- \dot{d} q_+}{4 \dot{d} r_n} E_1 [C_1(\Sigma) + C_1(\Delta) - C_3(\Sigma) - C_3(\Delta)],$$

$$g_{\xi,q} = - \frac{r_n \dot{p}_+ \dot{d} q_-}{4 \dot{d} r_n} (E_2 - E_0) [C_2(\Sigma) + C_2(\Delta)],$$

$$g_{\eta,q} = \frac{r_n q_+ \dot{d} q_-}{4 \dot{d} r_n} E_1 [C_1(\Sigma) + C_1(\Delta) - C_3(\Sigma) - C_3(\Delta)].$$

$$g_{A,\eta} = \frac{1}{A_+} \frac{\dot{d} A_+ q_-}{\dot{d} r_n} E_1 [C_0(\Sigma) + C_0(\Delta) - C_2(\Sigma) - C_2(\Delta)],$$

$$g_{\eta,A} = \frac{1}{A_-} \frac{\dot{d} A_- q_+}{\dot{d} r_n} E_1 [C_0(\Sigma) - C_0(\Delta) - C_2(\Sigma) + C_2(\Delta)],$$

Again, for  $r < r_c$ , modifications such as have already been pointed out must be made:  $(1/A_-)$   $(\dot{d} A_- / \dot{d} r_n)$  is replaced by  $(1/A_-)$   $(\dot{d} A_- / \dot{d} r_n) - (1/r_n)$ ,  $E_n$  by  $E_{n+1}$ , and  $\dot{p} E_n$  by  $\dot{p} E_{n+1} - (2/r_n) E_n$  in these expressions.

## μ-Mesonic Molecules. II. Molecular-Ion Formation and Nuclear Catalysis\*

STANLEY COHEN,† DAVID L. JUDD, AND ROBERT J. RIDDELL, JR.‡  
Lawrence Radiation Laboratory, University of California, Berkeley, California

(Received July 27, 1959)

The methods developed in the preceding paper are applied to the study of the behavior of μ mesons in liquid hydrogen. Numerically evaluated energy eigenvalues for the bound states of the various molecular-ion configurations are presented. Phase shifts and cross sections for the scattering of mesonic atoms from hydrogen and deuterium are given. It is shown that in the neighborhood of 0.2 ev the scattering of (dμ) atoms from protons exhibits a Ramsauer-Townsend effect with an anomalously small cross section occurring in this region. The existence of this effect provides an explanation for the appearance of "gaps" in the experimental observation of the catalytic process. The rate of exchange of mesons from protons to deuterons in pure deuterium is calculated along with the rates of formation of the (pμp)<sup>+</sup>, (pμd)<sup>+</sup>, and (dμd)<sup>+</sup> molecular ions. It is shown that the predominant mechanism for the formation of the molecular ions is dipole electron ejection. These results are shown to be in agreement with available experimental data. A semiphenomenological treatment of the (pd) nuclear reaction is also given. A rough estimate of the γ-emission process indicates that the dominant mode of emission is from the singlet proton spin states.

### I. INTRODUCTION

THE theoretical possibility that μ mesons could greatly enhance the nuclear fusion of hydrogen and deuterium so that it might be observed experimentally was first suggested by Frank<sup>1</sup> and later was estimated by Zeldovich.<sup>2</sup> Later, this fusion was in fact observed by Alvarez *et al.* in a liquid-hydrogen bubble chamber in a process in which the energy of fusion was given to the μ meson.<sup>3</sup> Still later, γ rays from

the reaction were detected by Ashmore *et al.* with counters, again in liquid hydrogen.<sup>4</sup> The process has been further investigated theoretically by Jackson<sup>5</sup> and by Skyrme,<sup>6</sup> who have also given phenomenological descriptions in which the reaction is assumed to proceed through the following steps:

(a) A fast μ meson is rapidly slowed down and captured to form a hydrogen (pμ) or deuterium (dμ) mesonic atom. (Because the experiments have been carried out with much more hydrogen than deuterium, it is much more likely that hydrogen mesonic atoms are formed.)

(b) The neutral mesonic hydrogen atom will then move about with thermal energy, colliding with the atoms of the liquid. It may then undergo elastic scatter-

\* This work was performed under the auspices of the U. S. Atomic Energy Commission.

† Present address: Argonne National Laboratory, Lemont, Illinois.

‡ On leave of absence to U. S. Atomic Energy Commission, Washington, D. C.

<sup>1</sup> F. C. Frank, *Nature* **160**, 525 (1947).

<sup>2</sup> Ya. B. Zeldovitch, *Doklady Akad. Nauk. S.S.S.R.* **95**, 493 (1954).

<sup>3</sup> L. W. Alvarez, H. Bradner, F. S. Crawford, Jr., J. A. Crawford, P. Falk-Vairant, M. L. Good, J. D. Gow, A. H. Rosenfeld, F. Solmitz, L. Stevenson, H. K. Ticho, and R. D. Tripp, *Phys. Rev.* **105**, 1127 (1957).

<sup>4</sup> Ashmore, Nordhagen, Strauch, and Townes, *Proc. Phys. Soc. (London)* **71**, 161 (1958).

<sup>5</sup> J. D. Jackson, *Phys. Rev.* **106**, 330 (1957).

<sup>6</sup> T. H. R. Skyrme, *Phil. Mag.* **2**, 910 (1957).