

noted, however, that the behavior of the s -wave scattering determined in this analysis is dependent upon the assumption that only one resonant p -wave state is present. The resonant parameters used in the analysis of the p -wave resonant state in Be^7 at an excitation energy of 7.58 MeV are shown in Table I along with the corresponding values for the mirror level in Li^7 as given by Gabbard. (The author wishes to thank Dr. F. Gabbard for permission to use the results of his calculations.)

ACKNOWLEDGMENTS

The author wishes to express his gratitude to Dr. C. A. Barnes for suggesting this work and for his continuous encouragement and discussions concerning it. The interest shown by Dr. W. A. Fowler and Dr. T. Lauritsen was greatly appreciated. The advice given by Dr. R. F. Christy on the theoretical analysis of the data was also very helpful. The author would like to thank Mrs. Barbara Zimmerman for her assistance in performing part of the calculations.

New Isomers of Astatine-212

W. BARCLAY JONES

Lawrence Radiation Laboratory, University of California, Berkeley, California

(Received 8 March 1963)

The ${}_{83}\text{Bi}^{209}(\alpha, n){}_{85}\text{At}^{212}$ reaction was investigated in the energy range of 17 to 25 MeV. Astatine-212 was observed to decay by alpha-particle emission. An alpha decay group of 7.60 and 7.66 MeV having a half-life of 0.305 sec and another group of 7.82 and 7.88 MeV having a half-life of 0.120 sec were observed. Relative excitation functions were obtained for both isomers. An energy-level diagram for the alpha decay of astatine-212 is proposed.

THE ${}_{83}\text{Bi}^{209}(\alpha, n){}_{85}\text{At}^{212}$ reaction was investigated at the Crocker Laboratory 60-in. cyclotron of the University of California. Previous work on this reaction¹⁻⁴ reports that At^{212} has a half-life of 0.20 sec, and emits alpha particles of 7.6 or 7.88 MeV.




FIG. 1. Spectrum of alpha energies for 24-MeV alpha particles on bismuth.

In this experiment, the alpha decay energies were measured with a phosphorus-diffused-junction counter having an energy resolution of 30 keV. The spectra were observed at selected intervals between cyclotron beam bursts. The spectrum taken at a bombarding energy of 24 MeV is shown in Fig. 1 (the small peaks at 5.63, 6.04, 6.28, 6.78, and 8.78 MeV are due to the calibration source, Th^{228}). The results of the experiment are summarized in Table I. The half-life associated with each of these alpha energies was measured individually by time analysis of each pulse height.

A search was made for a gamma transition between the states responsible for the 7.82- and 7.60-MeV alpha groups by means of detecting the conversion electrons; however, no such transition was observed. Less than 1% of the alpha activity could have a gamma decay in the energy range from 100 to 600 keV; however, a ≈ 63 -keV transition was observed with a half-life of ~ 0.13 sec.⁵

TABLE I. Alpha-decay energies and half-lives for At^{212} .

Alpha decay energy (MeV)	Half-life (sec)	Approximate relative abundance (%)	Hindrance factor
7.60	0.305	20	6200
7.66	0.305	80	1700
7.82	0.120	80	1600
7.88	0.120	20	9500

* Work supported by the U. S. Atomic Energy Commission.
¹ M. Weissbluth, T. M. Putnam, and E. Segrè (unpublished work) reported by D. Strominger, J. M. Hollander, and G. T. Seaborg, *Rev. Mod. Phys.* **30**, 585 (1958).
² M. M. Winn, *Proc. Phys. Soc. (London)* **A67**, 949 (1954).
³ J. C. Ritter and W. G. Smith, *Phys. Rev.* **128**, 1778 (1962).
⁴ R. G. Griffioen and R. D. Macfarlane, in *Lawrence Radiation Laboratory Report UCRL-10023*, 1962 (unpublished).

⁵ F. S. Stephens and R. M. Diamond, Lawrence Radiation Laboratory (private communication).




FIG. 2. Proposed energy-level diagram for alpha decay of At²¹².

The data indicate that the level structure might be that shown in Fig. 2.

The 4+ and 5+ assignments for Bi²⁰⁸ are from theoretical computations by Wahlborn,⁶ and are justified by experimental work.⁷ The shell-model configuration for nuclei in the region of At²¹² suggests that the neutron outside the closed shell⁸ should be $^2g_{9/2}$ and that the protons should be $(1h_{9/2})^3$; however, by the predictions of Pryce⁹ this would lead to a ground state of 0- and an isomeric state of 9-. The 0- to 4+ transition is forbidden, so the assignment of 0- or 4+ does not appear to be justified. Alternately, as in Bi²¹⁰, the lower spin state for At²¹² could be 1-.

Figure 3 shows the relative excitation functions for these two isomers, as well as the absolute cross section




FIG. 3. Excitation functions for Bi²⁰⁹(α, xn)At. The triangles refer to the 0.12-sec state; the circles refer to the 0.305-sec state. Cross section units for the two isomers of At²¹² are not the same. Data for the At²¹¹ cross section are taken from reference 10.

for Bi²⁰⁹($\alpha, 2n$)At²¹¹.¹⁰ However, neither the (α, n) cross sections relative to each other nor the absolute cross sections have been determined.

The hindrance factors were calculated from the empirical relation: $\log_{10}F = \log_{10}t_{1/2} - A_Z Q_{\text{eff}}^{-1/2} - B_Z$, where A_Z and B_Z are the arithmetical means of corresponding values for the two adjacent even atomic numbers.¹¹

⁶ S. Wahlborn, Nucl. Phys. **3**, 644 (1957).

⁷ P. Mukherjee and B. L. Cohen, Phys. Rev. **127**, 1284 (1962).

⁸ H. J. Mang, Phys. Rev. **119**, 1069 (1960).

⁹ M. H. L. Pryce, Proc. Phys. Soc. (London) **A65**, 773 (1952).

¹⁰ E. L. Kelly and E. Segrè, Phys. Rev. **75**, 999 (1949).

¹¹ C. J. Gallagher and J. O. Rasmussen, J. Inorg. Nucl. Chem. **3**, 333 (1957).