

Green Function Theory of Ferromagnetism*

HERBERT B. CALLEN

Laboratory for Research on the Structure of Matter, University of Pennsylvania, Philadelphia, Pennsylvania

(Received 26 December 1962)

A theory of ferromagnetism for general spin, approximately valid through the entire temperature range, is given. At low temperatures the magnetization agrees with the Dyson results, having no term in T^3 and having a term in T^4 equal to that found by Dyson in first Born approximation; terms arising from the approximations of the theory first appear in order $T^{3(2S+1)/2}$, so that a spurious T^3 term does appear for $S=1/2$, but for no other spin. Curie temperatures are within a few percent of the Brown and Luttinger estimates for spins greater than unity, and agree within 1% of the Domb and Sykes estimate of the large-spin limit. The susceptibility at high temperatures agrees with the Opechowski expansion to terms in $1/T^2$. The quasiparticle energies are renormalized by the energy at low temperature and by the magnetization at higher temperature. The Green function is decoupled by a physical criterion involving self-consistency of the decoupling at all temperatures. The Green function method is extended to higher spin by a technique of parametrizing the Green function and explicitly finding the functional dependence on this parameter by solution of an auxiliary differential equation.

1. INTRODUCTION

THE Heisenberg model of a ferromagnet has been theoretically analyzed by Dyson¹ by series expansion in powers of T , valid at low temperatures, and by Opechowski² by series expansion in $1/T$, valid at very high temperatures. We here develop an approximate theory which covers the entire temperature range, including the particularly interesting intermediate region in the neighborhood of the Curie temperature, and which agrees satisfactorily with the rigorous results at both very low and very high temperatures.

At low temperatures the magnetization has terms of order $T^{3/2}$, $T^{5/2}$, $T^{7/2}$ which agree with the Dyson results, it properly has no term in T^3 , and the term in T^4 is equal to that found by Dyson in first Born approximation. Terms arising from the approximation in the theory first arise in order $T^{3(2S+1)/2}$, so that the case of $S=1/2$ is an exception to the above statements, having a spurious T^3 term. Curie temperatures are quite close ($\approx 3\%$ for spin 2) to the values estimated by Brown and Luttinger³ by extrapolation of the high-temperature series expansion, except for very small values of spin (again the case of spin 1/2 is particularly unsatisfactory). The Curie temperatures agree within 1% with the estimate of Domb and Sykes⁴ for the high-spin limit. The susceptibility at high temperatures for all spin values agrees with the Opechowski² expansion to terms in $1/T^2$.

The quasiparticle energies are equivalent to simple spin-wave energies "renormalized" by a factor which is proportional to the thermodynamic energy at low temperatures, but which becomes proportional to the magnetization at higher temperatures.

An heuristic interpretation of the renormalization of quasiparticle energies has been given by Keffer and

Loudon.⁵ They point out that, at low temperatures where only long-wavelength spin waves are excited, the local magnetization direction varies slowly through the crystal. Excitation of an additional spin wave is analogous to excitation of a ripple relative to this slowly varying local magnetization. The effective exchange integral determining the energy of this ripple is influenced by the angle between neighboring spins in the slowly varying background medium. This angle also determines the thermodynamic energy; hence, the renormalization of the spin-wave energy by the thermodynamic energy. However, at higher temperatures, the thermally excited excitations have wavelengths comparable to the interspin distance, and the correlation distance in the background medium is as short, or shorter, than the wavelength of the particular excitation being considered. In this region our results indicate that the effective exchange integral is influenced by the angle relative to the average magnetization; that is, the spin-wave energies are renormalized by the average magnetization.

The Heisenberg ferromagnet with spin 1/2 was analyzed by Tyablikov⁶ using the technique of double-time temperature-dependent Green functions.^{7,8} Extension of the theory to higher spin has been achieved recently by Tahir-Kheli and ter Haar.⁹ The present theory differs from those applications of the Green function method in two respects. First, the decoupling of the higher order Green functions is guided by a plausible physical criterion. Second, the method of employing Green functions for general spin is simplified; the Green function is parametrized and the functional dependence on this parameter is found explicitly by

⁵ F. Keffer and R. Loudon, *J. Appl. Phys.* **32**, 25 (1961).

⁶ S. V. Tyablikov, *Ukr. Mat. Zh.* **11**, 287 (1959).

⁷ N. N. Bogolyubov and S. V. Tyablikov, *Dokl. Akad. Nauk SSSR* **126**, 63 (1959) [translation: *Soviet Phys.—Doklady* **4**, 604 (1959)].

⁸ A convenient review of Green functions and of Tyablikov's application of them to ferromagnetism is given by D. N. Zubarev, *Usp. Fiz. Nauk* **71**, 71 (1960) [translation: *Soviet Phys.—Usp.* **3**, 320 (1960)].

⁹ R. Tahir-Kheli and D. ter Haar, *Phys. Rev.* **127**, 88 (1962).

* Supported by the U. S. Office of Naval Research.

¹ F. J. Dyson, *Phys. Rev.* **102**, 1217, 1230 (1956).

² W. Opechowski, *Physica* **4**, 181 (1937); **6**, 1112 (1938).

³ H. A. Brown and J. M. Luttinger, *Phys. Rev.* **100**, 685 (1955).

⁴ C. Domb and M. F. Sykes, *Phys. Rev.* **128**, 168 (1962).

solution of an auxiliary differential equation. This single differential equation replaces the $2S$ simultaneous Green function equations in the method of Tahir-Kheli and ter Haar.⁹

2. THE GREEN FUNCTION

The problem to be considered is the statistical mechanics of the system described by the Hamiltonian

$$\mathcal{H} = -\mu H \sum_g S_g^z - \sum_{g,f} J(g-f) \mathbf{S}_g \cdot \mathbf{S}_f, \quad (1)$$

where μS is the magnetic moment per ion; H is the applied magnetic field (in the negative z direction); \mathbf{S}_g is the spin operator (in units of \hbar) for the ion at site g ; and $J(g-f)$ is the exchange integral between ions at sites g and f . The exchange integral is assumed to be a function only of the distance between ions; it is not restricted to nearest neighbors or to only positive values, but it is assumed that the ground state of the system is a ferromagnetically aligned state.

The temperature-dependent retarded Green function involving the two operators A and B , $\langle\langle A(t); B \rangle\rangle$, is defined by⁸

$$\langle\langle A(t); B \rangle\rangle \equiv -i\theta(t)\langle[A(t), B]\rangle, \quad (2)$$

where $A(t)$ is the Heisenberg operator at time t ; $\theta(t)$ is unity for positive t and zero for negative t ; square brackets denote a commutator; and single angular brackets denote an average with respect to the canonical density matrix of the system at temperature T . The Fourier transform of the Green function is a function of ω (or of $E = \hbar\omega$), and is denoted by $\langle\langle A; B \rangle\rangle_E$. It satisfies the equation of motion⁸

$$E\langle\langle A; B \rangle\rangle_E = (1/2\pi)\langle[A, B]\rangle + \langle\langle [A(t), \mathcal{H}]; B \rangle\rangle_E. \quad (3)$$

If this equation can be solved for $\langle\langle A; B \rangle\rangle_E$ one then extracts knowledge of the correlation function $\langle BA(t) \rangle$ from the relation⁸

$$\langle BA(t) \rangle = \lim_{\epsilon \rightarrow 0} i \int_{-\infty}^{\infty} \frac{\langle\langle A; B \rangle\rangle_{\hbar\omega+i\epsilon} - \langle\langle A; B \rangle\rangle_{\hbar\omega-i\epsilon}}{\exp(\hbar\omega/kT) - 1} e^{-i\omega t} d\omega. \quad (4)$$

Equations (3) and (4) are the only equations required for the application of the Green function method.

For reasons which will become evident subsequently, we consider the Green function

$$G_t^a(g, l) \equiv \langle\langle S_g^+(t); e^{aS_l^z} S_l^- \rangle\rangle, \quad (5)$$

where a is a parameter. The Fourier transform of this Green function, $G_E^a(g, l)$, satisfies the equation of motion [Eq. (3)]

$$EG_E^a(g, l) = \frac{1}{2\pi} \Theta(a) \delta_{g,l} + \langle\langle [S_g^+(t), \mathcal{H}]; e^{aS_l^z} S_l^- \rangle\rangle_E, \quad (6)$$

where

$$\Theta(a) \equiv \langle[S^+, e^{aS^z} S^-]\rangle. \quad (7)$$

The commutator of S_g^+ with the Hamiltonian, required in the last term of Eq. (6), is easily computed, giving

$$EG_E^a(g, l) = \frac{\Theta(a)}{2\pi} \delta_{g,l} + \mu H G_E^a(g, l) - 2 \sum_f J(g-f) \langle\langle S_g^z S_f^+ - S_f^z S_g^+; e^{aS_l^z} S_l^- \rangle\rangle_E. \quad (8)$$

The remaining problem is to express the higher order Green function on the right in terms of lower order Green functions, so that Eq. (8) can be explicitly solved for $G_E^a(g, l)$.

3. A DECOUPLING APPROXIMATION

The essential approximation in the methods of Tyablikov and of Tahir-Kheli and ter Haar consists of ignoring the fluctuations of S_g^z , replacing this operator by its average value:

$$\langle\langle S_g^z S_f^+; B \rangle\rangle \rightarrow \langle S^z \rangle \langle\langle S_f^+; B \rangle\rangle. \quad (\text{Tyablikov}) \quad (9)$$

This approximation results in the magnetization renormalization of quasiparticle energies, in disagreement with the low-temperature theory.

The decoupling approximation to be used here is most clearly described for the special case of spin 1/2. In that case we can write S_g^z in either of the following forms

$$S_g^z = S - S_g^- S_g^+, \quad (S = 1/2), \quad (10)$$

$$S_g^z = \frac{1}{2}(S_g^+ S_g^- - S_g^- S_g^+), \quad (11)$$

or, multiplying the first of these equations by an arbitrary parameter α and the second by $(1-\alpha)$ and adding,

$$S_g^z = \alpha S + \frac{1}{2}(1-\alpha) S_g^+ S_g^- - \frac{1}{2}(1+\alpha) S_g^- S_g^+. \quad (S = 1/2) \quad (12)$$

The Green function $\langle\langle S_g^- S_g^+ S_f^+; B \rangle\rangle$ is reasonably decoupled in the symmetric form¹⁰

$$\langle\langle S_g^- S_g^+ S_f^+; B \rangle\rangle \rightarrow \langle S_g^- S_g^+ \rangle \langle\langle S_f^+; B \rangle\rangle + \langle S_g^- S_f^+ \rangle \langle\langle S_g^+; B \rangle\rangle, \quad (13)$$

and similarly for the Green function $\langle\langle S_g^+ S_g^- S_f^+; B \rangle\rangle$. Thus, the identity (12) leads to

$$\langle\langle S_g^z S_f^+; B \rangle\rangle \rightarrow \langle S^z \rangle \langle\langle S_f^+; B \rangle\rangle - \alpha \langle S_g^- S_f^+ \rangle \langle\langle S_g^+; B \rangle\rangle. \quad (14)$$

If α is chosen as unity, the result corresponds to decoupling on the basis of identity (10); $\alpha = 0$ corresponds to decoupling on the basis of identity (11); $\alpha = -1$ corresponds to decoupling on the basis of the identity

¹⁰ The remaining "contraction," $\langle S_g^+ S_f^+ \rangle \langle\langle S_g^-; B \rangle\rangle$ vanishes because the operator $S_g^+ S_f^+$ is not diagonal in the total z component of spin.

$S_g^z = -S + S_g^+ S_g^-$. Thus, we are faced with the possibility of obtaining a correction to the Tyablikov decoupling with either a positive or negative sign, or no correction at all, or any intermediate value, depending on the choice of α . Clearly, a physical criterion is required at this point.

The operator $S^- S^+$ in Eq. (10) represents the deviation of S^z from $+S$. It is this operator $S^- S^+$ which is treated approximately when decoupling on the basis of Eq. (10). It, therefore, seems reasonable to use Eq. (10) when the deviation from $S^z = S$ is small; that is, when $\langle S^z \rangle \simeq S$.

Similarly, the operator $\frac{1}{2}(S^+ S^- - S^- S^+)$ in Eq. (11) represents the deviation of S^z from zero, and it, therefore, is reasonable to use Eq. (11) as the basis of decoupling when $\langle S^z \rangle \simeq 0$.

Both of the above observations are contained in the choice

$$\alpha = \langle S^z \rangle / S, \quad (S = 1/2) \quad (15)$$

for then Eq. (12) becomes

$$S_g^z = \langle S^z \rangle + \left[\frac{S - \langle S^z \rangle}{2S} S_g^+ S_g^- - \frac{S + \langle S^z \rangle}{2S} S_g^- S_g^+ \right]. \quad (16)$$

The operator in brackets, which is to be decoupled, represents the deviation of S^z from $\langle S^z \rangle$ and should be self-consistently small in all temperature regions.

Inserting the above value of α into the decoupling equation (19) gives

$$\begin{aligned} \langle \langle S_g^z S_f^+; B \rangle \rangle &\rightarrow \langle S^z \rangle \langle \langle S_f^+; B \rangle \rangle \\ &- \frac{\langle S^z \rangle}{S} \langle S_g^- S_f^+ \rangle \langle \langle S_g^+; B \rangle \rangle, \quad (S = 1/2). \end{aligned} \quad (17)$$

This is the basic decoupling approximation for spin 1/2; we now generalize it for higher spin.

The analog of Eq. (10), for general spin, is

$$S_g^z = S(S+1) - (S_g^z)^2 - S_g^- S_g^+, \quad (18)$$

whereas, Eq. (11) remains true. Decoupling as before, and neglecting¹¹ the fluctuations of $(S^z)^2$, we find in this case as well

$$\begin{aligned} \langle \langle S_g^z S_f^+; B \rangle \rangle &\rightarrow \langle S^z \rangle \langle \langle S_f^+; B \rangle \rangle \\ &- \alpha \langle S_g^- S_f^+ \rangle \langle \langle S_g^+; B \rangle \rangle, \end{aligned} \quad (19)$$

where α is the fractional contribution of the identity (18), and $(1-\alpha)$ is the contribution of the identity (11), to this result. Unfortunately, $S_g^- S_g^+$ is no longer the only operator treated approximately in decoupling equation (18), and the interpretation of the decoupled operator as being the deviation from $S^z = +S$ is no

longer true. Hence, the choice of α is no longer quite so evident. However, we determine it by the following requirements:

(a) For $S = 1/2$, α should reduce to our previous result, or to $\alpha = \langle S^z \rangle / S$.

(b) For $\langle S^z \rangle = 0$, α should vanish. This requirement follows from the fact that identity (11) retains its interpretation for arbitrary S .

(c) For $\langle S^z \rangle \simeq S$ we expect that S^z should have the form $S^z \simeq S - n$, where n is a deviation which is of order unity rather than of order S .

Requirement (c) implies that $\alpha \langle S_g^- S_f^+ \rangle$ should be of order unity, rather than of order S , at low temperatures. Now $(1/2S) \langle S_g^- S_g^+ \rangle$ is the spin deviation in lowest order¹²; similarly $\langle S_g^- S_f^+ \rangle$ will be of order $2S$ if f and g are closely coupled. Hence, we take

$$\alpha = (1/2S) \langle S^z \rangle / S \quad (20)$$

and it is clear that this satisfies all of the physical requirements above. Equations (19) and (20) characterize the basic decoupling approximation of the theory.

4. SOLUTION OF THE GREEN FUNCTION EQUATION

Inserting the decoupling approximation into the equation of motion (8) gives

$$\begin{aligned} EG_E^a(g, l) &= \frac{\Theta(a)}{2\pi} \delta_{g, l} + \mu HG_E^a(g, l) \\ &- 2 \langle S^z \rangle \sum_f J(g-f) [G_E^a(f, l) - G_E^a(g, l)] \\ &+ \frac{\langle S^z \rangle}{S^2} \sum_f J(g-f) [\langle S_g^- S_f^+ \rangle G_E^a(g, l) \\ &- \langle S_f^- S_g^+ \rangle G_E^a(f, l)]. \end{aligned} \quad (21)$$

These equations are a set of coupled equations for various pairs of sites (g, l) , (f, l) . Translational invariance dictates consideration of the spatial Fourier transforms

$$G_E^a(\mathbf{k}) \equiv \sum_{g-l} e^{-i(\mathbf{g}-l) \cdot \mathbf{k}} G_E^a(g, l), \quad (22)$$

$$J(\mathbf{k}) \equiv \sum_{g-l} e^{-i(\mathbf{g}-l) \cdot \mathbf{k}} J(g-l), \quad (23)$$

$$\psi(\mathbf{k}, a) \equiv \sum_{g-l} e^{-i(\mathbf{g}-l) \cdot \mathbf{k}} \langle e^{a S_l^z} S_l^- S_g^+ \rangle. \quad (24)$$

Here $\mathbf{g} \cdot \mathbf{k}$ denotes the vector product $\mathbf{R}_g \cdot \mathbf{k}$, where \mathbf{R}_g is the position vector of the g site. Equation (21) then implies

¹¹ The Green function $\langle \langle (S_g^z)^2 S_f^+; B \rangle \rangle$ also can be symmetrically decoupled, but the results are of the same form as those obtained by the simpler procedure above.

¹² The appearance of $2S$ here corresponds to the identification of S^+ and S^- , respectively, as $(2S)^{1/2} a^+$ and $(2S)^{1/2} a^-$ in the leading terms both of the Dyson and the Holstein-Primakoff transformations. Here a^+ and a^- are destruction and creation operators of elementary boson-type excitations.

$$EG_E^a(\mathbf{k}) = \frac{\Theta(a)}{2\pi} + \mu H G_E^a(\mathbf{k}) - 2\langle S^z \rangle [J(\mathbf{k}) - J(0)] G_E^a(\mathbf{k}) + \frac{\langle S^z \rangle}{NS^2} \sum_{\mathbf{k}'} [J(\mathbf{k}') - J(\mathbf{k}' - \mathbf{k})] \psi(\mathbf{k}', 0) G_E^a(\mathbf{k}) \quad (25)$$

or

$$G_E^a(\mathbf{k}) = \frac{\Theta(a)}{2\pi[E - E(\mathbf{k})]} \quad (26)$$

where

$$E(\mathbf{k}) = \mu H + 2\langle S^z \rangle [J(0) - J(\mathbf{k})] + \frac{\langle S^z \rangle}{NS^2} \sum_{\mathbf{k}'} [J(\mathbf{k}') - J(\mathbf{k}' - \mathbf{k})] \psi(\mathbf{k}', 0). \quad (27)$$

The correlation function which can be obtained from $G_E^a(\mathbf{k})$ by Eq. (4) is the quantity $\psi(\mathbf{k}, a)$, defined in Eq. (24). Inserting Eq. (26) in (4), and taking $t=0$, we thereby obtain

$$\psi(\mathbf{k}, a) = \Theta(a) / (e^{E(\mathbf{k})/kT} - 1). \quad (28)$$

Equation (28), together with Eq. (27) defining $E(\mathbf{k})$, Eq. (24) defining $\psi(\mathbf{k}, a)$, and Eq. (7) defining $\Theta(a)$, is the basic equation of the theory. It must, however, be augmented by an explicit relationship between ψ and Θ , and it is this step which complicates the problem for spin $> 1/2$.

The problem which distinguishes the simple case of spin $1/2$ from the more complicated case of higher spin becomes evident if we particularize our solution temporarily to $S = 1/2$. We also take $a=0$, and we note from Eqs. (24) and (10) that

$$\bar{\psi}(0) \equiv \frac{1}{N} \sum \psi(\mathbf{k}, 0) = \langle S_a^- S_a^+ \rangle = \frac{1}{2} - \langle S^z \rangle, \quad (S = \frac{1}{2}). \quad (29)$$

Thus, $\bar{\psi}(0)$ is simply related to the magnetization. $N\bar{\psi}(0)$ is the total number of spin reversals in the crystal, and $\psi(\mathbf{k}, 0)$ is the occupation number of the basic excitation of wave vector \mathbf{k} . Similarly, from Eq. (7)

$$\Theta(0) = 2\langle S^z \rangle. \quad (30)$$

Hence, Eqs. (28) and (30) determine $\psi(\mathbf{k}, 0)$ as a function of $\langle S^z \rangle$, and Eq. (29) provides a requirement of self-consistency which determines $\langle S^z \rangle$.

For higher spin values Eq. (29) is replaced by

$$\bar{\psi}(0) \equiv \frac{1}{N} \sum_{\mathbf{k}} \psi(\mathbf{k}, 0) = \langle S_a^- S_a^+ \rangle = S(S+1) - \langle (S^z)^2 \rangle - \langle S^z \rangle. \quad (31)$$

This introduces the new quantity $\langle (S^z)^2 \rangle$, which is not determined by Eq. (29). Tahir-Kheli and ter Haar,⁹ therefore, introduce the Green function $\langle\langle S_a^+; (S_a^-)^2 S_a^+ \rangle\rangle$ to evaluate this quantity, but its solution introduces $\langle (S^z)^3 \rangle$. By introducing $2S$ such Green functions, and

by recalling that $(S^z)^{2S+1}$ is related by an identity to lower powers of S^z , they are thereby able to obtain a solution. That method gets laborious very quickly, but they have given explicit solutions to $S=3$.

By exploiting the functional dependence of $\Theta(a)$ and $\bar{\psi}$ on the parameter a , which was inserted in Eq. (5) for just this purpose, both Θ and $\bar{\psi}$ can be explicitly related to $\langle S^z \rangle$ for arbitrary spin, in close analogy with the case of spin $1/2$.

5. RELATIONSHIP OF Θ TO $\langle S^z \rangle$

Calculation of the commutator defining $\Theta(a)$ [Eq. (7)] is facilitated by the identity

$$[S^+, (S^z)^n] = \{(S^z - 1)^n - (S^z)^n\} S^+ \quad (32)$$

which is easily corroborated for $n=1, 2$ and extended to higher n by mathematical induction. It follows that

$$[S^+, e^{aS^z}] = (e^{-a} - 1) e^{aS^z} S^+, \quad (33)$$

and thence,

$$\Theta(a) = 2\langle e^{aS^z} S^z \rangle + (e^{-a} - 1) \langle e^{aS^z} S^+ S^- \rangle. \quad (34)$$

Expressing $S^+ S^-$ in terms of S^z by the identity (18),

$$\Theta(a) = S(S+1)(e^{-a} - 1) \langle e^{aS^z} \rangle + (e^{-a} + 1) \langle e^{aS^z} S^z \rangle - (e^{-a} - 1) \langle e^{aS^z} (S^z)^2 \rangle. \quad (36)$$

Finally, it is convenient to introduce the quantity

$$\Omega(a) \equiv \langle e^{aS^z} \rangle \quad (36)$$

and the notation

$$D \equiv d/da. \quad (37)$$

Then the two quantities of interest can be written

$$\Theta(a) = S(S+1)(e^{-a} - 1)\Omega + (e^{-a} + 1)D\Omega - (e^{-a} - 1)D^2\Omega \quad (38)$$

and

$$\bar{\psi}(a) = \langle e^{aS^z} S^- S^+ \rangle = S(S+1)\Omega - D\Omega - D^2\Omega. \quad (39)$$

The relationship (28) between $\psi(\mathbf{k}, a)$ and $\Theta(a)$ can be recast in a more convenient form as well, by defining

$$\phi(\mathbf{k}) \equiv 1 / (e^{E(\mathbf{k})/kT} - 1), \quad (40)$$

whence,

$$\psi(\mathbf{k}, a) = \phi(\mathbf{k}) \Theta(a). \quad (41)$$

The self-consistency requirement on the Green function is now contained in the condition

$$\bar{\psi}(a) = \Phi \Theta(a), \quad (42)$$

where

$$\Phi \equiv \frac{1}{N} \sum_{\mathbf{k}} \phi(\mathbf{k}). \quad (43)$$

This condition should determine $\Theta(a)$, and thence $\langle S^z \rangle$, which is just $\frac{1}{2}\Theta(0)$, as we see from Eq. (25). It is more convenient to determine $\Omega(a)$, and thence to find $\Theta(a)$ by Eq. (38). In fact, inserting Eqs. (38) and

(39) into (42) gives a differential equation for $\Omega(a)$:

$$D^2\Omega + \frac{(1+\Phi)e^a + \Phi}{(1+\Phi)e^a - \Phi} D\Omega - S(S+1)\Omega = 0. \quad (44)$$

This differential equation is the analog of the set of $2S$ coupled equations of Tahir-Kheli and ter Haar.

To completely determine the solution we require two boundary conditions. From the definition (36) we observe

$$\Omega(0) = 1. \quad (45)$$

The second condition is provided by the operator identity

$$\prod_{p=-S}^S (S^z - p) = 0. \quad (46)$$

Taking the average of this equation it can be written in the form

$$\prod_{p=-S}^S (D - p)\Omega(a=0) \equiv \mathfrak{D}_S \Omega(0) = 0. \quad (47)$$

In the Appendix it is shown that the solution of the differential Eq. (44) satisfying the boundary conditions (45) and (47) is

$$\Omega(a) = \frac{\Phi^{2S+1} e^{-Sa} - (1+\Phi)^{2S+1} e^{(S+1)a}}{[\Phi^{2S+1} - (1+\Phi)^{2S+1}][(1+\Phi)e^a - \Phi]}, \quad (48)$$

from which $\langle S^z \rangle$ or $\Theta(a)$ can be found by differentiation.

6. THE FORMAL SOLUTION

For convenience and clarity we recapitulate the final form of the equations, preparatory to analyzing their low-temperature and high-temperature behavior, Curie temperature, etc.

Given a spin magnitude S , a temperature T , a magnetic field H , and an exchange interaction with Fourier components $J(\mathbf{k})$, the quasiboson occupation number $\phi(\mathbf{k})$ is

$$\phi(\mathbf{k}) = 1 / (e^{E(\mathbf{k})/kT} - 1), \quad (49)$$

where

$$E(\mathbf{k}) = \mu H + 2\langle S^z \rangle \left\{ [J(0) - J(\mathbf{k})] + \frac{\langle S^z \rangle}{NS^2} \sum_{\mathbf{k}'} [J(\mathbf{k}') - J(\mathbf{k}' - \mathbf{k})] \phi(\mathbf{k}') \right\}. \quad (50)$$

This is an implicit equation for $\phi(\mathbf{k})$, involving the unknown quantity $\langle S^z \rangle$. However, $\langle S^z \rangle$ is given in terms of

$$\Phi = \frac{1}{N} \sum \phi(\mathbf{k}) \quad (51)$$

by

$$\langle S^z \rangle = D\Omega(0) = \frac{(S-\Phi)(1+\Phi)^{2S+1} + (S+1+\Phi)\Phi^{2S+1}}{(1+\Phi)^{2S+1} - \Phi^{2S+1}}. \quad (52)$$

Thus, Eqs. (49) and (50) constitute a set of coupled equations which must be solved self-consistently for $\langle S^z \rangle$.

7. NEAREST-NEIGHBOR INTERACTION

For simple lattices with nearest-neighbor interaction, the formalism simplifies markedly. In this case the exchange $J(\mathbf{k})$ is

$$J(\mathbf{k}) = J \sum_{\delta} \exp(i\mathbf{k} \cdot \delta), \quad (53)$$

where δ goes over the nearest neighbors of a representative ion; we assume all magnetic ions are crystallographically equivalent. Consider the sum

$$\mathcal{K} \equiv \sum_{\mathbf{k}'} [J(\mathbf{k}') - J(\mathbf{k}' - \mathbf{k})] \phi(\mathbf{k}'), \quad (54)$$

which appears in Eq. (50) for $E(\mathbf{k})$. Then

$$\mathcal{K} = J \sum_{\delta} [1 - \exp(-i\mathbf{k} \cdot \delta)] \sum_{\mathbf{k}'} \exp(i\mathbf{k}' \cdot \delta) \phi(\mathbf{k}'). \quad (55)$$

The sum over \mathbf{k}' is clearly independent of δ , by symmetry, so that it can be replaced by $\frac{1}{z} \sum_{\delta'} \sum_{\mathbf{k}'} \times \exp(i\mathbf{k}' \cdot \delta) \phi(\mathbf{k}')$, where z is the number of nearest neighbors. We thus obtain

$$\mathcal{K} = J^{-1}(0) [J(0) - J(\mathbf{k})] \sum_{\mathbf{k}'} J(\mathbf{k}') \phi(\mathbf{k}'), \quad (56)$$

where $J(0) = zJ$ is the $k=0$ Fourier component of the exchange interaction. This equation was first pointed out by Michelene Bloch.¹³ The sum $\sum_{\mathbf{k}'} J(\mathbf{k}') \phi(\mathbf{k}')$ is a function only of the temperature, $\langle S^z \rangle$, and H (and, of course, of J and the lattice structure).

$$\frac{1}{NJ(0)} \sum_{\mathbf{k}} J(\mathbf{k}) \phi(\mathbf{k}) \equiv f(T, \langle S^z \rangle, H) \quad (57)$$

and

$$E(\mathbf{k}) = \mu H + 2\langle S^z \rangle [J(0) - J(\mathbf{k})] [1 + (\langle S^z \rangle / S^2) f]. \quad (58)$$

Thus, for simple lattices (including simple cubic, body-centered cubic, and face-centered cubic lattices in particular) and for nearest-neighbor interaction the simple spin-wave energies are renormalized by a factor depending only on the temperature (for zero field), independent of the wave vector \mathbf{k} . Equations (57) and (58), together with the definition of $\phi(\mathbf{k})$ in terms of $E(\mathbf{k})$, constitute a pair of coupled equations for the renormalization function f . For these lattices and nearest neighbor interaction the complete solution would be obtained explicitly by solving Eqs. (57) and (58) for f , thereby obtaining $\phi(\mathbf{k})$ or Φ in terms of T , $\langle S^z \rangle$ and H . Then eliminating Φ between this equation and Eq. (52) would give $\langle S^z \rangle$ as a function of T and H .

¹³ M. Bloch, Phys. Rev. Letters 9, 286 (1962).

8. THE LOW-TEMPERATURE REGION

The summation (or integral) over \mathbf{k}' involved in the calculation of Φ for $H=0$ occurs in simple spin wave theory and has been carried out by Dyson¹ and others by standard series expansion procedures appropriate to low temperatures. Thus, if

$$E(\mathbf{k}) = 2SR[J(0) - J(\mathbf{k})] \tag{59}$$

and if a reduced temperature is defined by

$$\tau \equiv 3kT/4\pi\delta JS\nu, \tag{60}$$

then¹

$$\Phi = \zeta\left(\frac{3}{2}\right)(\tau/R)^{3/2} + \frac{3}{4}\pi\nu\zeta\left(\frac{5}{2}\right)(\tau/R)^{5/2} + \pi^2\omega\nu^2\zeta\left(\frac{7}{2}\right)(\tau/R)^{7/2} + \dots \tag{61}$$

Similarly

$$f \equiv \frac{1}{N\delta J} \sum_{\mathbf{k}} J(\mathbf{k})\phi(\mathbf{k}) = \zeta\left(\frac{3}{2}\right)(\tau/R)^{3/2} - \frac{1}{4}\pi\nu\zeta\left(\frac{5}{2}\right)(\tau/R)^{5/2} + (\omega - \frac{5}{4})\pi^2\nu^2\zeta\left(\frac{7}{2}\right)(\tau/R)^{7/2} + \dots \tag{62}$$

The constants ν and ω depend upon the lattice structure and are defined by

$$\nu = 1, \quad \omega = 33/32, \quad \text{for simple cubic;} \tag{63}$$

$$\nu = \frac{3}{4} \times 2^{2/3}, \quad \omega = 281/288, \quad \text{for body-centered cubic;} \tag{64}$$

$$\nu = 2^{1/3}, \quad \omega = 15/16, \quad \text{for face-centered cubic.} \tag{65}$$

To apply these results to our case we take [compare Eqs. (59) and (58)]

$$R = \frac{\langle S^2 \rangle}{S} \left[1 + \frac{\langle S^2 \rangle}{S^2} f \right]. \tag{66}$$

The remaining equation is Eq. (52) for $\langle S^2 \rangle$ in terms of Φ ; expanding it in powers of Φ (which is small at low temperature), we find

$$\langle S^2 \rangle = S - \Phi + (2S+1)\Phi^{2S+1} - (2S+1)^2\Phi^{2S+2} + O(\Phi^{2S+3}). \tag{67}$$

To disentangle the coupled equations we first substitute Eq. (66) for R into Eq. (62), solving for f as a series in τ :

$$f = \zeta\left(\frac{3}{2}\right)(S\tau/\langle S^2 \rangle)^{3/2} - \frac{1}{4}\pi\nu\zeta\left(\frac{5}{2}\right)(S\tau/\langle S^2 \rangle)^{5/2} + (\omega - \frac{5}{4})\pi^2\nu^2\zeta\left(\frac{7}{2}\right)(S\tau/\langle S^2 \rangle)^{7/2} + \dots - \frac{3}{2}\zeta^2\left(\frac{3}{2}\right)S/\langle S^2 \rangle^2\tau^3 - \frac{5}{8}\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)(S^2/\langle S^2 \rangle^3)\tau^4 + \dots \tag{68}$$

Inserting this series into Eq. (61) for Φ , we find

$$\Phi = \zeta\left(\frac{3}{2}\right)(S\tau/\langle S^2 \rangle)^{3/2} + \frac{3}{4}\pi\nu\zeta\left(\frac{5}{2}\right)(S\tau/\langle S^2 \rangle)^{5/2} + \pi^2\omega\nu^2\zeta\left(\frac{7}{2}\right)(S\tau/\langle S^2 \rangle)^{7/2} + \dots - \frac{3}{2}\zeta^2\left(\frac{3}{2}\right)S/\langle S^2 \rangle^2\tau^3 - \frac{3}{2}\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)(S^2/\langle S^2 \rangle^3)\tau^4 + \dots \tag{69}$$

Finally, inserting this equation into Eq. (67), we find

$$\langle S^2 \rangle = S - \zeta\left(\frac{3}{2}\right)\tau^{3/2} - \frac{3}{4}\pi\nu\zeta\left(\frac{5}{2}\right)\tau^{5/2} - \pi^2\omega\nu^2\zeta\left(\frac{7}{2}\right)\tau^{7/2} + \dots - (3/2S)\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)\tau^4 + \dots + (2S+1)\zeta^{2S+1}\left(\frac{3}{2}\right)\tau^{3S+3/2} + (2S+1)^2\frac{3}{4}\pi\nu\zeta^{2S}\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)\tau^{3S+5/2} + \dots \tag{70}$$

Neglecting, for the moment, the terms in $\tau^{3S+3/2}$, $\tau^{3S+5/2}$ we see that the coefficients of $\tau^{3/2}$, $\tau^{5/2}$, $\tau^{7/2}$ are in exact agreement with the Dyson result. The term in τ^3 is absent, as required by Dyson. The term involving τ^4 is given by Dyson¹ as

$$- (3/2S)\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)Q\tau^4, \tag{71}$$

where Q , the ratio of this term to our τ^4 term, is

$$Q = 1 + \frac{4}{3}[GS - 1]^{-1} + \alpha/3S \tag{72}$$

with

$$G = 10, 16, 24 \quad \text{for sc, bcc, fcc,} \tag{73}$$

$$\alpha = 0.52, 0.39, 0.34 \quad \text{for sc, bcc, fcc.} \tag{74}$$

Thus, our τ^4 term corresponds to the leading term ($Q=1$) of the Dyson result; that is, to the result which Dyson finds in the first Born approximation.

The terms $\tau^{3S+3/2}$, $\tau^{3S+5/2}$... in Eq. (70) seem to be spurious results of the Green function approximation. For $S=1/2$ they give incorrect contributions to the τ^3 and τ^4 terms. For spin unity the first spurious contribution appears in the $\tau^{9/2}$ term, and for spin 3/2 it appears first in the τ^6 term; thereafter it moves rapidly to higher order in τ . Consequently the spurious terms are of consequence in the low-temperature region only for spin 1/2.

Finally, it is of interest to substitute Eq. (70) into Eqs. (69), (68), and (66), to obtain Φ , f , and the renormalization factor R explicitly as series in τ . We thus find

$$\Phi = S - \zeta\left(\frac{3}{2}\right)\tau^{3/2} - \frac{3}{4}\pi\nu\zeta\left(\frac{5}{2}\right)\tau^{5/2} - \pi^2\omega\nu^2\zeta\left(\frac{7}{2}\right)\tau^{7/2} + \dots - (3/2S)\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)\tau^4 + \dots, \tag{75}$$

$$f = \zeta\left(\frac{3}{2}\right)\tau^{3/2} - \frac{1}{4}\pi\nu\zeta\left(\frac{5}{2}\right)\tau^{5/2} + (\omega - 5/4)\pi^2\nu^2\zeta\left(\frac{7}{2}\right)\tau^{7/2} + \dots - (1/8S)\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)\tau^4 + \dots, \tag{76}$$

and

$$R = 1 - \pi\nu/S\zeta\left(\frac{5}{2}\right)\tau^{5/2} - 5/4S\pi^2\nu^2\zeta\left(\frac{7}{2}\right)\tau^{7/2} + \dots - 2/S^2\zeta\left(\frac{3}{2}\right)\tau^3 - 21/8S^2\pi\nu\zeta\left(\frac{3}{2}\right)\zeta\left(\frac{5}{2}\right)\tau^4 + \dots + \frac{2S+1}{S}\zeta^{2S+1}(3/2)\tau^{3(2S+1)/2} + \frac{3(2S+1)^2}{4S}\pi\nu\zeta^{2S}(3/2)\zeta(5/2)\tau^{3S+5/2}. \tag{77}$$

It will be recalled that R is the "renormalization

factor"; the ratio of the actual quasiparticle energies to the simple spin-wave energies. At low temperatures the leading temperature dependence of R arises in the $\tau^{5/2}$ term. Hence, the spin-wave energies are renormalized by a factor proportional to the thermodynamic energy rather than the magnetization in this temperature region.

9. THE CURIE TEMPERATURES, AND THE HIGH-TEMPERATURE REGION

Just below the Curie temperature (assuming $H=0$) the average magnetization $\langle S^z \rangle$ is small, and the mean number of excited quasiparticles is large. In fact, expanding Eq. (52) in powers of Φ^{-1} we find

$$\langle S^z \rangle = \frac{1}{3}S(S+1)\Phi^{-1} + O(\Phi^{-2}). \quad (78)$$

Furthermore, since $E(\mathbf{k})$ is proportional to $\langle S^z \rangle$ the exponential in the Bose distribution can be expanded, giving

$$\Phi = \frac{1}{N} \sum_{\mathbf{k}} \left\{ \frac{2\langle S^z \rangle}{kT} [J(0) - J(\mathbf{k})] \left[1 + \frac{\langle S^z \rangle}{S^2} f \right] \right\}^{-1}. \quad (79)$$

Multiplying Eq. (79) by $\langle S^z \rangle$, and replacing $\langle S^z \rangle \Phi$ by $S(S+1)/3$ [from Eq. (78)] we then have, in the limit $\langle S^z \rangle \rightarrow 0$,

$$\frac{S(S+1)}{3} = \frac{kT}{2J(0)[1 + (\langle S^z \rangle / S^2) f]} F(-1), \quad (80)$$

where $F(-1)$ denotes the summation

$$F(-1) \equiv \frac{1}{N} \sum_{\mathbf{k}} \frac{J(0)}{J(0) - J(\mathbf{k})}. \quad (81)$$

This summation has been evaluated by Watson;¹⁴ it has the values

$$F(-1) = 1.51638(\text{sc}); 1.39320(\text{bcc}); 1.34466(\text{fcc}). \quad (82)$$

Equation (80) determines the Curie temperature. However we must evaluate the limiting value of the quantity $(\langle S^z \rangle / S^2) f$ which appears in that equation. From Eq. (57) defining f , again expanding the Bose factor,

$$\frac{\langle S^z \rangle}{S^2} f = \frac{\langle S^z \rangle}{NS^2 J(0)} \sum_{\mathbf{k}} J(\mathbf{k}) \left\{ \frac{2\langle S^z \rangle}{kT} [J(0) - J(\mathbf{k})] \times \left[1 + \frac{\langle S^z \rangle}{S^2} f \right] \right\}^{-1}, \quad (83)$$

¹⁴ G. N. Watson, *Quart. J. Math.* **10**, 266 (1939). See also, M. Tikson, *J. Res. Natl. Bur. Std.* **50**, 177 (1953).

TABLE I. Curie temperatures (kT_c/J) for cubic lattices, nearest neighbor interaction.

S	Simple cubic			Body-centered cubic			Face-centered cubic ^a		
	Brown Lutt.	t. Haar	Cal-len	Brown Lutt.	t. Haar	Cal-len	Brown Lutt.	t. Haar	Cal-len
1/2	1.9	2.0	2.7	2.39	2.9	3.7	4.2	4.5	5.6
1	5.4	5.3	6.5	7.82	7.7	9.1	12.7	11.9	13.9
3/2	10.6	9.4	11.7	15.42	14.4	16.6	24.7	22.3	25.5
2	17.5	15.8	18.5	25.17	23.0	26.2	40.0	35.7	40.1
5/2	25.8	23.1	26.8	37.10	33.5	37.9	58.7	52.1	58.3
3	35.7	31.6	36.4	51.19	45.9	51.6	80.9	71.4	79.5

^a For the face-centered cubic, Domb and Sykes (reference 4) give

$$\lim_{S \rightarrow \infty} [kT_c / JS(S+1)] = 6.38 \text{ (Domb and Sykes),}$$

$$= 5.95 \text{ (Tahir-Kheli and ter Haar),}$$

$$= 6.45 \text{ (Callen).}$$

or

$$\frac{\langle S^z \rangle}{S^2} f \left[1 + \frac{\langle S^z \rangle}{S^2} f \right] = \frac{F(-1) - 1}{2S^2 J(0)} kT. \quad (84)$$

Eliminating $(\langle S^z \rangle / S^2) f$ between this equation and Eq. (80), we finally find the equation determining the Curie temperature.

$$\frac{kT_c}{J} = \frac{2\frac{1}{3}(S+1)}{9F^2(-1)} [(4S+1)F(-1) - (S+1)]. \quad (85)$$

In Table I we give the values of kT_c/J as estimated by Brown and Luttinger³ by extrapolation of the series expansion for the susceptibility in powers of $1/T$. We also list the values obtained by Tahir-Kheli and ter Haar,⁹ and the values calculated from Eq. (85). It will be noted that our values are higher than those of Brown and Luttinger, whereas those of Tahir-Kheli and ter Haar are lower. For spin 2 the deviation of our results from the Brown and Luttinger values is of the order of 3%, and the agreement improves with increasing spin. In fact, Domb and Sykes have recently published⁴ an estimate of the limiting value $kT_c / JS(S+1)$ for large S , for the face-centered cubic lattice. This estimate was obtained by a painstaking examination of the systematics of the extrapolation of high-temperature series. They obtain the value of 6.384, with which our value of 6.45 agrees within 1%.

Domb and Sykes⁴ also give estimates of kT_c/J for $S=1/2$ and $S=1$ only, for the face-centered-cubic lattice; their values are 4.07 and 11.95, respectively. These are lower than the Brown and Luttinger results, and further aggravate the disparity between our values and the estimated values for these small spins. It is apparent, both from the Curie temperatures and from the low-temperature results, that our approximations are more reliable for large spin, being particularly bad for spin 1/2. Fortunately, most cases of practical

interest are associated with spins greater than 1/2, for which the results appear to be quite reliable.

It should, perhaps, be noted that the Curie temperatures are quite sensitive to the decoupling parameter α , of Eq. (20). If the chosen value of α were to be multiplied by $(\langle S^z \rangle / S)^\epsilon$, where ϵ is any positive constant no matter how small, the Curie temperatures would become identical to those obtained by Tahir-Kheli and ter Haar, whereas the low-temperature and the high-temperature behavior of the theory would remain unaltered.

Finally, the high-temperature expansion of the susceptibility χ is of interest. We assume $\langle S^z \rangle$ small, and maintain only terms proportional to the applied magnetic field. Equation (78) remains valid, Eq. (79) contains the additional Zeeman term, and the analog of Eq. (80) becomes

$$\frac{S(S+1)}{3} = \frac{\chi kT}{\mu^2} \frac{1}{N} \sum_{\mathbf{k}} \left\{ 1 + \frac{2\chi}{\mu^2} [J(0) - J(\mathbf{k})] \times \left[1 + \frac{\langle S^z \rangle}{S^2} f \right] \right\}^{-1}. \quad (86)$$

Similarly, the analog of Eq. (83) is

$$\frac{\langle S^z \rangle}{S^2} f = \frac{\chi kT}{\mu^2} \frac{1}{S^2 J(0)} \frac{1}{N} \sum_{\mathbf{k}} J(\mathbf{k}) \left\{ 1 + \frac{2\chi}{\mu^2} \times [J(0) - J(\mathbf{k})] \left[1 + \frac{\langle S^z \rangle}{S^2} f \right] \right\}^{-1}. \quad (87)$$

It is convenient to denote

$$[1 + (\langle S^z \rangle / S^2) f] \equiv X \quad (88)$$

and to define a quantity χ_1 by

$$\chi \equiv \lim_{H \rightarrow 0} \frac{\mu \langle S^z \rangle}{H} \equiv \frac{\mu^2 S(S+1)}{3kT} (1 + \chi_1). \quad (89)$$

Then, multiplying Eq. (86) by $[1 + (2\chi/\mu^2)J(0)X]$ and Eq. (87) by $-(2\chi/\mu^2)S^2 J(0)X$ and adding, we find

$$X^2 - \frac{4S+1}{3S} X + \frac{kT}{2S^2 J(0)} \frac{\chi_1}{1 + \chi_1} = 0. \quad (90)$$

Furthermore, expanding the summand in Eq. (86) we find

$$1 = (1 + \chi_1) \left[1 - \frac{2J(0)}{\mu^2} X \chi + \frac{4J^2(0)}{\mu^4} X^2 \chi^2 \frac{\chi + 1}{\chi} + \dots \right]. \quad (91)$$

In this summation we have employed the identities⁹

$$F(1) = 1, \quad F(2) = (\chi + 1) / \chi, \quad (92)$$

where

$$F(n) \equiv \frac{1}{N} \sum_{\mathbf{k}} \left(\frac{J(0) - J(\mathbf{k})}{J(0)} \right)^n. \quad (93)$$

The quantity X can now be eliminated between Eqs. (90) and (91), enabling χ_1 to be evaluated in a series in $1/T$. In this way we find

$$\chi = \frac{\mu^2 S(S+1)}{3kT} \left[1 + \frac{T_M}{T} + \left(1 - \frac{2S-1}{3\chi S} \right) \left(\frac{T_M}{T} \right)^2 + O\left(\frac{1}{T^3} \right) \right], \quad (94)$$

where T_M is the Curie temperature of the molecular field theory:

$$kT_M \equiv \frac{2}{3} \chi JS(S+1). \quad (95)$$

The two leading terms in this expansion are in agreement with the values found⁹ by a rigorous expansion of the susceptibility in powers of $1/T$.

ACKNOWLEDGMENT

I am very grateful to Dr. Raza Tahir-Kheli, who investigated a number of the limiting processes, read the manuscript, critically discussed the theory through its development, and repeatedly offered valuable suggestions.

APPENDIX

It is easily corroborated that the solution of the differential equation (44), satisfying the boundary conditions (46) and (47), is

$$\Omega(a) = \frac{1}{1 + \Phi} \times \frac{\omega(-S, a) \mathfrak{D}_S \omega(S+1, 0) - \omega(S+1, a) \mathfrak{D}_S \omega(-S, 0)}{\mathfrak{D}_S \omega(S+1, 0) - \mathfrak{D}_S \omega(-S, 0)}, \quad (A1)$$

where

$$\omega(x, a) \equiv \frac{e^{ax}}{(1 + \Phi)e^a - \Phi}. \quad (A2)$$

The evaluation of the derivatives $\mathfrak{D}_S \omega(x, 0)$ is then required to reduce this result to Eq. (48). Consider

$$\mathfrak{D}_S \omega(x, a) = \prod_{-S}^S (D - p) \frac{e^{ax}}{(1 + \Phi)e^a - \Phi} \quad (A3)$$

$$= \prod_{-S}^S \left(y \frac{d}{dy} - p \right) \frac{y^x}{(1 + \Phi)y - \Phi}, \quad (A4)$$

where

$$y = e^a. \quad (A5)$$

Expanding in powers of y , and noting that $y(d/dy)y^n$

$=ny^n$, we find

$$\mathcal{D}_{S\omega}(x,a) = -\frac{1}{\Phi} \sum_{n=x}^{\infty} \left(\frac{1+\Phi}{\Phi}\right)^{n-x} y^n \prod_{p=S}^S (n-p). \quad (\text{A6})$$

We now take $a=0$ (or $y=1$), let $x=S+1$, and change variables from n to $m=n-S$, and from p to $r=S-p$. Then

$$\mathcal{D}_{S\omega}(S+1, 0) = \frac{-1}{\Phi} \sum_{m=0}^{\infty} \frac{(m+2S+1)!}{m!} \left(\frac{1+\Phi}{\Phi}\right)^m. \quad (\text{A7})$$

Similarly, taking $a=0$, letting $x=-S$ and changing variables from n to $m=n-S-1$, and from p to $r=S$

$+1-p$, we find

$$\mathcal{D}_{S\omega}(-S, 0) = -\frac{(1+\Phi)^{2S+1}}{\Phi^{2S+2}} \times \sum_{m=0}^{\infty} \frac{(m+2S+1)!}{m!} \left(\frac{1+\Phi}{\Phi}\right)^m. \quad (\text{A8})$$

From Eqs. (A7) and (A8), we note that the ratio of the relevant quantities is

$$\mathcal{D}_{S\omega}(-S, 0)/\mathcal{D}_{S\omega}(S+1, 0) = (1+\Phi)^{2S+1}/\Phi^{2S+1}. \quad (\text{A9})$$

Finally, inserting this ratio into Eq. (A1) gives Eq. (48).

Electron Number of the Nitrogen Atom in Mn_4N

MASAO KURIYAMA

Westinghouse Research Laboratories, Beulah Road, Churchill Borough, Pittsburgh, Pennsylvania,
and

Institute for Solid State Physics, University of Tokyo, Azabu, Minato-ku, Tokyo, Japan

AND

SUKEAKI HOSOYA AND TADASU SUZUKI*

Institute for Solid State Physics, University of Tokyo, Azabu, Minato-ku, Tokyo, Japan

(Received 26 December 1962)

In order to obtain some information about the effective electron number of nitrogen in Mn_4N , the x-ray atomic scattering factors of nitrogen and manganese were measured by Fe $K\alpha$ radiation. The experimental values of the scattering factor of nitrogen determined within a resultant probable error of 5% were close to the theoretical curve for N^0 or N^{1-} . This result contradicts the donor theory in which a nitrogen atom is supposed to donate electrons to the neighboring manganese atoms.

INTRODUCTION

RECENTLY, Takei *et al.*^{1,2} have shown by a neutron diffraction study that the magnetic structure of Mn_4N is ferrimagnetic and is explicable in terms both of the donor property of nitrogen and of the energy splittings of manganese atoms caused by their local environments. In the donor theory which was proposed by Guillaud,³ Wiener and Berger,⁴ and Juza and Puff,⁵ a nitrogen atom is supposed to donate one electron to each of the three face-centered manganese atoms.

In order to obtain some information about the effective electron number of nitrogen, the x-ray atomic scattering factors of nitrogen and manganese in Mn_4N

were measured by Fe $K\alpha$ radiation for two samples supplied separately by Mekata and Takei, with the compositions $\text{Mn}_4\text{N}_{0.98}$ and $\text{Mn}_4\text{N}_{1.00}$, respectively, the latter of which contained a very small amount of MnO .

In Mn_4N , manganese atoms occupy the sites of a face-centered cubic lattice with a nitrogen atom at the body-centered position of the unit cell. There are four types of structure factor as follows:

$$4f_{\text{Mn}} + f_{\text{N}} \text{ for all even indices,} \quad (1)$$

$$4f_{\text{Mn}} - f_{\text{N}} \text{ for all odd indices,} \quad (2)$$

$$f_{\text{N}} \text{ for mixed indices with } h+k+l = \text{even,} \quad (3)$$

$$-f_{\text{N}} \text{ for mixed indices with } h+k+l = \text{odd.} \quad (4)$$

The last two types of structure factor may be used for determining the values of the scattering factor of nitrogen in the range of small scattering angles, where the behavior of the scattering factor is more sensitively influenced by the total electron number of the relevant atom than in other angular ranges.

* Research fellow from College of General Education, University of Tokyo, Tokyo, Japan.

¹ W. J. Takei, G. Shirane, and B. C. Frazer, *Phys. Rev.* **119**, 122 (1960).

² W. J. Takei, R. R. Heikes and G. Shirane, *Phys. Rev.* **125**, 1893 (1962).

³ C. P. Guillaud, *Rev. Mod. Phys.* **25**, 119 (1953).

⁴ G. W. Wiener and J. A. Berger, *Trans. AIME* **203**, 360 (1955).

⁵ R. Juza and H. Puff, *Z. Elektrochem.* **61**, 810 (1957).