## Fission Fragment Kinetic Energies of Cf<sup>246</sup>, Cf<sup>248</sup>, and Cf<sup>254†</sup>

A. M. FRIEDMAN, J. W. MEADOWS, A. B. SMITH, P. R. FIELDS, J. MILSTED, AND J. F. WHALEN Argonne National Laboratory, Argonne, Illinois (Received 27 March 1963)

Fission fragment kinetic energy spectra of Cf<sup>246</sup>, Cf<sup>248</sup>, and Cf<sup>254</sup> were measured by comparison to Cf<sup>252</sup>. The most probable fragment energies were (in MeV):  $Cf^{246}$ ,  $84.8\pm1.3$  and  $110.8\pm1.5$ ;  $Cf^{248}$ ,  $81.6\pm0.9$  and  $107.1 \pm 0.8$ ; Cf<sup>254</sup>, 83.0  $\pm 2$  and  $103.1 \pm 2$ .

SURVEY of the kinetic energy of fission frag-A ments from spontaneous fission or neutroninduced fission shows an increasing amount of kinetic energy in the fragments as the fission parameter  $Z^2/A$ increases.<sup>1</sup> It appeared interesting to determine how the fission kinetic energy varied as the mass number varied for a given Z. A convenient series for this study was the californium isotopes Cf<sup>246</sup>, Cf<sup>248</sup>, Cf<sup>252</sup>, and Cf<sup>254</sup>. Cf<sup>252</sup> had already been studied<sup>1,2</sup> and recently data on the fission kinetics of Cf<sup>254</sup> has been reported.<sup>3</sup> In this paper the fission kinetic energy of Cf<sup>246</sup>, Cf<sup>248</sup>, and Cf<sup>254</sup> are reported and compared with Cf<sup>252</sup>.

Cf<sup>246</sup> and Cf<sup>248</sup> were prepared simultaneously by irradiating curium containing 98% Cm<sup>244</sup> and 2% Cm<sup>246</sup> with 41-MeV helium ions in the 60 in. Argonne cyclotron. The californium was isolated from the curium by a series of ionic exchange columns. The californium was prepared in a form suitable for measuring the kinetic energies of the fission fragments by volatilizing the oxide onto a quartz plate in a special vacuum evapora-

TABLE I. Most probable fission fragment kinetic energies (in MeV).

Isotope	E heavy	E light	E total	Mass ratio
Cf <sup>246</sup> Cf <sup>248</sup> Cf <sup>252</sup> <sup>a</sup> Cf <sup>254</sup>	$84.8\pm1.3$ $81.6\pm0.9$ $79.8\pm1$ $83.0\pm2$	$\begin{array}{c} 110.8 \pm 1.5 \\ 107.1 \pm 0.8 \\ 104.7 \pm 1 \\ 103.1 \pm 2 \end{array}$	$\begin{array}{c} 195.6{\pm}2.0\\ 188.7{\pm}1.3\\ 184.5{\pm}1.4\\ 186.1{\pm}2.8 \end{array}$	$\begin{array}{c} 1.31{\pm}0.03\\ 1.31{\pm}0.02\\ 1.31{\pm}0.03\\ 1.24{\pm}0.05\end{array}$

<sup>a</sup> Standard; see Ref. 2.

<sup>2</sup> J. C. Milton and J. S. Frazer, Phys. Rev. 111, 877 (1958). <sup>3</sup> R. Brandt, Lawrence Radiation Laboratory Report UCRL-10481, 1962 (unpublished).

tor. At the time the source was prepared 99.5% of the spontaneous fission events of the source was due to the 35-h Cf<sup>246</sup>; after about 30 days essentially all the spontaneous fissions observed were due to the longer lived Cf<sup>248</sup>.

A highly enriched sample of Cf<sup>254</sup> was prepared by utilizing the small electron-capture branching ratio  $(\sim 0.1\%)$  of the 38-h isomer of Es<sup>254</sup>. Es<sup>254</sup> was prepared by irradiating a sample of Cf<sup>252</sup> for 6 months in the Materials Testing Reactor. The Es<sup>254</sup> was chemically separated from the californium and fission products and allowed to decay. The products Cf<sup>250</sup>, formed from the  $\alpha$  decay of accompanying Fm<sup>254</sup>, and Cf<sup>254</sup> from the electron capture decay of Es<sup>254</sup> were isolated. The mass ratio of Cf<sup>250</sup> to Cf<sup>254</sup> was about 1000 to 1. These isotopes were then separated in the Argonne isotope separator. In the final product the Cf<sup>250</sup> made a negligible contribution to the total fissions. The separated Cf<sup>254</sup> was collected on aluminum and contained 1 spontaneous fission per min; this was used directly for energy measurements.

The fission fragment kinetic energy spectrum of each sample was measured with silicon diffuse junction detectors and compared to a Cf<sup>252</sup> standard at frequent intervals during each run. Due to the small amounts of activity it was only feasible to measure single-sided noncoincident spectra.

Table I lists the most probable kinetic energies found for the high- and low-energy peaks in each sample. The values used for the Cf<sup>252</sup> standard were  $104.7\pm1$  and 79.8±1 MeV.<sup>2</sup> A Gaussian analysis was used to determine the energies. The energies were normalized to the data taken by the time-of-flight measurements.<sup>2</sup> More accurate values would have been obtained by correcting the measured energies for the kinetic energies carried off by the neutrons; however, this was not done since  $\nu$  is unknown for the isotopes involved.

<sup>\*</sup> Based on work performed under the auspices of the U.S. Atomic Energy Commission.

<sup>&</sup>lt;sup>1</sup> A. B. Smith, P. Fields, A. Friedman, S. Cox, and R. Sjoblom, in Proceedings of the Second International Conference on the Peaceful Uses of Atomic Energy, Geneva, 1958 (United Nations, New York 1958).