

## ACKNOWLEDGMENTS

We would like to express our gratitude to Professor Jack Steinberger for his constant encouragement, advice, and help in this experiment, and Dr. Dan Tycko for the least-squares fitting program and several interesting discussions. The invaluable efforts of the Nevis Scanning and Measuring Staff are well appreciated.

Further thanks go to Dr. U. Nauenberg, Dr. C. Alff-Steinberger, Dr. C. Y. Chang, and S. Marateck for providing some of the events used in their experiments for this one. It is a pleasure to acknowledge the help of Dr. A. Prodell and of the 30-in. hydrogen-bubble-chamber operating crew. The assistance of the A.G.S. staff is also greatly appreciated.


Measurement of the  $K^0$  Mass and the  $K^0$ - $K^-$  Mass Difference\*

J. K. KIM, L. KIRSCH, AND D. MILLER†

*Columbia University, New York, New York*

(Received 21 July 1965)


From the decay of the  $K_1^0$  in a hydrogen bubble chamber, we have measured the mass of the  $K^0$  to be  $497.44 \pm 0.33$  MeV. From the reaction



we have measured the  $K^0$ - $K^-$  mass difference to be  $3.71 \pm 0.35$  MeV.

## I. INTRODUCTION

THE  $K^0$  mass was measured by two independent methods in the Columbia-BNL 30-in. hydrogen bubble chamber exposed to  $\bar{p}$  and  $K^-$  from the low-energy separated beam at the Brookhaven alternating gradient synchrotron (AGS). The first procedure was based on the study of a large sample of two-body decays of  $K^0$  mesons produced by stopping antiprotons. The second procedure was to determine  $\Delta$ , the  $K^0$ - $K^-$  mass difference, from the charge exchange reaction



followed by elastic scattering of the neutron.

A discussion is included of the tests used to determine an absolute scale factor for the conversion of curvature measurements to momenta, and to check the validity of the error assignments.

II.  $K^0$  MASS FROM DECAY

A sample of  $V$ 's was fitted to the three-constraint hypothesis  $K^0 \rightarrow \pi^+\pi^-$  using the measured directions of the neutral  $K^0$  meson and the measured momenta and directions of the pions. The fits were performed using the GRIND kinematics program.<sup>1</sup> Since the program is unable to treat the mass as a variable, we fit each event

using five values of the  $K^0$  mass, ranging from 496.5 to 498.5 MeV in steps of 0.5 MeV. Near the minimum, the curve of  $\chi^2$  versus  $K^0$  mass is a parabola with the minimum located at the best value of the mass. For each event, the lowest three values of  $\chi^2$  are fitted to a parabola to determine the value of  $\chi^2$  at the minimum, and if this value exceeds 5.0, the event is rejected. The accepted sample contains 2223 events. The best value of the  $K^0$  mass from these events is determined by fitting the curve  $\sum_{i=1}^{2223} \chi_i^2$  versus mass to a parabola as shown in Fig. 1. The result is  $m_{K^0} = 497.44 \pm 0.23$  MeV, where the error is obtained from the values of the mass for which  $\sum_i \chi_i^2 = (\sum_i \chi_i^2)_{\min} + 1.0$ . This error reflects the measurement and multiple-scattering errors assigned to each track.

We have two independent checks on the error assignments. The experimental width of the momentum distribution for a set of events produced at a fixed momentum should equal the width of the resolution function for the same sample. The resolution function is obtained by multiplying the assigned momentum error for each event by  $\sqrt{2}$  and then plotting a Gaussian ideogram about the fixed momentum. The pion momenta from the reactions  $K^-p \rightarrow \Sigma\pi$  and  $\bar{p}p \rightarrow \pi^+\pi^-$  are used to perform this test. Good agreement is achieved in both cases, indicating not only a correct magnitude, but also a correct momentum dependence for the errors. The curves for the reaction  $\bar{p}p \rightarrow \pi^+\pi^-$  are plotted in Fig. 2.

The second check is whether or not the errors which are determined by the fitting program based on the assigned errors are correct for the three-constraint fit.

\* Work supported in part by U. S. Atomic Energy Commission.

† Present address: Massachusetts Institute of Technology, Cambridge, Massachusetts.

<sup>1</sup> R. Böck, CERN Report 61-29, 1961 (unpublished).

This is checked by examining the  $K_1^0$  momentum ideograms after fitting the  $K_1^0$  decay from the reaction  $\bar{p}p \rightarrow K_1^0 K_2^0$ . Again the widths of the experimental distribution and the resolution function are comparable.

From a measurement of the momenta of pions produced in the reaction  $\bar{p}p \rightarrow \pi^+\pi^-$  we can experimentally determine the scale factor for the conversion of curvature measurements to momenta. The scale factor takes into account systematic effects such as optical distortions, uncertainties in the magnetic field, and approximations in the geometrical reconstruction program. Since this is a two-body decay occurring from an initial state of fixed energy and zero momentum, the outgoing prongs must be collinear and have unique momentum. The momentum distribution of 1632 pion tracks which dip less than  $60^\circ$ , and which are collinear within  $1.15^\circ$  were used to determine  $p_{\text{exp}}$ , the mean experimental value of the pion momentum from  $\bar{p}p \rightarrow \pi^+\pi^-$ . The expected value,  $p_{\text{theor}}$ , is based only upon the pion and proton mass. The scale factor,  $p_{\text{theor}}/p_{\text{exp}}$ , is applied to the momentum of all tracks used in a determination of the  $K^0$  mass. There is an uncertainty in this ratio due to the uncertainty in  $p_{\text{exp}}$ ;  $\delta p_{\text{exp}}/p_{\text{exp}} = 0.1\%$ . Each event, therefore, contributes an additional error,


$$\delta m_{K^0} = (\delta p_{\text{exp}}/p_{\text{exp}})(m_{K^0}/\sqrt{2}) \times \{1 - (E_K/E_1 E_2)(2m_\pi^2/m_K^4)(m_{K^0}^2 + 2m_\pi^2) + 2(m_\pi/m_K)^4(E_K/E_1 E_2)^2\}^{1/2},$$

where  $E_K$ ,  $E_1$ , and  $E_2$  are the total lab energies of the  $K^0$  and decay pions, respectively. The average contribution to the error in the  $K^0$  mass is 0.24 MeV, so that the result of this measurement is

$$m_{K^0} = 497.44 \pm 0.33 \text{ MeV.}$$

### III. $K^0$ MASS FROM $K^-$ CHARGE EXCHANGE

For a systematic study of low-energy  $K^-p$  cross sections, 60 000 pictures were scanned for all  $K^-$  interactions. Those events which were kinematically compatible with the hypothesis  $K^-p \rightarrow K^0n$  were rescanned. The scanners were given the predicted direction of the neutron as projected into each camera view, and the maximum length of the recoil proton track resulting from an  $n-p$  elastic scattering. They were instructed to accept any event within  $\pm 5^\circ$  of the predicted direction and within 15 cm of the production vertex. Twenty-five events met this criteria. Since the incident beam is inadequately shielded, many of these events are due to a background of unassociated recoil protons. The events were fitted to the hypothesis



where the momentum of the  $K^0$ , the incident neutron, and the outgoing neutron, and the direction of the outgoing neutron, are not observable. The program simul-




FIG. 1. Distribution of  $\chi^2$  versus  $K^0$  mass for 2223  $K^0 \rightarrow \pi^+\pi^-$  events.

taneously fits three vertices so that the problem has seven constraints.

Seven of the events fit with  $\chi^2 < 12$ , whereas among the remainder, none has  $\chi^2 < 55$ . Each of the seven acceptable events is fit to five values of the  $K^0$  mass ranging from 496.5 to 498.5 in steps of 0.5 MeV, using a  $K^-$  mass of 493.8 MeV. The three lowest values of  $\chi^2$  are used to fit a parabola to determine the best value of the  $K^0$  mass. Table I gives the value of the mass difference and its error for each event. The weighted




FIG. 2. The solid curve is a Gaussian ideogram of the momentum of 1632 pion tracks from the reaction  $\bar{p}p \rightarrow \pi^+\pi^-$ . The dashed curve is the resolution function for the same events. The vertical scale is in arbitrary units.

TABLE I. Values of  $K^0$ - $K^-$  mass difference for individual events.

Event number	$K^0$ - $K^-$ mass difference (MeV)
309 941	$5.37 \pm 1.12$
309 691	$3.60 \pm 0.69$
359 964	$2.44 \pm 3.13$
368 562	$3.92 \pm 0.64$
372 533	$4.08 \pm 1.12$
372 823	$3.95 \pm 1.24$
487 277	$2.25 \pm 0.87$
Weighted average	$\Delta = 3.71 \pm 0.35$ (MeV)

average is  $\Delta = 3.71 \pm 0.35$  MeV. Combining this result with the assumed value<sup>2</sup> of the  $K^-$  mass

$$m_{K^-} = 493.8 \pm 0.15 \text{ MeV}$$

gives  $m_{K^0} = 497.51 \pm 0.41$  MeV.

The determination of the mass from these events depends upon the  $K^-$  and pion momenta. Since both the  $\bar{p}$  and  $K^-$  exposures were made in the same chamber, the momenta were corrected in the same manner as discussed in Sec. II. The 0.1% momentum uncertainty results in less than 0.01-MeV change in the  $K^0$  mass.

<sup>2</sup> P. Schmidt, preceding paper, Phys. Rev. **140**, B1328 (1965).

The proton momentum is determined from a range-momentum table which has a scale factor known to 0.5%. The mass change due to this uncertainty is 0.03 MeV.

Both of our results are in agreement with the values of the mass obtained by Burnstein and Rubin,<sup>3</sup>  $m_{K^0} = 497.70 \pm 0.32$  MeV, and the value given in the compilation by Rosenfeld *et al.*,<sup>4</sup>  $m_{K^0} = 498.0 \pm 0.5$  MeV. The weighted average of these four values is  $\bar{m}_{K^0} = 497.62 \pm 0.18$  MeV.

#### ACKNOWLEDGMENTS

We would like to thank Professor J. Steinberger and Professor P. Franzini for many helpful discussions. We would also like to thank R. Goldberg for his assistance in the calibration of measurement errors. It is a pleasure to acknowledge the help of Dr. A. Prodel and of the 30-in. hydrogen-bubble-chamber operating crew. The assistance of the AGS staff is also greatly appreciated.

<sup>3</sup> R. A. Burnstein and H. A. Rubin, Phys. Rev. **138**, B895 (1965).

<sup>4</sup> A. H. Rosenfeld, A. Barbaro-Galtieri, W. H. Barkas, P. L. Bastien, J. Kirz, and M. Ross, Rev. Mod. Phys. **36**, 977 (1964).

## Faddeev Formalism for Four-Particle Systems\*

A. N. MITRA,† J. GILLESPIE,‡ AND R. SUGAR‡

National Science Foundation 1965 Summer Institute for Theoretical Physics,  
University of Wisconsin, Madison, Wisconsin

AND

NARGIS PANCHAPAKESAN

Department of Physics, University of Delhi, Delhi, India

(Received 28 July 1965)

From the four-particle Lippmann-Schwinger equation a set of equations of the Faddeev type with properly connected kernels has been obtained. The kernels contain only the two- and three-particle scattering amplitudes, and do not involve the potentials.

### I. INTRODUCTION

RECENTLY, there has been considerable interest in the three-particle problem<sup>1-5</sup> and in particular the Faddeev equation<sup>1</sup> for the three-body scattering

\* Work supported in part by the National Science Foundation and the U. S. Atomic Energy Commission.

† Permanent address: Department of Physics, University of Delhi, Delhi, India.

‡ Permanent address: Department of Physics, Columbia University, New York, New York.

<sup>1</sup> L. D. Faddeev, Zh. Eksperim. i Teor. Fiz. **39**, 1459 (1960) [English transl.: Soviet Phys.—JETP **12**, 1014 (1961)].

<sup>2</sup> C. Lovelace, Phys. Rev. **135**, B1225 (1964); references to Faddeev's other papers are cited here.

<sup>3</sup> S. Weinberg, Phys. Rev. **133**, B232 (1964).

amplitudes. The main advantages of the Faddeev approach are (1) only the off energy shell two-body scattering amplitudes rather than the potentials occur in the kernel, and (2) the square of the kernel is completely connected, consequently the kernel is likely to be square integrable since the troublesome delta functions arising from disconnected graphs are absent. Of course, the Faddeev method is only one of an infinite number of ways of obtaining  $L^2$  kernels.<sup>4</sup> Weinberg<sup>3</sup> has given a

<sup>4</sup> R. Sugar and R. Blankenbecler, Phys. Rev. **136**, B472 (1964).

<sup>5</sup> For a review of current methods for multiparticle systems see J. Gillespie, *Final State Interactions* (Holden-Day, Inc., San Francisco, California, 1965), Chap. 10.