

## Equivalent-Potential Calculation of $\pi\pi$ and $\pi K$ Scattering\*

JEROME FINKELSTEIN

Lawrence Radiation Laboratory, University of California, Berkeley, California

(Received 10 December 1965)

A method suggested by Balázs is used to calculate the  $\pi\pi$  and  $\pi K$  amplitudes in the resonance region. An input force corresponding to vector-meson exchange reproduces the vector mesons, and also produces  $2^+$  mesons at roughly the masses of the  $f^0(1250)$ ,  $f^{\prime}(1500)$ , and  $K^{**}(1405)$ . Inclusion of an estimate of Pomeron-chuk exchange in the input improves the output slightly. Regge trajectories and residues are computed, and their intercepts agree approximately with estimates based on high-energy experiments.

### I. INTRODUCTION

RECENTLY Balázs has proposed a method for constructing, from a given absorptive part in the crossed channel, a Schrödinger-equation potential that will reproduce the relativistic amplitude for scattering of spinless particles.<sup>1</sup> In this paper I present the results of some calculations of ( $\pi\pi$ ) and ( $\pi K$ ) scattering, using an approximation to this Balázs method; these calculations reproduce many generally accepted features of resonance-energy ( $\pi\pi$ ) and ( $\pi K$ ) scattering.

The Schrödinger-equation potential in this method is local but energy-dependent. Writing it as

$$V(r, q^2) = \frac{-1}{2\pi\mu} \int_{t_0}^{\infty} dt v(t, q^2) e^{-r\sqrt{t}/r},$$

where  $q$  is the magnitude of the momentum in the center-of-mass system of the  $s$  channel and  $\mu$  is the reduced mass, Balázs proposed an iterative method for constructing  $v(t, q^2)$  from the supposedly known absorptive part  $A_t$ ; the details of this construction are in Ref. 1. The scattering amplitude calculated from this Schrödinger equation would then be identical with the relativistic amplitude of which  $A_t$  was the absorptive part. In a problem with exchange potentials, this procedure is to be used for amplitudes of definite  $J$  parity. For the Balázs method to be exact would require that (a) the amplitude obey elastic unitarity at the energy in question, (b)  $v(t, q^2) \rightarrow 0$  as  $t \rightarrow \infty$ , and (c)  $A_t$  be known exactly.

The practical application of this derivation is to justify an approximation scheme, in which the lowest approximation consists of setting

$$v(t, q^2) = 2s^{-1/2} A_t^B(t, s),$$

where  $A_t^B$  is the Born approximation<sup>2</sup> to  $A_t$ , and  $s$  is the square of the energy in the center-of-mass system. The scattering amplitude calculated from the Schrödinger equation will be unitary, and in this sense this approximation is similar to the  $N/D$  approach. It differs from

the  $N/D$  approach in several interesting features: First, if  $A_t^B$  is constructed from elementary-particle exchange, there is no high-energy cutoff, even if the exchanged particles have spin.<sup>3</sup> And second, although the usual  $N/D$  calculation takes the Born approximation to the left-hand cut, solving the Schrödinger equation means in effect doing the entire Mandelstam iteration.<sup>4</sup> Thus in the Balázs approximation, contributions from all orders of this iteration are considered, even though none but the first has relativistic  $s$  dependence.

In Sec. II, I describe the potentials that I have used, and in Sec. III present the results obtained.

### II. CONSTRUCTION OF POTENTIALS

#### A. First ( $\pi\pi$ ) Potential

My first calculation of ( $\pi\pi$ ) scattering is an extension of an example given by Balázs.<sup>1</sup> He considered the force due to exchange of an elementary  $\rho$ , and obtained, in the small-width approximation,

$$A_t^B(t, s) = 4\pi\beta_{I1}(2l+1)P_l(z_t)\Gamma_{\text{in}}qR^2\delta(t-m^2), \quad (1)$$

where  $\beta$  is the isotopic spin-crossing matrix,  $l=1$  the spin and  $m$  the mass of the exchanged  $\rho$ ,  $qR^2 = \frac{1}{4}m^2 - m_\pi^2$ , and  $\Gamma_{\text{in}}$  is the input reduced width. The potential corresponding to (1) is

$$V(r, q^2) = -24\beta_{I1}\Gamma_{\text{in}}s^{-1/2}(s+2qR^2)r^{-1}e^{-mr}. \quad (2)$$

Balázs looked for the  $\rho$  resonance in the  $l=1$  amplitude obtained from (2), and found an approximately self-consistent solution with  $m=4.2 m_\pi$  and  $\Gamma=0.47$  (a  $\rho$  of mass 750 MeV and width 100 MeV has  $m=5.3 m_\pi$  and  $\Gamma=0.17$ ).

However, it turns out that approximate self-consistency is not sufficient to determine the value of these parameters, as within 10% there are many self-consistent solutions. In what is referred to below as calculation A, I use the potential given by (2), with the input  $\rho$  mass fixed at 750 MeV. I then adjust the input width to obtain the output  $\rho$  mass at the same energy; this solution is also self-consistent, and has  $\Gamma=0.46$  (cor-

\* This work was performed under the auspices of the U. S. Atomic Energy Commission.

<sup>1</sup> L. A. P. Balázs, Phys. Rev. **137**, B1510 (1965).

<sup>2</sup> That is,  $A_t^B$  represents the contribution of a few elastic partial waves in the  $t$  channel, which in the strip approximation (see Ref. 5) is the Born approximation to  $A_t$ .

<sup>3</sup> If the external particles have spin, a cutoff may be necessary. See L. A. P. Balázs, Phys. Rev. **139**, B1646 (1965), where he discusses the possible application of this method to ( $\pi N$ ) scattering.

<sup>4</sup> R. Blankenbecler, M. L. Goldberger, N. N. Khuri, and S. B. Trieman, Ann. Phys. (N. Y.) **10**, 62 (1960).

responding to a width of 270 MeV). This calculation has no cutoff and no strip width; with  $\Gamma_{\text{in}}$  determined, there are no other free parameters, and the entire (low-energy) ( $\pi\pi$ ) amplitude in all three isotopic-spin states can be calculated.

### B. Second ( $\pi\pi$ ) Potential

The results of calculation A, which are described in Sec. III, resembled experimental results sufficiently to encourage me to try to improve the potential by including some estimate of the contribution due to the exchange of the Pomeranchuk trajectory.

Chew<sup>5</sup> has discussed the generalized potential arising from exchange of a Regge trajectory in the  $t$  channel, where the generalized potential  $V^s(t, s)$  is

$$V^s(t, s) = \frac{1}{\pi} \int_{t_0}^{\infty} \frac{dt' A_t^B(t', s)}{t' - t}. \quad (3)$$

He writes  $V^s$  as a sum of partial waves in the  $t$  channel (even partial waves for trajectories of positive signature) and argues that only the lowest need be kept. He gives approximate expressions for the two lowest terms of the Pomeranchuk potential: The  $J=2$  part looks like the exchange of an elementary  $f^0$  multiplied by a "form factor" in  $t$ ; the  $J=0$  part, even though it has no associated particle, is more important, and is repulsive. The exact value of the  $J=0$  part depends on a strip parameter, which I have set equal to  $200 m_\pi^2$ .

I cannot make direct use of Chew's estimates, since they are intended for only small values of  $t$ , and hence do not accurately define  $A_t^B$  where I need it; instead, for each of these two partial waves I have replaced  $A_t^B$  by a delta function, chosen so that the  $V^s$  calculated from Eq. (3) has the same value and derivative at  $t=0$  as do Chew's estimates. It then turns out that the  $J=2$  term is almost completely negligible; the  $J=0$  term is substantially smaller, but has a slightly larger range, than the  $\rho$  potential used in calculation A. The complete potential is the sum of the  $\rho$  and the Pomeranchuk potentials<sup>6</sup>:

$$V(r, q^2) = V^\rho(r, q^2) + V_{J=0}^P(r, q^2) + V_{J=2}^P(r, q^2),$$

where, in units in which  $m_\pi^2 = 1$ ,

$$V^\rho(r, q^2) = -24\beta_{I1}\Gamma_{\text{in}}s^{-1/2}(s+12)e^{-5.3r}r^{-1},$$

$$V_{J=0}^P(r, q^2) = +180\beta_{I0}s^{-1/2}e^{-5.0r}r^{-1},$$

$$V_{J=2}^P(r, q^2) = -\beta_{I2}s^{-1/2}(12+0.94s+0.013s^2)e^{-5.1r}r^{-1}. \quad (4)$$

I again adjust  $\Gamma_{\text{in}}$  to get the output  $\rho$  mass at 750 MeV, and find  $\Gamma_{\text{in}} = 0.56$ . The results obtained from the potential given by (4) I call calculation B.

<sup>5</sup> G. F. Chew, Progr. Theoret. Phys. (Kyoto) Suppl., extra number, 118 (1965); Phys. Rev. 140, B1427 (1965).

<sup>6</sup> P. D. B. Collins (private communication) assures me that the effect of a Reggeized  $\rho$  potential is about the same as that of the elementary  $\rho$  potential which is used here.

Surely the expressions given in (4) have at best a tenuous connection with the correct Pomeranchuk potential, whatever that may be. However, if we accept Chew's arguments, these expressions should be reasonable qualitative estimates, and since they appear as fairly small corrections to the much better established  $\rho$  potential, might be expected to improve the results of calculation A. As shown in Sec. III, this apparently is the case.

### C. ( $\pi K$ ) Potential

The last calculation, which I call calculation C, is of the ( $\pi K$ ) scattering amplitude. I have not attempted here to put in the Pomeranchuk potential, but consider only vector-meson exchange, in analogy with calculation A. There are now two exchange forces,  $\rho$  exchange in the  $t$  channel and  $K^*(891)$  exchange in the  $u$  channel. With the assumption that the couplings  $\gamma_{\rho\pi\pi}$  and  $\gamma_{\rho K\bar{K}}$  are related to  $\gamma_{K^*K\pi}$  as predicted by  $SU_3$ , and with the physical values for the input masses used, the potential appropriate for  $I = \frac{1}{2}$  and angular momentum  $l$  is given by

$$\begin{aligned} V(r, q^2) &= V^\rho(r, q^2) + (-1)^l V^{K^*}(r, q^2), \\ V^\rho(r, q^2) &= -32\Gamma_{\text{in}}'s^{-1/2}(s+0.5)e^{-5.3r}r^{-1}, \\ V^{K^*}(r, q^2) &= -8\Gamma_{\text{in}}'s^{-1/2}(s+12)e^{-6.3r}r^{-1}. \end{aligned} \quad (5)$$

Here  $\Gamma_{\text{in}}'$ , the reduced  $K^*$  width, is determined by requiring the output mass of the  $K^*$  to be 891 MeV; this requirement gives  $\Gamma_{\text{in}}' = 0.57$ , corresponding to a width of 133 MeV.

Having constructed the potentials for the three calculations, I solved the Schrödinger equation numerically, to find the scattering amplitude and the Regge parameters at physical values of  $q^2$ . The computations were done on an IBM 7094 computer, in part with a program written by Burke and Tate.<sup>7</sup>

## III. RESULTS

In both calculations A and B, for  $I=0$  there are two Regge trajectories above  $l=0$  at threshold; these I identify with the  $P$  and  $P'$  trajectories, and the associated particles at  $l=2$  with the  $f^0(1250)$  and the  $f'^0(1500)$ .<sup>8</sup> For  $I=1$  there is one such trajectory, which I identify with the  $\rho$ ; for  $I=2$  the force is repulsive. The real parts of the trajectories continue to rise as the energy is increased, at least until  $s=400 m_\pi^2$ , and in fact do eventually go through higher physical values of  $l$ . However, one of the assumptions of the Balázs method is that the amplitude is elastic at the energy in question, and so it must lose its validity as the energy approaches the strip width. As none of these "recurrences" occurs below  $s=300 m_\pi^2$ , they could be considered spurious.

<sup>7</sup> P. G. Burke and C. Tate, Fortran Program TREGGE, Lawrence Radiation Laboratory Report No. UCRL-10384, 1962 (unpublished).

<sup>8</sup> V. E. Barnes *et al.*, Phys. Rev. Letters 15, 322 (1965).

TABLE I. Results for the  $(\pi\pi)$  amplitude. The force input to calculation A is  $\rho$  exchange, and to calculation B is  $\rho$  exchange and Pomeranchuk exchange.

	$\rho$ mass (MeV)	$\Gamma_\rho$	$f^0_{\text{mass}}$ (MeV)	$\Gamma_{f^0}$	$f^{0'}_{\text{mass}}$ (MeV)	$\Gamma_{f^{0'}}$
Calculation A	750	0.50	1070	0.50	1900	0.55
Calculation B	750	0.29	920	0.35	1435	0.95
Experiment	750 <sup>a</sup>	0.17 (100 MeV) <sup>a</sup>	1250 <sup>b</sup>	0.25 (100 MeV) <sup>c</sup>	1500 <sup>c</sup>	

<sup>a</sup> Ambiguities in the experimental values are not important to this paper; these numbers for the  $\rho$  come from C. Alif *et al.*, Phys. Rev. Letters 9, 322 (1962).

<sup>b</sup> W. Selove *et al.*, Phys. Rev. Letters 9, 272 (1962).

<sup>c</sup> See Ref. 7.

The masses and the reduced widths of the three output resonances are listed in Table I; the output reduced widths are computed by

$$\Gamma = \left[ \frac{\sqrt{s} \left( \frac{d\delta}{ds} \right)^{-1}}{8q^{2l+1}} \right]_{s=m_{\text{resonance}}^2},$$

and have units of  $(\text{GeV})^{2-2l}$ .

A calculation of  $s$ -wave scattering by this method is less reliable than a calculation of higher partial waves, for the usual reason that the  $s$  wave depends more strongly on the shorter-range parts of the potential; but it might be interesting anyway. The results of this calculation are the strange-looking phase shifts shown in Fig. 1; the fact that these phase shifts wander through  $90^\circ$  has no significance. A large but not necessarily resonant  $s$ -wave amplitude near the  $\rho$  mass has been previously suspected from the asymmetry in  $\rho^0$  decay.<sup>9</sup> The  $s$ -wave scattering length is  $-0.08 m_\pi^{-1}$  in calculation A and  $-0.18 m_\pi^{-1}$  in calculation B; the negative sign would be expected from the existence of trajectories above  $l=0$  at threshold, even though the forces are attractive.<sup>10</sup>

Because of the factor  $s^{-1/2}$  in the definition of the potential, the Balázs approximation must break down near  $s=0$ . Nevertheless it would be interesting to be able to compute the values of the Regge parameters at




FIG. 1.  $I=0$   $s$ -wave  $(\pi\pi)$  phase shift.

<sup>9</sup> M. M. Islam and R. Piñon, Phys. Rev. Letters 12, 310 (1964).

<sup>10</sup> See G. F. Chew, Phys. Rev. Letters 16, 60 (1966).

this point, for then they could be compared with the values obtained from experiments at high energy in the crossed channel. The procedure I have adopted is to calculate the Regge parameters  $\alpha$  and  $\ln\gamma = \ln(\beta/q^{2\alpha})$  above threshold, and attempt a straight-line extrapolation to  $s=0$ . In most cases this extrapolation seemed possible, and the results for the  $(\pi\pi)$  amplitude are given in Table II.

TABLE II. Values of the Regge parameters of the  $(\pi\pi)$  amplitude at  $s=0$ .

	$\alpha_\rho$	$\gamma_\rho^a$	$\alpha_P$	$\gamma_P$	$\alpha_{P'}$	$\gamma_{P'}$
Calculation A	0.7	b	1.3	b	0.7	b
Calculation B	0.45	0.05	1.3	0.0036	0.65	0.09
Other estimates	0.54 <sup>c</sup>	0.026 <sup>d</sup>	1.0 <sup>c</sup>	0.006 <sup>d</sup>	0.50 <sup>e</sup>	0.065 <sup>d</sup>

<sup>a</sup>  $\gamma$  is defined by  $\gamma = \beta/q^{2\alpha}$ , where  $q$  is in units of  $m_\pi$ .

<sup>b</sup> I was unable to make a reliable extrapolation from the physical region.

<sup>c</sup> R. J. N. Phillips and W. Rarita, Phys. Rev. 139, B1336 (1965).

<sup>d</sup> These values come from an analysis of high-energy  $\pi N$ ,  $NN$ , and  $\bar{N}N$  scattering, and use of the factorization theorem; see R. J. N. Phillips and W. Rarita, Phys. Rev. Letters 14, 502 (1965), and the Appendix to Heinz J. Rothe, Phys. Rev. 140, B1421 (1965).

The width of the  $K^*$  in calculation C is very nearly equal to the input width, which is about 2.5 times the experimental value. There is one other  $(\pi K)$  trajectory, with the quantum numbers of the  $K^{**}(1405)$ . The results for these two trajectories are presented in Table III. For the  $I=\frac{3}{2}(\pi K)$  amplitude the force is repulsive.

As can be seen from the tables, the general features of the experimental situation are reproduced quite well. Some of the closer agreements with experiment may well be fortuitous, but the over-all pattern, especially the appearance of the  $2^+$  mesons, could not be. The most glaring discrepancy of my results with experimental results is the fact that the input and output particle widths are too large, but this seems to be a common feature of most dynamical calculations.<sup>11</sup> The results of calculation B seem somewhat better than those of calculation A, although most of the differences are small. The output  $\rho$  width is improved substantially, destroying the consistency between input and output.

It is interesting to compare the above calculations, especially calculation A, with the  $N/D$  calculation of

<sup>11</sup> For example, F. Zachariasen, Phys. Rev. Letters 7, 112 (1961); J. R. Fulco, G. L. Shaw, and D. Y. Wong, Phys. Rev. 137, B1242 (1965).

TABLE III. Results for the ( $\pi K$ ) amplitude.

	$K^{*}_{\text{mass}}$ (MeV)	$\Gamma_{K^*}$	$\alpha_{K^*}$	$\gamma_{K^*}$ <sup>a</sup>	$K^{**}_{\text{mass}}$ (MeV)	$\Gamma_{K^{**}}$	$\alpha_{K^{**}}$	$\gamma_{K^{**}}$
Calculation C	891	0.55	0.4	0.14	1265	0.16	0.75	0.02
Experiment	891 <sup>b</sup>	0.22 <sup>b</sup> (50 MeV)			1405 <sup>b</sup>	0.12 <sup>b</sup> (95 MeV)		

<sup>a</sup> See footnote a, Table II.

<sup>b</sup> A. H. Rosenfeld *et al.*, Rev. Mod. Phys. **37**, 633 (1965).

Collins and Teplitz.<sup>12</sup> They used an input  $\rho$  with a width of 0.43, which is about the same as was used in calculation A, but their output  $\rho$  trajectory did not quite make it to  $l=1$ , and no trajectory rose above  $l=1.5$ . Thus the effective force in the present calculation is much stronger than in the  $N/D$  calculation, even though the input forces are similar. It seems reasonable that

<sup>12</sup> P. D. B. Collins and V. L. Teplitz, Phys. Rev. **140**, B663 (1965).

this is because the method used here does include contributions to the force from higher terms in the Mandelstam iteration. If this conjecture be correct, then the results presented here indicate that a calculation that actually performs the iteration might be expected to be very successful.

#### ACKNOWLEDGMENTS

I am grateful to Professor Geoffrey F. Chew for many helpful suggestions and discussions.

## Group Embedding for the Harmonic Oscillator

R. C. HWA\* AND J. NUYTS†

*The Institute for Advanced Study, Princeton, New Jersey*

(Received 23 December 1965)

The embedding of the algebra of the invariance group  $SU(n)$  for the  $n$ -dimensional harmonic oscillator in larger algebras is considered. Among the four classical algebras of rank  $n$ , only  $\mathfrak{su}(n+1)$  and  $\mathfrak{sp}(2n)$  are found possible for this purpose. Specific generators and their commutation relations are examined, and the general Casimir operators are constructed. It is found that the whole spectrum of the harmonic oscillator can be embedded in one representation of  $\mathfrak{su}(n+1)$ . Depending upon the value of a partition constant  $c$ , a finite portion of the spectrum can be embedded in the compact algebra  $su(n+1)$ ; the remainder is in the noncompact  $su(n,1)$ . In the case of  $\mathfrak{sp}(2n)$ , only the noncompact  $sp(n, R)$  can include the  $su(n)$  of the harmonic oscillator; moreover, the two useful representations of  $sp(n, R)$ , which are true representations of the universal covering group of  $Sp(n, R)$ , contain either all the even or all the odd levels of the spectrum.

### I. INTRODUCTION

IN the attempt to combine internal symmetries with the Poincaré group, it has been repeatedly proposed<sup>1-3</sup> that noncompact groups (or algebras) be used to describe not only the dynamical symmetries, but also the internal symmetries of elementary particles. The study of the problem has led to very beautiful and mysterious results on the one hand, but, despite a large freedom, to many difficulties on the other.

In order to learn about the problem of embedding groups (or algebras) into larger ones, it is useful to examine completely soluble models provided by non-relativistic classical and quantum-mechanical systems. The Kepler problem (or quantum-mechanically the hydrogen atom) and the harmonic oscillator are two standard examples.

It was shown<sup>4</sup> long ago that the Hamiltonian of the Kepler problem is invariant under the  $SO(4)$  group, which explains the "accidental" degeneracy of the hydrogen atom. More recently, it has been conjectured that this  $SO(4)$  group can be embedded in an  $SO(4,1)$  group, whose generators have been written down for the classical case.<sup>5</sup> This  $SO(4,1)$  group does not com-

\* Work supported by the National Science Foundation.

† Research sponsored by the Air Force Office of Scientific Research, Office of Aerospace Research, U. S. Air Force, under AFOSR 42-65.

<sup>1</sup> B. Kurşunoğlu, Phys. Rev. **135**, B761 (1964); A. Barut, Nuovo Cimento **32**, 234 (1964).

<sup>2</sup> R. Delbourgo, A. Salam, and J. Strathdee, Proc. Roy. Soc. (London) **A284**, 146 (1965); H. Bacry and J. Nuyts, Nuovo Cimento **37**, 1702 (1965); W. Rühl, Phys. Letters **14**, 346 (1965).

<sup>3</sup> Y. Dothan, M. Gell-Mann, and Y. Ne'eman, Phys. Letters **17**, 148 (1965).

<sup>4</sup> W. Pauli, Z. Physik **36**, 336 (1926); V. Fock, *ibid.* **98**, 145 (1935); V. Bargmann, *ibid.* **99**, 576 (1936).

<sup>5</sup> H. Bacry, Nuovo Cimento **41A**, 222 (1966); E. C. G. Sudarshan, N. Mukunda, and L. O'Raifeartaigh, Phys. Letters **19**, 322 (1965).