

THE PHYSICAL REVIEW

A journal of experimental and theoretical physics established by E. L. Nichols in 1893

SECOND SERIES, VOL. 149, No. 4

30 SEPTEMBER 1966

Observation on the Elemental Abundances of Low-Energy Cosmic Rays in July 1964

D. V. REAMES AND C. E. FICHEL

Goddard Space Flight Center, Greenbelt, Maryland

(Received 20 April 1966)

In a sounding rocket experiment flown at Fort Churchill, Manitoba in July 1964, nuclear emulsion detectors were used to study the composition and fluxes of primary cosmic-ray nuclei with charge $Z \geq 3$ above the atmosphere in the region of kinetic energy from 40 to 200 MeV/nucleon. The results show a flux ratio for light ($3 \leq Z \leq 5$) to medium ($6 \leq Z \leq 9$) nuclei of 0.16 ± 0.07 in the 40- to 100-MeV/nucleon region and a ratio for heavy ($Z \geq 10$) to medium nuclei of 0.42 ± 0.11 in the 70- to 160-MeV/nucleon interval. The elemental structure of the medium nuclei is found to contain $(40 \pm 7)\%$ carbon, $(10_{-8}^{+4})\%$ nitrogen, and $(48 \pm 7)\%$ oxygen. Extrapolating the fluxes back through approximately 2.5 g/cm^3 of interstellar hydrogen to the source, one finds a flux of nitrogen consistent with zero.

I. INTRODUCTION

MUCH of our present understanding of the origin of cosmic rays and their propagation through interstellar space results from the study of the abundances of multiply charged nuclei in the cosmic-ray flux at kinetic energies above a few hundred MeV/nucleon.¹⁻³ It is of particular interest to examine ratios of the fluxes of the different heavy nuclei (having approximately equal charge to mass ratio) since such flux ratios are insensitive at all energies to the somewhat poorly understood modulation mechanisms which affect the fluxes themselves. The flux ratios in the energy region below 200 MeV/nucleon should be sensitive to the amount of interstellar material traversed, owing to the differing energy-loss rates of the particles of different charge, in a way which depends upon the energy spectra of the particles at their source.

The existence of a measurable flux of heavy nuclei in the 20-200-MeV/nucleon range was demonstrated in a September 1963 sounding rocket experiment flown at Ft. Churchill, Manitoba using large sheets of nuclear emulsion as a detector.⁴ The experiment reported herein

was flown in June 1964 using an improved version of the original apparatus to extend the range of measurements to *all* charges $Z \geq 3$ and to unambiguously isolate only those particles of primary origin.

An additional advantage of a sounding rocket observation of cosmic radiation not shared by balloonborne experiments is that the observed composition is entirely free from secondary particles produced in interactions in the atmosphere. Though corrections for such interactions may be applied to the gross charge groups commonly investigated, the partial cross sections for interaction in air are not known on an element by element basis with sufficient precision to determine the detailed elemental composition of the cosmic radiation at the top of the atmosphere. Knowledge of the detailed charge structure of cosmic rays may provide additional information about the source and interstellar propagation of the radiation.

II. EXPERIMENTAL PROCEDURE

The nuclear emulsion detectors used in this experiment were stacks of $600 \mu \times 6.3 \text{ cm} \times 29 \text{ cm}$ Ilford G.5 pellicles with 18 pellicles per stack. The stacks were arranged in 3 trays which could be extended at a 17.5° angle with respect to the skin of the Aerobee 150 rocket with 2 stacks per tray. The trays were kept inside the payload until the sounding rocket had left the atmosphere, at which time they were extended. As the trays were extended, the top pellicle of each stack was displaced by 1 cm so that subsequent matching of

¹ W. R. Webber, *Handbuch der Physik*, edited by S. Flügge (Springer-Verlag, Berlin, 1966), Vol. 46.

² *Proceedings of the International Conference on Cosmic Rays, Jaipur, India* (Commercial Printing Press, Ltd., Bombay, India, 1963).

³ *Proceedings of the Ninth International Conference on Cosmic Rays* (The Institute of Physics and the Physical Society, London, 1966).

⁴ C. E. Fichtel, D. E. Guss, and K. A. Neelakantan, *Phys. Rev.* **138**, B732 (1965).




FIG. 1. Scatter plot of particle charges determined by the method described in the text using the measurements of two different observers. Note that all tracks accepted are represented here so that the energy or rigidity interval is a function of charge. Particles with a measured charge less than 2.5 have been suppressed.

the tracks would allow the isolation of those which entered during the exposure. The exposure lasted about 350 seconds, after which the trays were retracted and the top plate of each stack was restored to its initial position in preparation for re-entry into the atmosphere. Extension and retraction each took about 7 secs. The total amount of material above the top emulsion was 0.091 g/cm^2 (emulsion equivalent).

After processing, the top plate of each stack was area scanned for the tracks of all particles entering its surface within the dip angle ($< 65^\circ$) and ionization criteria. The minimum ionization accepted was set sufficiently low to insure that all $Z=3$ tracks would be selected and in practice this meant over half of the alpha particles were also included in the preliminary selection. These tracks were then followed into the stack and all tracks which failed to follow (i.e., were displaced by 1 cm with respect to the top pellicle) or failed to stop in the stack were rejected.

Charges of the particles were determined from delta-ray counting. If $n_\delta(R)$ is the number of delta rays per unit length a distance R from the track ending, then a plot of $\log N_\delta = \log \int_0^R n_\delta(R) dR$ versus $\log R$ for a given track produces a curve whose shape does not change with the charge (or mass) of the particle if one neglects the small effect of range extension caused by charge

pickup. For particles of the same charge to mass ratio one may write

$$\log N_\delta - \log Z = F(\log R + \log Z) \quad (1)$$

so that curves for particles of different charge are displaced diagonally from each other by the logarithm of their charge ratio. All charge identifications in this experiment were made using this technique. Delta rays were counted using a parallel-line reticle in the microscope and counting the rays which appeared to extend beyond the lines on either side of a track that was centered on a central line. A count using a narrow line spacing was first made to separate the alpha particles from the higher charges of interest. The narrow line spacing required that the delta rays extend at least 1.8μ from the center of the track. At least two independent counts were then made for each heavy track using a wider line spacing which selected delta rays extending at least 4.3μ . This spacing was more convenient for counting the heavier tracks. Owing partially to difficulties in counting the high delta-ray densities of the heavier tracks and partially to the effects of range extension mentioned previously, charges of the highly charged particles are underestimated by our method of analysis. To correct for these effects a "saturation" correction was applied which increased the

measured charges by an amount which varied from a few percent at charge 10 to about 20% at charge 26. The results and conclusion of this paper are relatively insensitive to the detailed nature of this kind of correction.

Figure 1 is a comparison of the charges of the particles accepted as determined by two different observers and indicates the charge resolution obtained.

The solid-angle considerations of this experiment are similar to those previously reported⁴ except that all particles with arrival directions making an angle of greater than 90° with respect to the magnetic field direction were excluded. These particles would be coming from below the rocket and might have resulted from interactions in the atmosphere. The probability of stopping a given particle is a function of its range in the detector, and an efficiency factor, as a function of range, can be compiled from geometrical considerations. This function is essentially flat for ranges from 2 to 12 mm, a lower limit of slightly over 1 mm being set by the requirement that particles enter the second plate in order to insure their arrival when the rocket was outside the atmosphere. The efficiency function used was found by numerically integrating $\mathbf{n} \cdot d\mathbf{A}d\Omega$ over the allowed area and solid angle for a given particle range where \mathbf{n} is the particle arrival direction. For particles of intermediate ranges the total allowed area-solid-angle factor was 1680 cm² sr. The upper limit on the energy was determined by the condition that the collection efficiency be greater than 20% of the maximum value possible i.e., that the range be less than about 2.5 cm.

Scanning efficiency was determined by a rescan of approximately 25% of the total area and was found to be between 96 and 99% for the different observers. Considerable effort was made in the original scan to insure that there be no bias against the light nuclei as evidenced by the large number of $Z=2$ nuclei included in the original sample. No additional light nuclei were located in the rescan and there is no evidence of a charge dependent scanning efficiency. Angular distributions were also searched unsuccessfully for evidence of scanning bias.

III. RESULTS AND DISCUSSION


A. General Features

Differential fluxes of the light L , medium M , and heavy H , nuclei collected into 30 MeV/nucleon energy bins are shown in Fig. 2. For comparison with the differential flux values of the M and H nuclei, we also show the smoothed α -particle spectra⁵⁻⁸ multiplied by

⁵ C. E. Fichtel, N. Durgaprasad, and D. E. Guss, in *Proceedings of the Ninth International Conference on Cosmic Rays* (The Institute of Physics and the Physical Society, London, 1966), Vol. 1, p. 371.

⁶ C. J. Waddington and P. S. Freier, in *Proceedings of the Ninth International Conference on Cosmic Rays* (The Institute

Fig. 2. Differential fluxes of light ($3 \leq Z \leq 5$), medium ($6 \leq Z \leq 9$) and heavy ($Z \geq 10$) nuclei in July 1964. Solid curves are the α -particle fluxes for the same period (see text) multiplied by 0.070 and 0.027 for comparison with the M and H nuclei, respectively.



the high-energy flux ratios, 0.070 and 0.027, respectively. No α -particle data taken in July 1964 have been reported for energies below 100 MeV/nucleon, and the curves in this energy region have been obtained by an appropriate averages of data taken before and after this period.^{7,8}

Ratios of fluxes of various constituent charge groups are shown in Table I together with the energy interval from which they are taken.

The theoretical implications of these ratios when compared with their higher energy values are of considerable interest and will be discussed in a subsequent paper.

B. Elemental Structure

Even a casual observation of Fig. 1 reveals considerably finer structure in the charge spectrum than that described by the charge groups discussed above. The elemental structure of the flux of medium nuclei be-

TABLE I. Observed particle ratios.

Charge group	Ratio	Energy interval (MeV/nucleon)
L/M	0.16 ± 0.07	40-100
$(10 \leq Z \leq 15)/M$	0.29 ± 0.09	70-160
$(16 \leq Z \leq 19)/M$	0.02 ± 0.02	70-160
$(Z \leq 20)/M$	0.11 ± 0.05	70-160
H/M	0.42 ± 0.11	70-160

of Physics and the Physical Society, London, 1966), Vol. 1, p. 339.

⁷ C. Y. Fan, G. Gloeckler, and J. A. Simpson, *J. Geophys. Res.* **70**, 3515 (1965).

⁸ C. Y. Fan, G. Gloeckler, and J. A. Simpson, *Proceedings of the Ninth International Conference on Cosmic Rays* (The Institute of Physics and the Physical Society, London, 1966), Vol. 1, p. 380.

TABLE II. Composition of the flux of medium nuclei.

Element	Cosmic-ray abundance (this work)		"Universal abundance"		Solar cosmic-ray abundances ^c
	At earth	At source (2.5 g/cm ²)	Cameron ^a	Suess and Urey ^b	
C	40±7%	~38	25.3%	11.2%	33±4%
N	10 ₋₆ ⁺⁴	~0	6.5	20.9	11 ₋₆ ⁺²
O	48±7	~62	68.1	67.9	56±7
F	2±2	~0	0.004	0.005	<2

^a Reference 9.^b Reference 10.^c Reference 11.

tween 40 and 160 MeV/nucleon is shown in Table II along with the "universal"^{9,10} and solar¹¹ abundances.

It is of interest that data reported from balloon-borne observations of higher energy cosmic rays indicate a richer nitrogen content than that reported here, while the OGO-I satellite experiments^{12,13} flown more recently confirm our result of $N/(C+N+O) \approx 10\%$. Even though there are many experiments at high energies which give higher values, some care must be exercised in drawing conclusions about the energy dependence of this ratio since there are several effects that might enhance the observed nitrogen abundance. Not the least consideration is the charge resolution obtained. Many of the balloon experiments were designed only to resolve charge groups and any broadening of the neighboring carbon and oxygen charge distributions could enhance the apparent number of nitrogen nuclei if not carefully eliminated. Further, the interaction of all nuclei with $Z \geq 8$ in the residual atmosphere above a balloon would enhance the abundance of N and C in a poorly known way, this effect being aggravated by particles collected during ascent of the balloon in many experiments. Experiments with high charge resolution such as those of Balasubrahmanyam and McDonald¹⁴ and of O'Dell *et al.*¹⁵ each find $N/(C+H+N) \approx 16\%$, though these experiments may still suffer from effects of the overlaying atmosphere.

Turning attention to the nuclei of even charge one can see from Table II that the O/C ratio of 1.2 ± 0.3 is still considerably smaller than the ratio from universal abundances but is consistent with solar abundances within statistics. Again our data are consistent with the OGO-I results, but apparently inconsistent with the ratios obtained in balloon experiments which typically give 0.64.¹⁵ As in the case of the nitrogen abundance it

remains to be proven that the discrepancy cannot result from fragmentation in the overlying atmosphere. One could, in principle, observe the effect of fragmentation in the atmosphere by comparing balloon results from different altitudes, but attempts to do so are frustrated by varying degrees of charge resolution and often by the presence of particles collected during ascent.

Above charge 9 we find the composition is characterized by the absence of nuclei of odd charge, with evidence of a higher proportion of Mg relative to Ne and Si when represented in terms of energy per nucleon though both statistics and charge resolution are poorer here than for the medium charge group. In addition to the well-known lack of particles of charges 15 to 19, we also see a gap between the two abundance peaks around charges 20 and 26. This gap is entirely consistent with universal abundances as are the low fluxes of odd-charged nuclei.

The principal motivation for studying the composition of the cosmic radiation is to learn as much as possible of the elemental composition of the source, and thereby to identify it and to study the processes of nucleogenesis in it. There is now considerable evidence¹ that the cosmic rays pass through 2 to 3 g/cm² of interstellar material before arriving at earth, and one would like to account for the effects of this material on the composition. Data on the reaction cross sections for protons (interstellar material) incident on various nuclei have been summarized by Badhwar *et al.*,^{16,17} and we have used these data to estimate the abundances of elements, particularly C, N, and O, at the source. The results of these calculations are shown in Table II, the most significant feature being the disappearance of the nitrogen on extrapolating backward through about 2.5 g/cm² of hydrogen. This effect results primarily from the fact that about half of the inelastic cross section for oxygen-proton reactions leads to nitrogen as a stable product. The relative abundances of the different species of medium nuclei are affected only very slightly by differing energy loss rates and given similar source spectra an energy-dependent composition can only be generated by assuming energy-dependent cross sections

⁹ A. G. W. Cameron, *Astrophys. J.* **129**, 676 (1959).¹⁰ H. E. Suess and H. C. Urey, *Rev. Mod. Phys.* **28**, 53 (1956).¹¹ S. Biswas and C. E. Fichtel, *Astrophys. J.* **139**, 941 (1964).¹² V. K. Balasubrahmanyam, D. E. Hage, G. H. Ludwig, and F. B. McDonald, in *Proceedings of the Ninth International Conference on Cosmic Rays* (The Institute of Physics and the Physical Society, London, 1966), Vol. 1, p. 427.¹³ G. M. Comstock, C. Y. Fan, and J. A. Simpson, in *Proceedings of the Ninth International Conference on Cosmic Rays* (The Institute of Physics and the Physical Society, London, 1966), Vol. 1, p. 383.¹⁴ V. K. Balasubrahmanyam and F. B. McDonald, *J. Geophys. Res.* **69**, 3289 (1964).¹⁵ F. W. O'Dell, M. M. Shapiro, and B. Stiller, *J. Phys. Soc. Japan* **17**, Suppl. AIII, 23 (1962).¹⁶ G. D. Badhwar, R. R. Daniel, and B. Vijayakshmi, *Progr. Theoret. Phys. (Kyoto)* **28**, 607 (1962).¹⁷ G. D. Badhwar and R. R. Daniel, *Progr. Theoret. Phys. (Kyoto)* **30**, 615 (1962).

for the production of C and N from O. Though incomplete, the existing measurements of similar reactions show no striking energy dependence or, if anything, an energy dependence which might be expected to increase the nitrogen production at low energies. If the fragmentation parameters are assumed to have their high-energy values and if the abundances measured in the low-energy experiments do reflect the cosmic-ray composition outside the atmosphere more correctly than do the balloon experiments, then we are left with the conclusions that the source might be almost entirely devoid of nuclei of odd charge with $Z \geq 3$ and that the abundance of N, like the abundances of Li, Be, and B, might be used as a measure of interstellar path length. In stating this result we would caution, however, that the measurement errors are sufficiently large that a source composition with abundances of medium nuclei similar to those given by Cameron cannot be excluded entirely.

IV. CONCLUSIONS

In conclusion, we find that the L/M ratio has attained the low value of 0.16 ± 0.07 in the 40 to 100 MeV/nucleon region while the H/M ratio has a value similar to that found at higher energies.

A detailed examination of the cosmic-ray charge structure shows a nitrogen abundance which is not inconsistent with that which would result from fragmentation of the observed $Z \geq 8$ nuclei in passing through ~ 2.5 g/cm² of hydrogen. The ratio of the fluxes of oxygen and carbon nuclei is found to be 1.2 ± 0.3 in reasonable agreement with satellite measurements in the vicinity of 100 MeV/nucleon but in disagreement with the much lower value found at high energies in balloon experiments. It is not at all clear whether this disagreement results from an energy dependence of unknown origin or from the effect of the residual atmosphere above the balloons.

Source Spectra and Composition of Cosmic Rays Implied by an Analysis of Interstellar and Interplanetary Travel

C. E. FICHEL AND D. V. REAMES

Goddard Space Flight Center, Greenbelt, Maryland

(Received 12 May 1966)

The implications of a simple set of assumptions related to galactic cosmic rays are examined and compared with the existing data. These assumptions are that (a) the multiply charged cosmic-ray nuclei all have the same spectral shape at the source, (b) the relative abundances of He³ and light nuclei ($3 \leq Z \leq 5$) are negligible at the source, and (c) the average amount of interstellar material traversed is independent of particle energy. The results show that within the present uncertainties of the experimental data and of the interaction cross sections, the data agree with the predictions without additional assumptions if the differential source spectra are relatively flat at low energies and the average interstellar path is 2.8 ± 0.4 g/cm². In particular, the nearly constant relative abundances of the helium, medium ($6 \leq Z \leq 9$), and ($Z \geq 10$) nuclei for different values of energy/nucleon can be explained, and the observed variation of the flux ratio of light to medium nuclei with energy/nucleon can be brought into fair agreement with predictions. The calculations also imply that, if the analysis does represent the true situation, then there is little or no solar modulation of the cosmic rays near solar minimum, and protons and helium nuclei have different source spectra.

I. INTRODUCTION

THE cosmic-ray energy spectra which are observed at the earth represent the source spectra after they have passed through interstellar matter and have been modulated within the solar system. Whereas presumably the solar-system modulation is primarily the result of electromagnetic fields, interstellar space is believed to contain enough material along the path of the particle to change appreciably the particle energy as well as the intensity of the radiation. In the latter case, it is normally assumed that the intensity in an energy interval is changed significantly only by fragmentation in interactions and by ionization loss, and not by acceleration nor by the complicated time-dependent

magnetic effects which probably cause the intensity variation in the solar system.

If the acceleration in interstellar space is negligible, it is possible to calculate the energy dependence of the relative abundances of various particle groups outside the solar system, assuming various source spectra, provided the cross section in interstellar space and the amount of material traversed are sufficiently well known. The exact nature of the solar modulation is not yet known, but the general belief is that it probably depends only on the velocity and charge-to-mass ratio of the particle. Therefore, although nuclei of the same charge-to-mass ratio but different charges will lose energy at different rates in interstellar space, the fluxes