

Possible Mechanism for the Acceleration of Ions in Some Astrophysical Phenomena

M. FRIEDMAN*

U.K.A.E.A. Research Group, Culham Laboratory, Abingdon, Berkshire, England

(Received 20 January 1969)

A model is suggested for the acceleration of ions, based on Petschek's theory of magnetic field annihilation. It is shown that ions can gain high energies in this model, and that energy distributions similar to those observed can be obtained. It is also shown that this model prevents electrons from gaining high energy, consistent with cosmic-ray observations. It is suggested that if this model is correct, we can obtain, from the observed ion energy distribution, a relation between the (average) plasma density and the (average) magnetic field in regions where the ions are accelerated.

1. INTRODUCTION

THE acceleration of charged particles is a common phenomenon in the universe. Tens of years of observation have provided us with a knowledge of the accelerated particles and their energy distribution. Many theories have been suggested to explain the mechanism of acceleration. In most of them the medium in which the particles are accelerated and from which they draw their energy is magnetoactive plasma. (The reader is referred to reviews by Parker¹ and by Ginzburg and Syrovatskii.²) Ginzburg and Syrovatskii³ have suggested that an acceleration mechanism should be universal, namely, that any particular astrophysical phenomenon in which particles are

accelerated and on which it is easy to make observations would serve as an example from which general laws may be drawn. One such example, it appears, would be the phenomenon of solar flares.

The author would like to suggest here a very simple version of Fermi-type mechanism^{4,5} for the acceleration of ions to very high energies such as are observed following intense solar activities. The model considered here is based on a modification of Petschek's wave theory of magnetic field annihilation. Details of the model are published elsewhere.^{6,7} Inherent in Petschek's model are stationary boundary regions (Fig. 1) separating much larger regions of antiparallel magnetic field B , which, with their trapped plasma, flow into the boundary with velocity U . It will be shown later that in the rest frame an electric field $E = -(1/c)(U \times B)$ exists in the plasma, from which a particular ensemble of ions can gain high energies.

We shall report here some preliminary computations (Secs. 3 and 5) which show that, with parameters based on the revised model of annihilation of magnetic fields, energy distributions of the accelerated ions can be obtained. The details of each energy distribution depend on quantities characteristic of the plasma, e.g., plasma density, magnetic field, etc. Towards the end of this article (Sec. 7), we discuss the consequences of using this model for cosmic-ray acceleration.

Speiser⁸ has suggested a somewhat similar model to account for the observed high-energy particles coming from the tail of the magnetosphere. In his model, charged particles entering the neutral sheet region (between antiparallel magnetic fields) can gain energy under the influence of an electric field. The energies of the particles depend on the spatial variations of the magnetic and electric fields as well as on the characteristic dimensions of the neutral sheet. Although his model explains satisfactorily the observed phenomenon, it




FIG. 1. Magnetic field and plasma-flow situation in Petschek's wave model.

* Present address: Department of Plasma Studies, Cornell University, Ithaca, N. Y.

¹ E. N. Parker, in *Nebulae and Interstellar Matter*, edited by B. M. Middlehurst and L. H. Aller (The University of Chicago Press, Chicago, 1958).

² V. L. Ginzburg and S. I. Syrovatskii, *The Origin of Cosmic Rays* (Pergamon Press, Inc., Oxford, 1964).

³ V. L. Ginzburg and S. I. Syrovatskii, *Usp. Fiz. Nauk* **88**, 485 (1966) [English transl.: *Soviet Phys.—Usp.* **9**, 223 (1966)].

⁴ E. Fermi, *Phys. Rev.* **75**, 1169 (1944).

⁵ E. Fermi, *Astrophys. J.* **119**, 1 (1954).

⁶ H. E. Petschek, in *Physics of Solar Flares*, AAS-NASA Symposium National Aeronautics and Space Administration Publication No. NASA-SP-50, 1963, p. 425 (unpublished).

⁷ M. Friedman and S. M. Hamberger, *Solar Phys.* (to be published).

⁸ T. W. Speiser, *J. Geophys. Res.* **72**, 3919 (1967).

fails to tie together the plasma parameters into a self-consistent picture which would enable the model to be extended to other phenomena in which acceleration occurs.

2. FIELD GEOMETRY OF THE MODEL

Petschek's basic idea is that a situation can exist in which oppositely magnetized plasma flows into two stationary slow shock fronts. Within the fronts the inflowing magnetic energy is largely converted into kinetic energy. The outflowing plasma is both heated and ejected from the shock region from which it escapes. Four regions exist in Petschek's model (Fig. 1):

- I: initial-flow regions;
- II: final-flow regions;
- III: shock-front regions;
- IV: diffusion region.

The plasma parameters in these four regions have been discussed elsewhere.^{6,7} Here we shall give only the quantities relevant to this discussion.

(a) The widths of regions IV and III are $2\delta^*$ and δ^* , respectively, where

$$\delta^* = c^2/4\pi\sigma V_A M_0. \quad (1)$$

Under the conditions of high current density in regions III and IV, the plasma conductivity σ is⁹

$$\sigma = \frac{1}{2}(M/m)^{1/3}(4\pi n e^2/m)^{1/2}, \quad (2)$$

where m and M are the masses of electrons and ions, respectively, and n is the plasma number density; M_0 is the Mach number in the flow region (region I),

$$M_0 = U_I/V_A, \quad (3)$$

where V_A is the Alfvén velocity and U_I the flow velocity of plasma in region I.

According to Petschek,⁶ the above situation exists only for

$$M_0 \lesssim 0.1. \quad (4)$$

(b) The Y dimension of region IV is

$$2Y^* = c^2/4\pi\sigma V_A M_0^2. \quad (5)$$

(c) The geometrical equations of the shock fronts (region III) can be written

$$\pm M_0 Y + X = 0, \quad (6)$$

for

$$|Y| > Y^*.$$

(d) An electric field $\mathbf{E} = (0, 0, E_z)$, which is constant in the four regions, is inherent in the model, and is given by

$$\mathbf{E} = -(\mathbf{U}_I \times \mathbf{B}_I)/c, \quad (7)$$

where \mathbf{B}_I is the magnetic field in region I.

⁹ S. M. Hamberger and M. Friedman, Phys. Rev. Letters 21, 674 (1968).

(e) From the Rankine-Hugoniot relations¹⁰ we can obtain the magnetic field in these regions. In regions where discontinuities in magnetic fields occur (regions III and IV), we shall assume linear variations of the magnetic field across these discontinuities.

The relations used are:

$$\begin{aligned} \text{region I: } \mathbf{B}_I &= (-2(Y/|Y|)M_0 B_{y0} f(Y), \\ &\quad -(X/|X|)B_{y0}, 0); \\ \text{region II: } \mathbf{B}_{II} &= (-Y/|Y|M_0 B_{y0}, 0, 0); \\ \text{region III: } \mathbf{B}_{III} &\text{ given by a linear interpolation} \quad (8) \\ &\quad \text{between } \mathbf{B}_I \text{ and } \mathbf{B}_{II}; \\ \text{region IV: } \mathbf{B}_{IV} &= (- (Y/|Y|)M_0 B_{y0}(1 + |X|/\delta^*) \\ &\quad \times f(Y), -(X/\delta^*)B_{y0}, 0), \end{aligned}$$

where B_{y0} is the magnetic field at infinity (i.e., remote from the shock regions), and $f(Y)$ is some monotonic function fulfilling the conditions

$$f(0) = 0, \quad f(Y > Y^*) = 1. \quad (9)$$

3. MOTION OF AN ION IN A PETSCHKEK-TYPE CONFIGURATION OF MAGNETIC AND ELECTRIC FIELDS

Given the magnetic field configuration in the different regions [Eqs. (8)], the motion of ions can be analyzed. We have found that it is very complicated to calculate analytically the trajectories of ions. However, one can get some qualitative information about the motion of the ions by looking at somewhat similar configurations to which analytical solution exists: (a) the hyperbolic configuration,¹¹ and (b) the perfect antiparallel magnetic field configuration.¹²

In both of these configurations the ions can have two distinct classes of orbit. In one class the ion does not cross the neutral plane ($B=0$). The ion gyrates and drifts in the magnetic field. The second class of orbit is one in which the ion crosses the neutral plane and moves in a "serpentine" path.

If we now superimpose on these configurations an electric field perpendicular to \mathbf{B} , only the ions having the second kind of orbit will accelerate and gain energy as they move along the Z direction.

Reverting to the configuration defined by Eq. (8), and assuming for the moment that only the Lorentz electric field exists, we can write the equation of motion of an ion in the general form

$$\frac{d}{dt} \frac{M\mathbf{V}}{[1 - (V/c)^2]^{1/2}} = e\mathbf{E} + e \frac{\mathbf{V} \times \mathbf{B}}{c}, \quad (10)$$

¹⁰ A. Kantrowitz and H. E. Petschek, in *Plasma Physics in Theory and Application*, edited by W. B. Kunkel (McGraw-Hill Book Co., Inc., New York, 1966), p. 147.

¹¹ E. Åstrom, *Tellus* 8, 260 (1956).

¹² A. A. Weiss and J. P. Wild, *Australian J. Phys.* 17, 28 (1964).




FIG. 2. Relation between the computed final energy of an ion and its initial Y coordinate. The plasma parameters are for case (i), $B_{y0}=500$ G, $n=2\times 10^{11}$ cm $^{-3}$, $M_0=0.1$, and for case (ii), $B_{y0}=500$ G, $n=5\times 10^{10}$ cm $^{-3}$, $M_0=0.05$, and $f(Y)\propto Y$.

where

$$\mathbf{E} = -(\mathbf{U}_I \times \mathbf{B}_I)/c \quad (11)$$

and \mathbf{B}_I is defined in Eq. (8).

Ions with velocity $\mathbf{V}=(V_x, V_y, V_z)$ will cross the neutral plane $X=0$ and gain energy provided the following approximate relation holds:

$$\rho_{Li} = \frac{(V_x^2 + V_z^2)^{1/2}}{eB_a/Mc} \gtrsim M_0 |Y|, \quad |Y| > Y^* \\ \gtrsim \delta^*, \quad |Y| < Y^* \quad (12)$$

where ρ_{Li} is the ion Larmor radius in an average magnetic field B_a . If the last relation does not hold, the ions will gyrate around a magnetic line of force without changing their average energy.




FIG. 3. Relation between the computed final energy of an ion and its initial Y coordinate. The plasma parameters are $B_{y0}=500$ G, $n=2\times 10^{11}$ cm $^{-3}$, $M_0=0.1$, and $f(Y)\propto Y^2$.

Numerical integrations of Eq. (10) for ions were performed for several conditions. The ions were taken to move initially in the positive X direction with velocity equal to the plasma velocity U in region I and with initial positions ($X=-\delta^*$, $Y=Y_0 < Y^*$, $Z=0$). Three sets of plasma conditions were assumed:

- (i) $B_{y0}=500$ G, $n=2\times 10^{11}$ cm $^{-3}$,
 $M_0=0.1$, $f(Y)\propto Y$;
- (ii) $B_{y0}=500$ G, $n=5\times 10^{10}$ cm $^{-3}$,
 $M_0=0.05$, $f(Y)\propto Y$;
- (iii) $B_{y0}=500$ G, $n=2\times 10^{11}$ cm $^{-3}$,
 $M_0=0.1$, $f(Y)\propto Y^2$.

The maximum energies achieved are shown in Fig. 2 for the cases (i) and (ii) and in Fig. 3 for case (iii), for different values of Y_0 .

From the detailed computations one can draw the following conclusions:

- (1) Ions can gain high energies.
- (2) The time during which an ion satisfies Eq. (12), and is therefore accelerated, is a sensitive function of the value of M_0 and of the form of $f(Y)$, which governs the spatial dependence of B .
- (3) The dependence of the final ion energy ϵ on its initial position Y_0 is given by

$$(a) Y_0 = A e^{-\epsilon/\epsilon_0}, \quad \text{for cases (i), (ii), } f(Y)\propto Y \quad (14)$$

$$(b) Y_0 = B \epsilon^{-q}, \quad \text{for cases (iii), } f(Y)\propto Y^2 \quad (15)$$

where A , B , ϵ_0 , and q are functions of the magnetic field B_{y0} , the plasma density n , the Mach number M_0 , and the spatial variations of the magnetic field components.

4. PLASMA EFFECTS ON ACCELERATED IONS

In the previous section we have demonstrated that ions can accelerate to high energies in the particular configuration of magnetic and electric fields existing in Petschek's model.

However, there are two obvious objections:

- (1) If all ions in a particular region accelerate, the momentum of the plasma cannot be preserved without introducing an additional electric field.
- (2) We have made the unjustifiable assumption that the medium, through which the particles are accelerated, does not influence the ion motions.

We must therefore examine the condition of the plasma in regions III and IV. It has been suggested⁷ that in these regions the plasma is subject to current-induced microinstability, one of the main effects of which is an enormous increase in the amplitude of electric field fluctuations in the plasma with wavelengths

of the order of a Debye distance $\lambda_D = (KT/4\pi ne^2)^{1/2}$, where KT is the mean thermal energy in the plasma.^{13,14}

By adopting the concept of "waves" for these fluctuations, one can describe these electric fields by their Fourier components

$$\mathbf{E}_f = \int \mathbf{E}_{k,\omega} \exp\{i[\omega(\mathbf{k})t + \mathbf{k} \cdot \mathbf{r}]\} d\mathbf{k}d\omega, \quad (16)$$

where ω, \mathbf{k} are the frequency and wave number, and $\omega(\mathbf{k})$ describes the relation between them. The chief effect of such waves is to increase the electrical resistivity of the plasma^{7,15} by scattering and randomizing the directed motion of the charged particles and hence to rise the temperature of the plasma.

We consider here only longitudinal waves, i.e.,

$$\mathbf{E}_{k,\omega} \parallel \mathbf{k}, \quad (17)$$

whose frequency will lie to a first approximation within two regions:

(a) near the electron plasma frequency,

$$\omega_{pe} = (4\pi ne^2/m)^{1/2}; \quad (18)$$

(b) between zero and the ion plasma frequency,

$$\omega_{pi} = (m/M)^{1/2}\omega_{pe}. \quad (19)$$

The change in the velocity of an ion \mathbf{V} due to its interaction with those fluctuations in a time of order one oscillation period will be

$$\delta\mathbf{V} = \frac{e}{M} \int \frac{\mathbf{E}_{k,\omega} \exp\{i[\omega(\mathbf{k})t + \mathbf{k} \cdot \mathbf{r}]\}}{i[\omega(\mathbf{k}) - \mathbf{k} \cdot \mathbf{V}]} d\mathbf{k}d\omega. \quad (20)$$

As we are interested in ions with high velocities, we can assume that either

$$\omega(\mathbf{k}) - \mathbf{k} \cdot \mathbf{V} = 0 \quad (21)$$

or

$$\omega(\mathbf{k}) - \mathbf{k} \cdot \mathbf{V} \ll 0. \quad (22)$$

Simple theoretical arguments¹⁶ show that for the type of waves considered the resonance condition (21) is not fulfilled for ions whose velocity lies in the range

$$(m/M)V_T < \mathbf{k} \cdot \mathbf{V} / |\mathbf{k}| < V_T,$$

where V_T is the mean random electron velocity.

When $V > V_T$, so that we can have $\omega(\mathbf{k}) - \mathbf{k} \cdot \mathbf{V} = 0$, the fluctuating electric field may scatter ions through a large angle in a single interaction. However, we shall

¹³ S. M. Hamberger, A. Malein, J. H. Adlam, and M. Friedman, *Phys. Rev. Letters* **19**, 350 (1967).

¹⁴ S. G. Alikhanov, N. I. Alinovskii, G. G. Dolgov-Savelev, B. G. Eiselevich, R. Kh. Kurtullaer, V. K. Malinovskii, Yu. E. Nesterikin, V. I. Pelskii, R. Z. Sagdeev, and V. N. Semenov, in *Proceedings of the Third International Conference on Plasma Physics, Novosibirsk, 1968*, Paper No. CN-24/A-1 (unpublished).

¹⁵ R. Z. Sagdeev, *Proc. Symp. Appl. Math.* **18**, 281 (1967).

¹⁶ A. A. Vedenov, E. P. Velikhov, and R. Z. Sagdeev, *Usp. Fiz. Nauk* **73**, 701 (1961) [English transl.: *Soviet Phys.—Usp.* **4**, 332 (1961)].

show that the effect of such scattering on the ion motion is very small and can be ignored.

(1) From Eq. (20) we see that

$$\delta\mathbf{V} \parallel \mathbf{k}, \quad (23)$$

so that after scattering, the quantity $\omega - \mathbf{k} \cdot \mathbf{V}$ will change from zero to approximately $\omega - \mathbf{k} \cdot \mathbf{V} + \mathbf{k} \cdot \delta\mathbf{V}$, which in general is different from zero.

(2) The fluctuations are not coherent and their interaction time with an ion is of the order

$$\tau \sim \omega_{pe}^{-1}. \quad (24)$$

Even if we take the situation of very strong turbulence, in which the electrical energy density in the fluctuations is of the same order as the thermal energy density of the particles nKT ,

$$E_f^2/8\pi \sim nKT,$$

the change in ion velocity will be

$$\delta V \sim (eE_f/M)\omega_{pe}^{-1} = (m/M)V_T, \quad (25)$$

which is still 10^3 times smaller than V .

From the above arguments we can conclude that once an ion reaches a velocity

$$V > (m/M)V_T, \quad (26)$$

the interaction of the type $\omega(\mathbf{k}) - \mathbf{k} \cdot \mathbf{V} = 0$ will not influence its motion.

When $\omega(\mathbf{k}) - \mathbf{k} \cdot \mathbf{V} \ll 0$, the scattering is not resonant and one has to take into account the effect of many small-angle scatterings on the ion motion.

Using the random-walk approximations, one can calculate the mean free path λ for an ion to lose all its forward momentum by these fluctuations:

$$\lambda = \lambda_0 V^2 / \langle \delta V^2 \rangle, \quad (27)$$

where λ_0 is the mean free path for a single scattering and $\langle \rangle$ denotes an average over many deflections. It can easily be shown that

$$\langle \delta V^2 \rangle = (e/M)^2 \langle E_f^2 \rangle / (\mathbf{k} \cdot \mathbf{V})^2, \quad (28)$$

so that

$$\lambda = \lambda_0 (M/e)^2 (\mathbf{k} \cdot \mathbf{V})^2 V^2 / \langle E_f^2 \rangle \quad (29)$$

or

$$\lambda < \frac{4\lambda_0 \epsilon^2}{\langle (eE_f)^2 \rangle / k^2}, \quad (30)$$

where ϵ is the kinetic energy of the ion,

$$\epsilon = \frac{1}{2} M V^2. \quad (31)$$

Let us define two parameters α and β such that

$$\lambda_0 = \beta \lambda_D, \quad \beta \geq 1 \quad (32)$$

$$\langle (eE_f)^2 \rangle / k^2 = (KT/\alpha)^2, \quad \alpha \geq 1 \quad (33)$$

$$\lambda < \lambda_D (\epsilon/KT)^2 \alpha^2 \beta. \quad (34)$$

The parameters α and β depend on the intensity of the fluctuating electric fields in the plasma and will be equal to 1 in the condition of a strong electrostatic plasma turbulence.

We have found before that during its acceleration an ion passes through regions III and IV many times. The condition that this ion will not be affected by these fluctuations can be expressed as

$$\delta^* \ll \lambda. \quad (35)$$

From Eqs. (34) and (35) we obtain a criterion for the minimum ion energy for which we can neglect the effect of the fluctuating fields on the ion motion:

$$\epsilon > \frac{1}{2}KT(\delta^*/\lambda_D)^{1/2}/\alpha\beta^{1/2}. \quad (36)$$

Since, in general, $\alpha, \beta > 1$, we relax the criterion to

$$\epsilon \gtrsim \frac{1}{2}KT(\delta^*/\lambda_D)^{1/2}. \quad (37)$$

For the case of a solar flare where $\delta^* \approx 20$ cm and $\lambda_D = 2 \times 10^{-1}$ cm, we require

$$\epsilon \gtrsim 5KT. \quad (38)$$

Since ions do not lose energy in small-angle scattering, and since they can be accelerated whatever their initial direction (as shown in Sec. 5), it can be argued that the mean effect of these fluctuations on the ensemble of accelerated ions will be negligible.

In the light of the above arguments, we assume that only those ions with energy

$$\epsilon \geq \gamma KT, \quad (39)$$

where

$$1 < \gamma < 10,$$

can be freely accelerated.

Thus it is clear that only those relatively few ions which have initial energies well above the average can be freely accelerated, and so remove momentum from the system. In fact, the momentum balance is maintained by the *bulk* of the plasma.

5. FINAL ENERGY DISTRIBUTIONS

In Sec. 3 we demonstrated that ions can accelerate to high energies in the particular configuration of magnetic and electric fields existing in Petschek's model, using the simplifying assumption that all ions have the same initial (directed) velocity. To examine a more realistic situation, we have integrated Eq. (10) for ions whose initial positions within the diffusion region and whose initial directions have both been randomly chosen. The initial energy of the ions has been taken as $4KT$, in accordance with Eq. (39).

Since we have random initial conditions for the ions, their final energies provide the required distribution.

The calculations have been performed for the follow-




FIG. 4. Computed energy distribution for ions. The plasma parameters are $B_{y0} = 500$ G, $n = 2 \times 10^{11}$ cm $^{-3}$, $M_0 = 0.1$, and $f(Y) \propto Y$.

ing flare conditions:

- (i) $B_y = 500$ G, $n = 2 \times 10^{11}$ cm $^{-3}$,
 $M_0 = 0.1$, $f(Y) \propto Y$;
(ii) $B_y = 500$ G, $n = 2 \times 10^{11}$ cm $^{-3}$,
 $M_0 = 0.1$, $f(Y) \propto Y^2$. (40)

The initial energy of the ions has been taken as 40 keV ($=4KT$, where KT is the thermal energy of electrons in solar flares found from the shock equations¹⁰).

Because of the very long computing time needed for such calculations, only 100–150 ion orbits were computed for each of the above conditions.

Although the statistics seem very poor, we can draw the following conclusions:

- (1) About 70% of the ions achieve energy >0.1 MeV.
- (2) 15% of all the ions have at some time a velocity component $V_y = 0$ for which we can use the calculations discussed in Sec. 3.
- (3) The energy distributions (Figs. 4 and 5) can be expressed as

$$\begin{aligned} N(>\epsilon) &= A'e^{-\epsilon/\epsilon_0}, & \text{for case (i)} \\ &= B'e^{-a'}, & \text{for case (ii)}. \end{aligned} \quad (41)$$




FIG. 5. Computed energy distribution for ions. The plasma parameters are $B_{y0} = 500$ G, $n = 2 \times 10^{11}$ cm $^{-3}$, $M_0 = 0.1$, and $f(Y) \propto Y^2$.

The constants ϵ_0' and q' are found to be the same as ϵ_0 and q from Eqs. (14) and (15) within the statistical error.

Because of the rather small number of ions used in the computation, the maximum ion energy obtained by this method is not spectacular, since there is a low statistical probability of choosing an ion which would reach a high energy [Eq. (41)].

If we make the assumption that the result (3) is not accidental, i.e., that $\epsilon_0' = \epsilon_0$ and $q' = q$, the results of Sec. 3 can be used to get the energy distribution of the "average" ions since $Y_0 \propto N(>\epsilon)$.

All ions with initial coordinate Y_1 and initial energy $4KT$ can be accelerated provided the relation [Eq. (12)] is fulfilled:

$$Y_1 < Y_0 = \frac{1}{M_0} \frac{(8kT/M)^{1/2}}{eB_a/Mc}, \quad (42)$$

where B_a is an average magnetic field.

The total number of particles N_i that satisfy the last condition is

$$N_i \sim 4L^2 Y_0 n, \quad (43)$$

where L is the characteristic dimension.

If we assume that the accelerated ions originate from the tail of a Maxwellian distribution with mean energy KT , then only 3% of N_i will be influenced appreciably by the steady electric field, of which only 70% will gain energy >0.1 MeV; then the total number of accelerated ions $N(>\epsilon)$ will be

$$N(>0.1 \text{ MeV}) \sim \frac{2 \times 10^{-1}}{M_0} \frac{(KT/M)^{1/2}}{L^2 n eB_a/Mc}. \quad (44)$$

For example, if we take as a typical solar flare condition $B_a \sim 100$ G, $n \sim 2 \times 10^{11} \text{ cm}^{-3}$, $L \sim 10^{10} \text{ cm}$, $KT \sim 10^4$ eV, then we find $N(>0.1 \text{ MeV}) \sim 10^{35} - 10^{36}$ for the number of particles with energy >0.1 MeV. If we use the calculated distribution Eq. (41) with $q' \sim 2$ (Fig. 5), we find that the number of ions with energy greater than 10 MeV will be

$$N(>10 \text{ MeV}) \sim 10^{31} - 10^{32}. \quad (45)$$

Krimigis¹⁷ has calculated that about 10^{31} ions with energy greater than 23 MeV are produced in a solar flare event.

The parameter $f(Y)$ controls the shape of the energy distribution. It is improbable that $f(Y)$ will depend on Y in the way we have assumed; neither it is probable that $f(Y)$ will be constant in time. Computations of the kind discussed in Sec. 3 were therefore performed, taking

$$f(Y) \propto Y^P, \quad 8 > P > 2. \quad (46)$$

The results were qualitatively similar to those for $P = 2$. If we assume that the temporal change of $f(Y)$ is slow

¹⁷ S. M. Krimigis, *J. Geophys. Res.* **70**, 2943 (1965).

compared with the time of acceleration (which in the case of solar flare $\sim 10^{-4}$ sec for an ion to reach energy ~ 10 MeV), the results obtained will remain the same.

6. ACCELERATION OF ELECTRONS

In the previous section we have considered only ion acceleration. This is because observations¹⁸ show that there are far fewer high-energy electrons than ions.

The mechanism that we have discussed may give an explanation of this phenomenon. We have shown that to be accelerated an ion should fulfill two conditions:

(1) It must be injected with sufficiently high energy that it will not be influenced by the fluctuating electric fields in the plasma.

(2) During the acceleration the condition (12):

$$\rho_{Li} = V_i / (eB_a/Mc) \gtrsim \delta^*, \quad |Y| < Y^* \\ \gtrsim M_0 |Y|, \quad |Y| > Y^* \quad (47)$$

where V_i is the ion velocity, should be fulfilled.

Similar conditions should be applied also to the electrons:

(1) Even if the electrons are injected with energy such that

$$\epsilon \gg KT,$$

there will be a strong interaction with the fluctuating electric fields which will randomize their velocities, in comparison to the weaker influence of the same phenomenon for ions. Only very few electrons in the remote part of the tail of the energy distribution will not be influenced strongly by the fluctuations.

(2) During the steady acceleration the condition

$$\rho_{Le} = V_e / (eB_a/mc) \gtrsim \delta^*, \quad |Y| \leq Y^* \\ \gtrsim M_0 |Y|, \quad |Y| \geq Y^* \quad (48)$$

where ρ_{Le} is the electron Larmor radius, should be fulfilled.

It can easily be seen that to suffer the same acceleration as an ion of velocity V_i , the electron must have velocity

$$V_e = (M/m)V_i.$$

For these reasons we should expect very few electrons in the high-energy particle spectrum resulting from this mechanism.

7. APPLICATION OF ACCELERATION MODEL TO COSMIC RAYS

Observations suggest that the occurrence of anti-parallel magnetic fields is not unique to the flare. For example, it has been suggested¹⁹ that the magnetic field in the Galaxy has opposite direction to the magnetic field in the Halo.

¹⁸ J. A. Earl, *Phys. Rev. Letters* **6**, 125 (1961).

¹⁹ *The Distribution and Motion of Interstellar Matter in Galaxies*, edited by L. Woltjer (W. A. Benjamin, Inc., New York, 1962).

If we speculate that a similar mechanism occurs in the boundary region between the Galaxy and the Halo, we can calculate the maximum energy that ions can reach.

Assuming

$$B_{y0} \sim 3 \times 10^{-6} \text{ G}, \quad n \sim 10^{-2} \text{ cm}^{-3} \quad (49)$$

in the boundary region, the maximum electric field is

$$E \approx M_0 B_{y0}^2 / (4\pi n M c^2)^{1/2} \sim 3 \times 10^{-8} \text{ V cm}^{-1}. \quad (50)$$

Taking the characteristic dimension L as

$$L \sim 5 \times 10^{22} \text{ cm},$$

the maximum energy an ion can achieve will be

$$\epsilon_{\max} = eEL \sim 10^{15} \text{ eV}. \quad (51)$$

The characteristic time for dissipation of the magnetic field by Petschek's mechanism is

$$\tau \sim \frac{L}{M_0 V_A} = \frac{L}{M_0} \left(\frac{4\pi n M}{B_{y0}^2} \right)^{1/2} \sim 10^{10} \text{ yr}, \quad (52)$$

which is the order of magnitude of the life of the Galaxy.

It can be shown, by using proper scaling laws for Eq. (10), that the results obtained above for the energy distribution will be the same in this case as for the flare.

The above argument implies that there is a limit to the energy an ion can gain in the Galaxy. However, let us speculate further and say that this sort of acceleration may occur outside the Galaxy. For example, taking the approximate value for magnetic field and plasma density with the characteristic dimension in the Meta-Galaxy,² energies $> 10^{19}$ eV can be reached.

The observed energy distribution of the ions in cosmic rays is²

$$N(>\epsilon) = C\epsilon^{-l}, \quad (53)$$

with $l \sim 1.7$. It is found that for certain ranges of energy, l deviates from this value. We can attribute these changes to different plasma parameters in places where acceleration occurs.

8. RELATION OF THIS ACCELERATION MECHANISM TO THE FERMI TYPE

The mechanism of acceleration discussed above can be classified as Fermi-type acceleration,^{4,5} since an ion gains energy by "collisions" with the two moving magnetized plasmas.

Gintzburg² has questioned the Fermi mechanism by calculating the increase in plasma density between

these moving regions. He finds that in order for an ion to gain an energy of 10 GeV, the plasma must be compressed 10^5 times, which seems unreasonable.

Using Petschek's model, it can be seen that situations of this sort are avoided by allowing the plasma to escape (Fig. 1).

The main criticism of the Fermi mechanism is that in order to make it efficient, ions should start with very high energies to overcome various energy-loss mechanisms (e.g., a proton should have energy ~ 200 MeV and an Fe nucleus $\gtrsim 300$ GeV).

Fermi obtains an energy distribution function for ions having an energy between ϵ and $\epsilon + d\epsilon$:

$$\pi(\epsilon)d\epsilon = \chi(Mc^2)^x \epsilon^{-(1+x)} d\epsilon, \quad (54)$$

where $\chi = \tau/(\Lambda/c)\beta^2$, Λ is the mean free path for nuclear interaction, βc is the velocity of the magnetized plasmas, and τ is the time of flight of an ion between the two magnetized plasmas. Fermi has suggested $\Lambda = 10^{25}$ cm for relativistic protons.

Comparing the last equation with the observed distribution function, one finds that

$$1 + \tau/(\Lambda/c)\beta^2 \approx 2.7, \quad (55)$$

which suggests $\tau \sim 1.7$ yr.

The model that we have introduced suggests that τ will be of the order $\delta^*/c \sim 0.1$ sec (for $B_y \sim 3.10^{-6}$ G, $n_e \sim 10^{-2}$ electrons cm^{-3}). This is very small compared to the nuclear interaction time $\Lambda/c \sim 10^{14}$ sec, so that nuclear interactions are not important in our model. The distribution functions that we have obtained have the right form and depend only on the plasma conditions in regions where the acceleration occurs. The ejection mechanism from the region, where ions are accelerated, is responsible for the shape of the distribution function. In the absence of any serious energy loss (compared with the rate of acceleration), this model requires no such high injection energy. Instead, the requirement for initial energy ϵ_0 is simply

$$\epsilon_0 > AKT, \quad (56)$$

where A is the atomic weight and KT the thermal energy of electrons in the plasma.

As τ is very small compared to Λ/c , it is possible to understand the existence of medium and heavy nuclei in the cosmic rays, a phenomenon which could not be explained by the original Fermi theory.

ACKNOWLEDGMENTS

Helpful discussions with Dr. R. J. Bickerton and Dr. S. M. Hamberger are gratefully acknowledged.