

LETTERS TO THE EDITOR

Prompt publication of brief reports of important discoveries in physics may be secured by addressing them to this department. Closing dates for this department are, for the first issue of the month, the twenty-eighth of the preceding month; for the second issue, the thirteenth of the month. The Board of Editors does not hold itself responsible for the opinions expressed by the correspondents.

The Presence of Neutral Oxygen in the Gaseous Nebulae

The most stable configuration, s^2p^4 , of neutral oxygen gives rise to three terms, 3P , 1D and 1S . McLennan¹ has shown that the green auroral line corresponds to the $^1D-^1S$ transition. Recently Frerichs² by a very exhaustive analysis of the oxygen spectrum has been able to predict the positions of the $^3P_2-^1D$ and $^3P_1-^1D$ transitions at $6299 \pm 5\text{\AA}$ and $6363 \pm 5\text{\AA}$ respectively while a private communication to Dr. Frerichs from Professor Paschen states that he has found these lines in the laboratory at 6300.00\AA and 6363.86\AA .

Wright's³ list of nebular lines shows weak unidentified lines at 6302\AA and 6364\AA . In agreement with the corresponding transitions in O_{III} the $^3P_2-^1D$ line is two or three times as intense as the $^3P_1-^1D$ line. The general behavior of the lines in various nebulae is in agreement with this identification.

The question at once arises as to why the $^1D-^1S$ line alone occurs under auroral conditions while the $^3P-^1D$ lines alone appear in the nebulae. The theory of the transition probability⁴ due to quadrupole radiation (as these lines are "forbidden" by the ordinary selection rules no dipole radiation exists)⁵ shows that the mean life of the 1D state before transition to the 3P level is much longer, since it is an intercombination line, than that of the 1S state before transition to the 1D state. On the other hand the number of atoms reaching the 1S state is much less than that reaching the 1D state. This follows from the fact that nearly all of the atoms reaching the 1S state make the $^1D-^1S$ transition and thus arrive at the 1D state in addition to those that reach there directly. Since the 1D state has lower energy than the 1S state the number reaching it directly is also greater than that reaching the 1S directly.

Under nebular conditions the mean time between impacts with other atoms is large compared with even the longest of these mean lives and consequently the intensity of the lines is determined largely by the number of atoms reaching the respective upper states. Consequently we should expect the $^1D-^1S$ transition to be much weaker than the $^3P-^1D$ transition. This is verified by the observation that in O_{III} and N_{III} , which have the same set of low terms, the $^1D-^1S$ line is 1/20 to 1/50 as intense as the $^3P-^1D$ lines. A corresponding ratio in O_I would give the $^1D-^1S$ line at 5577\AA an intensity much below the limit that could have been observed, as the $^3P-^1D$ lines are fairly weak.

On the other hand under auroral condition the collisions between atoms are so frequent that the atom in the 1D is almost always taken out by a collision of the second kind before it can radiate the $^3P-^1D$ lines. The $^1D-^1S$ transition, however, has a high enough probability so that even under auroral conditions it can take place before a collision of the second kind removes the atom from the 1S state.

I. S. BOWEN

California Institute,
Pasadena,
July 19, 1930.

¹ McLennan, Proc. Roy. Soc. **A120**, 327 (1928).

² Frerichs, Phys. Rev. in press.

³ Wright, Studies of the Nebulae, Lick Observatory, p. 242.

⁴ Huff and Houstoun, Phys. Rev. in press.

⁵ Frerichs and Campbell, Phys. Rev. **36**, 151 (1930).