

Letters to the Editor

PROMPT publication of brief reports of important discoveries in physics may be secured by addressing them to this department. Closing dates for this department are, for the first issue of the month, the eighteenth of the preceding month, for the second issue, the third of the month. Because of the late closing dates for the section no proof can be shown to authors. The Board of Editors does not hold itself responsible for the opinions expressed by the correspondents.

Communications should not in general exceed 600 words in length.

Radioactive Rb from Deuteron Bombardment of Sr

A. C. HELMHOLZ, *Department of Physics, Radiation Laboratory, University of California;*

CHARLES PECHER, *Fellow of the Belgian-American Education Foundation, Radiation Laboratory, University of California;*

AND

PERRY R. STOUT, *Department of Plant Nutrition, University of California, Berkeley, California*

May 12, 1941

ONE of us¹ has reported the production in large quantities of a long-lived radioactive Y from the bombardment of Sr with 16-Mev deuterons. At this bombarding energy one would expect an appreciable amount of Rb to be produced by the (d, α) reaction. We have found a radioactive Rb of half-life 19.5 ± 1 days, produced from this reaction in sufficient quantities for tracer work and with the added advantage of high specific activity.

To prove that the activity was Rb the following procedure was adopted. The filtrate remaining after Y and Sr had been precipitated as $Y(OH)_3$ and $SrCO_3$ was evaporated to dryness and the ammonium salts driven off. Fe, as carrier for the Y, and Sr were again added along with 10 mg of K. Fe and Sr were precipitated in alkaline Na_2CO_3 solution. After filtration, K and Rb were precipitated from the filtrate with $Na_3Co(NO_2)_6$. If the active material had been Na it would have remained largely in solution, but over 99 percent was precipitated. The precipitate was dissolved in HCl, 4 mg of Rb added, and Rb_2SnCl_6 precipitated. All the activity came with the precipitate. Sn was then removed as sulfide, and the whole activity of, in this case, 400 μ C remained in 4 mg of $RbCl$. Subsequent experiments showed that the activity could be separated quantitatively with Rb from equal molar mixtures of K, Sr, Ca and Mg. The activity must come from the Sr (d, α) Rb reaction since the Rb contamination of the target was negligible.

There are two "stable" isotopes of Rb, Rb^{85} and Rb^{87} . Snell² found an 18-min. and an 18-day period by the slow neutron bombardment of Rb. Our activity is undoubtedly his 18-day activity. The half-life of 19.5 days was obtained by following several samples over from one to four half-lives. However, there is some evidence for a weak longer

period activity, so that the half-life may be somewhat shorter. Of the two slow neutron activities, only that of Rb^{86} can be produced by the Sr (d, α) Rb reaction, since Sr^{88} is the heaviest stable isotope of Sr. Therefore this 19.5-day period can definitely be assigned to Rb^{86} . This agrees with the assignment of the 18-min. period to Rb^{88} on the basis of Th fission experiments.^{3,4}

Rb^{86} emits negative electrons and no strong γ -rays. From absorption measurements, the spectrum has an upper limit of 1.56 ± 0.05 Mev calculated from Feather's rule.

The yield of Rb^{86} from a 2000-microampere hour bombardment of Sr was 0.4 mC, as compared with 17 mC of the 55-day Sr^{89} from Sr^{88} (d, p) Sr^{89} , and 35 mC of the 100-day Y from Sr $(d, 2n)$ Y. The isotope responsible for the production of Y is uncertain, but if we assume that all the products come from Sr^{88} , the ratio of the cross sections for the $(d, 2n)$, (d, p) , and (d, α) reactions at 16 Mev is 29 : 10 : 0.1. Dr. Martin Kamen has kindly supplied us with yield data for Fe and Mg bombarded with deuterons. In Fe the reactions are Fe^{56} $(d, 2n)$ Co^{56} ,⁵ Fe^{58} (d, p) Fe^{59} , and Fe^{56} (d, α) Mn^{54} , and correcting for the reduced abundance of Fe^{58} , the ratio is 7 : 10 : 1.7. The Fe data are for 16-Mev deuterons. For Mg at 8 Mev the reactions are Mg^{26} (d, p) Mg^{27} and Mg^{24} (d, α) Na^{22} . The ratio here is ? : 10 : 200. We see that with increasing atomic number the (d, α) reaction drops off markedly as one would expect from the increase in the Coulomb barrier height. It is interesting to note that at 16 Mev, the $(d, 2n)$ reaction is one of the most probable, if not the most probable, reaction.

We wish to express our thanks to Professor E. O. Lawrence for his interest in this work and to the Research Corporation for financial support.

¹ Charles Pecher, *Phys. Rev.* **58**, 843 (1940).

² A. H. Snell, *Phys. Rev.* **52**, 1007 (1937).

³ A. Langsdorf, Jr., *Phys. Rev.* **56**, 205 (1939).

⁴ A. H. Aten, Jr., C. J. Bakker and F. A. Heyn, *Nature* **143**, 679 (1939).

⁵ The measured Co consists of two long-life activities, and since the assignment of these activities is uncertain, our estimate of the yield may be in error by a factor of 2 or 3.

Showers of Penetrating Particles

M. DAMY DE SOUZA SANTOS, P. A. POMPEIA AND G. WATAGHIN
Department of Physics, São Paulo University, São Paulo, Brazil
December 17, 1940

IN previous papers^{1,2} a report was given of measurements which gave evidence for the existence of showers of at least two particles of a range not smaller than 17 cm of lead and the size of the penetrating core of which is of the order of 0.2 sq. m.

In this paper we report further measurements made with a sixfold coincidence set with a resolving time of 1.8×10^{-6} sec. and with the multivibrator circuit developed by one of us.³ Some of the arrangements used (Fig. 1) were chosen in order to avoid knock-on showers and the secondary effects of the soft radiation which usually accompanies the penetrating rays.