

tests were made several times a day, for about one hour each time. A persistent increase of ionization extending over half a day or so could not have been missed.

The data observed at Fordham are given in Table I.

TABLE I. Total ionization on the roof of the Physics Building, Fordham University, New York, New York.

Time	Ionization	Time	Ionization
June 29 (mean)	6.28 <i>I</i>	July 7 11 A.M.	6.19 <i>I</i>
June 30 (mean)	6.07 <i>I</i>	July 7 8 P.M.	6.07 <i>I</i>
July 1 (mean)	6.07 <i>I</i>	July 8 8 A.M.	6.41 <i>I</i>
July 2 (mean)	6.32 <i>I</i>	July 8 10 A.M.	6.78 <i>I</i>
July 3 A.M.	5.96 <i>I</i>	July 8 11 A.M.	6.67 <i>I</i>
July 3 P.M.	6.08 <i>I</i>	July 8 noon	6.06 <i>I</i>
July 4 A.M.	6.26 <i>I</i>	July 8 1 P.M.	6.05 <i>I</i>
July 4 P.M.	5.93 <i>I</i>	July 8 7.30 P.M.	6.10 <i>I</i>
July 5 8 A.M.	5.95 <i>I</i>	July 9 8 A.M.	6.33 <i>I</i>
July 5 2 P.M.	6.20 <i>I</i>	July 9 3 P.M.	5.91 <i>I</i>
July 5 8 P.M.	6.47 <i>I</i>	July 10 10 A.M.	6.18 <i>I</i>
July 6 8 A.M.	6.24 <i>I</i>	July 10 8 P.M.	5.89 <i>I</i>
July 6 4 P.M.	6.00 <i>I</i>		

The total ionization *I* (in ion pairs produced per cc and sec. in pure nitrogen at N.P.T.) observed on the roof is, on the average, about 6 *I*, consisting of 2 *I* due to cosmic rays, 3 *I* local gamma-rays from the building, 0 to 0.5 *I* gamma-rays from the radioactive substances in the atmosphere and about 0.7 *I* residual ionization of the vessel itself.

The figures reported below are average values of from 30 to 60 min. continuous observation in each case. For the first few days mean values of the whole day are given.

Isolated higher values (as for instance on July 2 where a single value of 6.8 *I* was observed just for one hour) may be caused by cosmic-ray showers or bursts and are unimportant for judgment on any possible influence of the atom bomb cloud.

From the values shown in this table it is evident that no effect ascribable to the radioactive atom bomb cloud was noticed before July 8. On this date a slight increase of ionization (0.3 to 0.7 *I*) occurred and persisted for several hours. It is highly questionable, however, whether this effect was really produced by gamma-rays from the air, for two reasons: first, the general circulation in the United States, at levels of 7–12 km in the summer, is anticyclonic and in the eastern half of the country a northwesterly wind component would prohibit a radioactive air mass from reaching the New York area; second, from the weather maps (July 4 to July 8) the component wind velocity at 7 km, in the general direction from Houston to New York, was only 5 knots. Therefore the air mass present over Houston on July 5 would not have reached New York in much less than 200 hours. It is, of course, unfortunate that our observations were discontinued on July 10, but from the evidence presented here it is practically certain that no effects from the Bikini test were in evidence in the North Atlantic states, more than 7000 miles from the Bikini Islands.

We are indebted to Mr. Hormantas, Meteorological Office, LaGuardia Field for furnishing the meteorological data.

¹ G. Herzog, *Phys. Rev.* **70**, 227 (1946).

² See a forthcoming article of V. F. Hess, *Trans. Am. Geophys. Union* **27**, No. 5 (October, 1946).

Effort to Observe Anomalous Gamma-Rays Connected with Atomic Bomb Test of July 1, 1946

E. DOROTHY WEEKES

Petty Geophysical Engineering Company, San Antonio, Texas

AND

DONALD F. WEEKES

A & M College of Texas, College Station, Texas

September 17, 1946

A GAMMA-RAY anomaly associated with the atomic bomb test at Bikini Island on July 1, 1946, having been observed by Herzog¹ at Houston, Texas, on July 4 and 5, the absence of such an anomaly at College Station, 100 miles northwest of Houston, becomes a matter of interest.

Daily measurements of the background of stray radiation were made in a laboratory on the upper floor of a two-story brick building in College Station, during the following periods: June 25 to July 5; July 8 to 11; 16 to 19; and 23 to 26. The counting unit of an x-ray spectrometer which employs Geiger counter registration was used for the purpose. A rough indication of gamma-ray sensitivity was obtained by placing a four-ounce specimen of carnotite 15 inches from the longitudinal axis of the counter. This resulted in an increase of about 45 percent in the counting rate.

The observed background counting rates varied from 0.317 to 0.339 per sec., exceeding the average value of 0.329 by slightly more than the expected statistical fluctuation (0.0064 per sec.) on three dates, July 1, 2, and 18; but, even in these cases, the variations were too small to be regarded as significant.

Since the observations were all made at about the same time of day, it is unlikely that any local anomaly of as much as 24 hours duration occurred during the periods covered by the experiment. The observation of July 5 was concluded at 10:36 A.M., several hours after the maximum of the Houston anomaly but well before its expiration, and might have been expected to reveal the existence of this anomaly, had it been of regional extent.

¹ G. Herzog, *Phys. Rev.* **70**, 227 (1946).

Schwarzschild Interior Solution

NATHAN ROSEN AND BROWNIE NEWMAN

University of North Carolina, Chapel Hill, North Carolina

September 26, 1946

IN a recent paper,¹ Wyman gave an interesting discussion of the solution of the gravitational equations of general relativity for the case of a sphere of perfect fluid of constant density, using isotropic coordinates. He found that, for a given value of the density ρ , the mass m and the radius a were restricted by the conditions,

$$m \leq 0.4a, \quad a^2 \leq 0.27R^2, \quad (1)$$

with $R^2 = \frac{3}{8}\pi\rho$, instead of the conditions found for the Schwarzschild solution,²

$$m \leq 4a/9, \quad a^2 \leq 8R^2/9. \quad (2)$$