

an average would seem to be largely valid for distributions if not for end-point energies.

In view of the great difficulty of measuring the beta-spectrum of  $H^3$ , the concordance of adjusted data is remarkable. In terms of average deviations from the mean, it appears that the shape of the distribution function is now known to a precision of about 3 percent from 0.5 to 5.5 keV, about 1 percent from 6 to 11 keV, about 5 percent from 11.5 to 17.5 keV, and about 25 percent on up to within 2 percent of the end-point energy. Equally remarkable is the fact that the deviations of the composite from the theoretical spectrum are generally comparable with (in most cases less than) the corresponding average deviations of the measurements, except at the very lowest and highest energies where the measurements are very uncertain. A Fermi plot of the points in Figs. 1 and 2 is extremely satisfying from 2.5 to 18 keV; although the point at 0.9 keV is somewhat low, the points at 1.8 keV and above 18 keV are high (this last one about 100 percent).

\* Foothill Laboratory, Pasadena, California.

<sup>1</sup> C. S. Cook and L. M. Langer, Phys. Rev. **73**, 601 (1948).

<sup>2</sup> Curran, Angus, and Cockcroft, Phil. Mag. **40**, 53 (1949).

<sup>3</sup> Curran, Angus, and Cockcroft, Phys. Rev. **76**, 853 (1949).

<sup>4</sup> Byatt, Rogers, and Waltner, Phys. Rev. **75**, 909 (1949).

<sup>5</sup> G. C. Hanna and B. Pontecorvo, Phys. Rev. **75**, 983 (1949).

<sup>6</sup> E. R. Graves and D. I. Mayer, Phys. Rev. **76**, 183 (1949), and private communication.

## Natural Ferromagnetic Resonance

A. J. E. WELCH AND P. F. NICKS

Imperial College of Science and Technology, London, England

AND

ALAN FAIRWEATHER AND F. F. ROBERTS

Post Office Research Station, Dollis Hill, London, England

October 28, 1948

WHILE studying the magnetic properties of a number of magnesium ferrites, and related spinels, evidence was obtained in a few cases of a resonance with sufficiently small damping to produce a peak in the curve of permeability *versus* frequency for unmagnetized samples. Figure 1 shows the results




FIG. 1. Natural ferromagnetic resonance.  $\odot$   $\circ$   $\ominus$  are points measured on Mg ferrite sample.

found for a particular magnesium ferrite. In the region where the resonance occurred, the measurements were carried out in terms of standing-wave observations in a coaxial line containing the sample adjacent to the short-circuited end. Care was taken that higher modes, which might propagate within the sample, were not excited. The results may be interpreted in terms of an extension of the Landau-Lifshitz theory<sup>1</sup> as amended by Kittel.<sup>2</sup> It may be

deduced from Kittel's formula,

$$\frac{\chi}{\chi_0} = \frac{\omega_0^2 + \lambda^2 \mu_0 + j\omega\lambda}{\omega_0^2 + \lambda^2 \mu_0 - \omega^2 + j\omega\lambda(1 + \mu_0)}$$

that the real part of  $\chi$ ,  $Re(\chi)$ , can only become negative if  $\lambda < \omega_0$  but that it can exceed its static value  $\chi_0$  only if  $\lambda < \omega_0/\mu_0$ . Now, in fact, it has been found that all ferrites which are appreciably ferromagnetic show negative values of  $Re(\chi)$  over some part of the range 1000 Mc/sec. to 10,000 Mc/sec., but hitherto none has been found with the positive peak in  $Re(\chi)$ . Where the positive peak has occurred in our results it has been superimposed upon a basic curve of falling  $\chi$ , as seen from Fig. 1. The complete susceptibility can thus be regarded to a second approximation as the sum of two parts, each part represented by an equation of the above form. In the example given, the basic part contributes about three-fourths of the total low frequency susceptibility, has a resonance frequency,  $\omega_0/2\pi$ , of about 140 Mc/sec., and a damping ratio  $\lambda/\omega_0$  of about 0.5; the remaining one-fourth of the LF susceptibility has a resonance frequency of about 1500 Mc/sec. and a damping ratio of about 0.1. The observed shape of this resonance, however, appears to be much less sharp than that required by the formula; the addition of a third component enables a better fit to be obtained.

If Frenkel's form<sup>3</sup> of the absorption is used in place of Kittel's, the general frequency behavior is similar, but  $Re(\chi)$  is then negative for  $\omega > \omega_0$  for all values of the relaxation time  $\tau$ , while  $Re(\chi)/\chi_0$  exceeds unity at some frequency only if  $\tau > \pi/\omega_0$ . A slightly better fit to the observed points is obtainable with two components of Frenkel's form than of Kittel's, and the relaxation time for both components is near  $10^{-9}$  sec.

At low frequencies the formula gives a loss component which varies with frequency in the same way as an eddy current loss, and it has been found that the measured apparent eddy current loss of several ferrites is of the order of magnitude predicted by the formula from high frequency measurements.

It has also been observed that the application of a sufficiently strong transverse static magnetic field restores the 10,000 Mc/sec. susceptibility to about one-fourth the LF value. This suggests that the first term (which is dominant at low frequencies) is insensitive to external fields, whereas the second term behaves as implied in Kittel's analysis, i.e.,  $\omega_0$  increases with increase of the static field.

The interpretation of the two resonance frequencies in terms of internal fields would seem to imply rather smaller fields than have hitherto been suggested but the precise magnitudes involved would seem to depend considerably on the geometry assumed for them.

Acknowledgment is made to the Chief Scientist of the Ministry of Supply and to the Engineer-in-Chief of the General Post Office for permission to make use of the information contained in this letter.

<sup>1</sup> Landau and Lifshitz, Physik Zeits. Sowjetunion **8**, 153 (1935).

<sup>2</sup> C. Kittel, Phys. Rev. **73**, 155 (1948) and other papers.

<sup>3</sup> J. Frenkel, J. Phys. U.S.S.R. **9**, No. 4, 299 (1945).

## Beta-Spectrum and Decay Scheme of ${}_{85}\text{Tb}^{160}$

S. B. BURSON, K. W. BLAIR, AND D. SAXON

Argonne National Laboratory, Chicago, Illinois


December 22, 1949

THE  $\beta$ -spectrum of  $\text{Tb}^{160}$  has been investigated in the Argonne 180° spectrometer. The purified sample of  $\text{Tb}_4\text{O}_7$  from Oak Ridge was bombarded in the Argonne heavy water pile for about 90 days and allowed to cool for about one month. The activity of a portion of the sample was then followed for  $\sim 3$  months indicating a half-life of  $71 \pm 1$  day.

The source of  $\sim 0.1$  mg/cm<sup>2</sup> was prepared on a Nylon backing



TABLE I. Internal conversion electrons in  $Tb^{160}$ .

$E$	Line assignment	$\gamma$ -energy
$31.0 \pm 0.35$ keV	$K(\gamma_1)$	84.9 keV
$76.0 \pm 0.7$	$L(\gamma_1)$	85.1
$83.8 \pm 0.7$	$M(\gamma_1)$	85.3
$144.5 \pm 1.5$	$K(\gamma_2)$	198.4
$192.3 \pm 1.6$	$L(\gamma_2)$	201.4
$246.1 \pm 1.8$	$K(\gamma_3)$	300.0
$832 \pm 4.5$	$K(\gamma_4)$	886
$877 \pm 4.5$	$L(\gamma_4)$	886
$920 \pm 4.7$	$K(\gamma_5)$	964
$962 \pm 5.0$	$L(\gamma_5)$	971

FIG. 1. Momentum distribution for  $Tb^{160}$  beta-radiation.

of  $\sim 0.1$  mg/cm<sup>2</sup>. The counter window was of several layers of Nylon totaling 0.08 mg/cm<sup>2</sup>. A spectrometer resolution of 2 percent in  $H\rho$  was used.

The momentum plot of the spectrum is shown in Fig. 1. The lines in this plot may be assigned to conversion electrons from several gamma-rays of Dy as given in Table I.

FIG. 2. Kurie plot of  ${}_{65}Tb^{160}$  (71 days).FIG. 3. Proposed decay scheme for  ${}_{65}Tb^{160}$ .

The gamma-ray energies of 85, 200, and 300 keV agree well with those previously reported by Cork, Shreffler, and Fowler.<sup>1</sup> The gamma-ray of 212 keV reported by these authors was not observed.

The peaks on the momentum plot at approximately 640 and 690 gauss-cm corresponding to energies of  $34.8 \pm 0.8$  and  $39.8 \pm 1.0$  have been assigned to Auger electrons.

The Kurie plot of the Fermi theory is shown in Fig. 2. The plot appears to consist of three linear portions down to an energy of  $\sim 230$  keV below which there is an upward trend. Extrapolations of the linear portions of the plot give  $\beta$ -end points of  $860 \pm 10$ ,  $521 \pm 10$ , and  $396 \pm 10$  keV.

A search for positive radiation yielded no evidence for a positron decay. In view of this and the above energy values of the  $\beta$ 's and  $\gamma$ 's the decay scheme of Fig. 3 is proposed.

Values from the extrapolated linear portions of the Kurie plot were replotted on the momentum diagram (the dashed lines) with corrections being made at the extreme low energy end for the window absorption. Comparison of the areas under the curves corresponding to the three betas indicated the relative intensities of the three to be  $\beta_{860} : \beta_{521} : \beta_{396} :: 1.00 : 0.96 : 0.39$ .

<sup>1</sup> J. M. Cork, R. G. Shreffler and C. M. Fowler, Phys. Rev. **74**, 240 (1948).

## The Beta-Spectrum of $He^6$ \*

V. PEREZ-MENDEZ AND H. BROWN  
 Pupin Physics Laboratories, Columbia University, New York, New York  
 December 19, 1949

THE beta-ray spectrum of  $He^6$  has been investigated in a beta-spectrometer for gaseous emitters.<sup>1</sup>

This spectrum was first investigated by Bjerger,<sup>2,3</sup> using a cloud chamber. Subsequent absorption measurements by Sommers and Sherr<sup>4</sup> gave an energy end point of  $(3.5 \pm 0.6)$  Mev. Rustad<sup>5</sup> found an end point of  $(3.2 \pm 0.2)$  Mev, and a half-life of  $(0.84 \pm 0.02)$  sec. Holmes<sup>6</sup> gives a half-life of  $(0.823 \pm 0.013)$  sec. The spectrum is of considerable theoretical interest because of the low  $ft$  value and the high maximum energy.

The  $He^6$  is produced by a  $(n, \alpha)$  reaction by bombarding finely divided beryllium powder with fast neutrons from the cyclotron. In order to obtain high specific activity of  $He^6$ , it was necessary in this experiment to employ a modification of the internal probe technique used previously with gaseous targets. By placing the