

Scattering of 9.5-Mev Protons by Neon and Argon

R. G. FREEMANTLE, *University of Birmingham, Birmingham, England*D. J. PROWSE AND A. HOSSAIN, *University of Bristol, Bristol, England*

AND

J. ROTBLAT, *University of London, London, England*

(Received August 16, 1954)

The photographic emulsion technique was employed to study the scattering of 9.5-Mev protons by neon and argon. Energy levels of Ne^{20} at 1.58, 4.20, 4.95, and 5.62 Mev have been observed as well as of A^{40} at 1.48 Mev. Angular distributions have been determined for the protons elastically scattered by Ne^{20} and A^{40} , as well as for the inelastic groups from the 1.58-Mev level of Ne^{20} and the 1.48-Mev level of A^{40} .

THE experimental procedure in this work was similar to that described in the preceding paper.

The chamber was filled with neon of normal composition at a pressure of 6.2 cm. A typical histogram of the tracks observed at an angle of 80° is shown in Fig. 1; it clearly shows the presence of a number of groups corresponding to excitation levels in Ne^{20} . From the ranges of the elastically scattered particles we obtain 9.51 ± 0.01 Mev as the incident proton energy. The excitation energies obtained from measurements at a large number of angles are 1.58 ± 0.01 , 4.20 ± 0.01 , 4.95 ± 0.02 , and 5.62 ± 0.02 Mev. The first two agree within experimental error with those given by Ajzenberg and Lauritsen,¹ while the two states at 4.95 and 5.62 Mev replace the quoted level at 5.4 Mev; it would appear that previous measurements, on the reaction $\text{F}^{19}(d,n)\text{Ne}^{20}$, had not resolved this doublet.² The possibility that one or both of these levels might be due to Ne^{22} rather than to Ne^{20} has been ruled out by the variation of range with angle.

Figure 2 shows the angular distribution of the protons elastically scattered by neon (Curve A). It bears a marked similarity to both the carbon and oxygen

curves obtained at the same energy.³ The angular distribution of the protons arising from the 1.58-Mev level is shown in Fig. 3; it fits the equation:

$$\sigma(\phi) = 14.0P_0(\phi) - 10.4P_2(\phi) + 3.2P_4(\phi).$$

There is a slight tendency to asymmetry at about 60° which may just be significant. This distribution may be compared with that obtained by Burcham *et al.*³ for protons inelastically scattered from carbon, leading to




FIG. 1. Histogram of the tracks found at an angle of 80° (lab) in the neon experiment. A few tracks due to levels in Ne^{22} are visible but are not sufficient in number to enable accurate excitation energies to be given. All ranges should be reduced by 125 microns.

¹ F. Ajzenberg and T. Lauritsen, *Revs. Modern Phys.* **24**, 321 (1952).

² T. W. Bonner, *Proc. Roy. Soc. (London)* **174**, 339 (1940).




FIG. 2. Angular distributions of protons elastically scattered from (A) neon and (B) argon. $\sigma(\phi)$ is given in millibarns per steradian. The dashed lines are the curves to be expected from Rutherford scattering only.

³ Burcham, Gibson, Hossain, and Rotblat, *Phys. Rev.* **92**, 1266 (1953).




FIG. 3. Angular distribution of protons inelastically scattered from 1.58-Mev level of Ne^{20} . The solid line is the best fit in terms of even Legendre polynomials up to $P_4(\cos\phi)$. $\sigma(\phi)$ is given in millibarns per steradian.

the excited state at 4.4 Mev. We have fitted a series of Legendre polynomials to this carbon distribution and obtained:

$$\sigma(\phi) = 15.9P_0(\phi) + 10.1P_2(\phi) - 6.0P_4(\phi).$$

The constant terms in these two expressions are similar, but the coefficients of $P_2(\phi)$ while numerically indistinguishable from each other are of opposite sign. Since the spins and parities of the first excited states of C^{12} and Ne^{20} are both^{4,5} 2^+ and those of the ground states 0^+ , the difference between the angular distributions must be due to some other property of these states.

In the experiment with argon (at a gas pressure of 9.5 cm) we found in addition to the elastic group, one due to a level at 1.48 ± 0.02 Mev. This is probably identical with that reported by Heitler, May, and

⁴ Kraus, French, Fowler, and Lauritsen, *Phys. Rev.* **89**, 299 (1953).

⁵ J. Seed, *Phil. Mag.* **44**, 921 (1953).




FIG. 4. Angular distribution of protons inelastically scattered from the 1.48-Mev level of A^{40} . $\sigma(\phi)$ is given in millibarns per steradian.

Powell⁶ at 1.46 Mev. Further investigations of the other energy levels are being made.

The angular distribution of the elastically scattered protons is shown in Fig. 2 (Curve B). The nuclear effect is much less marked in this curve than in those obtained from nitrogen or neon. At 90° the cross-section is 4 times less than that of Rutherford scattering. The general shape is similar to the other curves obtained, but the minimum has moved up to 95° and the maximum up to 125° . The angular distribution of the protons leaving A^{40} in its excited state at 1.48 Mev is shown in Fig. 4; in contrast to those obtained for carbon, nitrogen, and neon it is not symmetrical. It bears a generic resemblance to the curve obtained by Davis⁷ for the inelastic scattering of 7.1-Mev protons from Be^9 .

⁶ Heitler, May and Powell, *Proc. Roy. Soc. (London)* **190**, 180 (1947).

⁷ K. E. Davis, *Phys. Rev.* **88**, 1433 (1952).