


## Canonical quantization for quantum plasmonics with finite nanostructures

V. Dorier, J. Lampart, S. Gu erin, and H. R. Jauslin\*

*Laboratoire Interdisciplinaire Carnot de Bourgogne, CNRS, UMR No. 6303,  
Universit  Bourgogne Franche-Comt , BP 47870, 21078 Dijon, France*

 (Received 10 August 2018; revised manuscript received 16 September 2019; published 14 October 2019)

The quantization of plasmons has been analyzed mostly under the assumption of an infinite-size bulk medium interacting with the electromagnetic field. We reformulate it for finite-size media, such as metallic or dielectric nanostructures, highlighting sharp differences. By diagonalizing the Hamiltonian by means of a Lippmann-Schwinger equation, we show the contribution of two sets of bosonic operators, one stemming from medium fluctuations and one from the electromagnetic field. The results apply to general models including dissipative and dispersive responses.

DOI: [10.1103/PhysRevA.100.042111](https://doi.org/10.1103/PhysRevA.100.042111)

### I. INTRODUCTION

Quantum effects in plasmonic systems are expected to open a whole branch of new perspectives both in fundamental physics and in technological applications [1–7], e.g., in quantum information, sensors, and generally in new types of circuits involving combinations of electronic and photonic effects. In the theories for quantum plasmonics there are two main approaches that have been widely used: (a) a phenomenological approach formulated in terms of quantum Langevin equations [8–11] and (b) microscopic oscillator models for the medium coupled to the electromagnetic field [12–15]. The latter are variations of models of the type first proposed by Hopfield [16]. The quantization of a model for a homogeneous bulk medium was first treated by Huttner and Barnett [12]. Extensions to inhomogeneous media were treated in [14,15]. The main criterion for the choice of the microscopic model is that, if one integrates the equations for the medium and one inserts the currents obtained into the macroscopic Maxwell equations, one should obtain the macroscopic Maxwell equations. The elementary quanta of the model are called plasmon-polariton.

Both approaches led to the following three basic relations: The diagonal plasmon-polariton Hamiltonian is of the form

$$\hat{H} = \int d^3r \int_0^\infty d\nu \hbar\nu \tilde{C}^\dagger(\vec{r}, \nu) \cdot \tilde{C}(\vec{r}, \nu), \quad (1)$$

where  $\tilde{C}^\dagger(\vec{r}, \nu)$  and  $\tilde{C}(\vec{r}, \nu)$  are bosonic creation and annihilation operators, respectively; the electric field is expressed in terms of the bosonic operators as

$$\begin{aligned} \vec{E}(\vec{r}) = & \sqrt{\frac{\hbar\mu_0}{\pi c^2}} \int_0^\infty d\nu \int d^3r' \nu^2 \epsilon_i^{1/2}(\vec{r}', \nu) G(\vec{r}, \vec{r}', \nu) \\ & \times \tilde{C}(\vec{r}', \nu) + \text{H.c.}, \end{aligned} \quad (2)$$

where  $\epsilon_i(\vec{r}', \nu)$  is the imaginary part of the dielectric coefficient and  $G(\vec{r}, \vec{r}', \nu)$  is a Green's tensor satisfying

$$\left[ \nabla \wedge \nabla \wedge - \epsilon(\vec{r}, \nu) \frac{\nu^2}{c^2} \right] \bar{G}(\vec{r}, \vec{r}', \nu) = \mathbb{1}_3 \delta(\vec{r} - \vec{r}'); \quad (3)$$

and the Green's tensor satisfies the identity

$$\frac{\nu^2}{c^2} \int d^3r \epsilon_i(\vec{r}) G^T(\vec{r}, \vec{r}_A) G^*(\vec{r}, \vec{r}_B) = \text{Im}G(\vec{r}_A, \vec{r}_B). \quad (4)$$

These three relations may be justified in the case of a bulk medium, i.e., when the dispersive and dissipative medium extends over all space  $\mathbb{R}^3$  [8–11,14,15].

In this article we show that within the microscopic oscillator models none of these three relations is valid for a finite medium, e.g., in particular for a system with finite metallic or dielectric nanostructures. We present a procedure to construct the complete diagonalization of the Hamiltonian (22) and the electric-field operator (25) adapted to finite media, which replace Eqs. (1) and (2). Although we do not use the Green's tensor (3) explicitly in our construction, we remark that for finite media a boundary term has to be added in Eq. (4), as described in [17], Appendix A.

It has been stated in several instances in the literature [18,19] that the formula (2) for the electric-field observable cannot be generally correct and that it has to be completed by including a term reflecting the contribution of the free electromagnetic field, while Eq. (2) describes only the electric field produced by the charges of the medium. A simple argument showing that (2) cannot be the complete formula for a finite medium is that if we take the limit where the size of the medium goes to zero, according to (2) the electric-field observable would disappear. There must be another contribution that converges to the free electric field in the limit of a vanishing medium.

Some extensions of Eq. (2) were proposed in Refs. [18,19], consisting in adding the free electric field  $\vec{E}_{\text{free}}$ ,

$$\vec{E}_{\text{tot}} = \vec{E}_{\text{free}} + \vec{E}_{\text{medium}}, \quad (5)$$

\*jauslin@u-bourgogne.fr

with  $\vec{E}_{\text{medium}}$  given by (2). This approach is not quite satisfactory, since  $\vec{E}_{\text{free}}$  was added without a justification from the diagonalization of the microscopic model. This might explain why this procedure was not widely adopted in the literature [20].

In this article we show that the electric-field operator has indeed two components: one that converges to the free electric field when the size of the medium goes to zero and another one that vanishes in that limit. However, our result shows that the first component is not just the free electric field, but a modification induced by the presence of the medium. The second component is not equal to Eq. (2), but to an expression that is modified due to the presence of the first term. We obtain this result by diagonalizing the microscopic model for a finite medium, as opposed to an infinite bulk medium as in Refs. [12–15]. Furthermore, we show that the diagonalized Hamiltonian is not of the form (1). It can be shown that, in the limit of vanishing coupling between the medium and the electromagnetic field, the operator (1) tends toward the Hamiltonian of the uncoupled medium and the Hamiltonian of the uncoupled electromagnetic field would be missing. Finally, we compare in perturbation theory the electric-field observable derived from our model with the one proposed in the literature.

## II. MICROSCOPIC PLASMON-POLARITON MODEL

We consider the microscopic model for the classical electromagnetic field interacting with a linear medium used in Ref. [15]. We consider a nonmagnetic medium for simplicity, but our construction can easily be extended to magnetic media and to the models of similar type considered, e.g., in Refs. [13,14]. It can be written in Hamiltonian form by choosing the pairs of canonically conjugate variables  $\vec{\Pi}_X(\vec{r}, \nu)$  and  $\vec{X}(\vec{r}, \nu)$  for the harmonic oscillators describing the medium, the vector potential  $\vec{A}(\vec{r})$ , and its canonically conjugate variable

$$\vec{\Pi}_A(\vec{r}) := -\vec{D}(\vec{r}) = -\epsilon_0 \vec{E}(\vec{r}) - \int_0^\infty d\nu \alpha(\vec{r}, \nu) \vec{X}(\vec{r}, \nu). \quad (6)$$

The Hamiltonian is

$$H = H_{\text{em}} + H_{\text{med}} + H_{\text{int}}, \quad (7)$$

with

$$H_{\text{em}} = \frac{1}{2} \int d^3r \left[ \frac{1}{\epsilon_0} \vec{\Pi}_A^2(\vec{r}) - \frac{1}{\mu_0} \vec{A}(\vec{r}) \cdot \Delta \vec{A}(\vec{r}) \right], \quad (8a)$$

$$H_{\text{med}} = \frac{1}{2} \int_{V_m} d^3r \int_0^\infty d\nu [\vec{\Pi}_X^2(\vec{r}, \nu) + \nu^2 \vec{X}^2(\vec{r}, \nu)], \quad (8b)$$

$$H_{\text{int}} = \frac{1}{\epsilon_0} \int_{V_m} d^3r \vec{\Pi}_A(\vec{r}) \cdot \int_0^\infty d\nu \alpha(\vec{r}, \nu) \vec{X}(\vec{r}, \nu) + \frac{1}{2\epsilon_0} \int_{V_m} d^3r \left[ \int_0^\infty d\nu \alpha(\vec{r}, \nu) \vec{X}(\vec{r}, \nu) \right]^2, \quad (8c)$$

where  $V_m$  is the volume occupied by the medium and  $\alpha(\vec{r}, \nu)$  is the coupling. It is related to the imaginary part  $\epsilon_i$  of the dielectric coefficient of the medium by  $\alpha^2(\vec{r}, \nu) = 2\epsilon_0 \nu \epsilon_i(\vec{r}, \nu) / \pi$ .

This microscopic model produces the macroscopic Maxwell equations when the medium variables are integrated [15].

We will use the real basis of transverse generalized eigenfunctions of the Laplacian

$$\vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r}) = \begin{cases} \frac{1}{2\pi^{3/2}} \vec{e}_\sigma(\vec{k}) \cos(\vec{k} \cdot \vec{r}), & \pi = c \\ \frac{1}{2\pi^{3/2}} \vec{e}_\sigma(\vec{k}) \sin(\vec{k} \cdot \vec{r}), & \pi = s, \end{cases} \quad (9)$$

where  $\sigma = \pm$  is the index for the basis of linear polarization and  $\vec{e}_\pm(\vec{k})$  are two real unit vectors orthogonal to  $\vec{k}$  and to each other. In the first step, we apply the simple canonical transformation  $(\vec{\Pi}_A, \vec{A}) \mapsto (\underline{p}, \underline{q})$ ,

$$\vec{\Pi}_A(\vec{r}) = \int d\kappa \sqrt{\epsilon_0 \omega_\kappa} \vec{\varphi}_\kappa(\vec{r}) q_\kappa, \quad (10a)$$

$$\vec{A}(\vec{r}) = - \int d\kappa \frac{1}{\sqrt{\epsilon_0 \omega_\kappa}} \vec{\varphi}_\kappa(\vec{r}) p_\kappa. \quad (10b)$$

Here we use the abbreviation for the labels  $\kappa := (\vec{k}, \sigma, \pi)$  and we define by  $\omega_\kappa = c|\vec{k}|$  the eigenfrequencies. The integral over  $d\kappa$  is an abridged notation for a combination of integrals and sums.

In these variables, the coupled Hamiltonian takes the form

$$H = \frac{1}{2} P \cdot P + \frac{1}{2} Q \cdot \Omega^2 Q, \quad (11)$$

where

$$P := \left[ \frac{p}{\vec{\Pi}_X} \right], \quad Q := \left[ \frac{q}{\vec{X}} \right], \quad (12)$$

$$\Omega^2 = \Omega_0^2 + V. \quad (13)$$

Here  $\Omega_0$  and  $\Omega$  are the frequency operators of the uncoupled and coupled systems, respectively, and  $V$  is the coupling operator, which can be explicitly extracted from Eq. (8c). Since  $V$  acts as a multiplication by  $\alpha(\vec{r}, \nu)$  and an integration over the volume of the medium, it is uniquely defined for every dielectric coefficient (hence for every material) and for every geometry of the medium.

## III. GENERAL QUANTIZATION PROCEDURE

The quantization of a Hamiltonian of the general form (11) and its expression in terms of creation and annihilation operators can be formulated in a systematic way [21,22] in terms of an orthonormal basis of generalized eigenfunctions of  $\Omega^2$ ,

$$\Omega^2 \psi_{\lambda, d^\lambda} = \lambda^2 \psi_{\lambda, d^\lambda}, \quad (14)$$

where  $d^\lambda$  is a label for the degeneracy indices. This quantization procedure can be thought of as a generalization of the usual quantization of the free electromagnetic field, in which the plane waves, which are eigenfunctions of the Laplacian, are replaced by the eigenfunctions of  $\Omega^2$ . The uncoupled  $\Omega_0^2$  has a continuous spectrum. Under some conditions on the coupling  $V$ , this is also the case for the coupled operator  $\Omega^2$ . The diagonal quantized Hamiltonian can then be written as

$$\hat{H} = \int_0^\infty d\lambda \sum_{d^\lambda} \hbar \lambda \hat{B}_{\psi_{\lambda, d^\lambda}}^\dagger \hat{B}_{\psi_{\lambda, d^\lambda}}. \quad (15)$$

The notation  $\sum_{d^\lambda}$  for the degeneracy is abridged notation for a combination of sums and integrals. The creation and annihilation operators act in the bosonic Fock space  $\mathfrak{F}^B(\mathcal{P})$  [21–23], with

$$\hat{B}_\psi^\dagger |\emptyset\rangle = |\psi\rangle, \quad (16)$$

where  $|\emptyset\rangle$  is the vacuum state and  $\psi$  are functions corresponding to the coordinates of the classical phase space  $\mathcal{P}$  that enters the definition of the Fock space. Their interpretation is that  $\hat{B}_\psi^\dagger$  creates one quantum on the phase-space mode  $\psi$ . The orthonormality relation of the generalized eigenfunctions of  $\Omega^2$  implies the canonical commutation relations. The canonical observables can be expressed in terms of the creation and annihilation operators as

$$\hat{Q} = \sqrt{\frac{\hbar}{2}} \int_0^\infty d\lambda \sum_{d^\lambda} \lambda^{-1/2} (\psi_{\lambda,d^\lambda} \hat{B}_{\psi_{\lambda,d^\lambda}} + \text{H.c.}), \quad (17a)$$

$$\hat{P} = -i\sqrt{\frac{\hbar}{2}} \int_0^\infty d\lambda \sum_{d^\lambda} \lambda^{1/2} (\psi_{\lambda,d^\lambda} \hat{B}_{\psi_{\lambda,d^\lambda}} - \text{H.c.}). \quad (17b)$$

#### IV. QUANTIZATION OF THE PLASMON-POLARITON MODEL

We can now apply the above general procedure to the model (7) and (8). As we show below, the (squared) frequency operator  $\Omega^2$  in general has a continuous spectrum with two distinct families of generalized eigenfunctions  $\psi^e$  and  $\psi^m$ ,

$$\Omega^2 \psi_{\omega,d^\omega}^e = \omega^2 \psi_{\omega,d^\omega}^e, \quad (18a)$$

$$\Omega^2 \psi_{v,d^v}^m = v^2 \psi_{v,d^v}^m, \quad (18b)$$

where the degeneracy indices are  $d^\omega := (\vartheta, \phi, \sigma, \pi)$ , with  $\vartheta$  and  $\eta$  the angles in spherical coordinates of  $\vec{k}$ , and  $d^v := (\vec{r}, j)$ , with  $j = 1, 2, 3$  labeling the three components of the fields.

The eigenfunctions have the same structure as Eq. (12), i.e., they are composed of two blocks:

$$\psi_{\omega,d^\omega}^e = \begin{bmatrix} u_{\omega,d^\omega}^e \\ v_{\omega,d^\omega}^e \end{bmatrix}, \quad \psi_{v,d^v}^m = \begin{bmatrix} u_{v,d^v}^m \\ v_{v,d^v}^m \end{bmatrix}. \quad (19)$$

The functions  $u$  in the upper block depend on the variables  $(\vec{k}, \sigma, \pi)$  associated with the electromagnetic field and the functions  $v$  in the lower block depend on the  $(v, \vec{r}, j)$  of the oscillators. In the limit where the coupling  $V$  vanishes, the  $\psi^e$  tend to eigenfunctions of the free electromagnetic field and the  $\psi^m$  tend to eigenfunctions of the uncoupled medium. They satisfy the Lippmann-Schwinger equations (see [24], Sec. XI.6, p. 98)

$$\psi_{\omega,d^\omega}^e = \phi_{\omega,d^\omega}^e - (\Omega_0^2 - \omega^2 \mp i0^+)^{-1} V \psi_{\omega,d^\omega}^e, \quad (20a)$$

$$\psi_{v,d^v}^m = \phi_{v,d^v}^m - (\Omega_0^2 - v^2 \mp i0^+)^{-1} V \psi_{v,d^v}^m, \quad (20b)$$

with  $\mp i0^+$  two possible ways to avoid the singularity in the complex plane. The functions  $\phi$  are eigenfunctions of the uncoupled operator  $\Omega_0^2$ ,

$$\phi_{\omega,d^\omega}^e(\omega', d^{\omega'}) = \begin{bmatrix} \delta(\omega - \omega') \delta_{d^\omega, d^{\omega'}} \\ 0 \end{bmatrix}, \quad (21a)$$

corresponding to a plane wave in Fourier representation, and

$$\phi_{v,\vec{r},j}^m(v', \vec{r}', j') = \begin{bmatrix} 0 \\ \delta(v - v') \delta(\vec{r} - \vec{r}') \delta_{j,j'} \end{bmatrix}, \quad (21b)$$

corresponding to an oscillator at position  $\vec{r}$  oriented in the direction  $\vec{e}_j$ . This gives a one-to-one correspondence between the coupled and the uncoupled generalized eigenfunctions. We will show below that the correspondence is unitary, which implies that the spectrum of  $\Omega^2$  has the same degeneracy structure as the one of the uncoupled operator  $\Omega_0^2$ .

The diagonalized Hamiltonian takes the form

$$\hat{H} = \int_0^\infty d\omega \sum_{d^\omega} \hbar\omega \hat{D}_{\omega,d^\omega}^\dagger \hat{D}_{\omega,d^\omega} + \int_0^\infty dv \int_{V_m} d^3r \hbar v \hat{C}_{v,\vec{r}}^\dagger \cdot \hat{C}_{v,\vec{r}}, \quad (22)$$

where we have introduced the notation

$$\hat{C}_{v,\vec{r},j} := \hat{B}_{\psi_{v,d^v}^m}, \quad (23a)$$

$$\hat{D}_{\omega,d^\omega} := \hat{B}_{\psi_{\omega,d^\omega}^e}. \quad (23b)$$

The expression (22) resembles the Hamiltonian (1) to which one would have added a free-field contribution. However, the creation and annihilation operators that appear in Eq. (22) are a deformation of the ones of the uncoupled model.

In the exterior of the medium, the last term in Eq. (6) vanishes and Eq. (10a) gives

$$\vec{E}(\vec{r}) = -\frac{1}{\sqrt{\epsilon_0}} \int d\kappa \omega_\kappa \vec{\varphi}_\kappa(\vec{r}) \hat{q}_\kappa. \quad (24)$$

Using Eq. (12), the electric-field observable can be written in terms of creation and annihilation operators on both sets of eigenmodes:

$$\begin{aligned} \vec{E}(\vec{r}) &= \vec{E}^e(\vec{r}) + \vec{E}^m(\vec{r}), \\ \vec{E}^e(\vec{r}) &= \int d\kappa \vec{\beta}_\kappa(\vec{r}) \int_0^\infty \frac{d\omega}{\sqrt{\omega}} \sum_{d^\omega} (u_{\omega,d^\omega}^e(\kappa) \hat{D}_{\omega,d^\omega} + \text{H.c.}), \\ \vec{E}^m(\vec{r}) &= \int d\kappa \vec{\beta}_\kappa(\vec{r}) \int_0^\infty \frac{dv}{\sqrt{v}} \sum_{d^v} (u_{v,d^v}^m(\kappa) \hat{C}_{v,d^v} + \text{H.c.}), \end{aligned} \quad (25)$$

with  $\vec{\beta}_\kappa(\vec{r}) := -\sqrt{\frac{\hbar}{2\epsilon_0}} \omega_\kappa \vec{\varphi}_\kappa(\vec{r})$ , and the sums over degeneracy indices correspond to the following combinations:

$$\sum_{d^\omega} := \frac{\omega^2}{c^3} \int_0^{\pi/2} d\vartheta \sin \vartheta \int_0^{2\pi} d\eta \sum_{\sigma=\pm} \sum_{\pi=c,s}, \quad (26)$$

$$\sum_{d^v} := \int_{V_m} d^3r \sum_{j=1}^3. \quad (27)$$

Equation (25) is the main result of this construction. In particular, it exhibits the appropriate convergence in the limit of zero coupling.

In summary, the electric-field observable at a point  $\vec{r}$  in the exterior of the medium can be written according to Eq. (25)

as the sum of two terms. When the coupling goes to zero, one can show that the first term  $\vec{E}^e$  goes to the free electric field

$$\vec{E}^e(\vec{r}) \rightarrow - \int d\omega \sum_{d^\omega} \sqrt{\frac{\hbar\omega}{2\epsilon_0}} \vec{\varphi}_{\omega,d^\omega}(\vec{r}) [\hat{D}_{\omega,d^\omega}^0 + \hat{D}_{\omega,d^\omega}^{0\dagger}] \quad (28)$$

and the second term  $\vec{E}^m$  vanishes. However, for an arbitrary coupling, the first term is not equal to the free electric field, but to a deformation due to the presence of the medium.

## V. STRUCTURE OF THE EIGENFUNCTIONS

The uncoupled operator  $\Omega_0^2$  has an absolutely continuous spectrum of infinite degeneracy, labeled by the indices  $d^\omega$  for the electromagnetic field and  $d^v$  for the oscillators of the medium. We will now explain why the spectrum of the coupled operator  $\Omega^2 = \Omega_0^2 + V$  has the same structure, including degeneracy, for the type of couplings that occur in plasmon-polariton models for a finite-size medium.

The methods and the intuition in the theory of continuous spectra are closely related to scattering theory. A central object in scattering theory is the Møller wave operator (see [24], Sec. XI, p. 18)

$$W_\pm(\Omega^2, \Omega_0^2) = \lim_{t \rightarrow \mp\infty} e^{i\Omega^2 t} e^{-i\Omega_0^2 t}. \quad (29)$$

If this operator is well defined, i.e., if the strong limit  $t \rightarrow \mp\infty$  exists, and is unitary we have

$$\Omega^2 = W_\pm \Omega_0^2 W_\pm^\dagger, \quad (30)$$

so  $\Omega^2$  is unitarily equivalent to  $\Omega_0^2$  and its spectrum must also be continuous with the same degeneracy. The generalized eigenfunctions are then given by  $\psi_{\lambda,d^\lambda} = W_\pm \phi_{\lambda,d^\lambda}$ . The unitarity of  $W_\pm$  ensures that the eigenfunctions satisfy the usual relations of orthogonality and completeness.

The coupling  $V$  in the plasmon-polariton models belongs to a class that has been thoroughly studied since pioneering works by Friedrichs [25–27]. The wave operator  $W_\pm$  is an integral operator that can be constructed using the methods of [28].<sup>1</sup> The choice of sign of  $W_\pm$  induces two possible sets of eigenfunctions verifying the Lippmann-Schwinger equation (20) either with  $-i0^+$  (for  $W_+$ ) or with  $+i0^+$  (for  $W_-$ ). The two sets of eigenfunctions are related by the unitary map  $S = W_+^\dagger W_-$ , which is the scattering operator [24]. Further analysis reveals that  $\Omega^2$  has no eigenvalues and a purely absolutely continuous spectrum, provided the coupling function  $\alpha(\nu)$  does not vanish for any  $\nu > 0$ . This assumption is satisfied for the usual models for dissipative and dispersive media. Consequently,  $W_\pm$  is unitary and this shows that the generalized eigenfunctions  $\psi_{\lambda,d^\lambda}$  have the structure we claim. Figure 1 is a sketch of this preservation of the spectral structure.

<sup>1</sup>Under appropriate continuity assumptions on  $\alpha(\nu)$  and with an ultraviolet cutoff of arbitrary size on the electromagnetic field, one can apply Theorem 4.2.1 in [28] if  $\alpha(\nu = 0) = 0$ . If  $\alpha(\nu = 0) \neq 0$  an adaptation of the argument yields an analogous result.




FIG. 1. Spectral structure of the plasmon-polariton model before and after diagonalization. The structure of two continuous spectra is preserved by the unitarity of the diagonalization transformation.

## VI. COMPARISON IN PERTURBATION THEORY

For a precise comparison between our construction and the results of Refs. [8–16, 18–20], one must be able to solve the Lippmann-Schwinger equations (20) in a given situation. A complete numerical solution, which includes the high degree of degeneracy of the model, is beyond the scope of the present work. We present here a preliminary result where the equations (20) are solved in first-order perturbation theory, i.e., where the volume of the medium and the dissipation are small. The details of the calculation are described in the Appendix. Once the obtained functions  $u^e$  and  $u^m$  are inserted in the electric-field operator (25), it reads

$$\vec{E}(\vec{r}) = \vec{E}_0(\vec{r}) + \sqrt{\frac{\hbar\mu_0}{\pi c^2}} \int_0^\infty d\nu \int_{V_m} d^3r' \nu^2 \epsilon_i^{1/2}(\nu, \vec{r}') \times [G_0^\pm(\nu; \vec{r}', \vec{r}) + F_{\text{nf}}(\nu; \vec{r}', \vec{r})] \vec{C}_{\nu, \vec{r}'} + \text{H.c.}, \quad (31)$$

with  $\vec{r}$  a position in the exterior of the medium. Here  $\vec{E}_0$  is of the same form as the electric-field operator in vacuum (28). It is however not strictly the same, since the annihilation operator  $\hat{D}$  acts on the plasmonic Fock space, which has a modified ground state. In addition,  $G_0^\pm$  are two possible Green's functions in vacuum

$$G_0^\pm(\vec{r}', \vec{r}) = c^2 \int d\vec{k} \sum_{\sigma, \pi} \frac{\vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r}') \otimes \vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r})}{\omega^2 - \nu^2 \mp i0^+} \quad (32)$$

and  $F_{\text{nf}}$  is a near-field term, which tends quickly to zero for  $\vec{r}$  far from the surface of the medium:

$$F_{\text{nf}}^{ij}(\vec{r}', \vec{r}) = \frac{-c^2}{4\pi \nu^2 |\vec{r} - \vec{r}'|^3} \left( \delta_{ij} - \frac{3(r'_i - r_i)(r'_j - r_j)}{|\vec{r} - \vec{r}'|^2} \right). \quad (33)$$

The two possible Green's tensors  $G_0^\pm$  can be obtained for a given choice of the Møller operator  $W_\pm$ . One can show that  $G_0^-$  satisfies the Sommerfeld radiation condition at infinity, with only outgoing waves to infinity, whereas  $G_0^+$  contains only incoming waves from infinity. It is customary to use the Sommerfeld radiation condition. Both representations can be used, however, since they are linked by a unitary transformation.

The expression (31) is obtained at first perturbative order in  $\sqrt{\epsilon_i}$ . The link with Eq. (2) can be done by also developing the Green's equation (3) to first order in  $\sqrt{\epsilon_i}$ , which corresponds



to replacing the Green's tensor in (2) by its vacuum expression  $G_0$ . We conclude that to first order in  $\sqrt{\epsilon_i}$ , the expression (31) takes the same form as the formulas proposed in Refs. [18,19], i.e., an electric field of the form (2) but with an extra term associated with the free field. There is also an additional near-field term in (31); this term diverges at the surface, and it is thus beyond the range of validity of perturbation theory.

It is worth mentioning that the free-field term  $\vec{E}_0$  is modified at higher orders of the perturbative development, which contradicts the idea that adding the contribution of the free field only is sufficient to achieve a complete quantization of the model.

## VII. CONCLUSION AND OUTLOOK

In summary, unlike in Eq. (1), the diagonalized coupled Hamiltonian (22) is of the same form as the uncoupled one: it has an absolutely continuous and positive spectrum with degeneracy labeled by the indices  $d^\omega$  and  $d^v$ .

Since the starting point for the diagonalization procedure in Refs. [14,15] is the assumption of a diagonal Hamiltonian of the form (1), the preceding arguments imply that the results of Refs. [14,15] cannot be applied to finite-size media, in particular to nanostructures, where the Hamiltonian has the form (22). The Fano-type diagonalization technique [29], which was adapted in Refs. [14,15], was conceived initially to treat a discrete mode coupled to a continuum. A characteristic phenomenon is that, under some hypothesis, the discrete mode “dissolves” in the continuum, as was first proven by Friedrichs [26] and rederived by Fano [29], and the coupled system has only a continuous spectrum. This is an intuition that may have led one to postulate a diagonal Hamiltonian of the form (1). However, in the present models for finite media we have two coupled continua instead of a discrete mode and a continuum, and the Fano-Friedrichs mechanism does not apply. Instead we need an analysis of what happens when two continua interact, which is what we presented in this article.

We note that the frequency operator  $\Omega$  is diagonalized by a unitary transformation, given by the wave operator  $W_+$ , which guarantees that the operators  $\vec{C}_{v,\vec{r}}$  and  $\hat{D}_{\omega,d^\omega}$  automatically satisfy the bosonic commutation relations. In the method of diagonalization used in Refs. [14,15], the commutation relations have to be imposed *a posteriori*, which is a technically nontrivial operation, in particular if one wants to extend it to include the two families of boson operators as in Eq. (22).

The relations (1)–(4) were constructed implicitly for bulk systems, but they have been widely used to analyze phenomena in systems with finite-size media, e.g., enhanced spontaneous emission; Purcell, Casimir, and Polder effects; and superradiant emission, among many others [11,30]. These applications have to be reassessed, taking into account the modified perspective including the missing terms, e.g., in the electric field. A natural question is whether the bulk formulas (1)–(4) can be a good approximation to the complete expressions in some particular regimes and beyond the perturbative approach described in Sec. VI.

For applications it will be necessary to develop or adapt efficient numerical methods to solve the Lippmann-Schwinger integral equations for media of different geometries (see, e.g.,

Ref. [31]), e.g., to treat the fields of nanoantennas [32,33]. An analytical and numerical study of these equations and the consequences for spontaneous emission is left for future work. The present results apply to a large class of open quantum systems coupled to a bath, featuring dispersion and dissipation.

## ACKNOWLEDGMENTS

This work was supported by the “Investissements d’Avenir” program, project ISITE-BFC/IQUINS (Grant No. ANR-15-IDEX-03), QUACO-PRC (Grant No. ANR-17-CE40-0007-01), and the EUR-EIPHI Graduate School (Grant No. 17-EURE-0002). We also acknowledge support from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 765075 (LIMQUET).

## APPENDIX: FIRST-ORDER PERTURBATION THEORY

We show here that a perturbation development to first order of the Lippmann-Schwinger equation results in the expression (31) of the electric-field operator. To lighten the notation, we show it only for  $G_0^-$ , which satisfies the Sommerfeld radiation condition.

The Lippmann-Schwinger equations (20) take the general form

$$\psi_{\lambda,d^\lambda}^{e/m} = \phi_{\lambda,d^\lambda}^{e/m} - R_0 V \psi_{\lambda,d^\lambda}^{e/m}, \quad (\text{A1})$$

where  $R_0(\lambda) = (\Omega_0^2 - \lambda^2 + i0^+)^{-1}$ . If the coupling  $V$  is small, Eq. (A1) can be developed as

$$\psi_{\lambda,d^\lambda}^{e/m} = [\mathbb{1} - R_0 V + (R_0 V)^2 - \dots] \phi_{\lambda,d^\lambda}^{e/m}. \quad (\text{A2})$$

In first order, we neglect all contributions of  $V^n$  with  $n > 1$ ; we thus have

$$\psi_{\lambda,d^\lambda}^{e/m} = \phi_{\lambda,d^\lambda}^{e/m} - R_0 V \phi_{\lambda,d^\lambda}^{e/m}. \quad (\text{A3})$$

How  $V$  acts on a general vector of the Hilbert space is deduced from the Hamiltonian of interaction (8c). It takes the form

$$V \begin{bmatrix} u(\kappa) \\ v(v, \vec{r}, j) \end{bmatrix} = \begin{bmatrix} (Bv)(\kappa) \\ (B^T u + Av)(v, \vec{r}, j) \end{bmatrix}, \quad (\text{A4})$$

with

$$(Bv)(\kappa) = \int dv \int_{V_m} d^3 r \omega \alpha(v, \vec{r}) \vec{\varphi}_\kappa(\vec{r}) \cdot \vec{v}(v, \vec{r}), \quad (\text{A5a})$$

$$(B^T u)(v, \vec{r}, j) = \alpha(v, \vec{r}) \int d\kappa \omega \vec{\varphi}_\kappa(\vec{r}, j) u(\kappa), \quad (\text{A5b})$$

$$(Av)(v, \vec{r}, j) = \alpha(v, \vec{r}) \int dv' \alpha(v', \vec{r}) v(v', \vec{r}, j), \quad (\text{A5c})$$

where we recall the concise notation  $\kappa = (\vec{k}, \sigma, \pi)$ . Note that we absorbed  $\epsilon_0$  by making the change of variable  $\alpha \mapsto \sqrt{\epsilon_0} \alpha$ . We then apply  $V$  to the uncoupled eigenfunctions  $\phi^{e/m}$  given by Eq. (21) and we insert it into Eq. (A3) to obtain the

solutions of the Lippmann-Schwinger equations in a small coupling regime. They take the form (19) with

$$u_{\kappa}^e(\kappa') = \delta(\kappa - \kappa'), \quad (\text{A6a})$$

$$v_{\kappa}^e(\nu, \vec{r}, j) = -\frac{\omega\varphi_{\kappa}(\vec{r}, j)\alpha(\nu, \vec{r})}{\nu^2 - \omega^2 + i0^+}, \quad (\text{A6b})$$

$$u_{\nu, \vec{r}, j}^m(\kappa) = -\frac{\omega\varphi_{\kappa}(\vec{r}, j)\alpha(\nu, \vec{r})}{\omega^2 - \nu^2 + i0^+}, \quad (\text{A6c})$$

$$v_{\nu, \vec{r}, j}^m(\nu', \vec{r}', j') = \delta(\nu - \nu')\delta(\vec{r} - \vec{r}')\delta_{j, j'} - \frac{\alpha(\nu, \vec{r})\alpha(\nu', \vec{r}')}{\nu^2 - \nu'^2 + i0^+}\delta(\vec{r} - \vec{r}')\delta_{j, j'}. \quad (\text{A6d})$$

Now that the eigenfunctions of  $\Omega^2$  have been found, we can use them to calculate the electric field in the exterior of the medium. We insert  $u^e$  and  $u^m$  into Eqs. (25) and we obtain, after some manipulations,

$$\vec{E}^e(\vec{r}) = \vec{E}^0(\vec{r}), \quad (\text{A7a})$$

$$\vec{E}^m(\vec{r}) = \sqrt{\frac{\hbar}{\pi\epsilon_0}} \int d\nu \int_{V_m} d^3r' \epsilon_i^{1/2}(\nu, \vec{r}') L(\nu; \vec{r}', \vec{r}) \vec{C}_{\nu, \vec{r}'} + \text{H.c.}, \quad (\text{A7b})$$

with

$$L(\vec{r}', \vec{r}) = \int d^3k \frac{\omega^2}{\omega^2 - \nu^2 + i0^+} \sum_{\sigma, \pi} \vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r}') \otimes \vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r}). \quad (\text{A8})$$

We express the integrals in spherical coordinates with Eq. (26) and we use the identity

$$\frac{1}{\omega^2 - \nu^2 \mp i0^+} = \frac{\mathcal{P}}{\omega^2 - \nu^2} \pm i\frac{\pi}{2\nu} \delta(\omega - \nu) \quad (\text{A9})$$

for  $\omega, \nu > 0$ , which gives

$$L(\nu; \vec{r}', \vec{r}) = \mathcal{P} \int_0^{\infty} d\omega \frac{\omega^2 g(\omega; \vec{r}', \vec{r})}{\omega^2 - \nu^2} - i\frac{\pi\nu}{2} g(\nu; \vec{r}', \vec{r}), \quad (\text{A10})$$

with

$$g(\omega; \vec{r}', \vec{r}) = \sum_{d^\omega} \vec{\varphi}_{\omega, d^\omega}(\vec{r}') \otimes \vec{\varphi}_{\omega, d^\omega}(\vec{r}). \quad (\text{A11})$$

The first term in (A10) can be rewritten as

$$\begin{aligned} \mathcal{P} \int d\omega \left( \frac{\omega^2}{\omega^2 - \nu^2} - 1 \right) g(\omega; \vec{r}', \vec{r}) + \int d\omega g(\omega; \vec{r}', \vec{r}) \\ = \mathcal{P} \int d\omega \frac{\nu^2 g(\omega; \vec{r}', \vec{r})}{\omega^2 - \nu^2} + \int d\omega g(\omega; \vec{r}', \vec{r}). \end{aligned}$$

Because of Eq. (A11) and the completeness of the eigenfunctions  $\vec{\varphi}_{\omega, d^\omega}$ , we have

$$\int d\omega g(\omega; \vec{r}', \vec{r}) = \delta_T(\vec{r} - \vec{r}'),$$

with the transverse  $\delta$  function defined as [34]

$$\begin{aligned} \delta_T^{ij}(\vec{r} - \vec{r}') &= \frac{2}{3} \delta_{ij} \delta(\vec{r} - \vec{r}') \\ &\quad - \frac{1}{4\pi |\vec{r} - \vec{r}'|^3} \left( \delta_{ij} - \frac{3(r_i - r'_i)(r_j - r'_j)}{|\vec{r} - \vec{r}'|^2} \right). \end{aligned} \quad (\text{A12})$$

Since  $\vec{r}'$  is defined only inside the medium whereas  $\vec{r}$  is taken in the exterior, the first term on the right-hand side vanishes. Hence we obtain

$$L(\nu; \vec{r}', \vec{r}) = \nu^2 \left[ \int d\omega \frac{g(\omega; \vec{r}', \vec{r})}{\omega^2 - \nu^2 + i0^+} + \frac{1}{c^2} F_{\text{nf}}(\nu; \vec{r}', \vec{r}) \right], \quad (\text{A13})$$

where

$$F_{\text{nf}}^{ij}(\vec{r}', \vec{r}) := \frac{-c^2}{4\pi\nu^2 |\vec{r} - \vec{r}'|^3} \left( \delta_{ij} - \frac{3(r_i - r'_i)(r_j - r'_j)}{|\vec{r} - \vec{r}'|^2} \right). \quad (\text{A14})$$

In (A13) we recognize the Green's function in vacuum

$$\begin{aligned} G_0^-(\nu; \vec{r}', \vec{r}) &= c^2 \int d\omega \frac{g(\omega; \vec{r}', \vec{r})}{\omega^2 - \nu^2 + i0^+} \\ &= c^2 \int d^3k \sum_{\sigma, \pi} \frac{\vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r}') \otimes \vec{\varphi}_{\vec{k}, \sigma, \pi}(\vec{r})}{\omega^2 - \nu^2 + i0^+}, \end{aligned} \quad (\text{A15})$$

and hence

$$L = \frac{\nu^2}{c^2} [G_0^- + F_{\text{nf}}]. \quad (\text{A16})$$

Introducing it back into Eq. (A7b) and using  $\epsilon_0\mu_0c^2 = 1$ , we obtain the electric field (31).

- [1] M. S. Tame, K. R. McEnery, S. K. Özdemir, J. Lee, S. A. Maier, and M. S. Kim, Quantum plasmonics, *Nat. Phys.* **9**, 329 (2013).
- [2] S. I. Bozhevolnyi and J. B. Khurgin, The case for quantum plasmonics, *Nat. Photon.* **11**, 398 (2017).
- [3] W. L. Barnes, A. Dereux, and T. W. Ebbesen, Surface plasmon subwavelength optics, *Nature (London)* **424**, 824 (2003).
- [4] E. Altewischer, M. P. van Exter, and J. P. Woerdman, Plasmon-assisted transmission of entangled photons, *Nature (London)* **418**, 304 (2002).
- [5] G. Di Martino, Y. Sonnefraud, M. S. Tame, S. Kéna-Cohen, F. Dieleman, K. Özdemir, M. S. Kim, and S. A. Maier, Observation of Quantum Interference in the Plasmonic Hong-Ou-Mandel Effect, *Phys. Rev. Appl.* **1**, 034004 (2014).

- [6] M.-C. Dheur, B. Vest, E. Devaux, A. Baron, J.-P. Hugonin, J.-J. Greffet, G. Messin, and F. Marquier, Remote preparation of single-plasmon states, *Phys. Rev. B* **96**, 045432 (2017).
- [7] L. Novotny and B. Hecht, *Principles of Nano-Optics* (Cambridge University Press, Cambridge, 2012).
- [8] T. Gruner and D.-G. Welsch, Correlation of radiation-field ground-state fluctuations in a dispersive and lossy dielectric, *Phys. Rev. A* **51**, 3246 (1995); Green-function approach to the radiation-field quantization for homogeneous and inhomogeneous Kramers-Kronig dielectrics, **53**, 1818 (1996); H. T. Dung, L. Knöll, and D.-G. Welsch, Three-dimensional quantization of the electromagnetic field in dispersive and absorbing inhomogeneous dielectrics, *ibid.* **57**, 3931 (1998); S. Scheel,

- L. Knöll, and D.-G. Welsch, QED commutation relations for inhomogeneous Kramers-Kronig dielectrics, *ibid.* **58**, 700 (1998); L. Knöll, S. Scheel, and D.-G. Welsch, in *Coherence and Statistics of Photons and Atoms*, edited by J. Peřina (Wiley, New York, 2001).
- [9] W. Vogel and D.-G. Welsch, *Quantum Optics* (Wiley-VCH, Weinheim, 2006).
- [10] S. Scheel and S. Y. Buhmann, Macroscopic quantum electrodynamics—Concepts and applications, *Acta Phys. Slovaca* **58**, 675 (2008).
- [11] S. Y. Buhmann, *Dispersion Forces I* (Springer, Berlin, 2012).
- [12] B. Huttner and S. M. Barnett, Quantization of the electromagnetic field in dielectrics, *Phys. Rev. A* **46**, 4306 (1992); S. M. Barnett, B. Huttner, and R. Loudon, Spontaneous Emission in Absorbing Dielectric Media, *Phys. Rev. Lett.* **68**, 3698 (1992).
- [13] L. G. Suttorp and M. Wubs, Field quantization in inhomogeneous absorptive dielectrics, *Phys. Rev. A* **70**, 013816 (2004).
- [14] L. G. Suttorp and A. J. van Wonderen, Fano diagonalization of a polariton model for an inhomogeneous absorptive dielectric, *Europhys. Lett.* **70**, 766 (2004).
- [15] T. G. Philbin, Canonical quantization of macroscopic electromagnetism, *New J. Phys.* **12**, 123008 (2010).
- [16] J. J. Hopfield, Theory of the contribution of excitons to the complex dielectric constant of crystals, *Phys. Rev.* **112**, 1555 (1958).
- [17] A. Drezet, Equivalence between the Hamiltonian and Langevin noise description of plasmon-polaritons in a dispersive and lossy inhomogeneous medium, *Phys. Rev. A* **96**, 033849 (2017).
- [18] O. Di Stefano, S. Savasta, and R. Girlanda, Mode expansion and photon operators in dispersive and absorbing dielectrics, *J. Mod. Opt.* **48**, 67 (2001).
- [19] A. Drezet, Quantizing polaritons in inhomogeneous dissipative systems, *Phys. Rev. A* **95**, 023831 (2017); Description of spontaneous photon emission and local density of states in the presence of a lossy polaritonic inhomogeneous medium, **95**, 043844 (2017).
- [20] M. Khanbekyan, L. Knöll, and D.-G. Welsch, Input-output relations at dispersing and absorbing planar multilayers for the quantized electromagnetic field containing evanescent components, *Phys. Rev. A* **67**, 063812 (2003).
- [21] S. de Bièvre, in *Contributions in Mathematical Physics: A Tribute to Gerard G. Emch*, edited by S. T. Ali and K. B. Sinha (Hindustan Book Agency, New Delhi, 2007), pp. 123–146.
- [22] S. de Bièvre, in *Large Coulomb Systems: Lecture Notes on Mathematical Aspects of QED*, edited by J. Dereziński and H. Siedentop, *Lecture Notes in Physics* Vol. 695 (Springer, Berlin, 2006), pp. 15–61.
- [23] M. Reed and B. Simon, *Methods of Modern Mathematical Physics* (Academic, San Diego, 1975), Vol. I.
- [24] M. Reed and B. Simon, *Methods of Modern Mathematical Physics* (Academic, San Diego, 1975), Vol. III.
- [25] K. O. Friedrichs, Über die spektralzerlegung eines integraloperators, *Math. Ann.* **115**, 249 (1937).
- [26] K. O. Friedrichs, On the perturbation of continuous spectra, *Commun. Pure. Appl. Math.* **1**, 361 (1948).
- [27] L. D. Faddeev, in *Fifty Years of Mathematical Physics: Selected Works of Ludwig Faddeev*, edited by M. Ge and A. J. Niemi, *World Scientific Series in 21st Century Mathematics* Vol. 2 (World Scientific, Singapore, 2016), pp.7–33.
- [28] D. R. Yafaev, *Mathematical Scattering Theory*, *Translations of Mathematical Monographs* Vol. 105 (American Mathematical Society, Providence, 1992).
- [29] U. Fano, Effects of configuration interaction on intensities and phase shifts, *Phys. Rev.* **124**, 1866 (1961).
- [30] D. Dzsotjan, A. S. Sørensen, and M. Fleischhauer, Quantum emitters coupled to surface plasmons of a nanowire: A Green's function approach, *Phys. Rev. B* **82**, 075427 (2010); J. Hakami, L. Wang, and M. S. Zubairy, Spectral properties of a strongly coupled quantum-dotmetal-nanoparticle system, *Phys. Rev. A* **89**, 053835 (2014); D. Dzsotjan, B. Rousseaux, H. R. Jauslin, G. C. des Francs, C. Couteau, and S. Guérin, Mode-selective quantization and multimodal effective models for spherically layered systems, *ibid.* **94**, 023818 (2016); A. Delga, J. Feist, J. Bravo-Abad, and F. J. Garcia-Vidal, Quantum Emitters Near a Metal Nanoparticle: Strong Coupling and Quenching, *Phys. Rev. Lett.* **112**, 253601 (2014); Theory of strong coupling between quantum emitters and localized surface plasmons, *J. Opt.* **16**, 114018 (2014); T. Hümmer, F. J. García-Vidal, L. Martín-Moreno, and D. Zueco, Weak and strong coupling regimes in plasmonic QED, *Phys. Rev. B* **87**, 115419 (2013).
- [31] K. Atkinson and W. Han, *Theoretical Numerical Analysis: A Functional Analysis Framework* (Springer, New York, 2009), pp. 473–549.
- [32] C. Sauvan, J. P. Hugonin, I. S. Maksymov, and P. Lalanne, Theory of the Spontaneous Optical Emission of Nanosize Photonic and Plasmon Resonators, *Phys. Rev. Lett.* **110**, 237401 (2013).
- [33] S. Franke, S. Hughes, M. K. Dezfouli, P. T. Kristensen, K. Busch, A. Knorr, and M. Richter, Quantization of Quasinormal Modes for Open Cavities and Plasmonic Cavity Quantum Electrodynamics, *Phys. Rev. Lett.* **122**, 213901 (2019).
- [34] A. M. Stewart, Longitudinal and transverse components of a vector field, *Sri Lankan J. Phys.* **12**, 33 (2011).