

Calculations of time-reversal-symmetry-violation sensitivity parameters based on analytic relativistic coupled-cluster gradient theory

Chaoqun Zhang, Xuechen Zheng, and Lan Cheng *Department of Chemistry, The Johns Hopkins University, Baltimore, Maryland 21218, USA*

(Received 8 April 2021; accepted 8 July 2021; published 26 July 2021)

We develop an analytic-gradient-based method for relativistic coupled-cluster calculations of effective electric field \mathcal{E}_{eff} with improved efficiency and robustness over the previous state of the art. The enhanced capability to calculate this time-reversal-symmetry-violation sensitivity parameter enables efficient screening of candidate molecules for the electron electric dipole moment (eEDM) search. As examples, the $|\mathcal{E}_{\text{eff}}|$ values of metal methoxides including BaOCH_3 , YbOCH_3 , and RaOCH_3 are shown to be as large as those of the corresponding fluorides and hydroxides, which supports the recent proposal of using these symmetric-top molecules to improve the sensitivity of eEDM measurements. The computational results also show that molecules containing late actinide elements, NoF , NoOH , LrO , and LrOH^+ , exhibit particularly large $|\mathcal{E}_{\text{eff}}|$ values of around 200 GV/cm.

DOI: [10.1103/PhysRevA.104.012814](https://doi.org/10.1103/PhysRevA.104.012814)

I. INTRODUCTION

The numerous discoveries at the Large Hadron Collider (LHC) of CERN included the observation of the Higgs boson, which completed the search for the fundamental particles in the Standard Model [1]. The exploitation of the 14-TeV collision energy of LHC has not observed fundamental particles associated with physics Beyond the Standard Model. One increasingly powerful method to search for new physics beyond the Standard Model is tabletop low-energy experiments based on precision spectroscopy of atoms and molecules [2–4]. In particular, the search for electron electric dipole moment (eEDM) through precision measurements of paramagnetic atoms and molecules has emerged as a promising route [2,4].

In paramagnetic atoms and molecules, the interaction between the eEDM (d_e) and the effective electric field $d_e\mathcal{E}_{\text{eff}}$ contributes to the atomic and molecular energy levels that are subject to spectroscopic interrogation. Although no nonzero eEDM has been reported, these measurements have set upper bounds to the eEDM value, which provides valuable information about the lower bounds for the energies of new fundamental particles. The sensitivity of the eEDM measurements is directly proportional to the effective electric field that the electrons experience. Paramagnetic atoms and molecules are sensitive to measurements of the eEDM because the $|\mathcal{E}_{\text{eff}}|$ values are far greater than applied laboratory electric fields [5]. Furthermore, since paramagnetic molecules exhibit orders of magnitude larger $|\mathcal{E}_{\text{eff}}|$ values than atoms [6], the recent work on diatomic molecules including YbF [7,8], ThO [9,10], and HfF^+ [11] has reduced the upper bound of eEDM by two orders of magnitude compared with a previous record set by the Thallium atom [12]. The present record of 1.1×10^{-29} e cm for the upper bound of eEDM obtained from

the measurements of ThO [10] corresponds to an energy scale of around 30 TeV for certain classes of new fundamental particles, which is beyond the reach of LHC. The ongoing experiments powered with new techniques to improve the precision [13–18] hold the promise to significantly improve the current limit. Furthermore, new schemes to use nearly degenerate vibrational and rotational states in laser-cooled polyatomic molecules have the potential to enhance the sensitivity by another several orders of magnitude [19–21].

The $|\mathcal{E}_{\text{eff}}|$ value, which represents the strength of an internal effective electric field in a paramagnetic atom or molecule, is not accessible to direct measurements. Electronic-structure calculations for this parameter [22–52] thus play an important role in the interpretation of experimental measurements and in the selection of candidate molecules. \mathcal{E}_{eff} is a “first-order property,” i.e., it corresponds to a first derivative of the electronic energy. Since \mathcal{E}_{eff} samples the electron density in the core region, it requires accurate treatments of both relativistic and electron-correlation effects to obtain accurate values. Relativistic Hartree-Fock (HF) and density-functional theory (DFT) calculations have proven useful in providing a good estimate of \mathcal{E}_{eff} [23–26,50], while accurate calculations of \mathcal{E}_{eff} have relied on wave-function-based electron-correlation methods including relativistic coupled-cluster (CC) methods [30,32,39,45] and multireference configuration interaction (CI) or CC methods [33,44]. The wave-function-based electron-correlation calculations to date have used unrelaxed formulations [30,32,33,39] or numerical differentiation of electronic energies [41,45,46]. The unrelaxed formulations involve an approximation of neglecting orbital-relaxation contributions. They also have a practical limitation of not being compatible with the standard noniterative treatment of triple excitations in the CC singles and doubles augmented with a noniterative triples [CCSD(T)] method [53,54]. Numerical differentiation can faithfully reproduce energy derivatives. However, the numerical-differentiation procedure for obtaining \mathcal{E}_{eff} is tedious due to the sensitivity of the numerical

*lcheng24@jhu.edu

results to the step size and the convergence of energy calculations [45,46]. These together with high computational cost of relativistic wave-function calculations render the calculations of \mathcal{E}_{eff} still a major challenge.

The tremendous efforts devoted to molecular structural optimization in quantum chemistry have established analytic gradients [55–59] as the standard tool for the calculations of first-order molecular properties. For example, a single analytic CC gradient calculation, which is two to three times as costly as a corresponding energy calculation, provides all first-order properties [58]. An analytic-gradient-based scheme is not only by far more efficient than numerical differentiation of electronic energies, but is also convenient to use. We report the development of an analytic-gradient-based method for the calculations of \mathcal{E}_{eff} using the relativistic exact two-component (X2C) CC analytic-gradient theory, hereby combining the analytic X2C gradient theory [60,61] and the recent development [62] of analytic first derivatives [53,54,63,64] for CC singles doubles (CCSD) [65] and CCSD with a noniterative triples [CCSD(T)] [66] methods with spin-orbit coupling included at the orbital level. The development of the present analytic-gradient-based method aims to significantly improve the efficiency, robustness, and convenience for the calculations of \mathcal{E}_{eff} to enable rapid and reliable screening of candidate molecules for use in the eEDM measurements.

II. THEORY

Relativistic electronic-structure calculations of effective electric field \mathcal{E}_{eff} are based on the Dirac Hamiltonian,

$$\hat{H} = \hat{H}_0 + d_e \hat{V}_{\text{eff}}, \quad (1)$$

with \hat{H}_0 and $d_e \hat{V}_{\text{eff}}$ representing the one-electron Dirac operator and the interaction between eEDM and the effective electric field [67],

$$\hat{H}_0 = \begin{pmatrix} \hat{V} & c\vec{\sigma} \cdot \vec{p} \\ c\vec{\sigma} \cdot \vec{p} & \hat{V} - 2c^2 \end{pmatrix}, \quad \hat{V}_{\text{eff}} = -2ic\beta\gamma_5 \hat{p}^2, \quad (2)$$

in which c is the speed of light, $\vec{\sigma}$ is a vector of Pauli spin matrices, \vec{p} is the momentum operator, \hat{V} is the nuclear attraction potential, and β and γ_5 are Dirac matrices given by

$$\beta = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}, \quad \gamma_5 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}. \quad (3)$$

\mathcal{E}_{eff} corresponds to the first derivative of the electronic energy with respect to d_e ,

$$\mathcal{E}_{\text{eff}} = \left. \frac{\partial E}{\partial d_e} \right|_{d_e=0}. \quad (4)$$

Since \hat{V}_{eff} involves the second derivatives of the small component wave function, \mathcal{E}_{eff} samples the wave function in the core region and is sensitive to treatments of relativistic and electron-correlation effects. The wave functions of the electronic states used in eEDM measurements, e.g., the $X^2\Sigma$ state of YbF and the $^3\Delta_1$ states of ThO, HfF⁺, and ThF⁺, are dominated by a single electron configuration. CC methods [68,69] can provide accurate treatments for dynamic correlation and are the methods of choice here.

The present scheme for the calculations of \mathcal{E}_{eff} uses the recent implementation of analytic first derivatives for X2C CCSD and CCSD(T) methods [62] and also the atomic-orbital-based algorithms [70] to enhance the computational efficiency. The calculation of \mathcal{E}_{eff} using the X2C-CC analytic-gradient theory involves a simple contraction between the relaxed one-electron density matrix $D^{\text{X2C-CC}}$ and the corresponding property integrals $[V_{\text{eff}}]^{\text{X2C}}$,

$$\mathcal{E}_{\text{eff}} = \left. \frac{\partial E_{\text{X2C-CC}}}{\partial d_e} \right|_{d_e=0} = \sum_{pq} D_{pq}^{\text{X2C-CC}} [V_{\text{eff}}]_{pq}^{\text{X2C}}. \quad (5)$$

We refer the readers to Ref. [62] for the calculations of the relaxed one-electron density matrix $D^{\text{X2C-CC}}$. We base the calculation of $[V_{\text{eff}}]^{\text{X2C}}$ on the analytic X2C gradient theory [60,61]. The X2C theory [71–73] uses the matrix representation of the one-electron Dirac equation,

$$\begin{pmatrix} h^{\text{LL}} & h^{\text{LS}} \\ h^{\text{SL}} & h^{\text{SS}} \end{pmatrix} \begin{pmatrix} C^{\text{L}} \\ C^{\text{S}} \end{pmatrix} = \begin{pmatrix} S & 0 \\ 0 & \frac{T}{2c^2} \end{pmatrix} \begin{pmatrix} C^{\text{L}} \\ C^{\text{S}} \end{pmatrix} E, \quad (6)$$

in which C^{L} and C^{S} are large- and small-component orbital coefficients in kinetically balanced basis sets [74],

$$\phi_i^{\text{L}} = C_{\mu i}^{\text{L}} f_{\mu}, \quad \phi_i^{\text{S}} = C_{\mu i}^{\text{S}} \frac{\vec{\sigma} \cdot \vec{p}}{2c} f_{\mu}, \quad (7)$$

h^{LL} , h^{LS} , h^{SL} , h^{SS} are the large-large, large-small, small-large, and small-small blocks of the Hamiltonian matrix,

$$h_{\mu\nu}^{\text{LL}} = V_{\mu\nu}, \quad h_{\mu\nu}^{\text{LS}} = h_{\mu\nu}^{\text{SL}} = T_{\mu\nu}, \quad (8)$$

$$h_{\mu\nu}^{\text{SS}} = \frac{1}{4c^2} \langle f_{\mu} | \vec{\sigma} \cdot \vec{p} V \vec{\sigma} \cdot \vec{p} | f_{\nu} \rangle - T_{\mu\nu}, \quad (9)$$

and S , T , and V represent overlap, kinetic energy, and nuclear attraction potential matrices. This four-component Hamiltonian matrix is block diagonalized [71],

$$\begin{pmatrix} h^{\text{LL}} & h^{\text{LS}} \\ h^{\text{SL}} & h^{\text{SS}} \end{pmatrix} \rightarrow \begin{pmatrix} h_+ & 0 \\ 0 & h_- \end{pmatrix}, \quad (10)$$

to decouple electronic and positronic degrees of freedom. The electronic block,

$$h_+ = R^\dagger [h^{\text{LL}} + h^{\text{LS}} X + X^\dagger h^{\text{SL}} + X^\dagger h^{\text{SS}} X] R, \quad (11)$$

$$C^{\text{S}} = X C^{\text{L}}, \quad R = (\tilde{S}^{-1} S)^{1/2}, \quad \tilde{S} = S + \frac{1}{2c^2} X^\dagger T X, \quad (12)$$

is used together with the untransformed two-electron Coulomb interaction in the subsequent many-electron treatment. We obtain $[V_{\text{eff}}]^{\text{X2C}}$ by differentiating h_+ , using a procedure developed in Ref. [75]. The calculation of $[V_{\text{eff}}]^{\text{X2C}}$ involves the derivatives of h^{LS} and h^{SL} , since V_{eff} appears on the LS and SL blocks,

$$\frac{\partial h_{\mu\nu}^{\text{LS}}}{\partial d_e} = \frac{\partial h_{\nu\mu}^{\text{SL}*}}{\partial d_e} = \langle f_{\mu} | \begin{pmatrix} -2icp^2 & 0 \\ 0 & -2icp^2 \end{pmatrix} \frac{\vec{\sigma} \cdot \vec{p}}{2c} f_{\nu} \rangle, \quad (13)$$

and the derivatives of the X and R matrices.

III. COMPUTATIONAL DETAILS

We have implemented the present analytic-gradient-based method for the calculations of effective electric field in the

CFOUR program package [76,77] and used it for all the calculations presented here. Our calculations used experimental equilibrium structures for HfF⁺ [78], TaN [79], ThF⁺ [80], ThO [81], BaF [82], BaOH [83], YbF [84], YbOH [85], a relativistic Fock-space CCSD bond length for HgF [86] to enable direct comparison with Ref. [38], and the spin-free X2C [73,75,87] CCSD(T) [66]/cc-pVTZ [88–91] structures for the other molecules, which are documented in detail in the Supplemental Material [92].

Unless otherwise stated, the calculations employed the X2C Hamiltonian [71,72,93] with the atomic mean-field [94] (AMF) spin-orbit integrals (the X2CAMF scheme) [95] and Gaussian nuclear model [96], and included the Gaunt term in the AMF approach. We used large uncontracted basis sets and correlated valence and a few additional shells of inner electrons. Calculations correlating all electrons were also carried out for a subset of the molecules to study inner-shell correlation effects. We mention that the basis-set errors beyond the uncontracted triple-zeta basis sets have been shown to be small [33,35,45,46]. The present calculations employed the uncontracted ANO-RCC basis sets [97–100] for heavy atoms, which are of augmented quadruple-zeta quality for W, Ta, Ba, Ra, Hg, and Lu and of augmented triple-zeta quality for Yb, Hf, and Th, except that we used the uncontracted cc-pVTZ basis sets [90] for No and Lr. For light elements, the uncontracted cc-pVTZ basis sets [88] were used in the calculations of RaOH, YbOCH₃, BaOCH₃, RaOCH₃, LuOH⁺, LrO, LrOH⁺, NoF, NoOH, while the uncontracted aug-cc-pVTZ basis sets were used for the other molecules. Although we used aug-cc-pVTZ basis sets in many calculations, the contributions from diffuse functions to $|\mathcal{E}_{\text{eff}}|$'s turned out to be negligible, e.g., they amount to less than 0.2% for BaOH, YbOH, and RaOH.

IV. RESULTS AND DISCUSSIONS

A widely used approach to assess the accuracy of the computed $|\mathcal{E}_{\text{eff}}|$ values is to compare other properties computed using the same method with the corresponding measured values. We have shown that the X2CAMF-CCSD(T) method provides accurate electric dipole moments and nuclear quadrupole-coupling constants for heavy-element containing molecules [62,95], and we expect it to provide accurate $|\mathcal{E}_{\text{eff}}|$ values. Taking advantage of the present extensive benchmark set, we also take complementary approaches to analyze the accuracy of the X2CAMF-CC calculations, by comparing the results with available calculations and by analyzing the remaining errors in the treatments of relativistic, electron-correlation, and nuclear-model effects.

The X2CAMF-CCSD(T) $|\mathcal{E}_{\text{eff}}|$ results agree well with the previous relativistic CC and CI calculations for all the molecules in Table I, with the discrepancies amounting to up to several percent of the total values. It has been reported that the X²Σ (4*f*¹⁴6*s*¹) states of YbF and YbOH can be perturbed by the 4*f*¹³6*s*² configuration [101–103], because the errors in the treatments of electron-correlation and basis-set effects both lead to underestimation of the relative energies between the 4*f*¹³6*s*² states and the X²Σ states [104]. It thus is necessary to use large basis sets to obtain accurate electron-correlation contributions [46,47]. We

TABLE I. The $|\mathcal{E}_{\text{eff}}|$ values (GV/cm) from the X2CAMF-HF, CCSD, and CCSD(T) calculations using the analytic energy-gradient theory. “fc ele” refers to the number of core electrons kept frozen together with virtual orbitals higher than 100 Hartree in the CC calculations.

	fc ele	HF	CCSD	CCSD(T)	Literature
HfF ⁺	48	25.8	22.8	22.5	22.5[43]/22.7[44]
WC	48	72.8	43.5	37.9	36[31]
TaN	48	59.3	39.6	34.8	34.9[36]/36.0[40]
ThF ⁺	62	42.7	36.6	36.6	37.3[35]/35.2[37]
ThO	62	98.2	83.3	79.8	75.2[33]/79.9[41]
BaF	30	6.5	6.4	6.3	6.52[51]/
BaOH	30	6.5	6.5	6.4	6.4[46]/6.2[49]
YbF	48	22.9	23.5	23.7	23.1[32]
YbOH	48	22.9	23.7	24.0	23.4[46]/23.8[47]/17.7[49]
RaF	48	55.0	54.9	54.2	52.9[34]/52.5[42]/50.9[50]
HgF	48	132.0	118.7	113.0	115.42[38]/116.37[45]

note that the present Kramers unrestricted calculations yield smaller electron-correlation contributions for YbF and YbOH than in previous studies [32,47], while the total CCSD values agree well with those in Refs. [32,46,47]. The present CCSD result for YbOH also agrees well with the Fock-space CCSD value in Ref. [46]. It has been shown that the CCSD(T) noniterative triples corrections to first-order properties of Yb-containing are not accurate representation of triples corrections [103]. The CCSD(T) triples corrections for YbF, YbOH, and YbOCH₃ in Tables I and II thus only serve as rough estimates for triples contributions in the present discussion.

Table II summarizes calculations for 10 molecules, for which no wave-function-based electron-correlation calculations have been reported. The present results for RaOH, LuO, NoF, and LrO are consistent with recent relativistic HF and DFT calculations [48,49]. The metal methoxides, BaOCH₃,

TABLE II. The $|\mathcal{E}_{\text{eff}}|$ values (GV/cm) from the X2CAMF-HF, CCSD, and CCSD(T) calculations using the analytic energy-gradient theory. “fc ele” refers to the number of core electrons kept frozen together with virtual orbitals higher than 1000 Hartree in the CC calculations.

	fc ele	HF	CCSD	CCSD(T)	Literature ^a
BaOCH ₃	28	6.5	6.4	6.3	–
YbOCH ₃	50	22.9	23.6	24.0	–
RaOCH ₃	46	55.0	55.0	54.2	–
RaOH	48	54.9	55.2	54.5	56.9 ^b , 52.3 ^c
LuO	28	36.1	33.7	32.4	37.4 ^b , 32.1 ^c
LuOH ⁺	28	32.3	29.8	29.2	–
NoF	70	185.4	192.4	191.9	200 ^b , 200 ^c
NoOH	70	185.2	192.4	191.7	–
LrO	70	303.3	263.9	246.5	323 ^b , 250 ^c
LrOH ⁺	70	268.6	259.5	255.1	–

^aZeroth-order regular approximation (ZORA) HF and DFT values [48,49].

^bHF values.

^cDFT values.

TABLE III. The errors of the X2CAMF scheme, the core-correlation contributions, and the finite nuclear size effects for $|\mathcal{E}_{\text{eff}}|$ (GV/cm). The percentages of the total values are enclosed in the parentheses.

	X2CAMF error ^a HF	Core correlation ^b CCSD	Finite nuclear size ^c CCSD
BaF	-0.03 (-0.5%)	0.01 (0.2%)	-0.04 (-0.5%)
YbF	-0.01 (-0.0%)	-0.06 (-0.2%)	-0.38 (-1.6%)
HfF ⁺	0.01 (0.0%)	0.40 (1.7%)	-0.47 (-2.0%)
RaF	0.14 (0.3%)	0.17 (0.3%)	-3.67 (-6.6%)
ThO	0.31 (0.3%)	0.34 (0.4%)	-6.27 (-7.4%)
ThF ⁺	0.50 (1.2%)	0.56 (1.5%)	-3.10 (-8.2%)
NoF	0.95 (0.5%)	-0.30 (-0.2%)	-35.04 (-18.0%)

^aThe differences between the Dirac-Coulomb-Gaunt and X2CAMF results.

^bThe differences between the correlation of all electrons and virtual orbitals below 10 000 Hartree and those in Tables I and II.

^cThe differences between using the Gaussian and pointlike nuclear models.

YbOCH₃, and RaOCH₃, possess $|\mathcal{E}_{\text{eff}}|$ values similar to those of the corresponding fluorides and hydroxides. For example, the X2CAMF-CCSD $|\mathcal{E}_{\text{eff}}|$ value for YbOCH₃ amounts to 23.6 GV/cm, very similar to the values of 23.5 and 23.7 GV/cm for YbF and YbOH. This is consistent with the chemical intuition that the unpaired electron in YbOCH₃ is localized at the Yb atom. These computational results support the recent proposal of using the nearly degenerate rotational states of these symmetric-top molecules to improve the sensitivity of eEDM measurements [19]. The $|\mathcal{E}_{\text{eff}}|$ values of 32.4 GV/cm and 29.0 GV/cm for LuO and LuOH⁺ are a little larger than those of YbF and YbOH and are similar to that of ThF⁺. The four small molecules containing late actinide elements, NoF, NoOH, LrO, and LrOH⁺, exhibit particularly large $|\mathcal{E}_{\text{eff}}|$ values of 191.9, 191.7, 246.5, and 255.1 GV/cm, respectively, because of the relativistic enhancement in the presence of these very heavy atoms.

The X2CAMF scheme uses the untransformed two-electron Coulomb interaction together with an AMF approximation to the two-electron spin-orbit integrals and the Gaunt term. This eliminates molecular relativistic atomic orbital two-electron integrals [95] and enables efficient electron-correlation calculations using atomic-orbital-based algorithms [70]. As shown in Table III, the errors of the X2CAMF values compared with the four-component results are small across the periodic table, e.g., the error amounts to -0.5% for BaF, 0.3% for RaF, 1.2% for ThF⁺, and 0.5% for NoF. We mention that the Gaunt-term contributions amount to around 1% for all the molecules studied here, except that it is around 3% for HfF⁺. We expect the remaining relativistic contributions from the Gauge term and quantum electrodynamics to be smaller than the Gaunt-term contributions.

The differences between the CC and HF results in Tables I and II represent the electron-correlation contributions. The electron-correlation contributions amount to more than half of the total values for TaN and WC. In contrast, they are less than 2% for MF, MOH, MOCH₃, with M = Ba, Yb, Ra. The rest of the molecules receive moderate yet important contributions from electron correlation, ranging from 5% to

30%. The magnitude of triples contributions are significantly smaller than singles and doubles contributions for most of the molecules studied here, except for some molecules exhibiting very small total correlation contributions such as BaOH, RaF, RaOH, and RaOCH₃. WC and TaN exhibit relatively large triples contributions of around 10%. We expect the high-level correlation contributions to be smaller than triples contributions. Finally, as shown in Table III, the correlation of the inner-shell core electrons makes minor contributions, amounting to up to a few percent.

The importance of the nuclear model increases rapidly for heavier elements, as demonstrated in Table III. While the difference between the Gaussian model and pointlike model is only -0.5% for BaF, it amounts to -7% for ThO and -18% for NoF. Since the Gaussian model is more realistic than the pointlike model, we expect the errors of the Gaussian model to be much smaller than the difference between the Gaussian and pointlike models.

Taking these error analyses into account, we conclude that the computational results in Tables I and II are accurate to within 10% except that we assign a 25% error estimate for NoF, NoOH, LrO, and LrOH⁺. Further improvement of the error estimate requires a study of the sensitivity of computed results to the sizes and function form of the finite nuclear model [49]. The current conservative error estimate still supports that the molecules containing late actinide elements possess extraordinarily large $|\mathcal{E}_{\text{eff}}|$ values. Therefore, given the structure of a molecule, the present analytic-gradient-based scheme only needs a single X2CAMF-CCSD(T) analytic-gradient calculation, which is of black-box nature, to provide a \mathcal{E}_{eff} value accurate enough for the initial screening of candidate molecules for eEDM measurements. One may improve the results by correcting the errors of the X2CAMF scheme and by including inner-shell correlation (Table III), e.g., the best values for YbF, HfF⁺, ThO, and ThF⁺ from the present calculations are obtained as 23.4, 22.9, 80.4, and 37.6 GV/cm, respectively, by combining the small corrections in Table III with the CCSD(T) results in Table I for all these molecules except that we adopt the CCSD result for YbF.

V. SUMMARY AND OUTLOOK

We report the development of an analytic-gradient-based method for the X2C-CCSD(T) calculations of effective electric field in paramagnetic molecules. Extensive benchmark calculations demonstrate the efficiency and accuracy of the present scheme. The extension of the present method to the calculations of other symmetry-violating parameters, e.g., the parameter associated with the measurements of nuclear magnetic quadrupole moment [22,36,40,105–107] is straightforward, by contracting the reduced density matrix with the corresponding property integrals. The present method thus provides significantly enhanced capabilities for the calculations of symmetry-violation sensitivity parameters in molecules. It will enable convenient, efficient, and reliable calculations of these parameters to help engineer new molecules suitable for the search of new physics via precision measurement. The X2C-CC analytic-gradient technique can also be extended to four-component theory as discussed in

Ref. [62] to enhance the efficiency of four-component CC calculations of these parameters.

ACKNOWLEDGMENTS

This work is supported by the National Science Foundation, under Grant No. PHY-2011794. L.C. is grateful to

Nicholas Hutzler (Pasadena), Phelan Yu (Pasadena), Kia Ng (Boulder), Anastasia Borschevsky (Groningen), Konstantin Gaul (Frankfurt), and Robert Berger (Frankfurt) for helpful comments on the manuscript and for stimulating discussions. All calculations were carried out at the Maryland Advanced Research Computing Center (MARCC).

C.Z. and X.Z. contributed equally to this work.

-
- [1] G. Aad *et al.*, *Phys. Lett. B* **716**, 1 (2012).
- [2] D. DeMille, J. M. Doyle, and A. O. Sushkov, *Science* **357**, 990 (2017).
- [3] M. S. Safronova, D. Budker, D. DeMille, D. F. J. Kimball, A. Derevianko, and C. W. Clark, *Rev. Mod. Phys.* **90**, 025008 (2018).
- [4] W. B. Cairncross and J. Ye, *Nat. Rev. Phys.* **1**, 510 (2019).
- [5] P. G. H. Sandars, *Phys. Lett.* **14**, 194 (1965).
- [6] P. G. H. Sandars, *Phys. Rev. Lett.* **19**, 1396 (1967).
- [7] J. J. Hudson, B. E. Sauer, M. R. Tarbutt, and E. A. Hinds, *Phys. Rev. Lett.* **89**, 023003 (2002).
- [8] J. J. Hudson, D. M. Kara, I. J. Smallman, B. E. Sauer, M. R. Tarbutt, and E. A. Hinds, *Nature (London)* **473**, 493 (2011).
- [9] J. Baron, W. C. Campbell, D. DeMille, J. M. Doyle, G. Gabrielse, Y. V. Gurevich, P. W. Hess, N. R. Hutzler, E. Kirilov, I. Kozryyev, B. R. O’Leary, C. D. Panda, M. F. Parsons, E. S. Petrik, B. Spaun, A. C. Vutha, and A. D. West, *Science* **343**, 269 (2014).
- [10] V. Andreev, D. G. Ang, D. DeMille, J. M. Doyle, G. Gabrielse, J. Haefner, N. R. Hutzler, Z. Lasner, C. Meisenhelder, B. R. O’Leary, C. D. Panda, A. D. West, E. P. West, X. Wu, and A. Collaboration, *Nature (London)* **562**, 355 (2018).
- [11] W. B. Cairncross, D. N. Gresh, M. Grau, K. C. Cossel, T. S. Roussy, Y. Ni, Y. Zhou, J. Ye, and E. A. Cornell, *Phys. Rev. Lett.* **119**, 153001 (2017).
- [12] B. C. Regan, E. D. Commins, C. J. Schmidt, and D. DeMille, *Phys. Rev. Lett.* **88**, 071805 (2002).
- [13] J. Lim, J. R. Almond, M. A. Trigatzis, J. A. Devlin, N. J. Fitch, B. E. Sauer, M. R. Tarbutt, and E. A. Hinds, *Phys. Rev. Lett.* **120**, 123201 (2018).
- [14] C. D. Panda, C. Meisenhelder, M. Verma, D. G. Ang, J. Chow, Z. Lasner, X. Wu, D. DeMille, J. M. Doyle, and G. Gabrielse, *J. Phys. B At. Mol. Opt. Phys.* **52**, 235003 (2019).
- [15] X. Wu, Z. Han, J. Chow, D. G. Ang, C. Meisenhelder, C. D. Panda, E. P. West, G. Gabrielse, J. M. Doyle, and D. DeMille, *New J. Phys.* **22**, 023013 (2020).
- [16] Y. Zhou, Y. Shagam, W. B. Cairncross, K. B. Ng, T. S. Roussy, T. Grogan, K. Boyce, A. Vigil, M. Pettine, T. Zelevinsky, J. Ye, and E. A. Cornell, *Phys. Rev. Lett.* **124**, 053201 (2020).
- [17] M. Verma, A. M. Jayich, and A. C. Vutha, *Phys. Rev. Lett.* **125**, 153201 (2020).
- [18] C. J. Ho, J. A. Devlin, I. M. Rabey, P. Yzombard, J. Lim, S. C. Wright, N. J. Fitch, E. A. Hinds, M. R. Tarbutt, and B. E. Sauer, *New J. Phys.* **22**, 053031 (2020).
- [19] I. Kozryyev and N. R. Hutzler, *Phys. Rev. Lett.* **119**, 133002 (2017).
- [20] B. L. Augenbraun, Z. D. Lasner, A. Frenett, H. Sawaoka, C. Miller, T. C. Steimle, and J. M. Doyle, *New J. Phys.* **22**, 022003 (2020).
- [21] N. R. Hutzler, *Quantum Sci. Technol.* **5**, 044011 (2020).
- [22] M. Kozlov, V. Fomichev, Y. Y. Dmitriev, L. Labzovsky, and A. Titov, *J. Phys. B At. Mol. Opt. Phys.* **20**, 4939 (1987).
- [23] Y. Y. Dmitriev, Y. G. Khait, M. Kozlov, L. Labzovsky, A. Mitrushenkov, A. Shtoff, and A. Titov, *Phys. Lett. A* **167**, 280 (1992).
- [24] M. G. Kozlov, A. V. Titov, N. S. Mosyagin, and P. V. Souchko, *Phys. Rev. A* **56**, R3326(R) (1997).
- [25] F. A. Parpia, *J. Phys. B At. Mol. Opt. Phys.* **31**, 1409 (1998).
- [26] H. Quiney, H. Skaane, and I. Grant, *J. Phys. B At. Mol. Opt. Phys.* **31**, L85 (1998).
- [27] B. Ravaine, S. G. Porsev, and A. Derevianko, *Phys. Rev. Lett.* **94**, 013001 (2005).
- [28] E. R. Meyer, J. L. Bohn, and M. P. Deskevich, *Phys. Rev. A* **73**, 062108 (2006).
- [29] E. R. Meyer and J. L. Bohn, *Phys. Rev. A* **78**, 010502(R) (2008).
- [30] L. V. Skripnikov, A. N. Petrov, and A. V. Titov, *J. Chem. Phys.* **139**, 221103 (2013).
- [31] J. Lee, J. Chen, L. V. Skripnikov, A. N. Petrov, A. V. Titov, N. S. Mosyagin, and A. E. Leanhardt, *Phys. Rev. A* **87**, 022516 (2013).
- [32] M. Abe, G. Gopakumar, M. Hada, B. P. Das, H. Tatewaki, and D. Mukherjee, *Phys. Rev. A* **90**, 022501 (2014).
- [33] T. Fleig and M. K. Nayak, *J. Mol. Spectrosc.* **300**, 16 (2014).
- [34] A. D. Kudashov, A. N. Petrov, L. V. Skripnikov, N. S. Mosyagin, T. A. Isaev, R. Berger, and A. V. Titov, *Phys. Rev. A* **90**, 052513 (2014).
- [35] L. V. Skripnikov and A. V. Titov, *Phys. Rev. A* **91**, 042504 (2015).
- [36] L. V. Skripnikov, A. N. Petrov, N. S. Mosyagin, A. V. Titov, and V. V. Flambaum, *Phys. Rev. A* **92**, 012521 (2015).
- [37] M. Denis, M. S. Nørby, H. J. A. Jensen, A. S. P. Gomes, M. K. Nayak, S. Knecht, and T. Fleig, *New J. Phys.* **17**, 043005 (2015).
- [38] V. S. Prasanna, A. C. Vutha, M. Abe, and B. P. Das, *Phys. Rev. Lett.* **114**, 183001 (2015).
- [39] S. Sasmal, H. Pathak, M. K. Nayak, N. Vaval, and S. Pal, *J. Chem. Phys.* **143**, 084119 (2015).
- [40] T. Fleig, M. K. Nayak, and M. G. Kozlov, *Phys. Rev. A* **93**, 012505 (2016).
- [41] L. V. Skripnikov, *J. Chem. Phys.* **145**, 214301 (2016).
- [42] S. Sasmal, H. Pathak, M. K. Nayak, N. Vaval, and S. Pal, *Phys. Rev. A* **93**, 062506 (2016).
- [43] L. V. Skripnikov, *J. Chem. Phys.* **147**, 021101 (2017).
- [44] T. Fleig, *Phys. Rev. A* **95**, 022504 (2017).
- [45] M. Abe, V. S. Prasanna, and B. P. Das, *Phys. Rev. A* **97**, 032515 (2018).

- [46] M. Denis, P. A. B. Haase, R. G. E. Timmermans, E. Eliav, N. R. Hutzler, and A. Borschevsky, *Phys. Rev. A* **99**, 042512 (2019).
- [47] V. S. Prasanna, N. Shitara, A. Sakurai, M. Abe, and B. P. Das, *Phys. Rev. A* **99**, 062502 (2019).
- [48] K. Gaul, S. Marquardt, T. Isaev, and R. Berger, *Phys. Rev. A* **99**, 032509 (2019).
- [49] K. Gaul and R. Berger, *Phys. Rev. A* **101**, 012508 (2020).
- [50] K. Gaul and R. Berger, *J. Chem. Phys.* **152**, 044101 (2020).
- [51] K. Talukdar, M. K. Nayak, N. Vaval, and S. Pal, *J. Phys. B At. Mol. Opt. Phys.* **53**, 135102 (2020).
- [52] R. Mitra, V. S. Prasanna, B. K. Sahoo, N. R. Hutzler, M. Abe, and B. P. Das, *Atoms* **9**, 7 (2021).
- [53] G. E. Scuseria, *J. Chem. Phys.* **94**, 442 (1991).
- [54] J. D. Watts, J. Gauss, and R. J. Bartlett, *J. Chem. Phys.* **98**, 8718 (1993).
- [55] P. Pulay, *Mol. Phys.* **17**, 197 (1969).
- [56] J. A. Pople, R. Krishnan, H. B. Schlegel, and J. S. Binkley, *Int. J. Quantum Chem. Symp.* **13**, 225 (1979).
- [57] T. Helgaker and P. Jørgensen, *Adv. Quant. Chem.* **19**, 188 (1988).
- [58] J. F. Stanton and J. Gauss, *Int. Rev. Phys. Chem.* **19**, 61 (2000).
- [59] P. Pulay, *WIREs Comp. Mol. Sci.* **4**, 169 (2014).
- [60] L. Cheng, S. Stopkowitz, and J. Gauss, *Int. J. Quantum Chem.* **114**, 1108 (2014).
- [61] M. Filatov, W. Zou, and D. Cremer, *Int. J. Quantum Chem.* **114**, 993 (2014).
- [62] J. Liu, X. Zheng, A. Asthana, C. Zhang, and L. Cheng, *J. Chem. Phys.* **154**, 064110 (2021).
- [63] A. C. Scheiner, G. E. Scuseria, J. E. Rice, T. J. Lee, and H. F. Schaefer III, *J. Chem. Phys.* **87**, 5361 (1987).
- [64] J. Gauss, J. F. Stanton, and R. J. Bartlett, *J. Chem. Phys.* **95**, 2623 (1991).
- [65] G. D. Purvis III and R. J. Bartlett, *J. Chem. Phys.* **76**, 1910 (1982).
- [66] K. Raghavachari, G. W. Trucks, J. A. Pople, and M. Head-Gordon, *Chem. Phys. Lett.* **157**, 479 (1989).
- [67] W. R. Johnson, D. S. Guo, M. Idrees, and J. Sapirstein, *Phys. Rev. A* **34**, 1043 (1986).
- [68] R. J. Bartlett and M. Musiał, *Rev. Mod. Phys.* **79**, 291 (2007).
- [69] T. D. Crawford and H. F. Schaefer III, *Rev. Comp. Chem.* **14**, 33 (2000).
- [70] J. Liu, Y. Shen, A. Asthana, and L. Cheng, *J. Chem. Phys.* **148**, 034106 (2018).
- [71] K. G. Dyall, *J. Chem. Phys.* **106**, 9618 (1997).
- [72] W. Kutzelnigg and W. Liu, *J. Chem. Phys.* **123**, 241102 (2005).
- [73] W. Liu and D. Peng, *J. Chem. Phys.* **131**, 1 (2009).
- [74] R. E. Stanton and S. Havriliak, *J. Chem. Phys.* **81**, 1910 (1984).
- [75] L. Cheng and J. Gauss, *J. Chem. Phys.* **135**, 084114 (2011).
- [76] J. F. Stanton, J. Gauss, L. Cheng, M. E. Harding, D. A. Matthews, and P. G. Szalay, "Computer code CFour: Coupled-cluster techniques for computational chemistry, a quantum-chemical program package", with contributions from A. A. Auer, R. J. Bartlett, U. Benedikt, C. Berger, D. E. Bernholdt, S. Blaschke, Y. J. Bomble, S. Burger, O. Christiansen, D. Datta, F. Engel, R. Faber, J. Greiner, M. Heckert, O. Heun, M. Hilgenberg, C. Huber, T.-C. Jagau, D. Jonsson, J. Jusélius, T. Kirsch, K. Klein, G. M. Kopper, W. J. Lauderdale, F. Lipparini, T. Metzroth, L. A. Mück, D. P. O'Neill, T. Nottoli, D. R. Price, E. Prochnow, C. Puzzarini, K. Ruud, F. Schiffmann, W. Schwalbach, C. Simmons, S. Stopkowitz, A. Tajti, J. Vázquez, F. Wang, J. D. Watts and the integral packages MOLEKULE (J. Almlöf and P. R. Taylor), PROPS (P. R. Taylor), ABACUS (T. Helgaker, H. J. Aa. Jensen, P. Jørgensen, and J. Olsen), and ECP routines by A. V. Mitin and C. van Wüllen. For the current version, see <http://www.cfour.de>.
- [77] D. A. Matthews, L. Cheng, M. E. Harding, F. Lipparini, S. Stopkowitz, T.-C. Jagau, P. G. Szalay, J. Gauss, and J. F. Stanton, *J. Chem. Phys.* **152**, 214108 (2020).
- [78] K. C. Cossel, D. N. Gresh, L. C. Sinclair, T. Coffey, L. V. Skripnikov, A. N. Petrov, N. S. Mosyagin, A. V. Titov, R. W. Field, E. R. Meyer, E. A. Cornell, and J. Ye, *Chem. Phys. Lett.* **546**, 1 (2012).
- [79] R. S. Ram, J. Liévin, and P. F. Bernath, *J. Mol. Spectrosc.* **215**, 275 (2002).
- [80] D. N. Gresh, K. C. Cossel, Y. Zhou, J. Ye, and E. A. Cornell, *J. Mol. Spectrosc.* **319**, 1 (2016).
- [81] G. Edvinsson and A. Lagerqvist, *Phys. Scr.* **30**, 309 (1984).
- [82] L. B. Knight, W. C. Easley, W. Weltner, and M. Wilson, *J. Chem. Phys.* **54**, 322 (1971).
- [83] S. Kinsey-Nielsen, C. R. Brazier, and P. F. Bernath, *J. Chem. Phys.* **84**, 698 (1986).
- [84] C. S. Dickinson, J. A. Coxon, N. R. Walker, and M. C. L. Gerry, *J. Chem. Phys.* **115**, 6979 (2001).
- [85] S. Nakhate, T. C. Steimle, N. H. Pilgram, and N. R. Hutzler, *Chem. Phys. Lett.* **715**, 105 (2019).
- [86] S. Knecht, S. Fux, R. van Meer, L. Visscher, M. Reiher, and T. Saue, *Theor. Chem. Acc.* **129**, 631 (2011).
- [87] K. G. Dyall, *J. Chem. Phys.* **115**, 9136 (2001).
- [88] T. H. Dunning Jr., *J. Chem. Phys.* **90**, 1007 (1989).
- [89] Q. Lu and K. A. Peterson, *J. Chem. Phys.* **145**, 054111 (2016).
- [90] R. Feng and K. A. Peterson, *J. Chem. Phys.* **147**, 084108 (2017).
- [91] J. G. Hill and K. A. Peterson, *J. Chem. Phys.* **147**, 244106 (2017).
- [92] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevA.104.012814> for a summary of the structural parameters used in the present calculations.
- [93] M. Iliaš and T. Saue, *J. Chem. Phys.* **126**, 064102 (2007).
- [94] B. A. Heß, C. M. Marian, U. Wahlgren, and O. Gropen, *Chem. Phys. Lett.* **251**, 365 (1996).
- [95] J. Liu and L. Cheng, *J. Chem. Phys.* **148**, 144108 (2018).
- [96] L. Visscher and K. G. Dyall, *At. Data Nucl. Data Tables* **67**, 207 (1997).
- [97] K. Fægri, *Theor. Chem. Acc.* **105**, 252 (2001).
- [98] B. O. Roos, R. Lindh, P.-Å. Malmqvist, V. Veryazov, and P.-O. Widmark, *J. Phys. Chem. A* **109**, 6575 (2005).
- [99] B. O. Roos, R. Lindh, P.-Å. Malmqvist, V. Veryazov, and P.-O. Widmark, *Chem. Phys. Lett.* **409**, 295 (2005).
- [100] B. O. Roos, R. Lindh, P.-Å. Malmqvist, V. Veryazov, P.-O. Widmark, and A. C. Borin, *J. Phys. Chem. A* **112**, 11431 (2008).
- [101] M. G. Kozlov, *J. Phys.* **30**, L607 (1997).
- [102] A. S. P. Gomes, K. G. Dyall, and L. Visscher, *Theor. Chem. Acc.* **127**, 369 (2010).

- [103] L. F. Pašteka, R. J. Mawhorter, and P. Schwerdtfeger, *Mol. Phys.* **114**, 1110 (2016).
- [104] M. Dolg, H. Stoll, and H. Preuss, *Chem. Phys.* **165**, 21 (1992).
- [105] O. Sushkov, V. Flambaum, and I. Khriplovich, *Zh. Eksp. Teor. Fiz.* **87**, 1521 (1984).
- [106] D. Maison, L. Skripnikov, V. Flambaum, and M. Grau, *J. Chem. Phys.* **153**, 224302 (2020).
- [107] M. Denis, Y. Hao, E. Eliav, N. R. Hutzler, M. K. Nayak, R. G. Timmermans, and A. Borschevsky, *J. Chem. Phys.* **152**, 084303 (2020).