





Quantum non-Markovianity elusive to interventions

Daniel Burgarth ^{1,*} Paolo Facchi ^{2,3,†} Davide Longro ^{2,3,‡} and Kavan Modi ^{4,§}

¹*Department of Physics and Astronomy, Center of Engineered Quantum Systems, Macquarie University, Sydney, New South Wales 2109, Australia*

²*Dipartimento di Fisica and MECENAS, Università di Bari, I-70126 Bari, Italy*

³*INFN, Sezione di Bari, I-70126 Bari, Italy*

⁴*School of Physics and Astronomy, Monash University, Clayton, Victoria 3800, Australia*



(Received 25 August 2021; accepted 3 November 2021; published 19 November 2021)

The non-Markovian nature of open quantum dynamics lies in the structure of multitime correlations, which are accessible by means of interventions. Here, by examining multitime correlations, we show that it is possible to engineer non-Markovian systems with only long-term memory but seemingly no short-term memory, so that their non-Markovianity is completely nondetectable by any interventions up to an arbitrarily large time. Our results raise the question about the assessability of non-Markovianity: in principle, non-Markovian effects that are perfectly elusive to interventions may emerge at much later times.

DOI: [10.1103/PhysRevA.104.L050404](https://doi.org/10.1103/PhysRevA.104.L050404)

Introduction. A key obstacle that lies in the path to quantum information processing is noise [1]. The conventional models for quantum noise, responsible for decoherence of qubits, make many simplifying assumptions. One of the key assumptions is that the noise is memoryless or Markovian [2]; this is widely known to be false, and an immense effort in understanding non-Markovian noise, both in general and in quantum information processors, has been initiated [3–6]. While non-Markovian noise is more complex than Markovian, it is *not* more detrimental. In fact, non-Markovian effects, which manifest as correlations in time, can be used to improve the functionality of the quantum information processor [7–9]. Thus, modeling and characterizing different varieties of non-Markovian noise is of strong interest.

The first challenge in this endeavor is to be able to differentiate between Markovian and non-Markovian noise in the quantum regime, which is not an easy task. Often, Markovian noise is associated with the exponential decay curves, e.g., a qubit that relaxes to the maximally mixed state exponentially fast. However, there are instances where a qubit exhibits an exponential decay, but nevertheless is undergoing a non-Markovian process [10,11]. A famous example is due to Lindblad, dubbed as a *shallow pocket* (SP), which has been scrutinized in detail recently in terms of dynamical decoupling [12,13], signaling [13], and multitime correlations [14]. (See Ref. [15] for a generalization of a SP.) In each case it is clear that the seemingly simple Markovian noise is, in fact, complex non-Markovian noise that can be exploited to prolong the coherence time of the system. On the other hand, there is a class of system-environment dynamics, generated

by Chebotarev-Gregoratti (CG) Hamiltonians [16] (also see Ref. [17]), that yield an exponential decay for the system but, unlike SP, do not allow for dynamical decoupling or other noise mitigation methods.

Recently, the matter took a turn for the worse; Ref. [18] introduced a class of models in which the decay of the system exhibits the tantalizingly Markovian exponential decay for a finite, but arbitrarily large, time window; then, abruptly, there is a departure from the exponential decay, i.e., the system's evolution displays non-Markovian features of the noise. These models have thus been termed as *hidden non-Markovian* (HNM) models. They occupy the space somewhere between SP and CG models. Importantly, given one of these processes, a snapshot [19] of the dynamics can fail to differentiate between them and will label each as Markovian [3,4]. This is an important space, as most real experiments exhibit the aforementioned exponential decay, and suggests that non-Markovianity may be unassessable.

One possibility is to examine how the system reacts to interventions. Indeed, since non-Markovian systems tend to respond positively to error mitigation methods such as dynamical decoupling, it would appear natural that even models with hidden non-Markovianity can be “stimulated” by interventions to already reveal this at much earlier time. For instance, with the right type of intervention we might be able to fill up the quantum environment with excitation and stimulate an earlier backflow. Multitime correlation measurements were also shown to reveal the failure of the quantum regression theorem, and therefore a subtle type of non-Markovianity [20], which may appear as a precursor.

In this Letter we show that the class of HNM processes are, in fact, genuinely Markovian within the finite window and genuinely non-Markovian outside of that temporal window. We do this by computing the multitime correlations, including those stemming from interventions. This is provably a set of *necessary and sufficient* quantum Markov conditions [21–23].

*daniel.burgarth@mq.edu.au

†paolo.facchi@ba.infn.it

‡davide.longro@ba.infn.it

§kavan.modi@monash.edu




FIG. 1. (a): Scheme of a quantum stochastic process for a system coupled with an environment, initially prepared in a global state $\varrho_0 = \rho \otimes \rho_{\text{vac}}$. At times t_1, \dots, t_4 , an intervention is performed on the system alone; between consecutive interventions, system and environment evolve freely. (b): Scheme of a qubit in a photon waveguide with a two-point interaction. The loop of length T creates a time-delayed coherent feedback and non-Markovian effects on the qubit evolution. (c): Free evolution of $|0, \text{vac}\rangle$: plot of the survival probability of the emitter $|a(t)|^2$ (top) and density plot of the wavefunction of the emitted boson $|\xi_t(x)|^2$ (bottom) for the model with double point interaction, with $\gamma T = \omega_0 T = 2$; darker shades correspond to larger values of $|\xi_t(x)|^2$. In both graphs, the dashed black line represents the time $t = T$ at which $|a(t)|^2$ ceases to be exponential and Eq. (10) ceases to hold. Plotted quantities are dimensionless.

We begin by revisiting the role of multitime correlations in quantum stochastic processes, and the basic properties of HNM models; then we compute such correlations for HNM models, showing that they vanish identically at small times. In other words, we show that there are non-Markovian processes whose non-Markovianity is *perfectly hidden* for a finite duration; non-Markovianity can be therefore undetectable at small times, even by taking into account the role of interventions, since the onset of all time correlations can be arbitrarily delayed.

Quantum multitime correlations. A classical stochastic process is a joint probability distribution of a random variable x in time. In practice, one usually considers the joint distribution of a discrete set of times, $\{t_k, \dots, t_0\}$, and corresponding probability distributions of $\mathbb{P}(x_k, \dots, x_0)$. A Markovian process satisfies the following condition: $\mathbb{P}(x_k | x_{k-1}, \dots, x_0) = \mathbb{P}(x_k | x_{k-1})$, i.e., the system, at a given time, is only conditionally dependent of its state at the previous time step. Practically speaking, this condition means that a Markov process is easy to work with. Nevertheless, such processes are not the norm, but rather a special case [24].

For multitime correlations in the quantum case, consider an initial system-bath (SB) state ϱ_0 , which undergoes an evolution U_0 and then an intervention A_1 is made on the system S alone. Next, the total state once again evolves due to U_1 , followed by a second intervention A_2 on S alone, and so on, up to a final intervention A_k performed following U_{k-1} ; see Fig. 1(a). The interventions $\{A_j\}$ are any physically implementable operation, which can be thought of as a generalized measurement with possible corresponding outcomes $\{x_j\}$. Mathematically, these are known as *instruments* [25]

and represented by a collection of completely positive maps $\mathcal{J} := \{A_{x_j}\}$ such that $\sum_{x_j} A_{x_j}$ is trace preserving.

The above machinery straightforwardly allows for the calculation of the probability to observe a sequence of quantum events (x_k, \dots, x_1) , corresponding to a choice of instruments $\{(\mathcal{J}_k, \dots, \mathcal{J}_1)\}$, as

$$\mathbb{P}(x_k, \dots, x_1 | \mathcal{J}_k, \dots, \mathcal{J}_1) = \text{tr}[A_{x_k} U_{k-1} \dots A_{x_1} U_0(\varrho_0)].$$

While the left-hand side is akin to a classical joint probability distribution, we have yet to identify the quantum stochastic process. We can do this by rewriting the right-hand side as

$$\text{tr}[A_{x_k} U_{k-1} \dots A_{x_1} U_0 \varrho_0] = \text{tr}[\Upsilon_{k:0} \mathbf{A}_{k:1}^T] \quad (1)$$

with $\mathbf{A}_{k:1} := A_{x_k} \otimes \dots \otimes A_{x_1}$ and $\Upsilon_{k:0} := \text{tr}_B[U_{k-1} \star \dots \star U_0 \varrho_0]$, where T denotes transposition and \star denotes the link product, defined as a matrix product on space B and a tensor product on space S [26]. The important feature here is the clear separation of the interventions $\mathbf{A}_{k:1}$ from the influences due to the bath, which are packaged in $\Upsilon_{k:0}$, which is the *Choi state* of the process [5].

The process tensor $\Upsilon_{k:0}$ is the quantum generalization of the joint classical probability distribution and unambiguously represents a quantum stochastic process [27–29]. It contains all accessible multitime correlations [9,30], including temporal entanglement [31–33], and Markovian processes are those satisfying the following property: any k -time process tensor factorizes as

$$\Upsilon_{k:0} = \Upsilon_{k:k-1} \otimes \dots \otimes \Upsilon_{2:1} \otimes \Upsilon_{1:0}. \quad (2)$$

Conversely, we can deduce a process to be non-Markovian by looking for correlations. For instance, in the SP model, while

the qubit dephases exponentially, an intervention of a Pauli σ_x operation reverses this process: therefore, in this model non-Markovianity isn't seen in two-time correlations, but it is seen in three-time correlations.

Hidden non-Markovianity models. We shall consider a qubit with ground state $|0\rangle$ and excited state $|1\rangle$, with excitation energy ω_0 , interacting with a one-dimensional bosonic bath on the real line. We write the Hamiltonian of the bath in the position representation, with annihilation and creation operators b_x, b_x^\dagger satisfying the commutation relations $[b_x, b_{x'}^\dagger] = \delta(x - x')$. Throughout this section, we shall use the following compact notation: for any wave function $\eta(x)$, we set

$$B^\dagger(\eta) = \int dx \eta(x) b_x^\dagger, \quad (3)$$

which represents the creation operator of a boson with wave function $\eta(x)$. The total Hamiltonian $H = H_0 + H_{\text{int}}$ is the sum of the free Hamiltonian

$$H_0 = \omega_0 |0\rangle\langle 0| \otimes \mathbf{1} + \mathbf{1} \otimes H_B, \quad (4)$$

and the interaction Hamiltonian

$$H_{\text{int}} = \sigma_+ \otimes B(g) + \sigma_- \otimes B^\dagger(g). \quad (5)$$

Above, H_B is the second quantization of the momentum operator $p = -i \frac{d}{dx}$ on the boson field, $\sigma_+ = \sigma_-^\dagger = |0\rangle\langle 1|$, and $g(x)$ is the form factor which encodes all information about the qubit-field interaction; different choices of the form factor can yield drastically distinct physics.

Local point interaction. A point interaction between the field and a qubit at $x = 0$ is obtained by setting $g(x) = \sqrt{\gamma} \delta(x)$, with coupling constant $\gamma > 0$, which gives $B(g) = \sqrt{\gamma} b_0$ [34]. This choice corresponds to a flat form factor in the frequency representation used in Ref. [18]. In the single-excitation sector, the model is exactly solvable; in particular, for every $t \in \mathbb{R}$ we have $e^{-itH} |1, \text{vac}\rangle = e^{-i\varepsilon_0 t} |1, \text{vac}\rangle$ and [35]

$$e^{-itH} |0, \text{vac}\rangle = a(t) |0, \text{vac}\rangle + B^\dagger(\xi_t) |1, \text{vac}\rangle. \quad (6)$$

Above, $|s, \text{vac}\rangle \equiv |s\rangle \otimes |\text{vac}\rangle$, $s = 0, 1$, with $|\text{vac}\rangle$ being the field vacuum state; besides, $a(t) = e^{-(i\varepsilon_0 + \gamma/2)t}$ with ε_0 being the dressed excitation energy of the qubit; finally, the boson wave function is given by $\xi_t(x) = \varphi_t(x)$, where

$$\varphi_t(x) = -i\sqrt{\gamma} a(t-x) 1_{[0,t]}(x), \quad (7)$$

with $1_\Omega(x)$ being the characteristic function of the set Ω , that is, $1_\Omega(x) = 1$ for $x \in \Omega$ and $= 0$ elsewhere. This is a simple model of an emission process: the decay of the qubit at $x = 0$ is associated with the creation of a photon which propagates along the positive direction of the x axis with unit velocity. The photon wave function exactly traces out the exponential decay of the qubit and is compactly supported in the interval $[0, t]$.

Consequently, the quantum evolution Λ_t of the qubit obtained by preparing the system in a state ρ and the field in the vacuum $\rho_{\text{vac}} = |\text{vac}\rangle\langle \text{vac}|$, letting them evolve for a time t , and tracing out the field, namely

$$\rho(t) \equiv \Lambda_t(\rho) = \text{tr}_B[e^{-itH} \rho \otimes \rho_{\text{vac}} e^{itH}], \quad (8)$$

satisfies the semigroup property $\Lambda_t \Lambda_s = \Lambda_{t+s}$ at all times, and yields a Markovian evolution: indeed, it is simply an amplitude-damping channel with decay rate γ [18].

Nonlocal point interactions. Now we consider a modification of the above setup. Namely, we allow for a nonlocal interaction of the qubit at two distinct points, say $x = 0$ and $x = T$, given by a form factor

$$g(x) = \sqrt{\frac{\gamma}{2}} [\delta(x) + \delta(x - T)], \quad (9)$$

so that $B(g) = \sqrt{\gamma/2} (b_0 + b_T)$; see Fig. 1(b). Physically, such a choice (up to a relative sign) has been used, see, e.g., [36], as a model of a qubit in a single-end waveguide with a perfect mirror at one end, with $T/2$ being the distance between the qubit and the mirror. This model was also studied in Ref. [18] as an example of non-Markovian model yielding exponential decay for the reduced dynamics of the qubit up to a time T .

Again, the single-excitation sector is fully solvable [35]. In particular, for $0 \leq t \leq T$, we have again an evolved state (6) with $a(t) = \exp[-(i\varepsilon_0 + \frac{\gamma}{2})t]$, and

$$\xi_t(x) = \frac{1}{\sqrt{2}} [\varphi_t(x) + \varphi_t(x - T)], \quad t \in [0, T], \quad (10)$$

where $\varphi_t(x)$ is as in Eq. (7). We may interpret this new situation as a two-point emission: at time $t = 0$, the qubit emits a photon at both positions $x = 0, T$, and each of its two spatially separated parts propagates separately in the positive direction of the x axis up to the time $t = T$; see Fig. 1(b). Notice that the two components, up to normalization, are two exact copies of the wave function of the one-point emission, and in particular the overall norm $\|\xi_t\|^2 = \|\varphi_t\|^2 = 1 - e^{-\gamma t}$ is the same as before. In this sense, at times smaller than T , this system behaves like the superposition of two identical, independent ‘‘copies’’ of the previous system. When $t = T$, one of the photon branches starts interfering back with the qubit and this simple picture ceases immediately to hold: $a(t)$ is no longer an exponential function, and $\xi_t(x)$ will no longer satisfy Eq. (10). This is clearly visible in the plots of $|a(t)|^2$ and $|\xi_t(x)|^2$ in Fig. 1(c).

As a result, the evolution of the qubit state $\rho(t)$, obtained in the same way as in Eq. (8) will be completely indistinguishable from the previous one, as long as $0 \leq t \leq T$. In fact, the form factor given by Eq. (9) is just one possible example: the same result would be obtained by taking into account any number of spatially separated nonlocal point interactions: until the various components of the photon wave function do not propagate to the next point and start interfering. We can thus construct a large family of quantum systems with the HNM property, as shown in Ref. [18]. In all such cases, we obtain non-Markovian processes which, as long as we examine their free dynamics, ‘‘look exactly like’’ a Markovian process at times $0 \leq t \leq T$, despite ultimately starting to show non-Markovian behavior at $t > T$.

This argument, however, still does not take into account multitime correlations of the qubit, which are accessible by means of external interventions on the system. The above result only accounts for correlations between two times, and in general, correlations at *all* orders should

be proved to vanish for assessing that a process is Markovian.

We will make a few remarks before going on and examining multitime correlations. The spin-boson model (4),(5), with a singular point interaction $g(x) = \sqrt{\gamma} \delta(x)$ and with boson Hamiltonian H_B the right shift generator, can be mapped into an SLH model [16,37], with $S = 1$, $L = \sigma_-$, and $H = \varepsilon_0 |0\rangle\langle 0|$. As such, it is associated with an Itô quantum stochastic differential equation [38], with a regression property for all multitime correlation functions, and thus with a strong quantum Markovianity property [4]. In this respect, the model with a two-point interaction (9), is a non-Markovian generalization of an SLH model, in which a feedback is considered; see Fig. 1(b). This is also linked with the theory of time-delayed coherent quantum feedback [39], where a similar scheme was considered. Interestingly enough, this theoretical model is an effective description of the experimental implementation of superconducting artificial giant atoms in waveguide electrodynamics [40]. Finally, we remark that a master equation hierarchy formalism for general systems interacting with an N -boson temporal wave packet was developed in Ref. [41].

Multitime correlations in HNM models. Now we turn to the main question of this Letter, that is, whether higher order correlations are present in HNM models. In other words, can intermediate interventions reveal hidden non-Markovianity already in the time interval $[0, T]$? Such interventions can cause the system to go in higher-excitation sectors, which are not solvable; the ‘indistinguishability’ between this model and the reference one may, in principle, be broken.

However, the system with form factor (9) satisfies a property which will be crucially employed here: given a time $t < T$ and any wave function $\eta(x)$ which satisfies

$$\eta(x) = 0, \quad \text{for } x \in [-t, 0] \cup [T - t, T], \quad (11)$$

then the following property holds [35] for $B^\dagger(\eta)$ in (3):

$$e^{-itH} B^\dagger(\eta) e^{itH} = B^\dagger(e^{-itp}\eta), \quad (12)$$

where $e^{-itp}\eta(x) = \eta(x - t)$ is the free evolution of $\eta(x)$. Physically, Eq. (12) can be interpreted as follows: since wave functions satisfying (11) are sufficiently far apart from the point interactions, they freely propagate without interfering. In particular, this implies that, for $t_1, t_2 \geq 0$ such that $t_1 + t_2 < T$, we have

$$e^{-it_2H} B^\dagger(\xi_{t_1}) e^{it_2H} = B^\dagger(e^{-it_2p}\xi_{t_1}), \quad (13)$$

with $\xi_t(x)$ being the boson wave function (10). This simple equation encodes a fundamental property of the system, which proves to be the key point of our discussion: as long as the total time of observation is less than T , the boson field *cannot carry information* about the non-Markovianity of the system. Local operations of any kind on the qubit cannot modify this simple picture: however we intervene on the qubit via a process as depicted in Fig. 1(a), with all interventions made at times $t_j < T$, all photons emitted in the process will evolve exactly as they would in the absence of coupling. Consequently, all multitime correlations vanish, and the process is genuinely Markovian up to a time T . Only at later times the field will ‘recognize’ the inherently non-Markovian structure of the coupling.

This argument is fully backed up by a lengthy but straightforward calculation which shows that, as a consequence of property (13), the process tensor $\Upsilon_{k,0}$ for all the HNM models (and, in particular, for the model discussed in the previous section) is *exactly the same*, and has the factor form (2), as long as $t_k \in [0, T]$. However, for $t_1 \in [0, T]$ and $t_2 > T$, the process tensor will display correlations. Therefore, all HNM models define a quantum process which is genuinely Markovian within a finite time window and (apart from the one-point interaction which is Markovian for all times) genuinely non-Markovian outside that window. The proof is shown in [35] for three-time correlations, and may be easily generalized for an arbitrary number of interventions.

What is surprising here is that these intricate memory structures stem from a rather simple time-independent Hamiltonian. This is a feature that may be used to engineer intricate temporal correlations, which we discuss in our concluding remarks.

Conclusions. When the Markovian properties of quantum noise were first investigated [10,11] the emphasis was on the natural properties of a given system. Nowadays, quantum information adds an important engineering perspective, which asks how systems behave differently in the context of design and control. This means that device characterization needs to test quantum systems under a wide range of interventions. We have shown here that it is ultimately impossible, in general, to fully conclude if a given system is truly Markovian, no matter how complex this characterization is. Using such a device under Markovian design assumptions can then lead to unexpected behavior at a later stage.

On the other hand, open systems quickly become too complex. Simple models like SP, CG, and now HNM allow us to form simple building block for complex processes and offer keen intuition about the structures of quantum stochastic processes. There are several features of the HNM of interest. First, this is an example of a process which has no short-term memory and only long-term memory. Moreover, this model could serve as the basis for constructing processes with only higher-order correlations, that is, a process where only correlations above four points in time are nonvanishing. Finally, HNM can serve as an ansatz for simulating processes with slow decaying correlations. To do this, we may add more loops to at the top in Fig. 1(b) and create a self-similar structure that will reduce the correlation strength geometrically after a delay of T . The slowly decaying correlation in time here will be akin to the highly common $1/f$ noise. Modeling processes with slowly decaying noise is thought to be hard for the same reasons as the tensor network representation of such correlation is nontrivial [42].

Acknowledgments. D.B. acknowledges funding by the ARC (Projects No. FT190100106, No. DP210101367, and No. CE170100009). P.F. and D.L. were partially supported by the Italian National Group of Mathematical Physics (GNFM-INdAM), by Istituto Nazionale di Fisica Nucleare (INFN) through the project ‘‘QUANTUM,’’ and by Regione Puglia and QuantERA ERA-NET Co-fund in Quantum Technologies (GA No. 731473), project PACE-IN. K.M. was supported

through Australian Research Council Future Fellowship No. FT160100073, Discovery Project No. DP210100597, and the

International Quantum U Tech Accelerator award by the U.S. Air Force Research Laboratory.

-
- [1] J. Preskill, Sufficient condition on noise correlations for scalable quantum computing, *Quant. Inf. Comput.* **13**, 181 (2013).
- [2] J. Helsen, I. Roth, E. Onorati, A. H. Werner, and J. Eisert, A general framework for randomized benchmarking, [arXiv:2010.07974](https://arxiv.org/abs/2010.07974).
- [3] Á. Rivas, S. F. Huelga, and M. B. Plenio, Quantum non-Markovianity: characterization, quantification and detection, *Rep. Prog. Phys.* **77**, 094001 (2014).
- [4] L. Li, M. J. Hall, and H. M. Wiseman, Concepts of quantum non-Markovianity: A hierarchy, *Phys. Rep.* **759**, 1 (2018).
- [5] S. Milz and K. Modi, Quantum stochastic processes and quantum non-markovian phenomena, *PRX Quantum* **2**, 030201 (2021).
- [6] H.-P. Breuer, E.-M. Laine, and J. Piilo, Measure for the Degree of Non-Markovian Behavior of Quantum Processes in Open Systems, *Phys. Rev. Lett.* **103**, 210401 (2009).
- [7] L. Viola, E. Knill, and S. Lloyd, Dynamical Decoupling of Open Quantum Systems, *Phys. Rev. Lett.* **82**, 2417 (1999).
- [8] C. Arenz, D. Burgarth, P. Facchi, and R. Hillier, Dynamical decoupling of unbounded hamiltonians, *J. Math. Phys.* **59**, 032203 (2018).
- [9] G. A. L. White, C. D. Hill, F. A. Pollock, L. C. L. Hollenberg, and K. Modi, Experimental non-Markovian process characterization and control on a quantum processor, *Nat. Commun.* **11**, 6301 (2020).
- [10] G. Lindblad, Response of Markovian and non-Markovian quantum stochastic systems to time-dependent forces, Report TRITA-TFY-79-9, Stockholm 1979.
- [11] L. Accardi, A. Frigerio, and J. T. Lewis, Quantum stochastic processes, *Pub. Res. Inst. Math. Sci.* **18**, 97 (1982).
- [12] C. Arenz, R. Hillier, M. Fraas, and D. Burgarth, Distinguishing decoherence from alternative quantum theories by dynamical decoupling, *Phys. Rev. A* **92**, 022102 (2015).
- [13] S. Milz, M. S. Kim, F. A. Pollock, and K. Modi, Completely Positive Divisibility Does Not Mean Markovianity, *Phys. Rev. Lett.* **123**, 040401 (2019).
- [14] P. Taranto, F. A. Pollock, and K. Modi, Memory strength and recoverability of non-markovian quantum stochastic processes, *npj Quantum Inf.* **7**, 149 (2021).
- [15] D. Burgarth, P. Facchi, and R. Hillier, Control of quantum noise: on the role of dilations, [arXiv:2010.00279](https://arxiv.org/abs/2010.00279).
- [16] M. Gregoratti, The hamiltonian operator associated with some quantum stochastic evolutions, *Commun. Math. Phys.* **222**, 181 (2001).
- [17] D. Burgarth, P. Facchi, M. Fraas, and R. Hillier, Non-Markovian noise that cannot be dynamically decoupled by periodic spin echo pulses, *SciPost Phys.* **11**, 27 (2021).
- [18] D. Burgarth, P. Facchi, M. Ligabò, and D. Lonigro, Hidden non-Markovianity in open quantum systems, *Phys. Rev. A* **103**, 012203 (2021).
- [19] M. M. Wolf, J. Eisert, T. S. Cubitt, and J. I. Cirac, Assessing Non-Markovian Quantum Dynamics, *Phys. Rev. Lett.* **101**, 150402 (2008).
- [20] G. Guarnieri, A. Smirne, and B. Vacchini, Quantum regression theorem and non-Markovianity of quantum dynamics, *Phys. Rev. A* **90**, 022110 (2014).
- [21] F. A. Pollock, C. Rodríguez-Rosario, T. Frauenheim, M. Paternostro, and K. Modi, Non-Markovian quantum processes: Complete framework and efficient characterization, *Phys. Rev. A* **97**, 012127 (2018).
- [22] F. A. Pollock, C. Rodríguez-Rosario, T. Frauenheim, M. Paternostro, and K. Modi, Operational Markov Condition for Quantum Processes, *Phys. Rev. Lett.* **120**, 040405 (2018).
- [23] F. Costa and S. Shrapnel, Quantum causal modelling, *New J. Phys.* **18**, 063032 (2016).
- [24] N. G. van Kampen, Remarks on non-Markov processes, *Braz. J. Phys.* **28**, 90 (1998).
- [25] E. Davies and J. Lewis, An operational approach to quantum probability, *Commun. Math. Phys.* **17**, 239 (1970).
- [26] G. Chiribella, G. M. D'Ariano, and P. Perinotti, Theoretical framework for quantum networks, *Phys. Rev. A* **80**, 022339 (2009).
- [27] S. Milz, F. Sakuldee, F. A. Pollock, and K. Modi, Kolmogorov extension theorem for (quantum) causal modelling and general probabilistic theories, *Quantum* **4**, 255 (2020).
- [28] P. Strasberg and M. G. Díaz, Classical quantum stochastic processes, *Phys. Rev. A* **100**, 022120 (2019).
- [29] S. Milz, D. Egloff, P. Taranto, T. Theurer, M. B. Plenio, A. Smirne, and S. F. Huelga, When Is a Non-Markovian Quantum Process Classical? *Phys. Rev. X* **10**, 041049 (2020).
- [30] G. A. L. White, F. A. Pollock, L. C. L. Hollenberg, C. D. Hill, and K. Modi, Non-Markovian Quantum Process Tomography, [arXiv:2106.11722](https://arxiv.org/abs/2106.11722).
- [31] C. Giarmatzi and F. Costa, Witnessing quantum memory in non-Markovian processes, *Quantum* **5**, 440 (2021).
- [32] S. Milz, C. Spee, Z.-P. Xu, F. A. Pollock, K. Modi, and O. Gühne, Genuine multipartite entanglement in time, *SciPost Phys.* **10**, 141 (2021).
- [33] G. A. L. White, F. A. Pollock, L. C. L. Hollenberg, C. D. Hill, and K. Modi, Diagnosing temporal quantum correlations with compressed non-Markovian calipers, [arXiv:2107.13934](https://arxiv.org/abs/2107.13934).
- [34] Strictly speaking, the sum of (4) and (5) is a well-defined (and self-adjoint) Hamiltonian only when $g(x)$ is square integrable, which is clearly not the case here; however, as discussed in [43,44], via a renormalization procedure (*singular coupling limit*) a self-adjoint operator can be obtained even in the cases that we are considering.
- [35] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevA.104.L050404> for the proof.
- [36] Y.-L. L. Fang, F. Ciccarello, and H. U. Baranger, Non-Markovian dynamics of a qubit due to single-photon scattering in a waveguide, *New J. Phys.* **20**, 043035 (2018).
- [37] J. Combes, J. Kerckhoff, and M. Sarovar, The SLH framework for modeling quantum input-output networks, *Adv. Phys.: X* **2**, 784 (2017).

- [38] R. L. Hudson and K. R. Parthasarathy, Quantum Ito's formula and stochastic evolutions, *Commun. Math. Phys.* **93**, 301 (1984).
- [39] A. L. Grimsmo, Time-Delayed Quantum Feedback Control, *Phys. Rev. Lett.* **115**, 060402 (2015).
- [40] B. Kannan, M. J. Ruckriegel, D. L. Campbell, A. Frisk Kockum, J. Braumüller, D. K. Kim, M. Kjaergaard, P. Krantz, A. Melville, B. M. Niedzielski, A. Vepsäläinen, R. Winik, J. L. Yoder, F. Nori, T. P. Orlando, S. Gustavsson, and W. D. Oliver, Waveguide quantum electrodynamics with superconducting artificial giant atoms, *Nature (London)* **583**, 775 (2020).
- [41] B. Q. Baragiola, R. L. Cook, A. M. Brańczyk, and J. Combes, n -photon wave packets interacting with an arbitrary quantum system, *Phys. Rev. A* **86**, 013811 (2012).
- [42] R. Orús, A practical introduction to tensor networks: Matrix product states and projected entangled pair states, *Ann. Phys.* **349**, 117 (2014).
- [43] D. Lonigro, P. Facchi, and M. Ligabò, The Friedrichs-Lee model and its singular coupling limit, *Proceedings (MDPI)* **12**, 17 (2019).
- [44] P. Facchi, M. Ligabò, and D. Lonigro, Spectral properties of the singular Friedrichs-Lee Hamiltonian, *J. Math. Phys.* **62**, 032102 (2021).