




Measurement disturbance tradeoffs in three-qubit unsupervised quantum classificationHector Spencer-Wood ^{1,*}, John Jeffers ² and Sarah Croke ¹¹*School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, Scotland*²*Department of Physics, University of Strathclyde, John Anderson Building, 107 Rottenrow, Glasgow G4 0NG, Scotland*

(Received 8 February 2022; revised 8 June 2022; accepted 10 June 2022; published 27 June 2022)

We consider measurement disturbance tradeoffs in quantum machine learning protocols which seek to learn about quantum data. We study the simplest example of a binary classification task in the unsupervised regime. Specifically, we investigate how a classification of two qubits, that can each be in one of two unknown states, affects our ability to perform a subsequent classification on three qubits when a third is added. Surprisingly, we find a range of strategies in which a nontrivial first classification does not affect the success rate of the second classification. There is, however, a nontrivial measurement disturbance tradeoff between the success rate of the first and second classifications, and we fully characterize this tradeoff analytically.

DOI: [10.1103/PhysRevA.105.062447](https://doi.org/10.1103/PhysRevA.105.062447)**I. INTRODUCTION**

The proliferation of huge datasets in modern science, technology, and society in general has spurred rapid developments in machine learning; a powerful set of techniques which seek to automate the drawing of inferences from data. A recent theoretical development has been to apply ideas from machine learning to the processing of quantum data [1], both in a supervised setting [2–4] and an unsupervised setting [5–7], for example, at the output of a quantum communication, sensing, or processing device. In the longer run, as quantum technologies develop further, such techniques may be expected to find use in, e.g., characterizing quantum channels and devices, including monitoring for malfunctions [6,7]. Indeed, in [6,7], the problem of determining a quantum change point was addressed. Here, the change point could be the result of some unknown error in a quantum device outputting quantum data. We note that another prominent line of research in quantum machine learning is that of using quantum processing techniques to aid and speed up machine learning when applied to classical data [1]. We, however, only consider the task of learning about quantum data, which requires rather different techniques.

Quantum data are fundamentally different to classical data, and learning strategies are therefore subject to different, peculiarly quantum limitations, which are not yet well explored. As an example, quantum data famously cannot be cloned [8,9], in stark contrast to the classical case. In addition, it is not possible to extract information about a quantum system without causing disturbance [10]. Measurement strategies must therefore be carefully chosen and generically (but not always) the globally optimal strategy for any learning task involves waiting until all data have been received and then performing a joint measurement over all systems [5,11–17].

Such considerations thus pose a problem unique to the quantum case: Can we learn about a subset of data without compromising performance on the dataset as a whole? We might expect a measurement-disturbance-type tradeoff between performance on the subset and performance on the whole dataset. In this paper we take the first steps towards understanding this tradeoff, studying the simplest case of unsupervised binary classification of qubit states, with three samples. A binary classification task is one in which the aim is to assign each sample provided to one of two possible classes, as accurately as possible. Unsupervised means that there is no labeled training data provided, and the user or algorithm must do as well as possible by comparing the data samples to each other. We give analytically the precise tradeoff between learning about the first two samples provided and learning about all three samples. This case is simple enough to allow analytic results, while rich enough to demonstrate the tradeoff. Surprisingly, for a range of strategies on the first two qubits, it is possible to avoid any reduction in performance on all three.

Our work is related to the problem of sequential observers extracting information about a system [18–21], however, so far, the literature has mostly considered the case in which sequential observers have access to the *same* system. Here, in the learning scenario, we are interested in how measurements on some part of a system (the first two subsystems in the example considered here) affect measurement on the whole. In addition, prior work considered the supervised learning case, in which a labeled training set was provided and used to induce a function to label test instances. Here it is known that, in the limit of many test instances, global measurements over training and test data are not required for optimal performance, and the training data may be measured in advance without access to the test data [11]. The unsupervised case is more complicated, as the algorithm seeks to both learn from and classify each instance provided.

In the remainder of this paper we will introduce the unsupervised binary classification problem, illustrate the measurement disturbance effect in the learning scenario, and

*h.spencer-wood.1@research.gla.ac.uk

quantify the tradeoff between learning about two samples and learning about all three.

II. BACKGROUND THEORY

We begin by noting the background theory and notation that will be used throughout this paper. First, as was alluded to earlier, we will be classifying qubits. A qubit $|\varphi\rangle$ is an element of the two-dimensional complex Hilbert space \mathbb{C}^2 such that $\langle\varphi|\varphi\rangle = 1$. It will be notationally convenient to consider our qubits as spin-1/2 particles. With this in mind, we can define the computational basis states of our qubits as follows:

$$\begin{aligned} |0\rangle &:= |s = \frac{1}{2}, m_s = \frac{1}{2}\rangle, \\ |1\rangle &:= |s = \frac{1}{2}, m_s = -\frac{1}{2}\rangle, \end{aligned} \quad (1)$$

where s, m_s denote the total spin and z -component of the total spin of the system, respectively. Now, the classification problem we will be considering is this: Given a number of qubits that can be in one of two unknown states $|\varphi_0\rangle, |\varphi_1\rangle$, how well can we assign a label, $|\varphi_0\rangle$ or $|\varphi_1\rangle$, to each of them? Being qubits, $|\varphi_k\rangle$ can be visualized as points on the Bloch sphere [22]. We can therefore explicitly write them as

$$|\varphi_k\rangle = \cos\frac{\theta_k}{2}|0\rangle + e^{i\phi_k}\sin\frac{\theta_k}{2}|1\rangle, \quad (2)$$

where $k \in \{0, 1\}$, and $\theta_k \in [0, \pi]$, $\phi_k \in [0, 2\pi)$ are the polar coordinates of a point on the Bloch sphere.

We will later see that a quantum classification can be formulated as a quantum measurement. A quantum measurement is mathematically equivalent to a positive operator-valued measure (POVM) [22]. We therefore define a measurement as a set of operators $\{\hat{\pi}_i\}$, called measurement operators, that satisfy

$$\hat{\pi}_i \geq 0 \quad \forall i, \quad (3a)$$

$$\sum_i \hat{\pi}_i = \hat{\mathbb{I}}, \quad (3b)$$

where $\hat{\mathbb{I}}$ denotes the identity operator. If a measurement $\{\hat{\pi}_i\}$ is performed on a state $\hat{\rho}_j$ and the measurement outcome is k (considered a correct measurement outcome if $k = j$), then the state $\hat{\rho}_j$ is updated as follows [22,23]:

$$\hat{\rho}_j \rightarrow \frac{\sqrt{\hat{\pi}_k}\hat{\rho}_j\sqrt{\hat{\pi}_k}^\dagger}{\text{Tr}(\hat{\pi}_k\hat{\rho}_j)}. \quad (4)$$

Although this is not the unique form of the allowed update rule for given $\hat{\pi}_k$, it is minimally disturbing and thus most appropriate for our purposes [24]. When a measurement $\{\hat{\pi}_i\}$ on a set of states $\{\hat{\rho}_i\}$ is performed, the probability of success is

$$P_{\text{succ}} = \sum_i p_i \text{Tr}(\hat{\pi}_i \hat{\rho}_i), \quad (5)$$

where p_i is the probability the input state is prepared in the state $\hat{\rho}_i$.

Finally, when considering multiple qubits, a basis that will turn out to be useful is the Schur basis [25]. The Schur basis states are denoted $|s, m_s\rangle|p_s\rangle$ where p_s is what we call the ‘‘path’’ degree of freedom. This basis is a consequence of

Schur-Weyl duality which says [25]

$$(\mathbb{C}^2)^{\otimes 2} \cong (\mathcal{Q}_1 \otimes \mathcal{P}_1) \oplus (\mathcal{Q}_0 \otimes \mathcal{P}_0) \quad (6)$$

for two qubits (i.e., two copies of the Hilbert space of a qubit) and

$$(\mathbb{C}^2)^{\otimes 3} \cong (\mathcal{Q}_{\frac{3}{2}} \otimes \mathcal{P}_{\frac{3}{2}}) \oplus (\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}) \quad (7)$$

for three qubits. Here, $\mathcal{Q}_s, \mathcal{P}_s \subset (\mathbb{C}^2)^{\otimes n}$ are the subspaces invariant under the action of the irreducible representations (irreps) of $SU(2)$ and S_3 , respectively. These irreps of $SU(2)$, S_3 , and therefore subspaces $\mathcal{Q}_s, \mathcal{P}_s$, respectively, can be labeled by total spin s since we are taking our qubits to be spin-1/2 particles. With all this in mind, $\{|s, m_s\rangle\}$ is a basis for \mathcal{Q}_s and $\{|p_s\rangle\}$ is a basis for \mathcal{P}_s . The reason we call $|p_s\rangle$ the path degree of freedom is that there exists a basis of \mathcal{P}_s , which corresponds to the different ways (or paths) by which a composite quantum system’s state develops a spin- s component via the spin addition of its constituent subsystems. This is the natural basis to work in for our problem because the states $|\varphi_k\rangle$ are completely unknown, so there is no preferred direction. This means that the states are maximally mixed within the subspaces \mathcal{Q}_s corresponding to the irreps of $SU(2)$, and all the information is contained within the path degree of freedom: the \mathcal{P}_s subspaces.

Explicitly, for two qubits, this basis relates to the computational basis as follows:

$$\begin{aligned} \mathcal{Q}_1 \otimes \mathcal{P}_1 &: |1, 1\rangle = |00\rangle, \\ &|1, 0\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle), \\ &|1, -1\rangle = |11\rangle, \\ \mathcal{Q}_0 \otimes \mathcal{P}_0 &: |0, 0\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle), \end{aligned} \quad (8)$$

and for three qubits

$$\begin{aligned} \mathcal{Q}_{\frac{3}{2}} \otimes \mathcal{P}_{\frac{3}{2}} &: \left| \frac{3}{2}, \frac{3}{2} \right\rangle = |000\rangle, \\ &\left| \frac{3}{2}, \frac{1}{2} \right\rangle = \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle), \\ &\left| \frac{3}{2}, -\frac{1}{2} \right\rangle = \frac{1}{\sqrt{3}}(|011\rangle + |101\rangle + |011\rangle), \\ &\left| \frac{3}{2}, -\frac{3}{2} \right\rangle = |111\rangle, \\ \mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}} &: \left| \frac{1}{2}, \frac{1}{2} \right\rangle|1\rangle = \frac{1}{\sqrt{6}}(|100\rangle + |010\rangle - 2|001\rangle), \\ &\left| \frac{1}{2}, -\frac{1}{2} \right\rangle|1\rangle = \frac{1}{\sqrt{6}}(-|011\rangle - |101\rangle + 2|110\rangle), \\ &\left| \frac{1}{2}, \frac{1}{2} \right\rangle|0\rangle = \frac{1}{\sqrt{2}}(|100\rangle - |010\rangle), \\ &\left| \frac{1}{2}, -\frac{1}{2} \right\rangle|0\rangle = \frac{1}{\sqrt{2}}(|101\rangle - |011\rangle). \end{aligned} \quad (9)$$

Here, we are using the abbreviation

$$|i_1 i_2 \dots i_n\rangle := |i_1\rangle \otimes |i_2\rangle \otimes \dots \otimes |i_n\rangle \quad (10)$$

for $i_k \in \{0, 1\}$. Note also that when $\dim(\mathcal{P}_s) = 1$ we do not include $|p_s\rangle$.

III. OPTIMAL CLASSIFICATION

We begin by considering the optimal classification of two and three qubits separately. Part of the reason for doing this explicitly is to introduce some of the ideas and notation required for when we perform two sequential classifications. These results were derived previously in [5,26].

A. Optimal classification of two qubits

The aim of a classification of two unknown qubits is to determine whether these two qubits are the same as or different from one another [5,26]. That is, the aim is to distinguish between $|\varphi_0\rangle|\varphi_0\rangle$, $|\varphi_0\rangle|\varphi_1\rangle$, $|\varphi_1\rangle|\varphi_0\rangle$, and $|\varphi_1\rangle|\varphi_1\rangle$ (actually, we only distinguish between $|\varphi_0\rangle|\varphi_0\rangle$ and $|\varphi_0\rangle|\varphi_1\rangle$ as we will see). Let us begin by considering, mathematically, the form of the two-qubit states. Since $|\varphi_k\rangle$ are unknown qubits, to our knowledge, they are equally likely to be located at *any* point on the Bloch sphere. We therefore describe the two possible two-qubit states as mixed states using density operators as follows:

$$\hat{\rho}_{ij} = \int |\varphi_i\rangle|\varphi_j\rangle\langle\varphi_i|\langle\varphi_j| d\varphi_0 d\varphi_1, \quad (11)$$

where $i = 0$, $j \in \{0, 1\}$, and the integral is with respect to the Haar measure and over the entire Bloch sphere. Note that we can always take $i = 0$. This is because, when averaging over the Bloch sphere, all information about whether each qubit is $|\varphi_0\rangle$ or $|\varphi_1\rangle$ is lost and all that remains is information about their relative positions on the Bloch sphere. This means that $\hat{\rho}_{00} = \hat{\rho}_{11}$ and $\hat{\rho}_{01} = \hat{\rho}_{10}$.

The explicit form of these two states, in the Schur basis [Eq. (8)], can be shown to be (see Appendix A 1 for more detail)

$$\hat{\rho}_{00} = \frac{1}{3}(|1, 1\rangle\langle 1, 1| + |1, 0\rangle\langle 1, 0| + |1, -1\rangle\langle 1, -1|), \quad (12a)$$

$$\hat{\rho}_{01} = \frac{1}{4}(|1, 1\rangle\langle 1, 1| + |1, 0\rangle\langle 1, 0| + |1, -1\rangle\langle 1, -1| + |0, 0\rangle\langle 0, 0|). \quad (12b)$$

We can therefore observe that, here, a quantum classification of two unknown qubits corresponds to a quantum measurement (in general, a POVM) that distinguishes between the two states $\hat{\rho}_{00}$, $\hat{\rho}_{01}$. The optimal measurement to do this is made up of the projectors onto the totally symmetric and antisymmetric subspaces invariant under $SU(2)$, respectively,

$$\hat{P}_+ = |1, 1\rangle\langle 1, 1| + |1, 0\rangle\langle 1, 0| + |1, -1\rangle\langle 1, -1|, \quad (13a)$$

$$\hat{P}_- = |0, 0\rangle\langle 0, 0|, \quad (13b)$$

where \hat{P}_+ (\hat{P}_-) is the outcome associated with measuring the state $\hat{\rho}_{00}$ ($\hat{\rho}_{01}$). Here we use the $+/-$ subscripts rather than 00/01 with the hope that this makes the notation later in this paper less confusing to read. This measurement can be motivated by realizing that $\hat{\rho}_{00}$ and $\hat{\rho}_{01}$ commute with one another, which means they have a common set of eigenstates.

So we take the optimal measurement operators \hat{P}_+ , \hat{P}_- to be the (sum of) projectors onto the eigenstates with the largest eigenvalues of $\hat{\rho}_{00}$, $\hat{\rho}_{01}$, respectively. In other words, it is the Holevo-Helstrom measurement for distinguishing between two quantum states [27]. Using Eq. (5), the maximal probability of successfully classifying two equally likely, unknown qubits is calculated as follows:

$$P_{\text{succ}} = \frac{1}{2}[\text{Tr}(\hat{P}_+ \hat{\rho}_{00}) + \text{Tr}(\hat{P}_- \hat{\rho}_{01})], \quad (14)$$

where the $1/2$ comes from the two states $\hat{\rho}_{00}$, $\hat{\rho}_{01}$ being equiprobable. This results in a success rate of

$$P_{\text{succ}} = \frac{5}{8} = 62.5\%. \quad (15)$$

B. Optimal classification of three qubits

Similarly to the two-qubit case, we begin by writing down the possible three-qubit states. In general, we once again express these states as

$$\hat{\rho}_{ijk} = \int |\varphi_i\rangle|\varphi_j\rangle|\varphi_k\rangle\langle\varphi_i|\langle\varphi_j|\langle\varphi_k| d\varphi_0 d\varphi_1, \quad (16)$$

where $i = 0$, $j, k \in \{0, 1\}$. As shown in Appendix A 2, using the Schur basis [Eq. (9)],

$$\hat{\rho}_{000} = \frac{1}{4}\hat{\mathbb{I}}_{\frac{3}{2}}, \quad (17a)$$

$$\hat{\rho}_{001} = \frac{1}{6}\hat{\mathbb{I}}_{\frac{3}{2}} + \frac{1}{6}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|, \quad (17b)$$

$$\hat{\rho}_{010} = \frac{1}{6}\hat{\mathbb{I}}_{\frac{3}{2}} + \frac{1}{24}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (|1\rangle - \sqrt{3}|0\rangle)(\langle 1| - \sqrt{3}\langle 0|), \quad (17c)$$

$$\hat{\rho}_{011} = \frac{1}{6}\hat{\mathbb{I}}_{\frac{3}{2}} + \frac{1}{24}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (|1\rangle + \sqrt{3}|0\rangle)(\langle 1| + \sqrt{3}\langle 0|), \quad (17d)$$

where $\hat{\mathbb{I}}_s$ is the identity operator on the subspace \mathcal{Q}_s . Note that $\hat{\mathbb{I}}_s$, $\hat{\mathbb{I}}_{s'}$ are orthogonal for $s \neq s'$. The optimal measurement that distinguishes the four states in Eq. (17) is

$$\hat{\pi}_{000} = \hat{\mathbb{I}}_{\frac{3}{2}}, \quad (18a)$$

$$\hat{\pi}_{001} = \frac{2}{3}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|, \quad (18b)$$

$$\hat{\pi}_{010} = \frac{1}{6}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (|1\rangle - \sqrt{3}|0\rangle)(\langle 1| - \sqrt{3}\langle 0|), \quad (18c)$$

$$\hat{\pi}_{011} = \frac{1}{6}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (|1\rangle + \sqrt{3}|0\rangle)(\langle 1| + \sqrt{3}\langle 0|). \quad (18d)$$

To motivate this, notice that $\hat{\rho}_{001}$, $\hat{\rho}_{010}$, $\hat{\rho}_{011}$ have S_3 permutation symmetry in their qubits. We can therefore require the optimal measurement to distinguish these three states to have this same symmetry. So, all we need to do is construct $\hat{\pi}_{000}$, $\hat{\pi}_{001}$ to optimally distinguish between $\hat{\rho}_{000}$, $\hat{\rho}_{001}$. From this, we can obtain $\hat{\pi}_{010}$, $\hat{\pi}_{011}$ via the S_3 symmetry mentioned. The construction of $\hat{\pi}_{000}$, $\hat{\pi}_{001}$ follows the same reasoning as that of two-qubit measurement in Eq. (13) aside from the factor of $2/3$ in $\hat{\pi}_{001}$, which is required for completeness. Therefore, using this measurement and Eq. (5), the maximal probability of successfully distinguishing the (equally likely) states in Eq. (17) is

$$P_{\text{succ}} = \frac{5}{12} \approx 41.7\%. \quad (19)$$

IV. MEASUREMENT DISTURBANCE

We thus consider what happens when we perform the optimal measurement (13) to classify two qubits, then add a third qubit and perform an optimal measurement on all three. In particular, we consider the case in which the outcome of the two-qubit measurement is known, and the measurement on all three is updated accordingly. After the first measurement is performed with outcome \hat{P}_j , as discussed earlier with Eq. (4), the two qubit states update as follows:

$$\hat{\rho}_{0i} \rightarrow \hat{\rho}_{0i}^j = \frac{\sqrt{\hat{P}_j} \hat{\rho}_{0i} \sqrt{\hat{P}_j}^\dagger}{\text{Tr}(\hat{P}_j \hat{\rho}_{0i})}. \quad (20)$$

Following this, we add a third qubit, however, it is convenient to instead think of the situation as beginning with three qubits, and performing the measurement in Eq. (13) on the first two. With this in mind, following a measurement outcome of \hat{P}_k , the three qubit states are found using

$$\hat{\rho}_{0ij}^k = \frac{(\sqrt{\hat{P}_k} \otimes \hat{\mathbb{1}}) \hat{\rho}_{0ij} (\sqrt{\hat{P}_k} \otimes \hat{\mathbb{1}})^\dagger}{\text{Tr}(\hat{P}_k \otimes \hat{\mathbb{1}} \hat{\rho}_{0ij})}, \quad (21)$$

where $\hat{\rho}_{0ij}$ are the states in Eq. (17) and $\hat{\mathbb{1}}$ denotes the identity operator on a single qubit.

Explicitly, the states are as follows:

$$\hat{\rho}_{000}^+ = \frac{1}{4} \hat{\mathbb{1}}_{\frac{1}{2}}, \quad (22a)$$

$$\hat{\rho}_{001}^+ = \frac{1}{6} \hat{\mathbb{1}}_{\frac{1}{2}} + \frac{1}{6} \hat{\mathbb{1}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|, \quad (22b)$$

$$\hat{\rho}_{01k}^+ = \frac{2}{9} \hat{\mathbb{1}}_{\frac{1}{2}} + \frac{1}{18} \hat{\mathbb{1}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|, \quad (22c)$$

$$\hat{\rho}_{000}^- = \hat{0} = \hat{\rho}_{001}^-, \quad (22d)$$

$$\hat{\rho}_{01k}^- = \frac{1}{2} \hat{\mathbb{1}}_{\frac{1}{2}} \otimes |0\rangle\langle 0| \quad (22e)$$

for $k = 0, 1$. For each of the first measurement outcomes \hat{P}_\pm , we can therefore find the optimal measurement to be made up of the following projectors:

$$\hat{\pi}_{000}^+ = \hat{\mathbb{1}}_{\frac{1}{2}}, \quad (23a)$$

$$\hat{\pi}_{001}^+ = \hat{\mathbb{1}}_{\frac{1}{2} \otimes \frac{1}{2}}, \quad (23b)$$

$$\hat{\pi}_{01k}^+ = \hat{0}, \quad (23c)$$

$$\hat{\pi}_{000}^- = \hat{\mathbb{1}}_{\frac{1}{2}}, \quad (23d)$$

$$\hat{\pi}_{001}^- = \hat{0}, \quad (23e)$$

$$\hat{\pi}_{01k}^- = \frac{1}{2} \hat{\mathbb{1}}_{\frac{1}{2} \otimes \frac{1}{2}}, \quad (23f)$$

where $\hat{\mathbb{1}}_{\frac{1}{2} \otimes \frac{1}{2}}$ denotes the identity on the subspace $\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}$. These measurements can be motivated by the fact that $\hat{\pi}_i^\pm$ projects its corresponding state $\hat{\rho}_i^\pm$ onto the components, which are larger, or the same as, the same components in all the other states.

Now, the probability of a successful second measurement is given by

$$\begin{aligned} P_{\text{succ}}^{\text{2nd}} &= \sum_{k \in \{+, -\}} \sum_{i, j \in \{0, 1\}} P(\hat{\rho}_{0ij}) P(\hat{P}_k, \hat{\pi}_{0ij}^k | \hat{\rho}_{0ij}), \\ &= \sum_k \sum_{i, j} P(\hat{\pi}_{0ij}^k | \hat{\rho}_{0ij}^k) P(\hat{P}_k | \hat{\rho}_{0ij}) P(\hat{\rho}_{0ij}) \\ &= \frac{1}{4} \sum_k \sum_{i, j} \text{Tr} \left[\hat{\pi}_{0ij}^k \left(\sqrt{\hat{P}_k} \otimes \hat{\mathbb{1}} \right) \hat{\rho}_{0ij} \left(\sqrt{\hat{P}_k} \otimes \hat{\mathbb{1}} \right) \right], \end{aligned} \quad (24)$$

where $P(\hat{\rho}_{0ij})$ is the probability that the system is prepared in the state $\hat{\rho}_{0ij}$ (this is 1/4 for all i, j), $P(\hat{P}_k, \hat{\pi}_{0ij}^k | \hat{\rho}_{0ij})$ denotes the probability that the first measurement outcome is k and the second is $0ij$ given that the state was prepared in the state $\hat{\rho}_{0ij}$, and $P(\hat{\pi}_{0ij}^k | \hat{\rho}_{0ij}^k) \equiv P(\hat{\pi}_{0ij}^k | \hat{\rho}_{0ij}, \hat{P}_k)$. We therefore find that the probability of a successful second classification is affected by an optimal first classification and is reduced to the following value:

$$P_{\text{succ}}^{\text{2nd}} = \frac{19}{48} \approx 39.6\%. \quad (25)$$

Although this is a small reduction in the success rate of the three-qubit measurement, it demonstrates the principle of measurement disturbance caused by the intermediate classification.

V. WEAKENING THE INTERMEDIATE MEASUREMENT

A. Weak two-qubit measurement

Our ultimate aim is to understand how a classification on two qubits affects our ability to perform a subsequent classification in general. So, instead of considering only the optimal measurement on two qubits, we interpolate between this and the weakest possible measurement: the identity measurement. This weakened measurement can be written as

$$\begin{aligned} \hat{\pi}_- &= \alpha \hat{P}_- + \beta \hat{\mathbb{1}}, \\ \hat{\pi}_+ &= \alpha \hat{P}_+ + (1 - \alpha - \beta) \hat{\mathbb{1}}, \end{aligned} \quad (26)$$

such that $\alpha \in [0, 1 - \beta]$ and $\beta \in [0, 1]$,

where the range of values α, β take come about due to the positivity condition of POVMs, given in Eq. (3a), as well as the convention we are adopting: we take the measurement outcome $\hat{\pi}_+$ ($\hat{\pi}_-$) to correspond to the measurement of the state $\hat{\rho}_{00}$ ($\hat{\rho}_{01}$). Note also that, by construction, this POVM is complete, as required [Eq. (3b)]. To reduce future work, note that we can change between the two situations corresponding to different measurement outcomes by performing the swaps

$$\begin{aligned} \alpha &\rightarrow -\alpha, \\ \beta &\rightarrow 1 - \beta. \end{aligned} \quad (27)$$

We conclude this subsection by noting that the probability of a successful two-qubit classification using the POVM in Eq. (26) is given by

$$P_{\text{succ}}^{\text{1st}} = \frac{1}{2} \left(1 + \frac{\alpha}{4} \right), \quad (28)$$

where the superscript is included in anticipation of the second classification introduced in the next subsection.

B. Adding a third qubit

As before, after the first classification of two qubits is performed, a third qubit, either $|\varphi_0\rangle$ or $|\varphi_1\rangle$, is added. To write down the resulting three-qubit state, just as in the case of an optimal intermediate measurement, it is convenient to think instead of the situation as an undisturbed three-qubit state $\hat{\rho}_{0ij}$

$$\hat{\rho}_{000}^- = \frac{1}{4}\hat{\mathbb{I}}_{\frac{3}{2}}, \quad (30a)$$

$$\hat{\rho}_{001}^- = \frac{1}{6}\hat{\mathbb{I}}_{\frac{3}{2}} + \frac{1}{6}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|, \quad (30b)$$

$$\hat{\rho}_{010}^- = \frac{4\beta}{6(\alpha + 4\beta)}\hat{\mathbb{I}}_{\frac{3}{2}} + \frac{1}{6(\alpha + 4\beta)}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (\sqrt{\beta}|1\rangle - \sqrt{3(\alpha + \beta)}|0\rangle)(\sqrt{\beta}\langle 1| - \sqrt{3(\alpha + \beta)}\langle 0|), \quad (30c)$$

$$\hat{\rho}_{011}^- = \frac{4\beta}{6(\alpha + 4\beta)}\hat{\mathbb{I}}_{\frac{3}{2}} + \frac{1}{6(\alpha + 4\beta)}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (\sqrt{\beta}|1\rangle + \sqrt{3(\alpha + \beta)}|0\rangle)(\sqrt{\beta}\langle 1| + \sqrt{3(\alpha + \beta)}\langle 0|), \quad (30d)$$

with probabilities (derived in Appendix B 1)

$$p_{000}^- = p_{001}^- = \frac{2\beta}{\alpha + 8\beta}, \quad (31a)$$

$$p_{010}^- = p_{011}^- = \frac{\alpha + 4\beta}{2(\alpha + 8\beta)}. \quad (31b)$$

To find $\hat{\rho}_{0ij}^+$, we can just perform the swaps in Eq. (27).

To achieve our aim of classifying the resulting three-qubit system, we construct a measurement $\{\hat{\pi}_i^-\}$ that distinguishes between the states $\{\hat{\rho}_i^-\}$ above (for additional detail, see Appendix B 2). To do this, first, notice that the totally symmetric components ($s = 3/2$) of $\hat{\rho}_{001}^-$, $\hat{\rho}_{01i}^-$ are strictly less than that of $\hat{\rho}_{000}^-$. Further, $\hat{\rho}_{000}^-$ has no $s = 1/2$ components. This motivates the fact that the optimal way to distinguish $\hat{\rho}_{000}^-$ from the other states is to take

$$\hat{\pi}_{000}^- = \hat{\mathbb{I}}_{\frac{3}{2}}, \quad (32)$$

while keeping the remaining measurement operators in $(\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}) \otimes (\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}})^*$, where V^* denotes the dual space of V . Next, note that in the $s = 1/2$ subspaces, $\hat{\rho}_{001}$, $\hat{\rho}_{010}$, $\hat{\rho}_{011}$ have a mirror symmetric form in their path degree of freedom (spanned by $|p_{\frac{1}{2}}\rangle = |0\rangle, |1\rangle$) as $p_{010}^- = p_{011}^-$ and the set is invariant under reflection about $|0\rangle$. Including $\hat{\pi}_{000}^-$ for completeness, the optimal measurement to distinguish these three states is known [28] and has the form

$$\hat{\pi}_{000}^- = \hat{\mathbb{I}}_{\frac{3}{2}}, \quad (33a)$$

$$\hat{\pi}_{001}^- = (1 - a_-^2)\hat{\mathbb{I}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|, \quad (33b)$$

$$\hat{\pi}_{010}^- = \frac{1}{2}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (a_-|1\rangle - |0\rangle)(a_- \langle 1| - \langle 0|), \quad (33c)$$

$$\hat{\pi}_{011}^- = \frac{1}{2}\hat{\mathbb{I}}_{\frac{1}{2}} \otimes (a_-|1\rangle + |0\rangle)(a_- \langle 1| + \langle 0|), \quad (33d)$$

where $a_- \in [0, 1]$ to preserve positivity. A closed form analytic expression for a_- in terms of the prior probabilities and overlaps of the states is given in [28], which we use below. Once again, to obtain $\{\hat{\pi}_i^+\}$, we just perform the swaps in Eq. (27).

that is updated by the intermediate measurement on the first two qubits as

$$\hat{\rho}_{0ij}^\pm = \frac{(\sqrt{\hat{\pi}_\pm} \otimes \hat{\mathbb{I}})\hat{\rho}_{0ij}(\sqrt{\hat{\pi}_\pm} \otimes \hat{\mathbb{I}})}{\text{Tr}(\hat{\pi}_\pm \otimes \hat{\mathbb{I}}\hat{\rho}_{0ij})}. \quad (29)$$

Explicitly, in the case when the measurement outcome on the first two qubits is $\hat{\pi}_-$, using similar techniques as those found in Eqs. (A13) and (A14) to find $\sqrt{\hat{\pi}_-} \otimes \hat{\mathbb{I}}$ in the Schur basis, the states $\hat{\rho}_{0ij}^-$ can be shown to be

To utilize [28] we first must define a prior probability for the states $\hat{\rho}_{001}^-$, $\hat{\rho}_{01i}^-$ when projected into the $s = 1/2$ subspace, and can then directly use the results of [28] to find the optimal value of the parameter a_- . Updating the prior probabilities gives

$$p^- = \frac{3\alpha + 4\beta}{6(\alpha + 2\beta)}, \quad (34)$$

which is derived in Appendix B 2. Using the analytical expression in [28] then gives (again more detail is given in Appendix B 2):

$$a_- = \begin{cases} \sqrt{\frac{\alpha + \beta}{3\beta}} & \text{if } \alpha \in [0, \min\{1 - \beta, 2\beta\}], \\ 1 & \text{if } \alpha \in (2\beta, 1 - \beta] \text{ with } 2\beta < 1 - \beta, \end{cases} \quad (35)$$

such that $\beta \in [0, 1]$ as always. Note that the conditions $2\beta < 1 - \beta$, $\beta \in [0, 1]$ can be rewritten as $\beta \in [0, \frac{1}{3})$. Similarly, when the outcome of the first measurement is $\hat{\pi}_+$, we arrive at

$$a_+ = \sqrt{\frac{1 - \alpha - \beta}{3(1 - \beta)}} \quad (36)$$

for all valid α, β . To achieve our aim and observe how the success probability of the first and second measurements compare to one another, we consider the two cases of Eq. (35).

I. Case 1: $\alpha \in [0, \min\{1 - \beta, 2\beta\}]$, $\beta \in [0, 1]$

Consider the first case in Eq. (35), that is, when

$$a_- = \sqrt{\frac{\alpha + \beta}{3\beta}}, \quad (37)$$

$$a_+ = \sqrt{\frac{1 - \alpha - \beta}{3(1 - \beta)}}.$$

Using Eq. (24) with $\hat{P}_\pm \rightarrow \hat{\pi}_\pm$, in this region, it is straightforward, albeit requiring a little algebra, to show that the

probability of a successful second classification stays constant at the optimal value for distinguishing three undisturbed qubits:

$$P_{\text{succ}}^{2\text{nd}} = \frac{5}{12} \quad (38)$$

for all $\alpha \in [0, \min\{1 - \beta, 2\beta\}]$, $\beta \in [0, 1]$.

2. Case 2: $\alpha \in (2\beta, 1 - \beta]$, $\beta \in [0, \frac{1}{3}]$

Considering now the second case in Eq. (35), let $a_- = 1$ and a_+ be as written in Eq. (36). Once again, using Eq. (24) with $\hat{P}_\pm \rightarrow \hat{\pi}_\pm$, after a little algebra, we find

$$P_{\text{succ}}^{2\text{nd}} = \frac{5}{12} - \frac{\beta}{12} - \frac{\alpha}{48} + \frac{1}{24}\sqrt{3\beta(\alpha + \beta)}. \quad (39)$$

Now, we want $P_{\text{succ}}^{2\text{nd}}$ to be its optimal value for each value of $P_{\text{succ}}^{1\text{st}}$. Since $P_{\text{succ}}^{1\text{st}}$ has the form given in Eq. (28) (linear in α alone), to do this, we hold α constant, and maximise $P_{\text{succ}}^{2\text{nd}}$ with respect to β . This occurs when

$$\beta = -\frac{3\alpha}{2} \quad \text{or} \quad \beta = \frac{\alpha}{2}. \quad (40)$$

The first option only holds when $\alpha = 0 \notin (2\beta, 1 - \beta]$. The second option corresponds to the boundary of the two scenarios in Eq. (35), that is, when $\alpha = 2\beta$. This tells us that for $\alpha > 2\beta$, there are no stationary points with respect to β , and we must therefore look to the boundaries of β : $\beta = 0$ or $\beta = 1 - \alpha$. However, the optimal boundary can be shown to be $\beta = 1 - \alpha$ when we notice that $P_{\text{succ}}^{2\text{nd}}$ is monotonically increasing with respect to β in the region $\alpha \in (2\beta, 1 - \beta]$, $\beta \in [0, \frac{1}{3}]$. This can be shown using the fact that there are no stationary points in this region, so it must therefore be monotonically increasing or decreasing, along with the fact there exists a point [e.g., $(\alpha, \beta) = (5/6, 1/6)$] in this region such that $\frac{\partial P_{\text{succ}}^{2\text{nd}}}{\partial \beta} > 0$. So, using $\beta = 1 - \alpha$ along with Eq. (28), we find the optimal probability of success in this region to be

$$P_{\text{succ}}^{2\text{nd}} = \frac{1}{12} + \frac{P_{\text{succ}}^{1\text{st}}}{2} + \frac{1}{24}\sqrt{3(5 - 8P_{\text{succ}}^{1\text{st}})}. \quad (41)$$

We can re-express the boundaries in $P_{\text{succ}}^{2\text{nd}}$ in terms of $P_{\text{succ}}^{1\text{st}}$ by noting that we would like Eq. (38) to be the success rate for as large a region as possible. This can be seen by noting that Eq. (39) can be rewritten as

$$P_{\text{succ}}^{2\text{nd}} = \frac{5}{12} - \frac{1}{48}(\sqrt{\alpha + \beta} - \sqrt{3\beta})^2 \quad (42)$$

and therefore is less than or equal to the optimal value of $5/12$. So, to make the region in which Eq. (38) is true as large as possible, we must maximize $\min\{2\beta, 1 - \beta\}$. That is, when $\beta = 1/3$ and so $\alpha \in [0, \frac{1}{3}]$. Therefore, using Eq. (28), we take $P_{\text{succ}}^{2\text{nd}}$ to be given by Eq. (38) when $P_{\text{succ}}^{1\text{st}} \in [0, \frac{7}{12}]$, and by Eq. (41) when $P_{\text{succ}}^{1\text{st}} \in (\frac{7}{12}, \frac{5}{8}]$.

To gain some intuition as to how the three qubit states and second measurement vary with the strength of the first measurement, we can plot their $s = 1/2$ path components, that is, their components when restricted to the subspace $\mathcal{P}_{\frac{1}{2}}$. Further, since $\hat{\rho}_{000}^\pm, \hat{\pi}_{000}^\pm$ are left invariant by the first measurement, no information is gained by considering them, so we only need look at the remaining states and measurement operators. Figure 1 shows how the states $\hat{\rho}_{001}^-, \hat{\rho}_{010}^-, \hat{\rho}_{011}^-$ and measurement operators $\hat{\pi}_{001}^-, \hat{\pi}_{010}^-, \hat{\pi}_{011}^-$ compare to one another for

various values of α, β . Note the mirror symmetry of the states and measurement operators in their $|0\rangle$ components as discussed earlier when constructing the optimal measurement of the three-qubit states. Further, the adjustment of the second measurement compensates for the disturbance caused by the first measurement in the region $\alpha \in [0, \min\{1 - \beta, 2\beta\}]$, $\beta \in [0, 1]$.

VI. RESULTS

Summarizing what we found, the tradeoff between the first and second classification is given by

$$P_{\text{succ}}^{2\text{nd}} = \begin{cases} \frac{5}{12} & \text{if } P_{\text{succ}}^{1\text{st}} \in [0, \frac{7}{12}], \\ \frac{1}{12} + \frac{P_{\text{succ}}^{1\text{st}}}{2} + \frac{1}{24}\sqrt{3(5 - 8P_{\text{succ}}^{1\text{st}})} & \text{if } P_{\text{succ}}^{1\text{st}} \in (\frac{7}{12}, \frac{5}{8}]. \end{cases} \quad (43)$$

A plot of this tradeoff can be seen in Fig. 2. Let us note some points of interest. First, when we require the second measurement to be optimal, the best first measurement occurs when $\alpha = 2/3$ and $\beta = 1 - \alpha = 1/3$. Here,

$$P_{\text{succ}}^{1\text{st}} = \frac{7}{12} \approx 58.3\%, \quad (44a)$$

$$P_{\text{succ}}^{2\text{nd}} = \frac{5}{12} \approx 41.7\%. \quad (44b)$$

So the success rate of the first measurement, under the requirement that $P_{\text{succ}}^{2\text{nd}}$ is optimal, ranges from $1/2$ to $7/12$. It is worth reiterating that the transition from optimal to suboptimal second-measurement success rate occurs at the boundary of the two cases in Eqs. (43) or (35). That is, given a first outcome of $\hat{\pi}_-$, when the second measurement stops varying with respect to α, β as can be seen in Fig. 1.

The next point to consider is when we optimize $P_{\text{succ}}^{1\text{st}}$. Here $\alpha = 1$ and $\beta = 1 - \alpha = 0$, which means that

$$P_{\text{succ}}^{1\text{st}} = \frac{5}{8} = 62.5\%, \quad (45a)$$

$$P_{\text{succ}}^{2\text{nd}} = \frac{19}{48} \approx 39.6\%, \quad (45b)$$

as was found in Secs. III A and IV. This limited success rate of the second measurement can perhaps be expected due to the fact the ρ_{01k}^- are parallel to one another as can be seen in Fig. 1(d).

VII. CONCLUSION

To summarize, we considered a base case in the tradeoff between two sequential unsupervised quantum learning tasks. In particular, we looked at the situation in which there were initially two qubits that could each be in one of two unknown quantum states. Once a binary classification of varying success rate, corresponding to a quantum measurement of varying strength, was performed, a third qubit was added and the optimal classification on all three qubits was then performed. We found that, although a binary classification of two unknown qubits causes measurement disturbance, which can degrade the performance of an optimal classifier on all three qubits, there is a large regime in which the performance remains unaffected. In this regime, the final measurement may be adjusted to fully mitigate the disturbance caused by the first measurement. That is, the success rate of the first classification




FIG. 1. Plots showing the effect that a measurement on the first two qubits of a system, with outcome $\hat{\pi}_-$ (for various values of α, β), has on the three-qubit states and measurement operators $\{\hat{\rho}_{001}^-, \hat{\rho}_{010}^-, \hat{\rho}_{011}^-\}$ and $\{\hat{\pi}_{001}^-, \hat{\pi}_{010}^-, \hat{\pi}_{011}^-\}$, respectively. Note that the states and measurement operators vary with the strength of the two-qubit measurement until (c) $\alpha = \frac{2}{3}, \beta = \frac{1}{3}$ after which, while the states continue to change, the measurement operators stay constant. (c) corresponds to the boundary between the constant and nonconstant regions in Fig. 2. Further, (a) corresponds to the case in which no measurement is performed on the first two qubits and (d) to when the optimal measurement is performed on the first two qubits.

can range from that of a guess, $P_{succ}^{1st} = 1/2$, to $P_{succ}^{1st} = 7/12$ without causing the success rate of the second classification to deviate from its optimal value of $P_{succ}^{2nd} = 5/12$. When P_{succ}^{1st} is further improved, however, P_{succ}^{2nd} decreases nonlinearly to a

success rate of $19/48$ as P_{succ}^{1st} increases to its optimal value of $5/8$.

This work provides an indication that sequential unsupervised classifications of quantum data can be performed. Further, depending on the strength of an earlier classification, a later classification's ability need not be compromised. Having said this, this work also highlights that there are nontrivial tradeoffs between sequential unsupervised quantum learning tasks, which, although small in this base case, may be more considerable in more complicated scenarios. Here we considered the simplest possible example of a quantum learning task in which a measurement disturbance tradeoff exists between performance on a subset of the data provided and performance on the entire dataset. We fully characterized this tradeoff. This is a peculiarly quantum effect due to fundamental features of quantum mechanics, which is not present in classical machine learning.

This is just the first step in exploring this tradeoff in learning tasks, and more work is required to fully understand the limitations imposed by quantum mechanics on sequential learning. For example, the next natural step would be to consider starting with n unknown qubits of two types and, following a classification of them, adding one or more extra qubits to be subsequently classified. Further, one could look




FIG. 2. Plot of the tradeoff between the success rates of the first binary classification of two qubits and the second binary classification when a third qubit is added. The probability of success of the first (second) measurement is denoted P_{succ}^{1st} (P_{succ}^{2nd}).

at the case in which there are a larger number of options of qubit (or d -dimensional qudit) to choose between. Another path to take could be the supervised analog of the content of this paper, with labeled qubits being given as a training set used to classify future ones. In addition, there are a range of learning scenarios, including partially or fully supervised learning, and reinforcement learning, in which similar effects may be explored. We leave these considerations for future work.

ACKNOWLEDGMENTS

The authors acknowledge The Engineering and Physical Sciences Research Council and the UK National Quantum Technologies Programme via the QuantIC Quantum Imaging Hub (EP/T00097X/1). S. Croke is supported by a Leverhulme Fellowship (RF-2020-397).

APPENDIX A: DERIVATION OF UNDISTURBED STATES

Let us first derive the two-qubit states in Eq. (12). To do this, we use some representation theory. For our purposes, we define the representation $(\mathbf{Q}_n, \mathcal{Q})$ of $SU(2)$ such that for any $U \in SU(2)$,

$$\mathbf{Q}_n(U)|i_1\rangle \cdots |i_n\rangle := U^{\otimes n}|i_1\rangle \cdots |i_n\rangle. \quad (\text{A1})$$

We also require the following representation of the symmetric group S_n :

$$\mathbf{P}(\sigma)|i_1\rangle \cdots |i_n\rangle := |i_{\sigma^{-1}(1)}\rangle \cdots |i_{\sigma^{-1}(n)}\rangle, \quad (\text{A2})$$

such that $\sigma \in S_n$. For instance, for $n = 3$, $\sigma = (123)$,

$$\mathbf{P}((123))|i_1 i_2 i_3\rangle = |i_3 i_1 i_2\rangle. \quad (\text{A3})$$

1. Two-qubit states

Beginning with $\hat{\rho}_{00}$, using the $SU(2)$ invariance of the Haar measure, notice that

$$\begin{aligned} \mathbf{Q}_2(U)\hat{\rho}_{00}\mathbf{Q}_2^\dagger(U) &= \mathbf{Q}_2(U)\left(\int |\varphi_0\varphi_0\rangle\langle\varphi_0\varphi_0|d\varphi_0\right)\mathbf{Q}_2^\dagger(U) \\ &= \int |\varphi_0\varphi_0\rangle\langle\varphi_0\varphi_0|d\varphi_0 = \hat{\rho}_{00}. \end{aligned} \quad (\text{A4})$$

So, by Schur's lemma, there exist bases, for example the Schur basis given in Eq. (8), such that

$$\hat{\rho}_{00} = \alpha_1 \hat{\mathbb{1}}_1 \oplus \alpha_0 \hat{\mathbb{1}}_0. \quad (\text{A5})$$

Recalling Eq. (6), note that the subscripts reference the total spin of the subspaces.

However, notice that for any qubit $|\varphi_0\rangle = a|0\rangle + b|1\rangle$,

$$|\varphi_0\varphi_0\rangle = a^2|00\rangle + ab(|01\rangle + |10\rangle) + b^2|11\rangle. \quad (\text{A6})$$

So, comparing with Eq. (8), $|\varphi_0\varphi_0\rangle$ lives entirely in $\mathcal{Q}_1 \otimes \mathcal{P}_1$. This implies that $\alpha_0 = 0$ and therefore, by the normalization of $\hat{\rho}_{00}$,

$$\begin{aligned} \hat{\rho}_{00} &= \frac{1}{3} \hat{\mathbb{1}}_1 \\ &= \frac{1}{3}(|1, 1\rangle\langle 1, 1| + |1, 0\rangle\langle 1, 0| + |1, -1\rangle\langle 1, -1|). \end{aligned} \quad (\text{A7})$$

Next, for $\hat{\rho}_{01}$, notice that with a slight abuse of notation,

$$\begin{aligned} \hat{\rho}_{01} &= \int |\varphi_0\varphi_1\rangle\langle\varphi_0\varphi_1|d\varphi_0d\varphi_1 \\ &= \int |\varphi_0\rangle\langle\varphi_0| \otimes |\varphi_1\rangle\langle\varphi_1|d\varphi_0d\varphi_1 \\ &= \int |\varphi_0\rangle\langle\varphi_0|d\varphi_0 \otimes \int |\varphi_1\rangle\langle\varphi_1|d\varphi_1 \\ &= \frac{1}{4} \hat{\mathbb{1}} \otimes \hat{\mathbb{1}}, \end{aligned} \quad (\text{A8})$$

where the last equality is obtained by the invariance of $\int |\varphi_i\rangle\langle\varphi_i|d\varphi_i$ under $\mathbf{Q}_1(SU(2))$ (or, more physically, due to each of the integrals describing a maximally mixed qubit) and the $1/4$ is required for normalization. Therefore, $\hat{\rho}_{01}$ is proportional to the identity on $(\mathbb{C}^2)^{\otimes 2}$ and hence, we can rewrite it as the identity in the Schur basis

$$\begin{aligned} \hat{\rho}_{01} &= \frac{1}{4} \hat{\mathbb{1}}_1 \oplus \hat{\mathbb{1}}_0 \\ &= \frac{1}{4}(|1, 1\rangle\langle 1, 1| + |1, 0\rangle\langle 1, 0| + |1, -1\rangle\langle 1, -1| \\ &\quad + |0, 0\rangle\langle 0, 0|). \end{aligned}$$

2. Three-qubit states

Similar arguments to the two-qubit case tell us that we can write the following:

$$\hat{\rho}_{000} = \alpha_{\frac{3}{2}} \hat{\mathbb{1}}_{\frac{3}{2}} \oplus \alpha_{\frac{1}{2}} \hat{\mathbb{1}}_{\frac{1}{2}} \oplus \alpha'_{\frac{1}{2}} \hat{\mathbb{1}}_{\frac{1}{2}} \quad (\text{A9})$$

due to its commutivity with all the elements of $\mathbf{Q}_3(SU(2))$. To motivate the presence of two copies of the $\mathcal{Q}_{\frac{1}{2}}$ space implied here, recall that the addition of three spin-half particles results in a system with *two* orthogonal spin-half components. To find $\alpha_{\frac{1}{2}}, \alpha'_{\frac{1}{2}}$, notice that for all $\sigma \in S_3$,

$$\mathbf{P}(\sigma)\hat{\rho}_{000}\mathbf{P}^\dagger(\sigma) = \hat{\rho}_{000}. \quad (\text{A10})$$

This implies $\hat{\rho}_{000}$ lives entirely within $(\mathcal{Q}_{\frac{3}{2}} \otimes \mathcal{P}_{\frac{3}{2}}) \otimes (\mathcal{Q}_{\frac{3}{2}} \otimes \mathcal{P}_{\frac{3}{2}})^*$, for if it did not, it would have a component within $(\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}) \otimes (\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}})^*$ and would therefore not be acted on trivially by $\mathbf{P}(S_3)$ since the irrep that $\mathcal{P}_{\frac{1}{2}}$ is invariant under is not trivial. It therefore follows that $\alpha_{\frac{1}{2}}, \alpha'_{\frac{1}{2}} = 0$ and

$$\hat{\rho}_{000} = \frac{1}{4} \hat{\mathbb{1}}_{\frac{3}{2}}, \quad (\text{A11})$$

where, again, $1/4$ is the normalization constant.

Now, for $\hat{\rho}_{001}$, using a similar technique to $\hat{\rho}_{01}$,

$$\begin{aligned} \hat{\rho}_{001} &= \int |\varphi_0\varphi_0\varphi_1\rangle\langle\varphi_0\varphi_0\varphi_1|d\varphi_0d\varphi_1 \\ &= \int |\varphi_0\varphi_0\rangle\langle\varphi_0\varphi_0|d\varphi_0 \otimes \int |\varphi_1\rangle\langle\varphi_1|d\varphi_1 \\ &= \frac{1}{2} \hat{\rho}_{00} \otimes \hat{\mathbb{1}} \\ &= \frac{1}{6}(|1, 1\rangle\langle 1, 1| + |1, 0\rangle\langle 1, 0| + |1, -1\rangle\langle 1, -1|) \\ &\quad \otimes \left(\left| \frac{1}{2}, \frac{1}{2} \right\rangle \left\langle \frac{1}{2}, \frac{1}{2} \right| + \left| \frac{1}{2}, -\frac{1}{2} \right\rangle \left\langle \frac{1}{2}, -\frac{1}{2} \right| \right), \end{aligned} \quad (\text{A12})$$

where the prefactors are determined using similar ideas to before. To rewrite this in the Schur basis of $(\mathcal{Q}_{\frac{3}{2}} \otimes \mathcal{P}_{\frac{3}{2}}) \otimes (\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}})$, we use the following [25]:

$$\begin{aligned} & |s, m\rangle|p\rangle \otimes \left| \frac{1}{2}, \pm \frac{1}{2} \right\rangle \\ & \rightarrow \sqrt{\frac{s \pm m + 1}{2s + 1}} \left| s + \frac{1}{2}, m \pm \frac{1}{2} \right\rangle |p, 0\rangle \\ & \mp \sqrt{\frac{s \mp m}{2s + 1}} \left| s - \frac{1}{2}, m \pm \frac{1}{2} \right\rangle |p, 1\rangle. \end{aligned} \quad (\text{A13})$$

For our case, $s = 1, m \in \{1, 0, -1\}$ and p is omitted since $\dim \mathcal{P}_1 = 1 = \dim \mathcal{P}_0$. Applying Eq. (A13) to Eq. (A12), we obtain

$$\hat{\rho}_{001} = \frac{1}{6} (\hat{\mathbb{I}}_{\frac{3}{2}} + \hat{\mathbb{I}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|), \quad (\text{A14})$$

where $\alpha = 1/6$ is found by again requiring $\text{Tr}(\hat{\rho}_{001}) = 1$.

Finally, we can find $\hat{\rho}_{010}, \hat{\rho}_{011}$. Noting that

$$\begin{aligned} \hat{\rho}_{011} &= \int |\varphi_0 \varphi_1 \varphi_1\rangle \langle \varphi_0 \varphi_1 \varphi_1| d\varphi_0 d\varphi_1 \\ &= \int |\varphi_1 \varphi_0 \varphi_0\rangle \langle \varphi_1 \varphi_0 \varphi_0| d\varphi_0 d\varphi_1 = \hat{\rho}_{100}, \end{aligned} \quad (\text{A15})$$

to obtain $\hat{\rho}_{010}, \hat{\rho}_{011}$, notice that we just have to permute the qubits in $\hat{\rho}_{001}$. To do this, first notice that $\hat{\mathbb{I}}_{\frac{3}{2}}$ is invariant under permutations of qubits since $\mathbf{P}(S_3)$ acts trivially on $\mathcal{P}_{\frac{3}{2}}$. Therefore, the only part of $\hat{\rho}_{001}$ affected by permutations is $\hat{\mathbb{I}}_{\frac{1}{2}} \otimes |1\rangle\langle 1|$.

Intuitively, we can guess the form of $\hat{\rho}_{010}, \hat{\rho}_{011}$ by the fact that,

$$\dots \hat{\rho}_{001} \xrightarrow{(123)} \hat{\rho}_{010} \xrightarrow{(123)} \hat{\rho}_{011} \xrightarrow{(123)} \hat{\rho}_{001} \dots, \quad (\text{A16})$$

where $(123) \in S_3$ is a three-cycle. Since permutations only have an affect on the path components of the states, it seems that $\hat{\rho}_{001}, \hat{\rho}_{010}, \hat{\rho}_{011}$ should be evenly distributed in the two-dimensional space $\mathcal{P}_{\frac{1}{2}}$. That is, since each state can be accessed by repeated application of the permutation (123), we would expect each state be accessible by the repeated application of some two-dimensional transformation (on $\mathcal{P}_{\frac{1}{2}}$). In particular, since $\hat{\rho}_{001}$ is known, we might guess that the remaining states could be found by rotating its $\mathcal{P}_{\frac{1}{2}}$ component by $2\pi/3$. This indeed results in the states given in Eq. (17).

More explicitly, we can derive $\hat{\rho}_{010}, \hat{\rho}_{011}$ using the following three steps.

- (1) Rewrite $\hat{\rho}_{001}$ in the computational basis [using Eq. (9)].
- (2) Permute the qubits $|i_1 i_2 i_3\rangle \rightarrow |i_2 i_3 i_1\rangle$ to obtain $\hat{\rho}_{010}$ in the computational basis.
- (3) Rewrite the state in the Schur basis given in Eq. (9).

APPENDIX B: DERIVATION OF UPDATED STATES AND MEASUREMENTS

1. Updated prior probabilities

Assuming the outcome of the measurement on the first two qubits is $-$, the probabilities of the disturbed states $\hat{\rho}_{000}^-, \hat{\rho}_{001}^-, \hat{\rho}_{010}^-, \hat{\rho}_{011}^-$ occurring are given by

$p_{000}^-, p_{001}^-, p_{010}^-, p_{011}^-$, respectively, such that

$$p_{0ij}^- := P(\hat{\rho}_{0ij} | \hat{\pi}_- \otimes \mathbb{1}). \quad (\text{B1})$$

Using Bayes' theorem, this can be written as

$$\begin{aligned} p_{0ij}^- &= \frac{P(\hat{\pi}_- \otimes \mathbb{1} | \hat{\rho}_{0ij}) P(\hat{\rho}_{0ij})}{P(\hat{\pi}_- \otimes \mathbb{1})} \\ &= \frac{P(\hat{\pi}_- | \hat{\rho}_{0i})}{4P(\hat{\pi}_-)}, \end{aligned} \quad (\text{B2})$$

where the second equality is obtained using the fact that $P(\hat{\rho}_{0ij}) = 1/4$, and that the third qubit is acted on only by the identity and therefore does not change any of the probabilities.

So, by noting that

$$\begin{aligned} P(\hat{\pi}_- | \hat{\rho}_{00}) &= \text{Tr}(\hat{\pi}_- \hat{\rho}_{00}) = \beta, \\ P(\hat{\pi}_- | \hat{\rho}_{01}) &= \text{Tr}(\hat{\pi}_- \hat{\rho}_{01}) = \frac{1}{4}(\alpha + 4\beta), \end{aligned} \quad (\text{B3})$$

and therefore

$$\begin{aligned} P(\hat{\pi}_-) &= P(\hat{\pi}_- | \hat{\rho}_{00}) P(\hat{\rho}_{00}) + P(\hat{\pi}_- | \hat{\rho}_{01}) P(\hat{\rho}_{01}) \\ &= \frac{1}{8}(\alpha + 8\beta), \end{aligned} \quad (\text{B4})$$

we find that

$$p_{000}^- = p_{001}^- = \frac{2\beta}{\alpha + 8\beta}, \quad (\text{B5a})$$

$$p_{010}^- = p_{011}^- = \frac{\alpha + 4\beta}{2(\alpha + 8\beta)}. \quad (\text{B5b})$$

2. Second measurement

Again, assuming the outcome of the first classification was $-$, recall that distinguishing $\hat{\rho}_{000}^-$ from the other states is done optimally by letting $\hat{\pi}_{000}^-$ be the projector onto the $s = 3/2$ space. This leads to the measurement that best distinguishes $\hat{\rho}_{001}^-, \hat{\rho}_{010}^-, \hat{\rho}_{011}^-$ being entirely contained in the $s = 1/2$ space $(\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}) \otimes (\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}})^*$. Further, since the $\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{Q}_{\frac{1}{2}}^*$ component of each of $\hat{\rho}_{001}^-, \hat{\rho}_{010}^-, \hat{\rho}_{011}^-$ is the identity, all the information about how they differ is contained in $\mathcal{P}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}^*$. So we can rephrase this as a state discrimination problem of the following states:

$$|\psi_{001}^-\rangle = \frac{\mathcal{N}_{001}^-}{\sqrt{6}} |1\rangle, \quad (\text{B6a})$$

$$|\psi_{010}^-\rangle = \frac{\mathcal{N}_{010}^-}{\sqrt{6(\alpha + 4\beta)}} (\sqrt{\beta} |1\rangle - \sqrt{3(\alpha + \beta)} |0\rangle), \quad (\text{B6b})$$

$$|\psi_{011}^-\rangle = \frac{\mathcal{N}_{011}^-}{\sqrt{6(\alpha + 4\beta)}} (\sqrt{\beta} |1\rangle + \sqrt{3(\alpha + \beta)} |0\rangle), \quad (\text{B6c})$$

where \mathcal{N}_{0ij}^- are normalization constants required so that we can think of this as a mirror-symmetric-state discrimination problem. Explicitly,

$$\begin{aligned} \mathcal{N}_{001}^- &= \sqrt{6}, \\ \mathcal{N}_{010}^- &= \sqrt{\frac{6(\alpha + 4\beta)}{4\beta + 3\alpha}} = \mathcal{N}_{011}^-. \end{aligned} \quad (\text{B7})$$

Now, in [28], the states to be discriminated were written as

$$|\psi_1\rangle = |1\rangle, \quad (\text{B8a})$$

$$|\psi_2\rangle = \cos\theta|1\rangle - \sin\theta|0\rangle, \quad (\text{B8b})$$

$$|\psi_3\rangle = \cos\theta|1\rangle + \sin\theta|0\rangle, \quad (\text{B8c})$$

such that $|\psi_{2,3}\rangle$ happen with probability $p_{2,3} = p$ and $|\psi_1\rangle$ with probability $p_1 = 1 - 2p$. So, we can let

$$\begin{aligned} \cos\theta &= \sqrt{\frac{\beta}{4\beta + 3\alpha}}, \\ \sin\theta &= \sqrt{\frac{3(\alpha + \beta)}{4\beta + 3\alpha}}, \end{aligned} \quad (\text{B9})$$

and $q_{010}^-, q_{011}^- = p^-, q_{001}^- = 1 - 2p^-$ where

$$q_{0ij}^- = P(\hat{\rho}_{0ij}|\hat{\pi}_-, s = 1/2) \quad (\text{B10})$$

is the probability of being in the state $|\psi_{0ij}^-\rangle$, and we add (with respect to [28]) a superscript to p to distinguish the two outcomes of the intermediate measurements.

So, using Eq. (B10), we can find $p^- = q_{0ij}^-$. Using Bayes' theorem, we find that

$$p^- = \frac{P(\hat{\pi}_- \otimes \mathbb{1}, s = \frac{1}{2}|\hat{\rho}_{0ij})P(\hat{\rho}_{0ij})}{P(\hat{\pi}_- \otimes \mathbb{1}, s = \frac{1}{2})}. \quad (\text{B11})$$

Note that requiring $s = 1/2$ is equivalent to projecting the state $\hat{\rho}_{0ij}$ onto the $s = 1/2$ space $(\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}}) \otimes (\mathcal{Q}_{\frac{1}{2}} \otimes \mathcal{P}_{\frac{1}{2}})^*$. Denoting this projector by $\hat{P}_{\frac{1}{2}}$,

$$\begin{aligned} P\left(\hat{\pi}_- \otimes \mathbb{1}, s = \frac{1}{2}|\hat{\rho}_{0ij}\right) &= P(\hat{\pi}_- \otimes \mathbb{1}, \hat{P}_{\frac{1}{2}}|\hat{\rho}_{0ij}) \\ &= \text{Tr}[\hat{P}_{\frac{1}{2}}(\sqrt{\hat{\pi}_-} \otimes \mathbb{1})\hat{\rho}_{0ij}(\sqrt{\hat{\pi}_-} \otimes \mathbb{1})\hat{P}_{\frac{1}{2}}]. \end{aligned} \quad (\text{B12})$$

The denominator can be found using

$$P\left(\hat{\pi}_- \otimes \mathbb{1}, s = \frac{1}{2}\right) = \sum_{ij} P(\hat{\pi}_- \otimes \mathbb{1}, \hat{P}_{\frac{1}{2}}|\hat{\rho}_{0ij})P(\hat{\rho}_{0ij}), \quad (\text{B13})$$

from which it follows that

$$p^- = \frac{3\alpha + 4\beta}{6(\alpha + 2\beta)}. \quad (\text{B14})$$

Now, according to [28], if

$$p \geq \frac{1}{2 + \cos\theta(\cos\theta + \sin\theta)}, \quad (\text{B15})$$

$a = 1$. Otherwise,

$$a = \frac{p \cos\theta \sin\theta}{1 - p(2 + \cos^2\theta)}. \quad (\text{B16})$$

Substituting p^- for p , a_- for a and our expressions for $\sin\theta, \cos\theta$ given in Eq. (B9), this can be restated in the following way: if

$$\alpha \geq 2\beta, \quad (\text{B17})$$

$a_- = 1$. Otherwise,

$$a_- = \sqrt{\frac{\alpha + \beta}{3\beta}}. \quad (\text{B18})$$

When coupled with the constraints on α, β given in Eq. (26), we obtain

$$a_- = \begin{cases} \sqrt{\frac{\alpha + \beta}{3\beta}} & \text{if } \alpha \in [0, \min\{1 - \beta, 2\beta\}], \\ 1 & \text{if } \alpha \in (2\beta, 1 - \beta] \text{ with } 2\beta < 1 - \beta, \end{cases} \quad (\text{B19})$$

such that $\beta \in [0, 1]$.

-
- [1] V. Dunjko and H. J. Briegel, Machine learning & artificial intelligence in the quantum domain: A review of recent progress, *Rep. Prog. Phys.* **81**, 074001 (2018).
- [2] J. A. Bergou and M. Hillery, Universal Programmable Quantum State Discriminator that is Optimal for Unambiguously Distinguishing between Unknown States, *Phys. Rev. Lett.* **94**, 160501 (2005).
- [3] M. Guță and W. Kottłowski, Quantum learning: Asymptotically optimal classification of qubit states, *New J. Phys.* **12**, 123032 (2010).
- [4] M. Sasaki, A. Carlini, and R. Jozsa, Quantum template matching, *Phys. Rev. A* **64**, 022317 (2001).
- [5] G. Sentís, A. Monràs, R. Muñoz-Tapia, J. Calsamiglia, and E. Bagan, Unsupervised Classification of Quantum Data, *Phys. Rev. X* **9**, 041029 (2019).
- [6] D. Akimoto and M. Hayashi, Discrimination of the change point in a quantum setting, *Phys. Rev. A* **83**, 052328 (2011).
- [7] G. Sentís, E. Bagan, J. Calsamiglia, G. Chiribella, and R. Muñoz-Tapia, Quantum Change Point, *Phys. Rev. Lett.* **117**, 150502 (2016).
- [8] W. K. Wootters and W. H. Zurek, A single quantum cannot be cloned, *Nature (London)* **299**, 802 (1982).
- [9] D. Dieks, Communication by EPR devices, *Phys. Lett. A* **92**, 271 (1982).
- [10] C. A. Fuchs and A. Peres, Quantum-state disturbance versus information gain: Uncertainty relations for quantum information, *Phys. Rev. A* **53**, 2038 (1996).
- [11] A. Monras, G. Sentís, and P. Wittek, Inductive Supervised Quantum Learning, *Phys. Rev. Lett.* **118**, 190503 (2017).
- [12] A. Peres and W. K. Wootters, Optimal Detection of Quantum Information, *Phys. Rev. Lett.* **66**, 1119 (1991).
- [13] S. Massar and S. Popescu, Optimal Extraction of Information from Finite Quantum Ensembles, *Phys. Rev. Lett.* **74**, 1259 (1995).
- [14] J. Calsamiglia, J. I. de Vicente, R. Muñoz-Tapia, and E. Bagan, Local Discrimination of Mixed States, *Phys. Rev. Lett.* **105**, 080504 (2010).
- [15] E. Chitambar and M.-H. Hsieh, Revisiting the optimal detection of quantum information, *Phys. Rev. A* **88**, 020302(R) (2013).
- [16] B. L. Higgins, A. C. Doherty, S. D. Bartlett, G. J. Pryde, and H. M. Wiseman, Multiple-copy state discrimination: Thinking globally, acting locally, *Phys. Rev. A* **83**, 052314 (2011).
- [17] S. Croke and S. M. Barnett, Difficulty of distinguishing product states locally, *Phys. Rev. A* **95**, 012337 (2017).

- [18] J. Bergou, E. Feldman, and M. Hillery, Extracting Information from a Qubit by Multiple Observers: Toward a Theory of Sequential State Discrimination, *Phys. Rev. Lett.* **111**, 100501 (2013).
- [19] M. Hillery and J. Mimih, Sequential discrimination of qudits by multiple observers, *J. Phys. A: Math. Theor.* **50**, 435301 (2017).
- [20] D. Fields, R. Han, M. Hillery, and J. A. Bergou, Extracting unambiguous information from a single qubit by sequential observers, *Phys. Rev. A* **101**, 012118 (2020).
- [21] L. Leppäjärvi and M. Sedlák, Postprocessing of quantum instruments, *Phys. Rev. A* **103**, 022615 (2021).
- [22] S. Barnett, *Quantum Information* (Oxford University Press, Oxford, 2009), Vol. 16.
- [23] *States, Effects, and Operations: Fundamental Notions of Quantum Theory*, edited by K. Kraus, A. Böhm, J. D. Dollard, and W. H. Wootters, Lecture Notes in Physics (Springer-Verlag, Berlin, Heidelberg, 1983), Vol. 190.
- [24] H. Barnum, [arXiv:quant-ph/0205155](https://arxiv.org/abs/quant-ph/0205155).
- [25] D. Bacon, I. L. Chuang, and A. W. Harrow, *The Quantum Schur and Clebsch-Gordan Transforms: I. Efficient Qudit Circuits* (Society for Industrial and Applied Mathematics, New Orleans, 2007), p. 1235–1244.
- [26] S. M. Barnett, A. Chefles, and I. Jex, Comparison of two unknown pure quantum states, *Phys. Lett. A* **307**, 189 (2003).
- [27] M. A. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (American Association of Physics Teachers, College Park, MD, 2002).
- [28] E. Andersson, S. M. Barnett, C. R. Gilson, and K. Hunter, Minimum-error discrimination between three mirror-symmetric states, *Phys. Rev. A* **65**, 052308 (2002).