

**Optimality of spatial search in graphs with infinite tail**

Weichen Xie\*

*Department of Mathematics, Clarkson University, Potsdam, New York 13699-5815, USA*Christino Tamon<sup>†</sup>*Department of Computer Science, Clarkson University, Potsdam, New York 13699-5815, USA*

(Received 23 January 2023; accepted 13 March 2023; published 20 March 2023)

Farhi and Gutmann [*Phys. Rev. A* **57**, 2403 (1998)] proved that a continuous-time analog of Grover search (also called spatial search) is optimal on complete graphs. We extend this result by showing that spatial search remains optimal on a complete graph even in the presence of an infinitely long path (or tail). If we view the latter as an external quantum system that has a limited but nontrivial interaction with our finite quantum system, this suggests that spatial search is robust against a coherent infinite one-dimensional probe. Moreover, we show that the search algorithm is *oblivious* in that it does not need to know whether the tail is present or not, and if so, where it is attached to.

DOI: [10.1103/PhysRevA.107.032416](https://doi.org/10.1103/PhysRevA.107.032416)**I. INTRODUCTION**

The celebrated quantum search algorithm of Grover [1] provides a provable quadratic speedup over any classical algorithm. Shortly thereafter, Farhi and Gutmann proposed an analog analogue of Grover's algorithm [2]. They defined the Grover search algorithm via a continuous-time quantum walk on a complete graph where the oracle or target vertex is marked by a suitably weighted self-loop. Remarkably, the Farhi-Gutmann algorithm achieved perfect fidelity on complete graphs of any size. In contrast, this property does not hold for Grover search (viewed as a discrete-time quantum walk) as it is inherently a bounded-error probabilistic algorithm.

This continuous-time search problem was later generalized by Childs and Goldstone [3] to arbitrary finite graphs, where it is known as the *spatial search* problem. A collection of different families of finite graphs had been studied in this context; for example, see Refs. [3–8]. But, to date, spatial search has not been studied on *infinite* graphs as it seems that the quantum walk will escape or diffuse to infinity before having a chance to localize on the marked vertex (or oracle). The goal of this work is to disabuse ourselves of this highly plausible intuition.

We consider infinite graphs which are obtained by attaching an infinite path (or tail) to a finite graph. This family of graphs with tails was explored by Golinskii [9]. In this work, we view the finite graph as our operational quantum system for performing quantum search and the tail as an external (possibly adversarial) infinite-dimensional quantum system which interacts with our finite system in a coherent manner.

The main question we explore in this work is this: Can spatial search still be performed optimally in the presence of an infinite-dimensional probe? We provide a positive answer to this question for complete graphs. This extends the result of Farhi-Gutmann [2] to the infinite setting. Moreover, the quantum search algorithm is *oblivious* as it does not need to know whether the infinite-dimensional probe is present (or not) and where it is attached (if present). Since we give our adversary the benefit of an infinite-dimensional quantum system, this serves only to strengthen the result.

Our technique relies on the theory of Jacobi operators (see Refs. [9,10]). The main idea is to decompose the adjacency operator  $A$  of our infinite graph using two pairwise orthogonal invariant subspaces (see Golinskii [9], Theorem 1.2) where the first one is finite dimensional while the second one is infinite dimensional. The next crucial observation is that spatial search takes place in the infinite-dimensional invariant subspace of  $A$ . Moreover, the action of  $A$  in this infinite-dimensional invariant subspace is given by a finite rank Jacobi matrix  $J$ . Finally, we show that the initial and target states of the spatial search are nearly confined in a two-dimensional subspace spanned by two *bound* states of  $J$ . This yields the claimed spatial search result. To the best of our knowledge, this is the first result which explores optimal spatial search on infinite graphs.

As outlined above, our argument is the standard argument for showing optimal spatial search in the finite setting (see Refs. [4,11,12]). Namely, we show that the initial and target states are spanned by two distinct eigenvectors of the perturbed adjacency matrix. A contribution of this work is to show that this argument holds in the infinite setting via bound states of the reduced adjacency operator. We believe that this argument might be useful in other settings.

Following a brief discussion of basic notation and terminology in Sec. II, we prove our main results in Secs. III and IV. Then, we justify the optimality of our search algorithm

\*xiew@clarkson.edu

†tino@clarkson.edu

in Sec. V. Finally, we conclude with some open questions in Sec. VI.

## II. PRELIMINARIES

We introduce the basic notation and terminology that we will use throughout. The set of all positive integers is denoted  $\mathbb{Z}^+$  and the set of all complex numbers with unit modulus is denoted  $U(1)$ . For vectors  $x, y$  where  $x = \zeta y$ , for some  $\zeta \in U(1)$ , we write  $x \equiv y$ . We adopt standard asymptotic notation:  $o(f_n)$  denotes any function  $g_n$  so that  $g_n/f_n \rightarrow 0$ ,  $O(f_n)$  denotes functions  $g_n$  for which  $g_n/f_n$  is bounded from above by a constant, and  $\Omega(f_n)$  denotes functions  $g_n$  where  $g_n/f_n$  is bounded from below by a constant, where in each case  $n \rightarrow \infty$ ; see Ref. [13]. In our case, the asymptotic parameter  $n$  corresponds to the size of a finite graph.

### A. Graphs and operators

We study undirected and connected graphs  $G = (V, E)$  with vertex set  $V$  and edge set  $E$ , respectively. The adjacency matrix  $A$  of  $G$  is a symmetric matrix whose  $(i, j)$  entry is 1 if  $(i, j) \in E$  and 0 otherwise. For a vertex  $u$ , let  $N(u) = \{v \in V : (u, v) \in E\}$  denote the set of neighbors of  $u$ . The degree of vertex  $u$ , denoted  $\deg(u)$ , is the cardinality of  $N(u)$ . The complete graph (or clique) on  $n$  vertices is denoted  $K_n$ . A rooted graph  $(G, r)$  is a graph  $G$  with a distinguished vertex  $r$  which we call the root. See Ref. [14] for further background on algebraic graph theory.

We allow countably infinite graphs, in which case  $V = \mathbb{Z}^+$  [15]. For example, the infinite path  $P_\infty$  has edges which are consecutive positive integers; its adjacency matrix is known as the free Jacobi matrix [9]. Related to these graphs, we associate a complex separable Hilbert space  $\mathcal{H} = \ell^2(V)$  equipped with the inner product  $\langle x, y \rangle = \sum_{u \in V} \bar{x}_u y_u$ , for vectors  $x, y \in \mathcal{H}$ . For  $\ell^2(V)$ , a standard basis is  $\{e_u : u \in V\}$ , where  $e_u$  is the unit vector corresponding to vertex  $u$ . An infinite graph  $\mathcal{G}$ , the adjacency operator  $A$  is a linear operator that maps the standard basis vector  $e_v$  to the vector associated with the neighboring vertices  $N(v)$ ; that is,  $Ae_v = \sum_{u \in N(v)} a_{u,v} e_u$ , or simply  $a_{u,v} = \langle e_u, Ae_v \rangle$ . Let  $\deg(\mathcal{G}) = \sup\{\deg(u) : u \in V\}$ . If  $\deg(\mathcal{G}) < \infty$ , then the adjacency operator  $A$  is a bounded self-adjoint operator [16].

The *spectrum* of a linear operator  $A$  is the set  $\sigma(A)$  of all complex numbers  $\lambda$  where  $\lambda I - A$  is not invertible. For a bounded and self-adjoint operator  $A$ , its spectrum can be classified further into the *point* spectrum  $\sigma_p(A)$  and the *continuous* spectrum  $\sigma_c(A)$ . The point spectrum consists of all eigenvalues  $\lambda \in \mathbb{C}$  of  $A$  such that  $Ax = \lambda x$  for some nonzero  $x \in \ell^2(V)$ . On the contrary, the values in  $\sigma_c(A)$  are not eigenvalues of  $A$  and have no corresponding eigenvectors in  $\ell^2(V)$ .

The spectral theorem [17,18] states that for a bounded self-adjoint operator  $A$  on a complex Hilbert space  $\mathcal{H}$ , there exists a unique resolution of the identity  $E$  on the Borel subsets of  $\sigma(A)$  so that  $A = \int_{\sigma(A)} \lambda dE(\lambda)$ . Moreover, if  $f$  is a bounded Borel function on  $\sigma(A)$ , then  $f(A) = \int_{\sigma(A)} f(\lambda) dE(\lambda)$ . We also use a decomposition induced by invariant subspaces (see Ref. [17], Theorem 3, Sec. 40) which states that if  $W_k$  ( $k =$




FIG. 1. Optimal spatial search on the infinite lollipop with the oracle hiding at vertex 1.

$1, 2, \dots, m)$  are pairwise orthogonal invariant subspaces of  $A$ , that is,  $\mathcal{H} = \bigoplus_{k=1}^m W_k$  and  $AW_k \subset W_k$ , for each  $k$ , then  $A = \sum_{k=1}^m A_k P_k$ , where  $P_k$  is the projection on  $W_k$  and  $A_k$  is the restriction of  $A$  to  $W_k$ .

### B. Spatial search

A continuous-time quantum walk on an infinite graph  $\mathcal{G}$  with bounded self-adjoint adjacency operator  $A$  is given by the unitary operator  $e^{-itA}$  [acting on the Hilbert space  $\ell^2(V)$ ]. Our focus is on infinite graphs obtained from a finite connected rooted graph  $(G, r)$  by attaching an infinite path  $P_\infty$  at the root vertex  $r$ ; denote the resulting infinite graph as  $\mathcal{G} = G(r, P_\infty)$ . As we use  $\{1, 2, \dots, n\}$  to label the vertices of a finite rooted graph of order  $n$ , we take the liberty to designate the last vertex  $n$  as the root (without loss of generality). These are the graphs with tails studied by Golinskii [9].

We say the infinite graph  $\mathcal{G} = G_n(P_\infty)$  has *optimal spatial search* (adopting Ref. [4]) if there is a real  $\gamma > 0$  so that for each vertex  $w$  of  $G_n$ , a continuous-time quantum walk on  $\mathcal{G}$  with a self-loop on  $w$  of weight  $\gamma$  will unitarily map the principal eigenvector  $z_1$  of  $G_n$  to the unit vector  $e_w$  with constant fidelity in time  $t = O(1/\epsilon_1)$ , where  $\epsilon_1 = |\langle e_w, z_1 \rangle|$ . That is,

$$|\langle e_w, e^{-it(A+\gamma P_w)} z_1 \rangle| = \Omega(1),$$

where  $A$  is the adjacency operator of  $\mathcal{G}$  and  $P_w$  is the projection onto the subspace spanned by  $e_w$ . It is customary to assume  $\epsilon_1 = o(1)$ , as otherwise we already have a constant overlap between the target state  $e_w$  and the initial state  $z_1$ .

## III. INFINITE LOLLIPOP IS OPTIMAL

For  $n \geq 2$ , consider the infinite *lollipop* graph  $\mathcal{L}_n = K_n(P_\infty)$ . The vertices of the infinite path  $P_\infty$  are labeled with  $n+1, n+2, \dots$ . To this lollipop graph  $\mathcal{L}_n$ , we place a self-loop of weight  $\gamma$  at vertex 1 (the oracle or target vertex) and denote the resulting infinite graph as  $\mathcal{L}_n(\gamma)$ . Please see Fig. 1.

The adjacency operator  $H$  of  $\mathcal{L}_n(\gamma)$  is obviously a bounded self-adjoint linear operator on  $\ell^2(\mathbb{Z}^+)$ . It can be written as an

infinite dimensional matrix under the standard basis  $\{e_j\}_{j=1}^\infty$  as

$$H = \begin{pmatrix} \gamma & 1 & \cdots & 1 \\ 1 & 0 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 0 & 1 \\ & & & 1 & 0 & 1 \\ & & & & 1 & 0 & 1 \\ & & & & & \ddots & \ddots & \ddots \end{pmatrix}.$$

Our goal is to prove the following infinite analog of the Farhi-Gutmann result.

*Theorem 1.* For  $\gamma = n + O(1)$ , we have

$$|\langle e_1, e^{-itH} z_1 \rangle| = \Omega(1),$$

for  $t = \pi/2\sqrt{n}$ , where  $z_1 = n^{-1/2} \sum_{j=1}^n e_j$ .

By a theorem of Golinskii (Ref. [9], Theorem 1.2), using a change of basis,  $H$  can be written in a block diagonal form. This new basis  $\{\tilde{e}_k\}_{k=1}^\infty$  is defined as follows. Let

$$\tilde{e}_k = e_k, \quad k = n, n + 1, \dots \tag{1}$$

and

$$\tilde{e}_{n-1} = \frac{1}{\sqrt{n-1}} \sum_{j=1}^{n-1} e_j. \tag{2}$$

The next orthogonal basis vector is then given by

$$\tilde{e}'_{n-2} = H\tilde{e}_{n-1} - \langle \tilde{e}_{n-1}, H\tilde{e}_{n-1} \rangle \tilde{e}_{n-1} - \langle \tilde{e}_n, H\tilde{e}_{n-1} \rangle \tilde{e}_n,$$

which, after normalization, yields

$$\tilde{e}_{n-2} = \frac{1}{\sqrt{(n-2)(n-1)}} [(n-1)e_1 - \tilde{e}_{n-1}]. \tag{3}$$

We take the remaining basis vectors to be the nonprincipal columns of the Fourier matrix of order  $n - 2$ . In particular, the basis vector  $\tilde{e}_k$  is defined as

$$\tilde{e}_k = \frac{1}{\sqrt{n-2}} \sum_{j=1}^{n-2} e^{\frac{2\pi i(j-1)k}{(n-2)}} e_j \quad (k = 1, \dots, n-3).$$

Since  $H\tilde{e}_{n-2} = (\gamma \frac{n-2}{n-1} - 1)\tilde{e}_{n-2} + \gamma \frac{\sqrt{n-2}}{n-1} \tilde{e}_{n-1}$ , the subspace  $\mathcal{S} = \text{span}\{\tilde{e}_{n-2}, \tilde{e}_{n-1}, \dots\}$  is  $H$  invariant. Under the new basis, the adjacency operator  $H$  becomes

$$H = \begin{pmatrix} -I_{n-2} & O \\ O & \hat{H} \end{pmatrix}$$

As will be clear soon, it suffices for us to restrict our focus on the operator  $\hat{H}$ , which is the operator  $H$  restricted to the subspace  $\mathcal{S} = \text{span}\{\tilde{e}_{n-2}, \tilde{e}_{n-1}, \dots\}$ . This is because the initial state  $z_1$  and the target state  $e_1$  of our spatial search problem both have nonnegligible overlap with  $\mathcal{S}$ .

TABLE I. For  $\gamma = n + O(1)$  and sufficiently large  $n$ , there are four sign changes of  $y_0(x)$ .

$x$	$-\infty$	$-1$	$\frac{1}{n}$	$1$	$+\infty$
$\gamma \sqrt{\frac{n-2}{n-1}} y_0$	$-\infty$	$\gamma + 1 > 0$	$-\frac{1}{n} + O(\frac{1}{n^2}) < 0$	$2n\gamma + O(n) > 0$	$-\infty$

It follows from the preceding analysis that the operator  $H$  under the basis  $\{\tilde{e}_{n-2}, \tilde{e}_{n-1}, \dots\}$  is given by

$$\hat{H} = \begin{pmatrix} \tilde{e}_{n-2} \\ \tilde{e}_{n-1} \\ \tilde{e}_n \\ \tilde{e}_{n+1} \\ \vdots \end{pmatrix} \begin{pmatrix} \gamma \frac{n-2}{n-1} - 1 & \gamma \frac{\sqrt{n-2}}{n-1} & & & \\ \gamma \frac{\sqrt{n-2}}{n-1} & n-2 + \frac{\gamma}{n-1} & \sqrt{n-1} & & \\ & \sqrt{n-1} & 0 & 1 & \\ & & 1 & 0 & 1 \\ & & & \ddots & \ddots & \ddots \end{pmatrix}. \tag{4}$$

This symmetric tridiagonal matrix is an *eventually free* or *finite rank* Jacobi matrix [9] whose full spectrum can be computed via the so-called *Jost* solution (see Ref. [10], Appendix). The Jost solution is a vector  $y(x) = [y_1(x), y_2(x), \dots]^T$  that satisfies the eigenequation of  $\hat{H}$  with eigenvalue of the form  $x + \frac{1}{x}$ , namely,

$$\hat{H}y(x) = \left(x + \frac{1}{x}\right)y(x), \tag{5}$$

and also the degree condition

$$\lim_{k \rightarrow \infty} x^{-k} y_k(x) = 1.$$

Given the special form of  $\hat{H}$ , we can set

$$y_k(x) = x^k, \quad k = 3, 4, \dots, \tag{6}$$

and use (5) to get the Jost polynomial

$$y_0(x) = \frac{1}{\gamma \sqrt{\frac{n-1}{n-2}}} \{ (2-n)x^4 + [(n-3)\gamma + 4 - 2n]x^3 + [(n-3)\gamma + 5 - 2n]x^2 + [3 - n - \gamma]x + 1 \}. \tag{7}$$

In order to compute the eigenvalues of  $\hat{H}$ , we need the following spectral theorem for finite rank Jacobi operators. We will only need information about the point spectrum, as will be clear soon.

*Theorem 2.* (Ref. [9], p. 8) Let  $J$  be an eventually free Jacobi matrix and  $y_0(x)$  be its Jost function. Then all roots of  $y_0(x)$  in the complex unit disk are real and simple,  $y_0(0) \neq 0$ . A real number  $\lambda_j$  is an eigenvalue of  $J$  if and only if

$$\lambda_j = x_j + \frac{1}{x_j}, \quad x_j \in (-1, 1), \quad y_0(x_j) = 0.$$

By choosing  $\gamma = n + O(1)$ , the roots of  $y_0(x)$  in the complex unit disk can be approximated consecutively. First note that there are four real roots for  $y_0(x)$  and two of them lie in the unit disk as indicated by Table I.

Denote the two roots within the unit disk as  $x_{\pm} = \frac{1}{n} + \delta_{\pm}$ . Hence,

$$0 = \gamma \sqrt{\frac{n-2}{n-1}} y_0(x_{\pm}) = -\frac{1}{n} + n^2 \delta_{\pm}^2 + o\left(\frac{1}{n}\right) + o(n^2 \delta_{\pm}^2).$$

In order for the left-hand side to attain 0 exactly, for all  $n$ , at least the two highest-order terms on the right-hand side should cancel perfectly; that is,  $o\left(\frac{1}{n}\right) = -\frac{1}{n} + n^2 \delta_{\pm}^2$ , which implies

$$x_{\pm} = \frac{1}{n} \pm \frac{1}{n^{3/2}} + o\left(\frac{1}{n^{3/2}}\right). \tag{8}$$

Therefore, the two distinct eigenvalues of  $\widehat{H}$  are given by

$$\lambda_{\pm} = n \pm \sqrt{n} + O(1). \tag{9}$$

The corresponding eigenvectors (or bound states) are given by  $y_{\pm} = [y_1(x_{\mp}), y_2(x_{\mp}), \dots]^T$ , where the entries are defined by the Jost polynomials

$$y_1(x_{\mp}) = \pm \frac{1}{\gamma} \sqrt{\frac{n-1}{n-2}} \times \frac{1}{n^{3/2}} + O\left(\frac{1}{n^3}\right), \tag{10}$$

$$y_2(x_{\mp}) = \frac{1}{\sqrt{n-1}} \times \frac{1}{n^2} \mp \frac{1}{\sqrt{n-1}} \frac{2}{n^{5/2}} + O\left(\frac{1}{n^{7/2}}\right), \tag{11}$$

$$y_k(x_{\mp}) = \frac{1}{n^k} \mp \frac{k}{n^{(2k+1)/2}} + O\left(\frac{1}{n^{k+1}}\right), \quad k = 3, 4, \dots \tag{12}$$

Now, we are ready to prove Theorem 1.

First, note that both  $e_1 = \sqrt{\frac{1}{n-1}} \tilde{e}_{n-2} + \sqrt{\frac{n-2}{n-1}} \tilde{e}_{n-1}$  and  $z_1 = \tilde{e}_{n-1}$  are completely in the invariant subspace  $\mathcal{S}$ . Thus, we can restrict our unitary evolution to  $\mathcal{S}$ . Moreover, as  $e_1$  overlaps almost completely with  $\tilde{e}_{n-1}$ , it suffices to consider  $\tilde{e}_{n-1}$  as the target state. Hence, the fidelity can be further approximated as

$$|\langle e_1, e^{-itH} z_1 \rangle| \geq \sqrt{\frac{n-2}{n-1}} |\langle \bar{e}_1, e^{-it\widehat{H}} \bar{e}_2 \rangle| + O\left(\frac{1}{n^{1/2}}\right),$$

where  $\bar{e}_k$  is a unit vector which is the  $k$ th basis vector for the invariant subspace  $\mathcal{S} = \text{span}\{\tilde{e}_{n-2}, \tilde{e}_{n-1}, \dots\}$ .

Notice that

$$\lim_{n \rightarrow \infty} \frac{\langle \bar{e}_k, y_+ \rangle \langle y_+, \bar{e}_k \rangle}{\langle y_+, y_+ \rangle} + \frac{\langle \bar{e}_k, y_- \rangle \langle y_-, \bar{e}_k \rangle}{\langle y_-, y_- \rangle} = 1, \quad k = 1, 2,$$

that is, both the initial and the target states lie in the two-dimensional invariant subspace spanned by the eigenvectors  $y_+$  and  $y_-$ . Thus, it again suffices to consider the fidelity in this subspace.

Straightforward calculation shows that when  $\gamma = n + O(1)$ , the fidelity satisfies  $|\langle e_1, e^{-itH} z_1 \rangle| = 1 + o(1)$  for time  $t = \frac{\pi}{2\sqrt{n}}$ . If we normalize the adjacency matrix of  $K_n$ , we obtain  $O(\sqrt{n})$  time (matching Grover search).

### IV. ORACLE AT THE EDGE OF INFINITY

In this section, we show that even when the oracle is placed at the attachment vertex of the infinite path, spatial search remains optimal on  $\mathcal{L}_n$ . See Fig. 2.

Together with Sec. III, this will show that the search algorithm is oblivious, as it does not need to know if the external




FIG. 2. Optimal spatial search still occurs on the infinite lollipop even if the oracle sits at gateway vertex  $n$ .

probe (tail) is present or not. This is because the two cases surprisingly require the same asymptotic time for optimal spatial search. The claim follows from a similar analysis as before.

The adjacency operator of  $\mathcal{L}_n(\gamma)$  is given as

$$H = \begin{pmatrix} 0 & 1 & \dots & 1 & & & & & & \\ 1 & 0 & \dots & 1 & & & & & & \\ \vdots & \vdots & \ddots & \vdots & & & & & & \\ 1 & 1 & \dots & \gamma & 1 & & & & & \\ & & & 1 & 0 & 1 & & & & \\ & & & & 1 & 0 & 1 & & & \\ & & & & & \ddots & \ddots & \ddots & & \end{pmatrix}.$$

First, we restrict our focus to the invariant subspace  $\mathcal{S} = \text{span}\{\tilde{e}_{n-1}, \tilde{e}_n, \dots\}$  since  $e_n = \tilde{e}_n \in \mathcal{S}$  and  $z_1 = \sqrt{\frac{n-1}{n}} \tilde{e}_{n-1} + \sqrt{\frac{1}{n}} \tilde{e}_n \in \mathcal{S}$ , where  $\tilde{e}_{n-1} = \frac{1}{\sqrt{n-1}} \sum_{j=1}^{n-1} e_j$  and  $\tilde{e}_k = e_k$ ,  $k \geq n$ . Under the new basis that spans  $\mathcal{S}$ , the operator  $H$  is a rank-2 Jacobi matrix

$$\widehat{H} = \begin{matrix} \tilde{e}_{n-1} \\ \tilde{e}_n \\ \tilde{e}_{n+1} \\ \vdots \end{matrix} \begin{pmatrix} n-2 & \sqrt{n-1} & & \\ \sqrt{n-1} & \gamma & 1 & \\ & 1 & 0 & 1 \\ & & \ddots & \ddots & \ddots \end{pmatrix}. \tag{13}$$

Hence, the following Jost polynomials are obtained:

$$y_0(x) = \frac{1}{\sqrt{n-1}} [-\gamma x^3 + (n-2)(\gamma-1)x^2 + (2-n-\gamma)x + 1], \tag{14}$$

$$y_1(x) = \frac{1}{\sqrt{n-1}} (x - \gamma x^2), \tag{15}$$

$$y_k(x) = x^k, \quad k = 2, 3, \dots \tag{16}$$

By choosing  $\gamma = n + O(1)$ , we can compute the two distinct roots for  $y_0$  that are within the interval  $[-1, 1]$ ,

$$x_{\pm} = \frac{1}{n} \pm \frac{1}{n^{3/2}} + o\left(\frac{1}{n^{3/2}}\right), \tag{17}$$

which yields the corresponding eigenvalues for  $\widehat{H}$ ,

$$\lambda_{\pm} = n \pm \sqrt{n} + O(1). \tag{18}$$

As shown previously, the corresponding eigenvectors are defined by the values of Jost polynomials:

$$y_1(x_{\mp}) = \pm \frac{1}{\sqrt{n-1}} \times \frac{1}{n^{3/2}} + O\left(\frac{1}{n^{5/2}}\right), \quad (19)$$

$$y_2(x_{\mp}) = \frac{1}{n^2} \mp \frac{2}{n^{5/2}} + O\left(\frac{1}{n^3}\right), \quad (20)$$

$$y_k(x_{\mp}) = \frac{1}{n^k} \mp \frac{k}{n^{2k+1/2}} + O\left(\frac{1}{n^{k+1}}\right), \quad k = 3, 4, \dots \quad (21)$$

Following the same argument, we can see that for time  $t = \frac{\pi}{2\sqrt{n}}$ , the fidelity satisfies  $|\langle e_n, e^{-itH} z_1 \rangle| = 1 + o(1)$ .

## V. OPTIMALITY

We show that the time bound obtained in Theorem 1 (and in Sec. IV) is optimal. To this end, we generalize an argument of Farhi and Gutmann [2] to a class of infinite graphs with tails.

For a finite graph  $G_n$  on  $n$  vertices, take the cone  $\widehat{G}_n = K_1 + G_n$ , which is obtained by adding a new vertex (called the conical vertex) and connecting it to all vertices of  $G_n$ . Then, attach a tail to the conical vertex and denote this infinite graph as  $\widehat{G}_n(P_{\infty})$ . Notice that we recover the infinite lollipop when  $G_n$  is a clique.

*Theorem 3.* Let  $G_n$  be a  $(n, d)$ -regular graph, where  $d = \omega(\sqrt{n})$  for some  $\delta > 0$ , and let  $\mathcal{H}_0$  be the adjacency operator of  $\widehat{G}_n(P_{\infty})$ . Let  $z_1 \in \ell^2$  be natural embedding of the principal eigenvector of  $G_n$ . Suppose there is a time  $t_0$  so that for some  $\gamma \in \mathbb{R}$  and  $\zeta \in U(1)$ , and for a vertex  $w$  of  $G_n$ , we have

$$\|e^{-it_0(\mathcal{H}_0 + \gamma P_w)} z_1 - \zeta e_w\|^2 = o(1).$$

Then,  $\gamma t_0 = \Omega(1/\epsilon_1)$ , where  $\epsilon_1 = |\langle e_w, z_1 \rangle|$ , provided  $1/\gamma\epsilon_1 \in o(1)$ .

The largest eigenvalue in  $\sigma_p(\mathcal{H}_0)$  has a unique bound eigenstate  $\beta_1$  which satisfies  $\|\beta_1 - z_1\| = o(1)$ . This can be shown using similar techniques as in previous sections.

We call a time-dependent state  $\psi(t) \in \ell^2(\mathbb{Z}^+)$  exponentially decaying if there is a positive integer  $M$  so that for all  $m \geq M$  we have  $|\langle e_m, \psi(t) \rangle| = O(\kappa^m)$ , for some  $\kappa < 1$ , for all  $t$ .

*Proof.* (of Theorem 3) Let  $\mathcal{H}_w = \mathcal{H}_0 + \gamma P_w$ . It suffices to show the claim under the assumption  $\|e^{-it\mathcal{H}_w} \beta_1 - \zeta e_w\|^2 = o(1)$ , since  $\|\beta_1 - z_1\| = o(1)$  and by using the triangle inequality for squared norm (see Ref. [19], Eq. 18.5).

Following Ref. [2], we compare two Schrödinger evolutions given by

$$\begin{aligned} \psi'_w(t) &= -i\mathcal{H}_w \psi_w(t), & \psi_w(0) &= \beta_1, \\ \psi'_0(t) &= -i\mathcal{H}_0 \psi_0(t), & \psi_0(0) &= \beta_1. \end{aligned}$$

Note  $\psi_0(t) \equiv \beta_1$ , for all  $t$ , as  $\beta_1$  is an eigenstate of  $\mathcal{H}_0$ .

For simplicity, we assume that spatial search achieves perfect fidelity, namely,  $\psi_w(t_0) \equiv e_w$ . The general case is handled using triangle inequality for squared norm.

The key quantity is  $M(t) := \|\psi_w(t) - \psi_0(t)\|^2$ . First, notice that

$$M(t_0) = 2(1 - \text{Re}\langle e_w, \beta_1 \rangle) \geq 2(1 - \epsilon_1). \quad (22)$$

Furthermore, we have  $M'(t) = -2\text{Re}\langle \psi_w(t), \psi_0(t) \rangle'$ . Given that the inner product is an infinite series, the existence of its derivative requires uniform convergence. As  $\psi_0(t) \equiv \beta_1$  is exponentially decaying (by virtue of being a Jost solution), we have  $|\langle \psi_w(t) \rangle_m \langle \beta_1 \rangle_m| \leq |\langle \beta_1 \rangle_m|$  and

$$\sum_{m=1}^{\infty} |\langle \beta_1 \rangle_m| \leq C_M + \sum_{m \geq M} \kappa^m = C_M + \frac{\kappa^M}{1 - \kappa} < \infty$$

for a constant  $C_M$ . Thus,  $\langle \psi_w(t), \beta_1 \rangle$  is uniformly convergent (see Titchmarsh [13], 1.11).

Since  $\mathcal{H}_w$  is a finite-rank Jacobi matrix and  $\psi'_w(t) = -i\mathcal{H}_w \psi_w(t)$ , we see that  $\langle \psi'_w(t), \psi_0(t) \rangle$  is also uniformly convergent. This allows us to take the derivative of  $\langle \psi_w(t), \psi_0(t) \rangle$  by termwise differentiation (see Titchmarsh [13], 1.72), i.e.,

$$\begin{aligned} \langle \psi_w(t), \psi_0(t) \rangle' &= \langle \psi_w(t), \psi'_0(t) \rangle + \langle \psi'_w(t), \psi_0(t) \rangle \\ &= -i\langle \psi_w(t), \mathcal{H}_0 \psi_0(t) \rangle + i\langle \mathcal{H}_w \psi_w(t), \psi_0(t) \rangle. \end{aligned}$$

So, we obtain

$$M'(t) = 2\gamma \text{Im}\langle P_w \psi_w(t), \psi_0(t) \rangle \leq 2\gamma \|P_w \psi_w(t)\|.$$

Thus,  $M'(t) \leq 2\gamma\epsilon_1$ , which further implies

$$M(t_0) = \int_0^{t_0} M'(t) dt \leq 2\gamma\epsilon_1 t_0.$$

By combining the lower and upper bounds on  $M(t_0)$ , we get  $\gamma t_0 \geq (1 - \epsilon_1)/\epsilon_1 = \Omega(1/\epsilon_1)$  as  $\epsilon_1 = o(1)$ .

Theorem 3 justifies the optimal time  $1/\epsilon_1$  used to define spatial search. For the infinite lollipop, we have  $\epsilon_1 = 1/\sqrt{n}$ , and thus,  $\gamma t_0 = \Omega(\sqrt{n})$ . Since the search algorithm uses  $\gamma = n + O(1)$ , we get  $t_0 = \Omega(1/\sqrt{n})$ , which matches the bound achieved by Theorem 1. Hence, the algorithm is optimal.

## VI. CONCLUSION

In this work, we proved optimal spatial search occurs on cliques even in the presence of an infinite path. This generalized a known result of Farhi and Gutmann [2] to the infinite setting. We view this as a first step in showing that optimal spatial search is robust against an adversary modeled as an infinite-dimensional external quantum probe. Interesting directions for future work include extending the result to multiple tails or to tails induced by more general Jacobi matrices and strengthening the lower bound to other families of infinite graphs.

Our work was motivated by a question posed in Ref. [20]. We mention that discrete-time quantum walk on graphs with tails was studied in Ref. [21].

## ACKNOWLEDGMENTS

Work started during the workshop ‘‘Graph Theory, Algebraic Combinatorics, and Mathematical Physics’’ at Centre de Recherches Mathématiques (CRM), Université de Montréal. C.T. would like to thank CRM for its hospitality and support during his sabbatical visit. W.X. was supported by NSF Grant No. DMS-2212755. We thank Pierre-Antoine Bernard and Luc Vinet for discussions.

- [1] L. Grover, Quantum Mechanics Helps in Searching for a Needle in a Haystack, *Phys. Rev. Lett.* **79**, 325 (1997).
- [2] E. Farhi and S. Gutmann, Analog analogue of a digital quantum computation, *Phys. Rev. A* **57**, 2403 (1998).
- [3] A. Childs and J. Goldstone, Spatial search by quantum walk, *Phys. Rev. A* **70**, 022314 (2004).
- [4] S. Chakraborty, L. Novo, and J. Roland, Optimality of spatial search via continuous-time quantum walks, *Phys. Rev. A* **102**, 032214 (2020).
- [5] A. Glos and T. Wong, Optimal quantum walk search on Kronecker graphs with dominant or fixed regular initiators, *Phys. Rev. A* **98**, 062334 (2018).
- [6] J. Janmark, D. Meyer, and T. Wong, Global Symmetry is Unnecessary for Fast Quantum Search, *Phys. Rev. Lett.* **112**, 210502 (2014).
- [7] H. Tanaka, M. Sabri, and R. Portugal, Spatial search on Johnson graphs by continuous-time quantum walk, *Quantum Inf. Process.* **21**, 74 (2022).
- [8] T. Wong, Quantum walk search on Johnson graphs, *J. Phys. A: Math. Theor.* **49**, 195303 (2016).
- [9] L. Golinskii, Spectra of infinite graphs with tails, *Linear Multilinear A.* **64**, 2270 (2016).
- [10] D. Damanik and B. Simon, Jost functions and Jost solutions for Jacobi matrices, II. Decay and analyticity, *Int. Math. Res. Not.* **2006**, 19396 (2006).
- [11] S. Chakraborty, L. Novo, A. Ambainis, and Y. Omar, Spatial Search by Quantum Walk is Optimal for Almost all Graphs, *Phys. Rev. Lett.* **116**, 100501 (2016).
- [12] A. Chan, C. Godsil, C. Tamon, and W. Xie, Of shadows and gaps in spatial search, *Quantum Inf. Comput.* **22**, 1110 (2022).
- [13] E. C. Titchmarsh, *The Theory of Functions*, 2nd ed. (Oxford University Press, New York, 1939).
- [14] C. Godsil and G. Royle, *Algebraic Graph Theory* (Springer, New York, 2001).
- [15] B. Mohar and W. Woess, A survey on spectra of infinite graphs, *Bull. London Math. Soc.* **21**, 209 (1989).
- [16] B. Mohar, The spectrum of an infinite graph, *Linear Algebra Appl.* **48**, 245 (1982).
- [17] N. I. Akhiezer and I. M. Glazman, *Theory of Linear Operators in Hilbert Space* (Dover, New York, 2013).
- [18] W. Rudin, *Functional Analysis*, 2nd ed. (McGraw-Hill, New York, 1991).
- [19] S. Sternberg, *A Mathematical Companion to Quantum Mechanics* (Dover, New York, 2019).
- [20] P.-A. Bernard, C. Tamon, L. Vinet, and W. Xie, Quantum state transfer in graphs with tails, [arXiv:2211.14704](https://arxiv.org/abs/2211.14704).
- [21] N. Konno, E. Segawa, and M. Štefaňák, Relation between quantum walks with tails and quantum walks with sinks on finite graphs, [arXiv:2105.03111](https://arxiv.org/abs/2105.03111).