





Rényi entanglement entropy after a quantum quench starting from insulating states in a free boson system

Daichi Kagamihara ^{1,2,*}, Ryui Kaneko ^{1,*}, Shion Yamashika ², Kota Sugiyama,² Ryosuke Yoshii,³ Shunji Tsuchiya,² and Ippei Danshita ¹

¹*Department of Physics, Kindai University, Higashi-Osaka, Osaka 577-8502, Japan*

²*Department of Physics, Chuo University, Bunkyo, Tokyo 112-8551, Japan*

³*Center for Liberal Arts and Sciences, Sanyo-Onoda City University, Yamaguchi 756-0884, Japan*



(Received 25 July 2022; accepted 24 January 2023; published 6 March 2023)

We investigate the time-dependent Rényi entanglement entropy after a quantum quench starting from the Mott-insulating and charge-density-wave states in a one-dimensional free boson system. The second Rényi entanglement entropy is found to be the negative of the logarithm of the permanent of a matrix consisting of time-dependent single-particle correlation functions. From this relation and a permanent inequality, we obtain rigorous conditions for satisfying the volume-law entanglement growth. We also succeed in calculating the time evolution of the Rényi entanglement entropy in extremely large systems by brute-force computations of the permanent. We discuss possible applications of our findings to the real-time dynamics of noninteracting bosonic systems.

DOI: [10.1103/PhysRevA.107.033305](https://doi.org/10.1103/PhysRevA.107.033305)

I. INTRODUCTION

The concept of entanglement is indispensable for understanding quantum many-body physics these days. A pure quantum many-body state is entangled when it cannot be represented by a product state [1]. The entanglement entropy quantifies the degree of entanglement and is a valuable probe for characterizing states of quantum many-body systems. For example, in critical systems, the entanglement entropy exhibits the universal scaling with the size of a subsystem; the universal coefficient is determined by the corresponding conformal field theory [2–7]. Topologically ordered states, which cannot be described by conventional order parameters, would be characterized by the topological entanglement entropy [8–11].

The von Neumann entanglement entropy is a standard reference value to quantify the entanglement. When a system possesses a pure state $|\psi\rangle$ and can be divided into two subsystems A and B, the von Neumann entanglement entropy is defined as $S_{\text{vN}} = -\text{Tr}_A \hat{\rho}_A \ln \hat{\rho}_A$, where $\hat{\rho}_A = \text{Tr}_B \hat{\rho}$ is the reduced density matrix of $\hat{\rho} = |\psi\rangle\langle\psi|$ and $\text{Tr}_{A(B)}$ is the trace over the basis of subsystem A (B). The Rényi entanglement entropy is another quantity, which behaves similarly to the von Neumann entanglement entropy [12,13], and is defined as $S_\alpha = [\ln \text{Tr}_A(\hat{\rho}_A^\alpha)]/(1 - \alpha)$. The von Neumann entanglement entropy can be regarded as the $\alpha \rightarrow 1$ limit of the Rényi entanglement entropy.

The entanglement entropy is not merely an ideal quantity in theory, but it is also measurable experimentally. The protocol for measuring the Rényi entanglement entropy was first

proposed in 2012 [13,14]. In Ref. [13], the authors considered the real-time dynamics of cold atoms in optical lattices, which can be realized in experiments [15], and suggested preparing two copies of the same state. The Rényi entanglement entropy can be evaluated by controlling the tunnel coupling between these copies and by measuring the parity of the atom numbers. The second Rényi entanglement entropy has been indeed observed during a quench dynamics of one-dimensional (1D) cold atomic gases in an optical lattice [16,17].

Previous theoretical studies have actively discussed the dynamics of entanglement entropy after a quantum quench in connection with information propagation and thermalization [6,7,18–30]. While the dynamics of entanglement entropy in 1D lattice systems can be accurately analyzed by means of numerical methods based on matrix product states [20–25], the tractable time scale is rather limited in general due to the linear growth of the entanglement entropy in time [18]. In the case of 1D systems described by fermionic quasiparticles, such as the transverse-field Ising model, when the system is quenched to a noninteracting parameter region, long-time dynamics can be investigated analytically [18,29,31,32]. In particular, when the initial state is a Gaussian state, i.e., a ground or thermal state of some free (quadratic) Hamiltonian, the time-evolved state remains Gaussian. Then, the time evolution of the entanglement entropy can be evaluated from single-particle correlation functions. The fermionic Gaussian states include simple product states such as the Mott-insulating (MI) and charge-density-wave (CDW) states, which can also be prepared in experiments.

By contrast, studies of the entanglement growth during the quench dynamics for (soft-core) bosons are very few even in the case that the system is quenched to the noninteracting point. This is partly because a simple product state, which is often used as an initial state of quench dynamics in experiments [16,17], is not a Gaussian state for bosonic

*These authors contributed equally to this work.

†kagamihara@phys.kindai.ac.jp

‡rkaneko@phys.kindai.ac.jp

systems. Starting from product states such as the MI and CDW states, single-particle correlation functions are analytically calculable [33–35]. However, it is not straightforward to calculate the entanglement entropy from these correlation functions because a time-evolved state is not a Gaussian state.

This situation raises the following questions:

(i) Can we get an analytical form of the Rényi entanglement entropy concerning real-time dynamics of noninteracting bosonic systems?

(ii) Supposing we obtain the analytical form, can we rigorously obtain conditions under which the volume-law scaling is satisfied during the real-time evolution?

(iii) Can we numerically evaluate the Rényi entanglement entropy in systems much larger than the best currently available methods can handle?

To answer these questions, we take the 1D soft-core Bose-Hubbard model as the simplest playground. When the system is quenched to the noninteracting Hamiltonian, we obtain the analytical form of evaluating the second Rényi entanglement entropy. It can be expressed by the expectation value of the shift (SWAP) operator [13,14,16,17] and is given as the negative of the logarithm of the permanent of a time-dependent matrix consisting of single-particle correlation functions. We also give the condition for the volume-law scaling of the Rényi entanglement entropy by using a permanent inequality [36]. In addition, we obtain the long-time evolution of the Rényi entanglement entropy in extremely large systems by numerically computing the matrix permanent. Although direct calculations of the permanent require exponential-time cost in general, accessible sizes are found to be much larger than the exact diagonalization and matrix-product-state methods can deal with. Last but not least, we propose that the infinity norm of rows of the matrix consisting of the correlation function offers an entropy-density-like value and would give a practical bound for the Rényi entanglement entropy. This value would free us from exponential-time computations of the permanent, as long as we are interested in a qualitative behavior of the entanglement entropy growth rather than the value itself.

This paper is organized as follows: In Sec. II, we present the 1D Bose-Hubbard model and introduce two initial states for the quench dynamics. In Sec. III, we calculate the time-evolved states after the sudden quench and derive the analytical form of the Rényi entanglement entropy. In Sec. IV, we summarize some interesting properties of the matrix consisting of the correlation function introduced in Sec. III and show the condition for the volume-law scaling of the Rényi entanglement entropy. We describe some examples of the application of the condition to quench dynamics in our model. In Sec. V, we directly compute the permanent of the matrix consisting of the correlation function and obtain the time evolution of the Rényi entanglement entropy. We also compare our results with other reference values. In addition, we introduce an entropy-density-like value, which can be calculated in polynomial time, and discuss a bound for the Rényi entanglement entropy. In Sec. VI, we draw our conclusions and discuss possible applications to several problems on real-time dynamics of free boson systems. Throughout this paper, we

set $\hbar = 1$, take the lattice constant to be unity, and consider the zero-temperature dynamics, for simplicity.

II. ONE-DIMENSIONAL BOSE-HUBBARD MODEL

We consider the quench dynamics in the 1D Bose-Hubbard model under the open boundary condition. The Hamiltonian is defined as

$$\hat{H} = -J \sum_{j=1}^{L-1} (\hat{b}_j^\dagger \hat{b}_{j+1} + \text{H.c.}) + \sum_{j=1}^L \Omega_j \hat{n}_j + \frac{U}{2} \sum_{j=1}^L \hat{n}_j (\hat{n}_j - 1). \quad (1)$$

Here the symbols \hat{b}_j and \hat{n}_j denote the boson annihilation and number operators, respectively. The strength of the hopping and the interaction are given as J and U , respectively, and Ω_j denotes the single-particle potential. The number of sites is represented as L . This model quantitatively describes 1D Bose gases in optical lattices when the lattice potential is sufficiently deep.

We focus on the quench from insulating states to the noninteracting ($U = 0$) and homogeneous ($\Omega_j = 0$) point. As initial states, we specifically choose the MI state at unit filling, which is represented as

$$|\psi^{\text{MI}}\rangle = \prod_{j=1}^L \hat{b}_j^\dagger |0\rangle, \quad (2)$$

and the 010101... CDW state at half filling, which is described as

$$|\psi^{\text{CDW}}\rangle = \prod_{j=2,4,\dots}^L \hat{b}_j^\dagger |0\rangle, \quad (3)$$

where $|0\rangle$ is the vacuum state of \hat{b}_j and L is taken as an even number. The MI state is the ground state of the Bose-Hubbard model at unit filling for the large- U limit and can be prepared in experiments via a slow ramp-up of the optical lattice potential [16,17,35,37]. The CDW state is the ground state of the Bose-Hubbard model at half filling when $\Omega_j = \Omega(-1)^{j+1}$, $\Omega/J \gg 1$, and $U/J \gg 1$. It can be prepared in experiments with use of a secondary optical lattice whose lattice constant is twice as large as that of the primary lattice [38].

III. EVALUATING THE SECOND RÉNYI ENTANGLEMENT ENTROPY USING SHIFT OPERATORS

We first consider the time evolution of the many-body wave function. Since the matrix representation of the single-particle Hamiltonian after the quench is tridiagonal, we easily find the single-particle energy

$$\epsilon_k = -2J \cos\left(\frac{k\pi}{L+1}\right) \quad (4)$$

and corresponding eigenstate

$$x_{k,l} = \sqrt{\frac{2}{L+1}} \sin\left(\frac{k\pi}{L+1}l\right), \quad (5)$$

where $k, l = 1, 2, \dots, L$. The time-evolved many-body states $|\psi(t)\rangle = e^{-i\hat{H}t}|\psi(0)\rangle$ are given as

$$|\psi^{\text{MI}}(t)\rangle = \prod_{j=1}^L \left[\sum_{l=1}^L y_{j,l}(t) \hat{b}_l^\dagger \right] |0\rangle, \quad (6)$$

$$|\psi^{\text{CDW}}(t)\rangle = \prod_{j=2,4,\dots}^L \left[\sum_{l=1}^L y_{j,l}(t) \hat{b}_l^\dagger \right] |0\rangle, \quad (7)$$

where all information about real-time dynamics is encoded in

$$y_{j,l}(t) = \sum_{k=1}^L x_{k,j} e^{-i\epsilon_k t} x_{k,l}. \quad (8)$$

The second Rényi entanglement entropy can be obtained by utilizing the expectation value of the shift (SWAP) operator \hat{V} [13,14,16,17]. Let us suppose that we have two copies of the state $|\psi(t)\rangle$, which we call copies 1 and 2, and that the total wave function is given by the product state of the two copies,

$$|\psi_{\text{copy}}(t)\rangle = |\psi(t)\rangle \otimes |\psi(t)\rangle. \quad (9)$$

We divide the system into two subsystems A and B. Here subsystem A contains $j = 1, 2, \dots, L_A$ sites in this paper. Let us consider the shift operator \hat{V}_A which swaps states in subsystem A. The expectation value of \hat{V}_A in terms of $|\psi_{\text{copy}}(t)\rangle$ is related to the reduced density matrix $\hat{\rho}_A$ as

$$\langle \psi_{\text{copy}}(t) | \hat{V}_A | \psi_{\text{copy}}(t) \rangle = \text{Tr}'_A (\hat{\rho}_A \otimes \hat{\rho}_A \hat{V}_A) = \text{Tr}_A \hat{\rho}_A^2, \quad (10)$$

where Tr'_A stands for the trace over the basis of subsystem A of copies 1 and 2. As long as \hat{V}_A acts on a product state of copies 1 and 2, such as Eq. (9), the shift operator transforms the creation operator as

$$\hat{V}_A \hat{b}_j^\dagger \hat{V}_A^{-1} = \begin{cases} \hat{c}_j^\dagger & (j \in \text{A}) \\ \hat{b}_j^\dagger & (j \in \text{B}), \end{cases} \quad (11)$$

$$\hat{V}_A \hat{c}_j^\dagger \hat{V}_A^{-1} = \begin{cases} \hat{b}_j^\dagger & (j \in \text{A}) \\ \hat{c}_j^\dagger & (j \in \text{B}), \end{cases} \quad (12)$$

where operators \hat{b} and \hat{c} respectively correspond to boson operators for copies 1 and 2. For derivation of Eqs. (11) and (12), see Appendix A. Making use of these relations, we can evaluate the second Rényi entanglement entropy in an elementary way. For example, for the MI state, it can be evaluated as

$$S_2 = -\ln \langle \psi_{\text{copy}}^{\text{MI}}(t) | \hat{V}_A | \psi_{\text{copy}}^{\text{MI}}(t) \rangle, \quad (13)$$

$$\begin{aligned} |\psi_{\text{copy}}^{\text{MI}}(t)\rangle &= \left\{ \prod_{j=1}^L \left[\sum_{l=1}^L y_{j,l}(t) \hat{b}_l^\dagger \right] \right\} \\ &\times \left\{ \prod_{j=1}^L \left[\sum_{l=1}^L y_{j,l}(t) \hat{c}_l^\dagger \right] \right\} |0\rangle^{\otimes 2}, \end{aligned} \quad (14)$$

$$\begin{aligned} \hat{V}_A |\psi_{\text{copy}}^{\text{MI}}(t)\rangle &= \left\{ \prod_{j=1}^L \left[\sum_{l=1}^{L_A} y_{j,l}(t) \hat{c}_l^\dagger + \sum_{l=L_A+1}^L y_{j,l}(t) \hat{b}_l^\dagger \right] \right\} \\ &\times \left\{ \prod_{j=1}^L \left[\sum_{l=1}^{L_A} y_{j,l}(t) \hat{b}_l^\dagger + \sum_{l=L_A+1}^L y_{j,l}(t) \hat{c}_l^\dagger \right] \right\} \\ &\times |0\rangle^{\otimes 2}. \end{aligned} \quad (15)$$

Since both $|\psi_{\text{copy}}^{\text{MI}}(t)\rangle$ and $\hat{V}_A |\psi_{\text{copy}}^{\text{MI}}(t)\rangle$ are many-boson states and their wave functions are symmetric under the permutation of \hat{b} bosons (and \hat{c} bosons as well), the expectation value $\langle \psi_{\text{copy}}(t) | \hat{V}_A | \psi_{\text{copy}}(t) \rangle$ can be rewritten by the permanent of single-particle correlation functions Z and \tilde{Z} :

$$\langle \psi_{\text{copy}}(t) | \hat{V}_A | \psi_{\text{copy}}(t) \rangle = \text{perm} \begin{pmatrix} \tilde{Z} & Z \\ Z & \tilde{Z} \end{pmatrix}, \quad (16)$$

where Z (\tilde{Z}) is a single-particle overlap matrix between single-particle states $e^{-i\hat{H}t} \hat{b}_j^\dagger |0\rangle^{\otimes 2}$ and $\hat{V}_A e^{-i\hat{H}t} \hat{c}_l^\dagger |0\rangle^{\otimes 2}$ ($e^{-i\hat{H}t} \hat{b}_j^\dagger |0\rangle^{\otimes 2}$ and $\hat{V}_A e^{-i\hat{H}t} \hat{b}_l^\dagger |0\rangle^{\otimes 2}$). Using the fact that $Z + \tilde{Z} = I$, where I is an identity matrix, we obtain the analytical expression of the Rényi entanglement entropy:

$$S_2 = -\ln \text{perm} A_Z, \quad (17)$$

$$A_Z = \begin{pmatrix} I - Z & Z \\ Z & I - Z \end{pmatrix}. \quad (18)$$

Here the element $z_{j,l}$ of the Hermitian matrix Z is given as

$$z_{j,l}^{\text{MI}} = \sum_{m=1}^{L_A} y_{j,m}^*(t) y_{l,m}(t) \quad (j, l = 1, 2, \dots, L), \quad (19)$$

$$z_{j,l}^{\text{CDW}} = \sum_{m=1}^{L_A} y_{2j,m}^*(t) y_{2l,m}(t) \quad (j, l = 1, 2, \dots, L/2). \quad (20)$$

Note that a somewhat similar formula for the Rényi entanglement entropy given by the permanent has been proposed for excited states in the static system [39].

In the following, \tilde{L} denotes the size of the square matrix A_Z . It is given by $2N$, where N is the number of particles. For example, $\tilde{L} = 2L$ (L) for the MI (CDW) state. Hereafter we will mainly consider the Rényi entanglement entropy for a bipartition of the system into two half chains ($L_A = L/2$).

IV. ANALYTICAL RESULTS

In this section, we analytically evaluate the system size L dependence of the Rényi entanglement entropy S_2 by examining the permanent of the matrix A_Z . We will discuss the condition for S_2 to satisfy the volume-law scaling. Then, we apply the obtained volume-law condition to the quench dynamics of the present case.

A. Remarks on the matrices Z and A_Z

Let us first summarize the characteristics of the matrices Z and A_Z . The purpose of this section is to show $\|A_Z\|_2 = 1$ for both MI and CDW initial states. Here the operator 2-norm is defined as $\|A\|_2 := \sup_{\|x\|_2 \leq 1, x \in \mathbb{C}^M} \|Ax\|_2$ with M being the

size of a square matrix A [on the right-hand side of the equation, $\|\mathbf{x}\|_p := (\sum_j |x_j|^p)^{1/p}$ is an L^p norm of a vector \mathbf{x}], or the largest singular value of the matrix A . We will utilize this fact to obtain the condition for the volume-law entanglement growth in the next section.

For the quench from the MI state, the matrix Z^{MI} is a complex orthogonal projection matrix, satisfying $(Z^{\text{MI}})^2 = Z^{\text{MI}} = (Z^{\text{MI}})^\dagger$ (see Appendix B). Therefore, all the eigenvalues are either 0 or 1. For the quench from the CDW state, the matrix Z^{CDW} is a principal submatrix of the Hermitian matrix Z^{MI} ; i.e., it can be obtained from Z^{MI} by removing $L/2$ rows and the same $L/2$ columns. Using Cauchy's interlace theorem [40,41], we can show that all the eigenvalues of Z^{CDW} are bounded by the largest eigenvalue 1 and the smallest eigenvalue 0 of Z^{MI} . As a result, $0 \leq \|Z^{\text{CDW}}\|_2 \leq \|Z^{\text{MI}}\|_2 = 1$.

All the eigenvalues of the matrix A_Z can be obtained from those of Z . Let us write the eigenvalues of Z as $\epsilon_k^{(Z)}$ and the eigenvectors of Z as $|\epsilon_k^{(Z)}\rangle = (v_{k,1}, v_{k,2}, \dots, v_{k,n})^T$ for $k = 1, 2, \dots, n$ with $n (= \tilde{L}/2)$ being the length of the square matrix Z . Then, half of all the eigenvalues of A_Z are $\epsilon_k^{(A_Z)} = 1$, and the corresponding eigenvectors are $|\epsilon_k^{(A_Z)}\rangle = (v_{k,1}, v_{k,2}, \dots, v_{k,n}, v_{k,1}, v_{k,2}, \dots, v_{k,n})^T$. The remaining half are $\epsilon_{k+n}^{(A_Z)} = 2\epsilon_k^{(Z)} - 1$, and the corresponding eigenvectors are $|\epsilon_{k+n}^{(A_Z)}\rangle = (v_{k,1}, v_{k,2}, \dots, v_{k,n}, -v_{k,1}, -v_{k,2}, \dots, -v_{k,n})^T$. Therefore, $\epsilon_k^{(A_Z)}, \epsilon_{k+n}^{(A_Z)} \in [-1, 1]$ because $\epsilon_k^{(Z)} \in [0, 1]$. Thus, the operator 2-norm of the matrix A_Z satisfies $\|A_Z\|_2 = 1$.

For the quench from the MI state, the matrix A_Z^{MI} becomes a unitary matrix, which can be shown by the relations $(Z^{\text{MI}})^2 + (I - Z^{\text{MI}})^2 = Z^{\text{MI}} + (I - Z^{\text{MI}}) = I$ and $Z^{\text{MI}}(I - Z^{\text{MI}}) = 0$. This unitarity also ensures $\|A_Z^{\text{MI}}\|_2 = 1$. For the quench from the CDW state, the matrix A_Z^{CDW} is not a unitary matrix in general; however, $\|A_Z^{\text{CDW}}\|_2 = 1$ still holds.

Note that the elements of matrix A_Z satisfy

$$\sum_l a_{j,l} = 1, \quad \sum_j a_{j,l} = 1, \quad (21)$$

for any rows j and columns l , which is a part of the definition of the doubly stochastic matrix while the non-negativity condition $a_{j,l} \geq 0$ is absent. (Here the matrix A_Z is complex and satisfies $0 \leq |a_{j,l}| \leq \max(|a_{j,l}|) \leq \|A_Z\|_2 = 1$.) The permanent of the doubly stochastic matrix has been intensively studied [42–49], while little is known about the permanent of a general complex matrix so far.

B. Condition for volume-law entanglement entropy

To quantify the \tilde{L} dependence of the entanglement entropy, we utilize the inequality [36]

$$|\text{perm}A| \leq C^M \exp \left\{ -10^{-5} \times \left[1 - \frac{g_A(M)}{C} \right]^2 M \right\}, \quad (22)$$

which holds for an arbitrary $M \times M$ complex matrix A and an arbitrary nonzero constant C satisfying $C \geq \|A\|_2$. The function $g_A(M)$ is defined as

$$g_A(M) := \frac{1}{M} \sum_{j=1}^M \|r_j\|_\infty \quad (23)$$

with r_j 's being rows of a matrix A and $\|\mathbf{x}\|_\infty := \max_j |x_j|$, which satisfies $0 \leq g_A(M) \leq \|A\|_2$.

We apply inequality (22) to our case given by Eqs. (17) and (18). From Eq. (10), $\text{perm}A_Z$ equals $\text{Tr}_A \hat{\rho}_A^2$, implying $\text{perm}A_Z > 0$. In addition, because $\|A_Z\|_2 = 1$, we can choose $C = 1$ as the tightest bound. Then, the inequality is simplified as

$$\text{perm}A_Z \leq \exp\{-10^{-5} \times [1 - g_{A_Z}(\tilde{L})]^2 \tilde{L}\}, \quad (24)$$

where

$$g_{A_Z}(\tilde{L}) = \frac{1}{N} \sum_{j=1}^N \max_{l=1}^N (|z_{j,l}|, |\delta_{j,l} - z_{j,l}|). \quad (25)$$

Note that $|\text{perm}A| \leq 1$ holds for any unitary matrix A [50], as is the case with the quench from the MI state. Even if A is nonunitary, as is the case with the quench from the CDW state, $|\text{perm}A| \leq (\|A\|_2)^M$ holds [51]. Inequality (22) gives a much tighter constraint on the permanent of A than these two inequalities.

Consequently, the Rényi entanglement entropy satisfies

$$S_2 \geq 10^{-5} \times [1 - g_{A_Z}(\tilde{L})]^2 \tilde{L}. \quad (26)$$

This inequality rigorously guarantees that when

$$\lim_{\tilde{L} \rightarrow \infty} [1 - g_{A_Z}(\tilde{L})] \neq 0, \quad (27)$$

the Rényi entanglement entropy shows the volume-law scaling. In other words, when Eq. (27) holds, the area-law scaling (and the area-law scaling with a logarithmic correction, as is often the case in critical systems [2–5]) is prohibited. If $\lim_{\tilde{L} \rightarrow \infty} [1 - g_{A_Z}(\tilde{L})] = 0$ ($\tilde{L} \propto L$), inequality (26) becomes meaningless and either the area-law or volume-law scaling is allowed. From the volume-law condition given in Eq. (27), we expect that the value $1 - g_{A_Z}(\tilde{L})$ (≥ 0) could be used as an entropy-density-like value, which we will discuss in Sec. V E.

C. Application to the quench dynamics

Here we give two examples which violate the volume-law condition in Eq. (27). One is a product state and the other is a time-evolved state after a short time. Neither of these is expected to follow the volume-law scaling, and in the following we show that they indeed break the condition in Eq. (27). Here, we take a state starting from the MI state as an example.

The first case is a product state at $t = 0$, which apparently has zero entanglement entropy. This fact implies the violation of the condition in Eq. (27). A straightforward calculation on the matrix Z^{MI} gives a matrix

$$Z^{\text{MI}} = \begin{pmatrix} I_{N/2} & 0_{N/2} \\ 0_{N/2} & 0_{N/2} \end{pmatrix}. \quad (28)$$

The matrix A_Z becomes just a permutation matrix, which is a square matrix whose every row and column contains a single 1 with 0s elsewhere. In fact, $g_{A_Z}(\tilde{L}) = \|A_Z\|_2 = 1$ if and only if the matrix A_Z is a permutation matrix [36]. Therefore, the product state always breaks the condition irrespective of L . On the other hand, we can directly calculate the Rényi entanglement entropy by the permanent. The permanent of a permutation matrix is unity, and thus the entanglement entropy for a product state is zero, as expected.




FIG. 1. Time dependence of the absolute value of the correlation function $|z_{j,l}|$ from the MI initial state with $L = 256$ at (a) $tJ/L = 0$, (b) $tJ/L = 0.05$, (c) $tJ/L = 0.1$, and (d) $tJ/L = 0.25$. At $t = 0$, the correlation function $z_{j,l}$ is given by Eq. (28). For $0 < tJ < L/4$, $z_{j,l}$ is approximately given by Eq. (29) and the length of the sides of the square where $|z_{j,l}| > 10^{-10}$ holds is nearly equal to $2\beta \approx 4tJ$.

The second case is a time-evolved state with a short time, $t \ll L/v_C$, where v_C is the propagation velocity of correlations. In the present 1D free boson system, v_C is equivalent to the maximum quasiparticle velocity v_{\max} . The velocity is given by the maximal group velocity and $v_{\max} = 2J$ [34,35] (when $\hbar = 1$ and the lattice constant is chosen to be unity) in the present case. Because the single-particle correlation function extends up to a distance $v_C t$, it is likely that the entanglement entropy does not grow with increasing L when $L > v_C t$. Therefore, the time-evolved state with fixed $t (\ll L/v_C)$ is expected to follow the area-law scaling, implying the breaking of the volume-law condition in Eq. (27).

In this situation, we can approximate the matrix Z^{MI} as

$$Z^{\text{MI}} \approx \begin{pmatrix} I_{N/2-\beta} & 0 & 0 \\ 0 & Z'_{2\beta} & 0 \\ 0 & 0 & 0_{N/2-\beta} \end{pmatrix}, \quad (29)$$

where β is a positive integer and is roughly proportional to $2tJ$. $Z'_{2\beta}$ would be a dense matrix with the size $2\beta \times 2\beta$. Figure 1 shows the time dependence of the absolute value of Z^{MI} . We see that Z^{MI} actually has the matrix structure given in Eq. (29) when $tJ \ll L/4$. For the derivation of Eq. (29) and the specific value of β , see Appendix C.

Using Eq. (29), we can verify the breaking of the volume-law condition of Eq. (27). Substituting Eq. (29) into $g_{A_Z}(\tilde{L})$ in Eq. (25), we obtain

$$g_{A_Z}(\tilde{L}) = \frac{N - 2\beta}{N} + \frac{1}{N} \sum_{j \in \Delta} \max_{l \in \Delta} (|z_{j,l}|, |\delta_{j,l} - z_{j,l}|), \quad (30)$$




FIG. 2. Time dependence of the Rényi entanglement entropy for the quench from the (a) MI and (b) CDW states. For a short time ($tJ \lesssim L/4$), the Rényi entanglement entropy S_2 exhibits an increase proportional to time t . After a long time ($tJ \gtrsim L/4$), S_2 is nearly saturated at the value proportional to the system size L .

where $\Delta = [N/2 - \beta, N/2 + \beta]$. Because $0 \leq |z_{j,l}| \leq 1$, $1 - g_{A_Z}(\tilde{L})$ is bounded like

$$0 \leq 1 - g_{A_Z}(\tilde{L}) \leq \frac{2\beta}{N}. \quad (31)$$

Since β depends on $2tJ$ but not on the system size L , by taking the thermodynamic limit $L(\propto N) \rightarrow \infty$, we conclude

$$\lim_{L \rightarrow \infty} [1 - g_{A_Z}(\tilde{L})] = 0. \quad (32)$$

We note that the breaking of the volume-law condition does not directly mean the area-law scaling, as mentioned before. Because we do not have the criterion on the area-law scaling of the entanglement entropy at this stage, we rely on the numerical calculation to check the area-law scaling. The numerical results of the Rényi entanglement entropy will be shown in the next section. Calculated Rényi entanglement entropies with a short time $tJ \ll L$ shown in Fig. 2 take almost the same value with increasing L , implying that the entanglement entropy of this state would obey the area-law scaling.

Another less rigorous evidence of the area-law entanglement scaling can be seen from the permanent formula. Substituting the approximated expression on Z^{MI} in Eq. (29) into Eq. (18), we obtain the permanent of the matrix A_Z as

$$\begin{aligned} \text{perma}_{A_Z} &\approx \text{perma}_{Z'} \\ &= \text{perm} \begin{pmatrix} I_{2\beta} - Z'_{2\beta} & Z'_{2\beta} \\ Z'_{2\beta} & I_{2\beta} - Z'_{2\beta} \end{pmatrix}. \end{aligned} \quad (33)$$

Applying the same discussion in Sec. IV B with replacing the matrix size \tilde{L} with 2β , we expect that the entanglement entropy would be constant when $1 - g_{A_{Z'}} = 0$ or proportional to 2β when $1 - g_{A_{Z'}} \neq 0$. Strictly speaking, this argument does not exclude the possibility that the entanglement entropy is also proportional to the system size L . However, since the condition for the volume-law scaling in Eq. (27) itself ensures the proportionality of the entanglement entropy with respect to \tilde{L} , it is unlikely that the volume-law scaling of entanglement would be satisfied if the proportionality to \tilde{L} is replaced by that to β .

V. NUMERICAL RESULTS

In this section, we will numerically evaluate the permanent to obtain the Rényi entanglement entropy. In general, permanent calculations require an exponentially long time. However, we can practically obtain the Rényi entanglement entropy for systems larger than the exact diagonalization method can handle and can perform longer simulations than the method based on matrix product states, even by performing a brute-force permanent calculation.

The advantages of getting the Rényi entanglement entropy by the permanent calculation are the following:

(i) We do not need Hamiltonian eigenstates, which require much memory cost. This is the main reason why our method enables us to access larger systems than the exact diagonalization method can handle.

(ii) Without explicit time evolution, we can directly calculate the Rényi entanglement entropy at a given time, which allows us parallel computations.

(iii) In a system of soft-core bosons, there is no upper limit to the number of bosons at any site, but it is common to set a realistic limit when performing numerical calculations. With the present method we have proposed, we do not have to care about the size of such a local Hilbert space limitation.

A. Summary of algorithm

Here we briefly review the algorithm for the permanent calculation. The permanent of an $M \times M$ matrix A is defined as

$$\text{perm}A = \sum_{\sigma \in \text{Sym}(M)} \prod_{j=1}^M a_{j,\sigma(j)}, \quad (34)$$

where $\text{Sym}(M)$ is the symmetric group, i.e., over all permutations of numbers $1, 2, \dots, M$. Since straightforward calculations require $M! \times M$ arithmetic operations, we should use a more efficient algorithm. The best known algorithms so far are the Ryser formula [52–54] and Balasubramanian-Bax-Franklin-Glynn (BBFG) formula [54–58]. Both take $O(M2^{M-1})$ computation time. Hereafter we mainly use the BBFG formula for the permanent calculation. It is given by

$$\text{perm}A = \frac{1}{2^{M-1}} \sum_{\delta} \left(\prod_{m=1}^M \delta_m \right) \prod_{l=1}^M \sum_{j=1}^M \delta_j a_{j,l}, \quad (35)$$

where $\delta = (\delta_1, \delta_2, \dots, \delta_M) \in \{\pm 1\}^M$ with $\delta_1 = 1$. Although the computation time using the above straightforward BBFG formula is $O(M2^{2M-1})$, it can be reduced to $O(M2^{M-1})$ by

utilizing a specific ordering of the binary numeral system, known as Gray code [59,60]. Our numerical source code is based on the PYTHON program in “The Walrus” library [61].

The current feasible matrix size is up to $\sim 50 \times 50$ [62–64]. For the quench from the MI (CDW) state, the size of the matrix A_Z is $2L \times 2L$ ($L \times L$). Therefore, in principle, we may handle $L \lesssim 25$ ($L \lesssim 50$) for the MI (CDW) case. Here we present our results for $L \leq 20$ ($L \leq 40$) for the quench from the MI (CDW) state. Although we may be able to utilize the symmetry of the matrix A_Z to accelerate permanent computations [65], we stick to the original BBFG formula. Even without improving the original algorithm, it allows us to calculate the permanent for systems much larger than the exact diagonalization method can deal with. (Note that, for example, exact diagonalization calculations for $L = 14$ from the MI state to $U/J = 3.01$ have been reported [23].) As for the size at which the permanent is computable, the Rényi entanglement entropy can be obtained at any given time, allowing for longer simulations than the methods based on matrix product states.

B. Time dependence of Rényi entanglement entropy

We examine the time dependence of the Rényi entanglement entropy (see Fig. 2). For a short time ($tJ \lesssim L/4$), S_2 grows linearly with t . After $tJ \gtrsim L/4$, S_2 is almost saturated and reaches the value nearly proportional to L . S_2 exhibits oscillations whose period grows with L . These observations are consistent with the fact that the t -linear growth of the entanglement entropy terminates at $t \sim L/(2v_{\text{max}})$, where $v_{\text{max}} = 2J$ is the maximum quasiparticle velocity [27,34,35].

To see this behavior more clearly, we rescale the time tJ and the Rényi entanglement entropy S_2 in the unit of the system size L (see Fig. 3). All the lines nearly overlap for $tJ/L \lesssim 1$ when $L \gtrsim 10$ ($L \gtrsim 20$) for the quench from the MI (CDW) state. The deviation from the thermodynamic limit appears to be smaller for the CDW state because the feasible size is larger than the MI state.

C. Comparison with the Rényi entanglement entropy estimated from the Gaussian state

The MI or CDW state quenched to $U = 0$ evolves to a Gaussian state after a long time in the thermodynamic limit ($L \gg tJ \gg 1$) [33,66,67]. The entanglement entropy of the Gaussian state can be calculated from the eigenvalues of the matrix, consisting of $\langle \hat{b}_j^\dagger \hat{b}_l \rangle$ and $\langle \hat{b}_j \hat{b}_l \rangle$ [18,29,31,32,68]. After diagonalizing the matrix in bosonic systems, we obtain the eigenvalues, which correspond to the expectation values of the mode occupation numbers n_μ . The Rényi entanglement entropy S_α of order α can be described by n_μ as [18,29,31,32,68]

$$S_\alpha = \frac{1}{\alpha - 1} \sum_{\mu} \ln [(n_\mu + 1)^\alpha - n_\mu^\alpha]. \quad (36)$$

The time-evolved state is not a Gaussian state for $tJ, L < \infty$ in general. However, it is not outrageous to extract reference values using the Gaussian state, which exhibits the same single-particle correlation functions of the (non-Gaussian) time-evolved state for a finite time and finite sizes. At least for the MI quench, as we see below, single-particle




FIG. 3. Rescaled time dependence of the Rényi entanglement entropy for the quench from the (a) MI and (b) CDW states. The Rényi entanglement entropy and the time are rescaled by L . The dashed lines correspond to the Rényi entanglement entropy densities S_2^{Gaussian}/L estimated from Eq. (36). The dashed line is independent of L for the MI state, while it is obtained for a sufficiently large size ($L = 1024$) for the CDW state. The insets show the long-time behavior of the Rényi entanglement entropy densities.

correlations are independent of time and size. As a result, the Rényi entanglement entropy of Eq. (36) obtained from the mode occupation numbers n_μ for any $tJ, L < \infty$ gives the true Rényi entanglement entropy for $L \gg tJ \gg 1$. Hereafter we use the symbol S_2^{Gaussian} to denote the Rényi entanglement entropy estimated from Eq. (36).

For the quench starting from the MI state, the matrix consisting of the correlation function is already diagonal; $\langle \hat{b}_j^\dagger \hat{b}_l \rangle = \delta_{j,l}$ and $\langle \hat{b}_j \hat{b}_l \rangle = 0$ ($j, l = 1, 2, \dots, L_A$) for all L and tJ . The mode occupation numbers, $n_\mu = 1$ ($\mu = 1, 2, \dots, L_A$), are independent of time and sizes. Therefore, the Rényi entanglement entropy of subsystem A, whose size is $L_A = L/2$, is given by $S_2^{\text{Gaussian}} = \ln 3 \times L/2 \approx 0.5493 \times L$. Indeed, numerically obtained S_2 for $tJ \gg L$ fluctuates around S_2^{Gaussian} [see Figs. 3(b) and 4(b)], while a t -linear growth is not reproduced in S_2^{Gaussian} for a short time ($tJ \ll L$).

For the quench starting from the CDW state, we numerically evaluate the Rényi entanglement entropy for a subsystem size $L_A = L/2$. In this case, $\langle \hat{b}_j^\dagger \hat{b}_l \rangle$ depends on time, and therefore, S_2^{Gaussian} oscillates in time. Again, numerically obtained S_2 for $tJ \gg L$ fluctuates around S_2^{Gaussian} [see Figs. 3(b) and 4(b)], whereas a t -linear growth in a short time is absent for S_2^{Gaussian} . The Rényi entanglement entropy in the thermodynamic limit for a long time is estimated to be $S_2^{\text{Gaussian}}/L = 0.31(1)$ for the Gaussian state, and S_2 is expected to converge to this value for $L \gg tJ \gg 1$.




FIG. 4. Comparisons among the Rényi entanglement entropy S_2 (solid lines), that estimated from Eq. (36), S_2^{Gaussian} (dashed lines), and the Page value S_2^{Page} (dotted lines) for the (a) MI and (b) CDW states. Note that S_2^{Gaussian} and S_2^{Page} of $L = 10$ for the MI state take almost the same value. We do not show error bars of S_2^{Page} because they are invisibly small.

D. Comparison with Page value

The Bose-Hubbard model is nonintegrable (integrable) for $|U| > 0$ ($U = 0$). The system is thermalized when it is quenched to the parameter region $|U| > 0$ [69,70]. The entanglement entropy would be nearly saturated at that of the random state vector, which is known as the Page value S_2^{Page} [71]. For the Bose-Hubbard model, an analytical expression for the Page value has not been obtained yet. We can obtain the Page value within the statistical error bars by numerically taking an average of random state vectors [25,72]. When taking this average, we should directly use the Hilbert space of the Bose-Hubbard model. The dimension of the Hilbert space grows exponentially when increasing the system size. Therefore, we can estimate the Page value only for rather small system sizes. In this paper, we calculate S_2^{Page} up to $L = 12$ ($L = 16$) at unit filling (half filling).

For the $U = 0$ quench, the time-evolved state is not thermalized. The Rényi entanglement entropy, in this case, would deviate from the Page value. To examine whether we can tell the difference between the entanglement entropy of the thermalized state and that of the state quenched to $U = 0$, we compare the Rényi entanglement entropy at $U = 0$ with the Page value. We calculate the Page value by averaging 1024 random samples.

As for the quench from the MI state, the Page value S_2^{Page} at unit filling is close to the Rényi entanglement entropy S_2 for a longer time ($tJ \gtrsim L/4$) and for all systems ($L \leq 12$)




FIG. 5. System size dependence of the Rényi entanglement entropy density for (a) the MI initial state and (b) the CDW initial state. Circles represent the time average of the Rényi entanglement entropy density of the time-evolved state S_2/L from $tJ = 1$ to $tJ = 10^4$. Diamonds are the time average of S_2^{Gaussian}/L from $tJ = 1$ to $tJ = 10^4$. Squares are the Page value. The solid and dotted lines are, respectively, linear fits of S_2/L and S_2^{Page}/L using five largest systems. The dashed line in (a) represents S_2^{Gaussian}/L , which is independent of the system size L . The dashed line in (b) shows a linear fit of S_2^{Gaussian}/L using data satisfying $L \geq 20$. We do not show error bars because they are small compared to symbol sizes.

that we have considered [see Fig. 4(a)]. When increasing the system size L , we see that the Page value becomes greater than the Rényi entanglement entropy S_2 at a longer time. This observation indicates that when taking the thermodynamic limit, the Page value converges to the value larger than S_2 . We examine the system-size dependence and extrapolate these values, as well as S_2^{Gaussian} , to the thermodynamic limit, as shown in Fig. 5(a). The Rényi entanglement entropy S_2 in the thermodynamic limit is very close to that of the Gaussian state, which is expected from previous studies [33,66,67], although we see a small deviation due to finite-size effects. On the other hand, in the thermodynamic limit, the Page value is greater than S_2 and S_2^{Gaussian} . Thus, it is expected that a jump occurs between the entropies of $U = 0$ and $|U| > 0$ quenches after long-time evolution for sufficiently large system sizes, although they take similar values in a small system. In this respect, we compare the Rényi entanglement entropy of $U = 0$ with that of $U > 0$ and confirm the presence of a jump in Appendix D.

For the quench from the CDW state at half filling, the Rényi entanglement entropy S_2 is slightly smaller than the Page value unlike the MI case [see Fig. 4(b)]. The deviation seems to be more enhanced with increasing system sizes. It

is likely that the Rényi entanglement entropy density would be smaller than the density of the Page value in the thermodynamic limit although the Rényi entanglement entropy itself satisfies the volume-law scaling, as can be seen from Fig. 3(b). We investigate the system-size dependence and extract the Rényi entanglement entropy density in the thermodynamic limit, as shown in Fig. 5(b). From the same discussion as in the MI case, there is a jump between the entropies of $U = 0$ and $|U| > 0$ quenches after long-time evolution, which is discussed in Appendix D.

The time-averaged Rényi entanglement entropy density S_2/L of the $U = 0$ quench is smaller than the Page value S_2^{Page}/L , which is also the value S_2/L expected for the $|U| > 0$ quench, as shown in Fig. 5. This would be understood from the viewpoint of the number of states and the integrability. When the state is thermalized, the entanglement entropy is saturated at the Page value and the state would be characterized by the thermal distribution. Therefore, the entanglement entropy as well as the Page value would be identified as the thermal entropy, described by the logarithm of the number of states. When we denote ζ as the number of states per site, the number of states in subsystem A is given by ζ^{L_A} and the Page value S_2^{Page}/L would be approximated by $[\ln(\zeta)]/2$ (where $L_A = L/2$). On the other hand, after a long-time evolution, the state quenched to $U = 0$ relaxes to a state characterized by the generalized Gibbs ensemble (GGE), which is the Boltzmann (thermal) distribution taking into account not only the internal energy but also a set of conserved quantities [73]. In this case, the time-averaged Rényi entanglement entropy can be seen as the logarithm of the number of states in the GGE [74]. Due to the presence of the exponentially large number of conserved quantities, the number of states in the GGE would be ζ^{L_A}/η^{L_A} , where $\eta (> 1)$ represents the number of conserved charges per site. Consequently, the entropy density of the GGE, S_2/L , is estimated as $[\ln(\zeta/\eta)]/2$. Comparing the Page value and the entropy density of the GGE, we see that S_2/L is always smaller than S_2^{Page}/L .

E. Entropy-density-like value and practical bound for the Rényi entanglement entropy

From the argument in Sec. IV B, we obtain the rigorous lower bound for the second Rényi entanglement entropy density, which is given as

$$\frac{S_2}{\tilde{L}} \geq 10^{-5} \times [1 - g_{A_z}(\tilde{L})]^2. \quad (37)$$

Moreover, the tighter bound was conjectured to be $\text{per}M \leq \exp\{-\text{const} \times M[1 - g_A(M)]\}$, where M is the size of matrix A [36], which gives the lower bound of the Rényi entanglement entropy as

$$\frac{S_2}{\tilde{L}} \geq \text{const} \times [1 - g_{A_z}(\tilde{L})]. \quad (38)$$

We briefly note that the conjectured bound gives the same condition for the volume-law entanglement growth as Eq. (27). These inequalities lead us to expect that $\tilde{s}(\tilde{L}) = 1 - g_{A_z}(\tilde{L})$ determines the qualitative behavior of the Rényi entanglement entropy density and serves as an entropy-density-like value.




FIG. 6. Comparisons between the rescaled Rényi entanglement entropy S_2/\tilde{L} and $\tilde{s}(\tilde{L})$ estimated from the infinity norm of rows of the matrix A_Z for the quench from the (a) MI ($\tilde{L} = 2L$) and (b) CDW ($\tilde{L} = L$) states. The value $\tilde{s}(\tilde{L} \rightarrow \infty)$ is obtained for a sufficiently large size ($L = 1024$). Note that $\tilde{s}(\tilde{L})$'s are the same for the MI and CDW states as shown in Appendix E.

$\tilde{s}(\tilde{L})$ is obtained from g_{A_Z} in Eq. (25):

$$\tilde{s}(\tilde{L}) = 1 - \frac{1}{N} \sum_{j=1}^N \max_{l=1}^N (|z_{j,l}|, |\delta_{j,l} - z_{j,l}|). \quad (39)$$

As shown in Appendix E, the entropy-density-like value can be expressed in a simple form:

$$\tilde{s}(\tilde{L}) = \frac{1}{2} - \frac{1}{N} \sum_{j=1}^N \left| z_{j,j} - \frac{1}{2} \right|. \quad (40)$$

Unlike the permanent of matrix A_Z , which requires costly calculation, $\tilde{s}(\tilde{L})$ is easy to compute numerically and analytically. Hence if $\tilde{s}(\tilde{L})$ has a similar tendency with S_2/\tilde{L} , it would be a helpful quantity to qualitatively capture features of the entanglement entropy density.

To see the behavior of $\tilde{s}(\tilde{L})$ in the thermodynamic limit, we compare $\tilde{s}(\tilde{L})$ for a much larger system $L = 1024$ with S_2/\tilde{L} for $\tilde{L} \leq 40$ in Fig. 6. We have confirmed that $\tilde{s}(\tilde{L})$ converges well when $L \gtrsim 256$ and, therefore, regarded $\tilde{s}(\tilde{L})$ with $L = 1024$ as that in the thermodynamic limit, $\tilde{s}(\tilde{L} \rightarrow \infty)$. Both $\tilde{s}(\tilde{L} \rightarrow \infty)$ and S_2/\tilde{L} are found to behave in a qualitatively similar way including the period of oscillations. Thus, we confirm that $\tilde{s}(\tilde{L} \rightarrow \infty)$ certainly captures the qualitative behavior of the Rényi entanglement entropy density.

While Eq. (38) means that $\text{const} \times \tilde{s}(\tilde{L})$ serves as a lower bound of S_2/\tilde{L} , we observe in Fig. 6 that the rescaled Rényi entanglement entropy S_2/\tilde{L} would be practically bounded from above by the entropy-density-like value in the thermodynamic




FIG. 7. Comparisons between the rescaled Rényi entanglement entropy S_2/\tilde{L} and $\tilde{s}(\tilde{L})$ estimated from the infinity norm of rows of the matrix A_Z for the quench from the (a) MI ($\tilde{L} = 2L$) and (b) CDW ($\tilde{L} = L$) states for $L = 4, 8$, and 12 . In both cases, the Rényi entanglement entropy density S_2/\tilde{L} seems to be bounded by $2\tilde{s}(\tilde{L})$. Note that $\tilde{s}(\tilde{L})$'s are the same for the MI and CDW states as shown in Appendix E.

limit $\tilde{s}(\tilde{L} \rightarrow \infty)$. This observation leads us to expect that $\tilde{s}(\tilde{L})$ would play a role of an upper bound of S_2/\tilde{L} for any system sizes and motivates us to examine how $\tilde{s}(\tilde{L})$ bounds the Rényi entanglement entropy S_2/\tilde{L} from above. For this purpose, we compare these for some finite L 's, as shown in Fig. 7. We find a region where the expected inequality $S_2/\tilde{L} < \tilde{s}(\tilde{L})$ is slightly violated. Even in this case, for the MI quench, S_2/\tilde{L} is bounded by $2\tilde{s}(\tilde{L})$ from above and they well overlap in the time range $tJ/(2L) \lesssim 0.02$ [see Fig. 7(a)]. Likewise, in the case of the CDW quench, $2\tilde{s}(\tilde{L})$ still gives an upper bound for S_2/\tilde{L} [see Fig. 7(b)]. From these observations, we speculate that some constant value times $\tilde{s}(\tilde{L})$ would practically give an upper bound of the Rényi entanglement entropy.

To sum up this section, we have observed

$$\frac{S_2}{\tilde{L}} \lesssim \tilde{s}(\tilde{L} \rightarrow \infty) \quad \text{for } \tilde{L} \gg 1, \quad (41)$$

$$\frac{S_2}{\tilde{L}} \lesssim \text{const} \times \tilde{s}(\tilde{L}) \quad \text{for finite } \tilde{L}. \quad (42)$$

These results imply that $\bar{s}(\tilde{L})$, which can be obtained from the infinity norm of rows of the matrix A_Z , would be a helpful guide in qualitatively estimating the Rényi entanglement entropy at least in the present case. We expect that the current discussion is applicable to other initial conditions and Hamiltonians.

VI. SUMMARY AND OUTLOOK

We have studied the time evolution of the Rényi entanglement entropy in a 1D free boson system. We have focused on the quench dynamics of the 1D Bose-Hubbard model at the noninteracting point starting from the Mott-insulating and charge-density-wave initial states. We have obtained the analytical form of the second Rényi entanglement entropy by calculating the expectation value of the shift operator. The Rényi entanglement entropy was found to be the negative of the logarithm of the permanent of the matrix whose elements are time-dependent single-particle correlation functions. Using a permanent inequality, we have rigorously proven that the Rényi entanglement entropy satisfies the volume-law scaling under a certain condition. We have also numerically obtained the long-time evolution of the Rényi entanglement entropy by direct computations of the permanent. Although it requires exponential time in general, the present approach is superior to the best currently available methods such as the exact diagonalization and matrix-product-state methods. The feasible system size is about twice the size that the conventional method can handle [23]. Since our method enables us to compute the Rényi entanglement entropy at any time, the reachable time is also much longer than the conventional methods.

The procedure presented in this paper can be extended to systems containing long-range hopping parameters and those with randomness. Real-time dynamics of such complex quantum many-body systems of free fermions have attracted much attention recently, whereas those of free bosons are yet to be explored because of their computational difficulties. Our method would be helpful for studying such bosonic counterparts. Typical examples include (i) noninteracting higher-dimensional systems (see Refs. [75–77] for correlation-spreading dynamics with an interaction quench in two dimensions), (ii) Anderson localization with long-range hopping (see Refs. [78,79] for free fermions), (iii) localization in disorder-free or correlated-disorder systems such as the Aubry-André model [80] (see Refs. [78,81] for free fermions), and (iv) Lindblad dynamics of free bosons (see Refs. [82–86] for free fermions). Our formula would also be useful in studying the entanglement properties of mixtures of bosons and fermions (e.g., fermion system in cavities [87] and Bose-Fermi-Hubbard systems [88]).

We have introduced the entropy-density-like value using the infinity norm of rows of the matrix consisting of the correlation function and have numerically demonstrated that it well captures the features of the Rényi entanglement entropy density. We have also discussed a practical bound of the entanglement entropy, i.e., that of the matrix permanent, which usually requires exponential time computations. Our findings on the practical bound would also stimulate mathematical research on the permanent of general complex matrices and

research in the field of quantum computing involving boson-sampling techniques [89,90].

ACKNOWLEDGMENTS

The authors acknowledge fruitful discussions with S. Goto and Y. Takeuchi. This work was financially supported by JSPS KAKENHI (Grants No. 18H05228, No. 19K14616, No. 20H01838, No. 21H01014, and No. 21K13855), by Grant-in-Aid for JSPS Fellows (Grant No. 22J22306), by JST CREST (Grant No. JPMJCR1673), by MEXT Q-LEAP (Grant No. JPMXS0118069021), and by JST FOREST (Grant No. JP-MJFR202T).

APPENDIX A: DERIVATION OF EQUATIONS (11) AND (12)

Although we apply Eqs. (11) and (12) to the product state of the same state living in copies 1 and 2, such as Eq. (9), in the main part, it holds even if states of copies 1 and 2 are different. To prove this, we first Schmidt-decompose the state of copy 1 (2) as

$$|\psi\rangle^{1(2)} = \sum_l s_l^{1(2)} |\phi_l\rangle^{1(2),A} |\varphi_l\rangle^{1(2),B}, \quad (\text{A1})$$

where $|\phi_l\rangle^{1(2),A}$ and $|\varphi_l\rangle^{1(2),B}$ are orthonormal states of subsystems A and B, respectively. $s_l^{1(2)}$ is the Schmidt coefficient. The product state of copies 1 and 2 is given by

$$|\psi_{\text{prod}}\rangle = |\psi\rangle^1 \otimes |\psi\rangle^2, \quad (\text{A2})$$

where we assume that copies 1 and 2 can have different states.

We consider how $\hat{V}_A \hat{b}_j^\dagger \hat{V}_A^{-1}$ acts on the state given by Eq. (A2). Note that $\hat{V}_A^{-1} = \hat{V}_A$ holds because of $\hat{V}_A^2 = \hat{I}$, where \hat{I} is an identity operator. When $j \in B$, $\hat{V}_A \hat{b}_j^\dagger \hat{V}_A^{-1} |\psi_{\text{prod}}\rangle = \hat{b}_j^\dagger |\psi_{\text{prod}}\rangle$ trivially holds. In the case of $j \in A$, the action of $\hat{V}_A \hat{b}_j^\dagger \hat{V}_A^{-1}$ on the product state is given by

$$\begin{aligned} & \hat{V}_A \hat{b}_j^\dagger \hat{V}_A^{-1} |\psi_{\text{prod}}\rangle \\ &= \hat{V}_A \hat{b}_j^\dagger \hat{V}_A^{-1} \sum_{l,m} s_l^1 s_m^2 |\phi_l\rangle^{1,A} |\varphi_l\rangle^{1,B} |\phi'_m\rangle^{2,A} |\varphi'_m\rangle^{2,B} \\ &= \hat{V}_A \hat{b}_j^\dagger \sum_{l,m} s_l^1 s_m^2 |\phi'_m\rangle^{1,A} |\varphi_l\rangle^{1,B} |\phi_l\rangle^{2,A} |\varphi'_m\rangle^{2,B} \\ &= \hat{V}_A \sum_{l,m} s_l^1 s_m^2 (\hat{b}_j^\dagger |\phi'_m\rangle^{1,A}) |\varphi_l\rangle^{1,B} |\phi_l\rangle^{2,A} |\varphi'_m\rangle^{2,B} \\ &= \sum_{l,m} s_l^1 s_m^2 |\phi_l\rangle^{1,A} |\varphi_l\rangle^{1,B} (\hat{c}_j^\dagger |\phi'_m\rangle^{2,A}) |\varphi'_m\rangle^{2,B} \\ &= \hat{c}_j^\dagger |\psi_{\text{prod}}\rangle. \end{aligned} \quad (\text{A3})$$

Thus, we proved Eq. (11) [and Eq. (12) in the same manner] as long as it acts on the product state of copies 1 and 2.

APPENDIX B: MORE ON THE PROPERTY OF THE MATRIX Z^{MI}

The matrix X , having the element $x_{k,l}$ in Eq. (5), is unitary and diagonalizes the single-particle Hamiltonian matrix H , namely, $HX = XE$, with the matrix E being a diagonal matrix consisting of all the corresponding eigenvalues. The matrix

$D = e^{-iEt}$ ($t \geq 0$) is also unitary, and the matrix Y , having the component $y_{j,l}$ in Eq. (8), is given as $Y = X^\dagger D X$. Because $Y Y^\dagger = Y^\dagger Y = I$, Y is also unitary.

The Hermitian matrix Z^{MI} can be represented as

$$Z^{\text{MI}} = (Y P_{L/2} Y^\dagger)^T, \quad P_{L/2} = \begin{pmatrix} I_{L/2} & 0_{L/2} \\ 0_{L/2} & 0_{L/2} \end{pmatrix}, \quad (\text{B1})$$

where $I_{L/2}$ and $0_{L/2}$ are, respectively, an $L/2 \times L/2$ identity matrix and an $L/2 \times L/2$ zero matrix. Since $P_{L/2}$ is a projection matrix, satisfying $P_{L/2}^2 = P_{L/2}$, we immediately obtain $(Z^{\text{MI}})^2 = Z^{\text{MI}}$. Therefore, the matrix Z^{MI} is a projection matrix. Note that this argument holds for any Hermitian single-particle Hamiltonian matrix H , in particular, containing long-range hopping parameters with randomness.

APPENDIX C: DERIVATION OF EQUATION (29)

The purpose of this Appendix is to prove Eq. (29) for a time-evolved state with a short time, $t \ll L/v_C$. Corresponding to the limit in Eq. (27), we consider the thermodynamic limit, $L \rightarrow \infty$ with a fixed time t in the following.

Since we consider the case $tJ \ll L$ and the thermodynamic limit, the summation in Eq. (8) can be replaced with the integral. In this case, the single-particle wave function $y_{j,l}$ and the correlation function $z_{j,l}$ can be expressed as

$$y_{j,l} = (-i)^{j-l} [J_{j-l} - (-1)^l J_{j+l}], \quad (\text{C1})$$

$$z_{j,l} = i^j (-i)^l \sum_{m=1}^{L_A} [J_{j-m} J_{l-m} + J_{j+m} J_{l+m} - (-1)^m J_{j-m} J_{l+m} - (-1)^m J_{j+m} J_{l-m}], \quad (\text{C2})$$

where we use an abbreviation $J_n = J_n(2tJ)$ with $J_n(x)$ being the Bessel function of the first kind.

We note that the Bessel function is exponentially small when $|n| > (e/2)x$ because of the inequality of the Bessel function,

$$|J_n(x)| \leq \frac{(x/2)^{|n|}}{|n|!} \approx \frac{e^{-(|n|+1/2)\ln(2|n|/ex)}}{\sqrt{\pi ex}}, \quad (\text{C3})$$

where $x \geq 0$ and n is an integer. Note that $J_{-n}(x) = (-1)^n J_n(x)$ holds. When deriving the right-hand side of Eq. (C3), we assume that $|n|$ is large and use Stirling's approximation. Using this fact, we can simplify the correlation function $z_{j,l}$. Let β be the smallest positive integer for which $|J_\beta(2tJ)|$ is negligibly small. When we specifically demand $|J_\beta(2tJ)| < \varepsilon$, β is approximately given by

$$\beta \gtrsim \frac{e}{2}(2tJ) - \ln \sqrt{\pi e(2tJ)\varepsilon} + \dots, \quad (\text{C4})$$

where we assume that $2tJ \gg 1$ and $\beta \gg 1$. By definition, $J_n(2tJ)$ for $|n| > \beta$ is exponentially small.

We evaluate $z_{j,l}$ in detail using the aforementioned properties of the Bessel function. When (i) $j > L_A + \beta$, (ii) $j < L_A - \beta$, (iii) $l > L_A + \beta$, and (iv) $l < L_A - \beta$, we can approximate $z_{j,l}$ to a simple form. When (i) $j > L_A + \beta$, all terms in the summation of Eq. (C2) are exponentially small because the smallest index ($j - m$ or $j + m$) of the Bessel functions with varying m is $j - L_A (> \beta)$. Thus, we

can approximate the correlation function as $z_{j,l} \approx 0$. A similar argument holds for case (iii) by replacing j with l .

Before moving to the next case, we note that due to the unitarity of $y_{j,l}$, the correlation function $z_{j,l}$ can be regarded by $\delta_{j,l}$ when the upper limit of the summation (L_A) in Eq. (C2) can be replaced by L . This can be approximately achieved when $\sum_{m=L_A+1}^L y_{j,m}^* y_{l,m}$ is negligibly small. [Compare the definition of $z_{j,l}$ in Eq. (19).] When (ii) $j < L_A - \beta$, the index with the smallest absolute value of the Bessel functions in $\sum_{m=L_A+1}^L y_{j,m}^* y_{l,m}$ with varying m is $j - (L_A + 1) (< -\beta)$ and we can approximate $\sum_{m=L_A+1}^L y_{j,m}^* y_{l,m} \approx 0$. Therefore, we can regard $z_{j,l}$ as $\delta_{j,l}$. For case (iv), we can obtain the same result by replacing j with l .

As a result, Z^{MI} has a structure shown in Fig. 1. Denoting the dense matrix part that does not fall under the above simplifications as $Z'_{2\beta}$ (whose matrix size is $2\beta \times 2\beta$), we obtain Eq. (29).

APPENDIX D: QUENCH TO FINITE U FROM THE MI AND CDW INITIAL STATES

As shown in Figs. 4 and 5, when the system is quenched to the noninteracting point ($U = 0$) starting from the MI and CDW initial state, the Rényi entanglement entropy after the long-time evolution deviates from the Page value. When the system is quenched to finite U , it is expected that thermalization occurs and the Rényi entanglement entropy approaches the Page value after the long-time evolution. These facts indicate that there is a jump between saturated values of the Rényi entanglement entropy for the $U = 0$ and finite- U quenches.

To confirm this, we calculate the time evolution of the Rényi entanglement entropy for the quench to finite U starting from the MI and CDW states by the exact diagonalization method [91,92] and compare it with that for the $U = 0$ quench, as shown in Figs. 8(a) and 8(b). In both cases, we observe the jump between the Rényi entanglement entropies of $U = 0$ and small but finite U . We also find that the Rényi entanglement entropies for finite U converge to the Page value. Thus, we can distinguish whether the state is thermalized (for $U > 0$) or not (for $U = 0$) from the Rényi entanglement entropy in a sufficiently large system (see Figs. 4 and 5).

APPENDIX E: PROPERTIES OF THE ENTROPY-DENSITY-LIKE VALUE

In this Appendix, we show the derivation of Eq. (40) and analytically prove the agreement of the entropy-density-like value $\tilde{s}(\tilde{L})$ of the MI initial state and 010101... CDW initial state with the same system size L as seen in Figs. 6 and 7.

To show Eq. (40), we first point out that there is an upper bound of elements of the matrix Z . This can be followed by the fact that $2Z^{\text{MI}} - I$ is a unitary matrix. Indeed, the relation

$$\begin{aligned} (2Z^{\text{MI}} - I)^\dagger (2Z^{\text{MI}} - I) &= (Y^\dagger)^T (2P_{L/2} - I) Y^T (Y^\dagger)^T (2P_{L/2} - I) Y^T = I \end{aligned} \quad (\text{E1})$$

holds from Eq. (B1). The unitarity of $2Z^{\text{MI}} - I$ ensures $\|Z^{\text{MI}} - I/2\|_2 = 1/2$, which leads to

$$\max_{j,l} \left(\left| z_{j,l}^{\text{MI}} - \frac{1}{2} \delta_{j,l} \right| \right) \leq \left\| Z^{\text{MI}} - \frac{1}{2} I \right\|_2 = \frac{1}{2}. \quad (\text{E2})$$




FIG. 8. Comparisons of the time evolution of the Rényi entropies for the quench to $U/J = 0, 0.2,$ and 0.4 from the (a) MI and (b) CDW initial states. The dashed line shows the Page value as the expected entanglement entropy when thermalization occurs. For finite U/J , the Rényi entanglement entropy is calculated from the exact diagonalization method [91,92]. To reduce the memory cost, it is calculated from a wave function utilizing $U(1)$ symmetry [93], associated with the conservation of the total particle number [94–97].

In particular, considering the case $j \neq l$ and recalling that all elements of Z^{CDW} are embedded in Z^{MI} as discussed in Sec. IV A, we obtain

$$\max_{j,l,j \neq l} (|z_{j,l}^{\text{MI}}|) \leq \frac{1}{2}, \quad \max_{j,l,j \neq l} (|z_{j,l}^{\text{CDW}}|) \leq \frac{1}{2}. \quad (\text{E3})$$

The function $\max_{l=1}^N (|z_{j,l}|, |\delta_{j,l} - z_{j,l}|)$ in the definition of $\tilde{s}(\tilde{L})$ in Eq. (39) always picks up the diagonal part of the matrix

Z because either $z_{j,j}$ or $1 - z_{j,j}$ is greater than $1/2$. Therefore, we get

$$\max_{l=1}^N (|z_{j,l}|, |\delta_{j,l} - z_{j,l}|) = \left| z_{j,j} - \frac{1}{2} \right| + \frac{1}{2}. \quad (\text{E4})$$

Substituting Eq. (E4) into Eq. (39), we immediately obtain Eq. (40).

Next, we show why the entropy-density-like values $\tilde{s}(\tilde{L})$ of the MI initial state and $010101 \cdots$ CDW initial state with the same system size L coincide. To this end, we first examine the relation between the matrices Z of the MI and CDW states. By definition, even index elements of Z^{MI} are simply related to elements of Z^{CDW} through

$$z_{2j,2l}^{\text{MI}} = z_{j,l}^{\text{CDW}} \quad (j, l = 1, 2, \dots, L/2). \quad (\text{E5})$$

Note that the sizes of Z^{MI} and Z^{CDW} are, respectively, $L \times L$ and $L/2 \times L/2$. Odd index elements of Z^{MI} are also related to elements of Z^{CDW} . The eigenfunction of the single-particle Hamiltonian $x_{k,j}$ is also the eigenfunction of the parity operator, satisfying $x_{k,L+1-j} = (-1)^{k+1} x_{k,j}$. This leads to $y_{L+1-j,l} = y_{j,L+1-l}$ and

$$z_{j,l}^{\text{MI}} + z_{L+1-j,L+1-l}^{\text{MI}} = \delta_{j,l}, \quad (\text{E6})$$

where we assume that $L_A = L/2$. Using Eqs. (E5) and (E6), we obtain

$$z_{2j-1,2l-1}^{\text{MI}} = \delta_{j,l} - z_{L/2+1-j,L/2+1-l}^{\text{CDW}}, \quad (\text{E7})$$

for $j, l = 1, 2, \dots, L/2$. Using the relations between the matrices Z of the MI and CDW states, Eqs. (E5) and (E7), and dividing the summation of $\tilde{s}(\tilde{L})$ in Eq. (40) of the MI initial state into the even and odd index parts, we conclude $\tilde{s}^{\text{MI}}(\tilde{L}) = \tilde{s}^{\text{CDW}}(\tilde{L})$.

The coincidence of $\tilde{s}(\tilde{L})$ between the MI and $010101 \cdots$ CDW initial states seen in Figs. 6 and 7 is rather special. This is true only for the MI and $010101 \cdots$ CDW initial states with $L_A = L/2$. When either condition is broken, such as considering $001001 \cdots$ CDW states or $L_A = L/4$, the coincidence disappears.

-
- [1] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2010).
- [2] C. Holzhey, F. Larsen, and F. Wilczek, *Nucl. Phys. B* **424**, 443 (1994).
- [3] P. Calabrese and J. Cardy, *J. Stat. Mech.* (2004) P06002.
- [4] P. Calabrese and J. Cardy, *J. Phys. A: Math. Theor.* **42**, 504005 (2009).
- [5] G. Vidal, J. I. Latorre, E. Rico, and A. Kitaev, *Phys. Rev. Lett.* **90**, 227902 (2003).
- [6] J. Eisert, M. Cramer, and M. B. Plenio, *Rev. Mod. Phys.* **82**, 277 (2010).
- [7] N. Laflorencie, *Phys. Rep.* **646**, 1 (2016).
- [8] A. Kitaev and J. Preskill, *Phys. Rev. Lett.* **96**, 110404 (2006).
- [9] M. Levin and X.-G. Wen, *Phys. Rev. Lett.* **96**, 110405 (2006).
- [10] Y. Zhang, T. Grover, and A. Vishwanath, *Phys. Rev. B* **84**, 075128 (2011).
- [11] S. V. Isakov, M. B. Hastings, and R. G. Melko, *Nat. Phys.* **7**, 772 (2011).
- [12] K. Życzkowski, *Open Syst. Inf. Dyn.* **10**, 297 (2003).
- [13] A. J. Daley, H. Pichler, J. Schachenmayer, and P. Zoller, *Phys. Rev. Lett.* **109**, 020505 (2012).
- [14] D. A. Abanin and E. Demler, *Phys. Rev. Lett.* **109**, 020504 (2012).
- [15] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, *Nature (London)* **415**, 39 (2002).
- [16] R. Islam, R. Ma, P. M. Preiss, M. Eric Tai, A. Lukin, M. Rispoli, and M. Greiner, *Nature (London)* **528**, 77 (2015).
- [17] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, *Science* **353**, 794 (2016).

- [18] P. Calabrese and J. Cardy, *J. Stat. Mech.* (2005) P04010.
- [19] R. Yoshii, S. Yamashika, and S. Tsuchiya, *J. Phys. Soc. Jpn.* **91**, 054601 (2022).
- [20] G. D. Chiara, S. Montangero, P. Calabrese, and R. Fazio, *J. Stat. Mech.* (2006) P03001.
- [21] J. H. Bardarson, F. Pollmann, and J. E. Moore, *Phys. Rev. Lett.* **109**, 017202 (2012).
- [22] A. Bauer, F. Dorfner, and F. Heidrich-Meisner, *Phys. Rev. A* **91**, 053628 (2015).
- [23] S. Goto and I. Danshita, *Phys. Rev. B* **99**, 054307 (2019).
- [24] A. M. Läuchli and C. Kollath, *J. Stat. Mech.* (2008) P05018.
- [25] M. Kunimi and I. Danshita, *Phys. Rev. A* **104**, 043322 (2021).
- [26] C. Rylands, B. Bertini, and P. Calabrese, *J. Stat. Mech.* (2022) 103103.
- [27] V. Alba and P. Calabrese, *Proc. Natl. Acad. Sci. USA* **114**, 7947 (2017).
- [28] R. Yao and J. Zakrzewski, *Phys. Rev. B* **102**, 104203 (2020).
- [29] M. Fagotti and P. Calabrese, *Phys. Rev. A* **78**, 010306(R) (2008).
- [30] S. Yamashika, D. Kagamihara, R. Yoshii, and S. Tsuchiya, *arXiv:2209.13340*.
- [31] V. Alba and P. Calabrese, *SciPost Phys.* **4**, 017 (2018).
- [32] I. Frérot and T. Roscilde, *Phys. Rev. B* **92**, 115129 (2015).
- [33] M. Cramer, C. M. Dawson, J. Eisert, and T. J. Osborne, *Phys. Rev. Lett.* **100**, 030602 (2008).
- [34] P. Barmettler, D. Poletti, M. Cheneau, and C. Kollath, *Phys. Rev. A* **85**, 053625 (2012).
- [35] M. Cheneau, P. Barmettler, D. Poletti, M. Endres, P. Schauß, T. Fukuhara, C. Gross, I. Bloch, C. Kollath, and S. Kuhr, *Nature (London)* **481**, 484 (2012).
- [36] R. Berkowitz and P. Devlin, *Isr. J. Math.* **224**, 437 (2018).
- [37] Y. Takasu, T. Yagami, H. Asaka, Y. Fukushima, K. Nagao, S. Goto, I. Danshita, and Y. Takahashi, *Sci. Adv.* **6**, eaba9255 (2020).
- [38] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, *Nat. Phys.* **8**, 325 (2012).
- [39] J. Zhang and M. A. Rajabpour, *J. Stat. Mech.* (2021) 093101.
- [40] S.-G. Hwang, *Am. Math. Mon.* **111**, 157 (2004).
- [41] S. Fisk, *Am. Math. Mon.* **112**, 118 (2005).
- [42] M. Marcus, *Am. Math. Mon.* **67**, 215 (1960).
- [43] H. S. Wilf, *Can. J. Math.* **18**, 758 (1966).
- [44] R. Merris, *Am. Math. Mon.* **80**, 791 (1973).
- [45] S. Friedland, *Ann. Math.* **110**, 167 (1979).
- [46] H. Minc, *Permanents, Encyclopedia of Mathematics and Its Applications* (Cambridge University Press, Cambridge, UK, 1984), Vol. 6.
- [47] G.-S. Cheon and I. M. Wanless, *Lin. Algebra Appl.* **403**, 314 (2005).
- [48] M. Laurent and A. Schrijver, *Am. Math. Mon.* **117**, 903 (2010).
- [49] L. Gurvits and A. Samorodnitsky, in *2014 IEEE 55th Annual Symposium on Foundations of Computer Science* (IEEE, New York, 2014), pp. 90–99.
- [50] M. Marcus and M. Newman, *Ann. Math.* **75**, 47 (1962).
- [51] L. Gurvits, in *Mathematical Foundations of Computer Science 2005*, edited by J. Jędrzejowicz and A. Szepletowski (Springer, Berlin, 2005), pp. 447–458.
- [52] H. J. Ryser, *Combinatorial Mathematics* (American Mathematical Society, Providence, RI, 1963), Vol. 14.
- [53] R. A. Brualdi and H. J. Ryser, *Combinatorial Matrix Theory* (Cambridge University Press, Cambridge, UK, 1991).
- [54] D. G. Glynn, *Eur. J. Comb.* **31**, 1887 (2010).
- [55] K. Balasubramanian, Ph.D. thesis, Indian Statistical Institute-Kolkata (1980).
- [56] E. Bax and J. Franklin, CalTech Report No. CS-TR-96-04 (1996).
- [57] E. T. Bax, *Finite-Difference Algorithms for Counting Problems* (California Institute of Technology, Pasadena, CA, 1998).
- [58] D. G. Glynn, *Des. Codes Cryptogr.* **68**, 39 (2013).
- [59] F. Gray, U.S. Patent No. 2,632,058 (1953).
- [60] A. Nijenhuis and H. S. Wilf, *Combinatorial Algorithms: For Computers and Calculators* (Elsevier, Amsterdam, 1978).
- [61] B. Gupt, J. Izaac, and N. Quesada, *J. Open Source Software* **4**, 1705 (2019).
- [62] A. Neville, C. Sparrow, R. Clifford, E. Johnston, P. M. Birchall, A. Montanaro, and A. Laing, *Nat. Phys.* **13**, 1153 (2017).
- [63] J. Wu, Y. Liu, B. Zhang, X. Jin, Y. Wang, H. Wang, and X. Yang, *Natl. Sci. Rev.* **5**, 715 (2018).
- [64] P.-H. Lundow and K. Markström, *J. Comput. Phys.* **455**, 110990 (2022).
- [65] S. Chin and J. Huh, *Sci. Rep.* **8**, 6101 (2018).
- [66] A. Flesch, M. Cramer, I. P. McCulloch, U. Schollwöck, and J. Eisert, *Phys. Rev. A* **78**, 033608 (2008).
- [67] M. Cramer, A. Flesch, I. P. McCulloch, U. Schollwöck, and J. Eisert, *Phys. Rev. Lett.* **101**, 063001 (2008).
- [68] I. Frérot and T. Roscilde, *Phys. Rev. Lett.* **116**, 190401 (2016).
- [69] G. Biroli, C. Kollath, and A. M. Läuchli, *Phys. Rev. Lett.* **105**, 250401 (2010).
- [70] S. Sorg, L. Vidmar, L. Pollet, and F. Heidrich-Meisner, *Phys. Rev. A* **90**, 033606 (2014).
- [71] D. N. Page, *Phys. Rev. Lett.* **71**, 1291 (1993).
- [72] A. Russomanno, M. Fava, and R. Fazio, *Phys. Rev. B* **102**, 144302 (2020).
- [73] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, *Phys. Rev. Lett.* **98**, 050405 (2007).
- [74] Y. O. Nakagawa, M. Watanabe, H. Fujita, and S. Sugiura, *Nat. Commun.* **9**, 1635 (2018).
- [75] G. Carleo, F. Becca, L. Sanchez-Palencia, S. Sorella, and M. Fabrizio, *Phys. Rev. A* **89**, 031602(R) (2014).
- [76] K. Nagao, M. Kunimi, Y. Takasu, Y. Takahashi, and I. Danshita, *Phys. Rev. A* **99**, 023622 (2019).
- [77] R. Kaneko and I. Danshita, *Commun. Phys.* **5**, 65 (2022).
- [78] R. Modak and T. Nag, *Phys. Rev. Res.* **2**, 012074(R) (2020).
- [79] R. Singh, R. Moessner, and D. Roy, *Phys. Rev. B* **95**, 094205 (2017).
- [80] S. Aubry and G. André, *Ann. Isr. Phys. Soc.* **3**, 18 (1980).
- [81] T. Devakul and D. A. Huse, *Phys. Rev. B* **96**, 214201 (2017).
- [82] X. Cao, A. Tilloy, and A. De Luca, *SciPost Phys.* **7**, 024 (2019).
- [83] O. Alberton, M. Buchhold, and S. Diehl, *Phys. Rev. Lett.* **126**, 170602 (2021).
- [84] T. Minato, K. Sugimoto, T. Kuwahara, and K. Saito, *Phys. Rev. Lett.* **128**, 010603 (2022).
- [85] M. Block, Y. Bao, S. Choi, E. Altman, and N. Y. Yao, *Phys. Rev. Lett.* **128**, 010604 (2022).
- [86] T. Müller, S. Diehl, and M. Buchhold, *Phys. Rev. Lett.* **128**, 010605 (2022).
- [87] F. P. M. Méndez-Córdoba, J. J. Mendoza-Arenas, F. J. Gómez-Ruiz, F. J. Rodríguez, C. Tejedor, and L. Quiroga, *Phys. Rev. Res.* **2**, 043264 (2020).

- [88] J. J. Mendoza-Arenas and B. Buča, [arXiv:2106.06277](#).
- [89] S. Aaronson and A. Arkhipov, in *Proceedings of the Forty-Third Annual ACM Symposium on Theory of Computing* (Association for Computing Machinery, New York, 2011), pp. 333–342.
- [90] S. Aaronson and T. Hance, *Quantum Inf. Comput.* **14**, 541 (2014).
- [91] P. Weinberg and M. Bukov, *SciPost Phys.* **2**, 003 (2017).
- [92] P. Weinberg and M. Bukov, *SciPost Phys.* **7**, 020 (2019).
- [93] J.-H. Jung and J. D. Noh, *J. Korean Phys. Soc.* **76**, 670 (2020).
- [94] J. Schnack, P. Hage, and H.-J. Schmidt, *J. Comput. Phys.* **227**, 4512 (2008).
- [95] J. Zhang and R. Dong, *Eur. J. Phys.* **31**, 591 (2010).
- [96] A. Szabados, P. Jeszenszki, and P. R. Surján, *Chem. Phys.* **401**, 208 (2012).
- [97] D. Raventós, T. Graß, M. Lewenstein, and B. Juliá-Díaz, *J. Phys. B: At. Mol. Opt. Phys.* **50**, 113001 (2017).