




Multi-sideband interference structures by high-order photon-induced continuum-continuum transitions in helium

D. Bharti ^{1,*} H. Srinivas,¹ F. Shobeiry,¹ A. T. Bondy,^{2,3} S. Saha ^{3,4} K. R. Hamilton ⁵
R. Moshhammer,¹ T. Pfeifer,¹ K. Bartschat,³ and A. Harth^{1,6,†}

¹Max-Planck-Institute for Nuclear Physics, D-69117 Heidelberg, Germany

²Department of Physics, University of Windsor, Windsor, Ontario, Canada N9B 3P4

³Department of Physics and Astronomy, Drake University, Des Moines, Iowa 50311, USA

⁴Department of Physics, Panskura Banamali College (Autonomous), Panskura, West Bengal 721152, India

⁵Department of Physics, University of Colorado at Denver, Denver, Colorado 80204, USA

⁶Center for Optical Technologies, Aalen University, D-73430 Aalen, Germany



(Received 21 September 2023; accepted 9 January 2024; published 12 February 2024)

Following up on a previous paper on two-color photoionization of Ar($3p$) [D. Bharti *et al.*, *Phys. Rev. A* **103**, 022834 (2021)], we present measurements and calculations for a modified three-sideband (3-SB) version of the reconstruction of attosecond beating by interference of two-photon transitions (RABBITT) configuration applied to He($1s$). The 3-SB RABBITT approach allows us to explore interference effects between pathways involving different orders of transitions within the continuum. The relative differences in the retrieved oscillation phases of the three sidebands provide insights into the continuum-continuum transitions. The ground state of helium has zero orbital angular momentum, which simplifies the analysis of oscillation phases and their angle dependence within the three sidebands. We find qualitative agreement between our experimental results and the theoretical predictions for many cases but also observe some significant quantitative discrepancies.

DOI: [10.1103/PhysRevA.109.023110](https://doi.org/10.1103/PhysRevA.109.023110)

I. INTRODUCTION

The reconstruction of attosecond beating by interference of two-photon transitions (RABBITT) is a widely-employed technique to characterize attosecond pulse trains and measure attosecond time delays in photoionization processes (e.g., Refs. [1–3]). In the standard scheme, photoionization by various spectral harmonics in an attosecond extreme ultraviolet (XUV) pulse train (the pump photons) results in multiple discrete peaks (main peaks) within the photoelectron signal. Simultaneously, the presence of a time-delayed infrared (IR) field (the probe photons) creates an additional photoelectron peak between every two main peaks, referred to as “sidebands.” The photoelectron yield within these sidebands oscillates as a function of the time delay, and this oscillatory pattern can be utilized to determine the relative photoionization time delay. However, in order to accurately determine photoionization time delays using the RABBITT method, it is essential to consider the contribution arising from the continuum-continuum transitions induced by the probe

pulse. This complexity is simplified by breaking down the phases of the two-photon transition matrix element into the sum of the Wigner phase, associated with the single-photon ionization process, and the continuum-continuum phase denoted as ϕ_{cc} . Details can be found, for example, in Refs. [4–6].

In 2019, Harth *et al.* [7] introduced a variant of the RABBITT scheme known as three-sideband (3-SB) RABBITT, in which the interaction with the probe pulse results in the creation of not just one but three sidebands between every two main peaks. In the context of the 3-SB RABBITT approach, Bharti *et al.* [8] extended the “decomposition approximation” to determine the phase of the N th-order above-threshold-ionization (ATI) matrix element. In this extension, the ϕ_{cc} contributions from each participating transition are simply summed. In the 3-SB scheme, one important consequence of the decomposition approximation is the prediction that the oscillation phases in the three sidebands formed between the same pair of main peaks should be identical, except for an additional π phase shift in the central sideband. Numerical calculations performed for atomic hydrogen revealed slight deviations from this expectation, and these deviations consistently diminished with an increase in kinetic energy. Experiments on atomic hydrogen, of course, are extremely challenging due to the difficulties in creating a dense H target.

A proof-of-principle experimental realization of the 3-SB scheme was reported by Bharti *et al.* [9] for an argon target. As a noble gas with the first ionization potential of ≈ 15.8 eV for the $3p$ electron, this target was experimentally favorable due to the fact that low-order (7, 9, 11, ...) 515-nm harmonics

*bharti@mpi-hd.mpg.de

†anne.harth@hs-aalen.de

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Open access publication funded by the Max Planck Society.

of the frequency-doubled fundamental field (1030 nm) in the XUV pulse train were able to ionize the atom. However, ionization of the 3s electron is also possible, resonances occur at relatively low ejected-electron energies of 10–15 eV due to the possible inner-shell promotion of the 3s electron, and the numerical treatment is very challenging. A further complication arises from the fact that already the XUV step alone promotes the 3p electron to two different angular momentum states (*s* and *d*). All this, together with the uncertainties regarding the detailed time dependence of the electric fields, resulted in only qualitative agreement between the experimental data and theoretical predictions based on the *R*-matrix with time dependence (RMT) approach.

In this paper, we report the outcomes of a 3-SB RABBITT experiment performed on helium. The ground state of helium has zero orbital angular momentum, and hence the single-photon ionization induced by an XUV pulse results in the creation of a photoelectron with a single orbital angular momentum $\ell = 1$. This simplifies the interpretation of the photoelectron interference patterns resulting from the interaction of the photoelectron with the IR photons. Due to its higher ionization potential of approximately 24.6 eV, however, helium presents further experimental challenges compared to argon, as it requires relatively high-energy XUV photons to create a sufficient number of main peaks. On the other hand, from a theoretical perspective, helium is significantly easier to handle than argon. In fact, since the remaining 1s electron is tightly bound, a single-active-electron (SAE) approach may be sufficient to explain the basic features.

This paper is organized as follows. We begin with a brief review of the basic idea behind the 3-SB setup in Sec. II. This is followed by a description of the experimental apparatus in Sec. III and the accompanying theoretical SAE and RMT approaches in Sec. IV. We first show angle-integrated data in Sec. V A before focusing on the angle dependence of the RABBITT phases in the three sidebands of each individual group in Sec. V B. We finish with a summary and an outlook in Sec. VI.

II. THE 3-SB SCHEME FOR HELIUM

In this section, we briefly review the 3-SB scheme introduced in Ref. [7] and the analytical treatment presented in Ref. [8], as applied to the 3-SB RABBITT experiment in general and then in our particular case of the helium target.

The basic scheme is illustrated in Fig. 1. The active electron originates from an *s* orbital, and the absorption of an XUV photon with a frequency corresponding to either H_{q-1} or H_{q+1} transitions it to a *p* orbital. This transition gives rise to the main photoelectron peaks labeled M_{q-1} and M_{q+1} , respectively. The frequencies H_{q-1} and H_{q+1} correspond to the $(q-1)$ and $(q+1)$ odd harmonics, respectively, of the pulse used for generating the XUV train via high-order harmonic generation (HHG) [10,11]. Additional transitions within the continuum, resulting from either the absorption or emission of probe photons, lead to the emergence of three sidebands situated between two main peaks. These transitions to the sidebands can traverse different angular momentum channels in compliance with the dipole selection rule. We designate the trio of sidebands positioned between M_{q-1} and M_{q+1} accord-




FIG. 1. 3-SB transition diagram with all angular momentum channels, illustrating only the lowest-order pathways from emission and absorption processes necessary for yield oscillations in each sideband. I_p is the ionization threshold, and the dashed line just above it labels the special case of the threshold sideband. See text for details. The solid and dashed lines in the transitions illustrate a propensity rule [13–15], with the solid line indicating a more probable transition compared to the dashed line.

ing to their energy positions within the group, labeling them as $S_{q,l}$, $S_{q,c}$, and $S_{q,h}$, where the second subscript designates the lower, central, and higher sidebands, respectively. All pathways leading to the same sideband interfere to produce the net photoelectron yield of the sideband. Changing the time delay between the XUV and IR pulses adds a dynamic phase to the interference of absorption and emission paths, leading to oscillations in the sideband yield.

As shown in Ref. [9], the general form of the signal in the sidebands is given by

$$S_{q,j}(\tau, \theta) = A_{q,j}(\theta) + B_{q,j}(\theta) \cos(4\omega\tau - \phi_{R,q,j}(\theta)) \quad (1)$$

with j standing for (*l*, *c*, *h*). Each signal is characterized by a constant term *A*, an oscillation amplitude *B*, and a RABBITT phase ϕ_R . As seen from the above equation, each of these parameters depends on the sideband group *q*, the location of the sideband within that group (*l*, *c*, *h*), and the detection angle θ . In angle-integrated detection mode, the angle dependence is averaged over, but the general form of the equation remains unchanged. In both the angle-differential and angle-integrated cases, the signal oscillates as a function of the delay τ with an angular frequency of four times the angular frequency ω of the fundamental IR field. Detailed expressions of the above parameters in terms of transition amplitudes can be found in Refs. [8,9].

III. EXPERIMENT

Since the experimental methodology was described in detail in Ref. [9], we only provide a brief summary of the most important settings for the current experiment. The fiber-based laser utilized in the experiment emits 50 fs (full width at half maximum) ultrashort infrared pulses centered around 1030 nm with a pulse energy of 1.2 mJ. The pulse is divided into two arms of a Mach-Zehnder-type interferometer using a holey mirror, which reflects approximately 85% of the beam in the pump arm and allows the remainder to pass through into the probe arm. Within the pump arm, a barium borate crystal is employed to produce the second harmonic (515 nm) of the laser pulse with an efficiency of 25–30%. The fundamental IR beam is subsequently filtered via a dichroic mirror, and the second harmonic is focused onto an argon gas jet to produce an XUV pulse train through HHG. The driving 515-nm beam was spatially filtered out from the beam line, and the resulting XUV beam then passed through a 150-nm-thick aluminum filter. Meanwhile, the IR beam within the probe arm was directed through a retroreflector mounted on a piezoelectric-translation stage, before it was spatially and temporally recombined with the XUV beam. Both the XUV and IR beams were focused inside a reaction microscope (ReMi) [12] onto a supersonic gas jet of the target species.

The relative strength of the harmonics was determined by analyzing the photoelectron spectra generated solely by the XUV beam in different gases. Beyond the helium ionization threshold, four harmonics (H_{11} , H_{13} , H_{15} , and H_{17}) were observed, with their strength decreasing as the photon energy increased. Below the ionization threshold, H_7 is positioned at the low-energy transmission edge of an aluminum filter, resulting in significant attenuation of this harmonic and effectively elimination of all harmonics below H_7 . The harmonic H_9 , whose energy is also below the ionization energy, is crucial for generating oscillations in the yield of the threshold sideband (see the dashed line just above the ionization threshold in Fig. 1). This threshold sideband (S_{th}) belongs to the so-called uRABBITT scheme [16–22].

We adjusted the IR probe intensity by utilizing an iris in the probe arm. RABBITT measurements were performed at three IR probe peak intensities: $I_1 \approx 5 \times 10^{11}$ W/cm², $I_2 \approx 7 \times 10^{11}$ W/cm², and $I_3 \approx 1.2 \times 10^{12}$ W/cm². The IR and XUV fields were linearly polarized parallel to each other and along the spectrometer axis of the ReMi. The latter enables the reconstruction of the three-dimensional momenta of the electrons and corresponding ions (in coincidence) created during the photoionization process [12]. Furthermore, the ReMi is capable of selecting electrons emitted in different directions, thereby enabling the construction of photoelectron angular distributions (PADs).

The XUV-IR temporal delay was sampled at regular intervals of $T_0/60$, where $T_0 = 3.44$ fs is the period of the IR pulse. The delay was scanned over a range equivalent to one and a half times the optical cycle of the IR pulse. The XUV-IR beam line was actively stabilized [9,23] to achieve a stability of approximately 30 as over a data acquisition period of more than 10 hours.

IV. THEORY

The numerical approaches to model the experiment were also described in detail in previous publications. Hence, we again limit ourselves to a brief summary with references given at the appropriate spots.

A. The SAE approach

We employed the same SAE model as Birk *et al.* [24] and Meister *et al.* [25]. Specifically, we used the one-electron potential

$$V(r) = -\frac{1}{r} - \left(\frac{1}{r} + 1.3313\right) \exp(-3.0634r), \quad (2)$$

where r is the distance from the nucleus, to calculate the valence orbitals. The difference of excitation energies compared to the recommended excitation from the NIST database [26] is less than 0.2 eV even in the worst-case scenario.

For both the XUV pulse train and the fundamental IR, we used temporal fields based on the measured spectrum of the IR pulse and on the measured XUV-only photoionization spectrum considering the experimental resolution. This gives reasonably accurate information about the relative strength of the harmonics. Additionally, we incorporated an estimated attochirp into the theoretical XUV pulse train.

Since both the XUV pulse train and the fundamental IR are linearly polarized along the same direction, the initial state can be propagated very efficiently and accurately. Specifically, we used an updated version of the code described by Douguet *et al.* [27]. In contrast to the heavier noble gases, the SAE approach is expected to be suitable for the helium target, as long as obvious two-electron correlation effects, e.g., autoionizing resonances, are not affecting the process significantly. This is, indeed, the case for the present study.

B. The RMT approach

As a second method, we employed the general RMT method [28]. To calculate the necessary time-independent basis functions and dipole matrix elements for the present work, we set up the simplest possible model, a nonrelativistic one-state approach. This model, labeled RMT below, enables efficient calculations whose predictions can be readily compared with both the experimental data and those from the SAE approach to make a first assessment regarding the likely quality of the theoretical predictions.

We took the same pulse as in the SAE calculation. Instead of the numerical orbital employed in the SAE calculation, which is close to the Hartree-Fock orbital of the ground-state configuration, however, we took the known $1s$ orbital of He^+ . While this is not optimal to obtain the very best ground-state energy, this disadvantage is mitigated almost completely by the continuum-continuum terms in the R -matrix Hamiltonian. Using this orbital will, however, be advantageous in future studies, where we plan to include additional states in the close-coupling expansion to check the convergence and sensitivity of the numerical predictions. In contrast to SAE, exchange effects between the two electrons are treated explicitly within the R -matrix box.

V. RESULTS AND DISCUSSION

We first present in Sec. V A our angle-integrated RABBITT results. This is followed by a discussion of angle-differential measurements in Sec. V B. For the latter case, we will concentrate on only two sideband groups, S_{12} and S_{14} , where we have a sufficient amount of data to conduct the angle-resolved investigation of the RABBITT phase.

A. Angle-integrated RABBITT scans

Figure 2 shows the results from three angle-integrated RABBITT measurements recorded at the probe intensities $I_1 \approx 5 \times 10^{11}$ W/cm² (top), $I_2 \approx 7 \times 10^{11}$ W/cm² (center), and $I_3 \approx 12 \times 10^{11}$ W/cm² (bottom). The logarithmic-scale color map in the lower part of each panel shows the RABBITT trace obtained by subtracting the delay-integrated signal from the original trace to highlight the oscillations. The upper part of the panels displays the XUV-only photoelectron spectrum (gray line) and the delay-integrated photoelectron spectrum (red line), both normalized to their peak values and plotted on a logarithmic scale.

Looking at the RABBITT traces, we see that the central sideband displays the clearest oscillation at all applied IR intensities. This was expected since both the absorption and the emission paths that populate this sideband are of the same order. In the lower and higher sidebands, the two interfering terms are of different (second and fourth) order, and hence the contrast in the oscillation is reduced compared to the central sideband.

Next, the RABBITT traces demonstrate that the oscillation contrast is generally better for the higher sideband in each group than for the lower one. This is due to the rapid decrease of the main peaks with increasing energy, which makes the amplitude of the two interfering transition paths more balanced in the case of the higher sideband. In $S_{12,h}$, for example, the upper main peak (M_{13}) is much weaker than the lower main peak (M_{11}), resulting in the magnitude of a three-photon transition from the stronger lower main peak ($M_{11} + 3\omega$) becoming comparable to the magnitude of a one-photon transition from the weaker upper main peak ($M_{13} - \omega$). Hence the interference of these two terms leads to a strong delay-dependent oscillation. In contrast, for the lower sideband $S_{12,l}$, the magnitude of a one-photon transition from the lower main peak ($M_{11} + \omega$) is much stronger than the magnitude of a three-photon transition from the already weaker upper main peak ($M_{13} - 3\omega$), thus resulting in small contrast of the oscillation.

Additionally, at the lowest applied IR intensity (top panel of Fig. 2), the highest-energy main peak M_{17} is almost entirely depleted, but it becomes repopulated as the intensity is increased. The same is seen for the main peak M_{15} , which first becomes weaker in the presence of the IR pulse but then gains strength when the intensity is increased. Looking at the S_{16} group, we observe that the contrast in the lower sideband ($S_{16,l}$) is very weak at the lowest applied intensity, but it gradually improves as the IR intensity increases. On the other hand, the contrast of the oscillation in the higher sideband $S_{16,h}$ deteriorates with increasing IR intensity.

When the kinetic energy or the IR intensity is increased, higher-order transitions should be accounted for as well.




FIG. 2. Results from the angle-integrated RABBITT measurements taken at peak IR intensities $I_1 \approx 5 \times 10^{11}$ W/cm² (top), $I_2 \approx 7 \times 10^{11}$ W/cm² (center), and $I_3 \approx 12 \times 10^{11}$ W/cm² (bottom). The logarithmic color map in each of the lower panels displays the angle-integrated RABBITT traces, while the upper panel shows the XUV-only (gray line) and the delay-integrated photoelectron spectrum (red line) normalized to the peak value. The dashed line in the top panels is the (relative) signal predicted by the RMT model. The main lines and the sideband groups are also indicated. The right y axis shows the phases of the delay-dependent yield oscillations of the sidebands obtained from our fitting method, along with their estimated fitting errors. Since the absolute phase is not known experimentally, we set it to zero for the center sideband in the S_{12} group. Also, a phase $\pm\pi$ was added to the center sideband to simplify the comparison with the other two. The solid circles represent the experimental data while the open symbols are the theoretical SAE (circles) and RMT (boxes) predictions.

Figure 3 shows a selection of many-order transition pathways leading to the lower sideband of the S_{16} group and the interference schemes contributing to the oscillations of the yield. At the low IR intensity, the oscillation in the photoelectron yield of the lower sideband is predominantly influenced by




FIG. 3. Selected interference schemes for 4ω oscillations in the lower sideband. Scheme T_A dominates at low probe intensity. The oscillation generated by scheme T_A is out of phase by π compared to the oscillations produced by schemes T_B , T_C , T_D , and T_E .

the interference scheme T_A . However, with increasing kinetic energy or IR intensity, the involvement of higher-order transitions featuring M_{15} and M_{13} becomes significant in shaping the oscillations of the yield.

In order to determine the phases of the oscillation from each sideband, the photoelectron spectrum was first integrated within an energy window of 0.8 eV across the peak and was then fitted to a cosine function of the form $A + B \cos(4\omega\tau - \phi_R)$. For the lower and higher sidebands, the trivial π phase was removed from the obtained phases. As the absolute experimental phase is unknown, the retrieved phases from the three measurements were shifted to align the data point for $S_{12,c}$ at zero for comparison of the relative phases. The phases of the oscillations obtained from the fitting process, along with their corresponding fitting errors, are shown in the upper panels of Fig. 2 and also listed in Table I.

Several noteworthy results can be seen from Table I:

(1) After accounting for the additional phase of π , the phases in the S_{12} and S_{14} groups are nearly the same in all three sidebands of the respective group. This is in excellent agreement with the “decomposition approximation” [8].

(2) Relative to S_{12} , the phases in the S_{14} group are larger by ≈ 0.5 rad.

(3) The lower sideband in S_{16} exhibits a significantly different phase compared to the other two. This is likely due to the contribution of the higher-order interference schemes (T_B , T_C , T_D , and T_E in Fig. 3) dominating over the lowest-order interference (T_A). Since these interference schemes, involving higher-order pathways, produce oscillations at the same frequency but π out of phase with that arising from the lowest-order interference scheme T_A , their contribution in the yield oscillation results in the observed change of $\approx \pi$. It should also be noted that the oscillation due to T_E includes the spectral-phase difference of H_{15} and H_{13} , while the schemes T_A , T_B , T_C , and T_D contain the spectral phase difference of H_{17} and H_{15} .

Regarding the agreement between experiment and theory, we see that both SAE and RMT agree with the experimental findings of nearly the same phases for each member of the S_{12} and S_{14} groups. Furthermore, the theoretical phases within S_{14} are systematically larger than those observed in the experiment. This is simply due to an overestimate of the chirp that was included in the description of the XUV pulse. Since RABBITT calculations are very time consuming, we decided not to repeat them with only the chirp being reduced. There is also good qualitative agreement in the fact that $S_{16,c}$ and $S_{16,h}$ have about the same RABBITT phase, while that of $S_{16,l}$ is very different.

Finally, there is significant quantitative disagreement between experiment and the two theories, as well as between the two theoretical predictions themselves, for the threshold sideband S_{th} , which is the upper member of the S_{10} group and the only one above the ionization threshold. However, there is general agreement in the qualitative finding of a strong intensity dependence. This strong sensitivity to the intensity is caused by near-resonant interactions with Rydberg states.

B. Angle-differential RABBITT scans

To examine the angle dependence of the RABBITT phases, we analyzed the photoelectrons emitted at various angles relative to the polarization axis of the driving light fields. This involved segregating the data into angle-differential data sets, which were generated by integrating the photoelectron yield over 10° angular windows at 5° intervals. The phase of the oscillation in each sideband was extracted from each angle-differential data set using the same method as outlined in the angle-integrated case. The photoelectron spectra from

TABLE I. Angle-integrated RABBITT phase (in radians) obtained from the fitting procedure to the measurements at three IR intensities.

Probe intensity	S_{10} (S_{th})			S_{12}			S_{14}			S_{16}		
	S_h	S_l	S_c	S_h	S_l	S_c	S_h	S_l	S_c	S_h		
I_1	-1.96 ± 0.02	0.11 ± 0.02	0.00 ± 0.01	0.08 ± 0.02	0.52 ± 0.06	0.52 ± 0.01	0.55 ± 0.01	-2.18 ± 0.49	0.67 0.08	0.60 ± 0.06		
I_2	-1.29 ± 0.01	0.31 ± 0.02	0.00 ± 0.01	0.07 ± 0.01	0.58 ± 0.04	0.56 ± 0.01	0.58 ± 0.01	-2.46 ± 0.11	0.56 0.04	0.54 ± 0.07		
I_3	-0.64 ± 0.01	0.32 ± 0.02	0.00 ± 0.01	0.07 ± 0.01	0.73 ± 0.02	0.53 ± 0.11	0.59 ± 0.01	-2.53 ± 0.05	0.61 0.03	0.11 ± 1.40		




FIG. 4. Angle-dependent phase retrieved from the S_{12} group in the RABBITT scans for IR peak intensities of (a) $I_1 \approx 5 \times 10^{11} \text{ W/cm}^2$, (b) $I_2 \approx 7 \times 10^{11} \text{ W/cm}^2$, and (c) $I_3 \approx 12 \times 10^{11} \text{ W/cm}^2$.

each sideband were integrated over a 0.6-eV energy window centered on the peak, and the resulting delay-dependent signal was fitted to a cosine function to obtain the phase.

Figures 4 and 5 illustrate the angle dependence of the retrieved phases from the three sidebands in the S_{12} and S_{14} groups, respectively, for different IR intensities: I_1 [Figs. 4(a) and 5(a)], I_2 [Figs. 4(b) and 5(b)], and I_3 [Figs. 4(c) and 5(c)]. For better comparison and since we do not know the absolute experimental phase, a common phase was added to the three sidebands of each group to shift the central sideband phase to zero at the first data point (5°) for each intensity. The statistical uncertainty of the data above 60° was insufficient to retrieve a meaningful oscillation phase at all, and the estimated fitting errors suggest some caution regarding the extracted phases at the larger angles, especially at the lowest intensity. The figures demonstrate a consistent angle-dependent RABBITT phase in the central sidebands of both the S_{12} and S_{14} groups across all three probe intensities. In contrast, the lower and the higher sidebands exhibit some dependence on both their group and the IR intensity.

We first discuss the angle-dependent oscillation phase in the central sideband while considering only the lowest-order transitions depicted in Fig. 1. According to the selection rules for electric dipole transitions, a three-photon transition to the

central sideband results in the creation of a combination of two partial waves corresponding to angular momentum states of p ($\ell=1$) and f ($\ell=3$), where $m=0$ is conserved. The amplitudes of these partial waves vary with the angle of electron emission and are determined by the spherical harmonics $Y_{\ell,0}(\theta)$. In our specific experimental setup, where both beams are linearly polarized along the same direction, we can substitute the spherical harmonics with Legendre polynomials denoted as $P_\ell(\theta)$. Consequently, the interference of all partial waves generated during the absorption and emission processes will result in a signal of the form

$$S_c(\tau, \theta) = A_c(\theta) + B_c(\theta) \cos(4\omega\tau - \phi_{R,c}(\theta)) \quad (3a)$$

$$= A_c(\theta) + a_{11}P_1^2(\theta) \cos(4\omega\tau - \phi_{11}) \quad (3b)$$

$$+ a_{33}P_3^2(\theta) \cos(4\omega\tau - \phi_{33}) \quad (3c)$$

$$+ P_1(\theta)P_3(\theta)[a_{13} \cos(4\omega\tau - \phi_{13}) + a_{31} \cos(4\omega\tau - \phi_{31})]. \quad (3d)$$

Here, $A_c(\theta)$ represents the signal that does not oscillate during the temporal delay scan. Each oscillation term is characterized by an amplitude determined by the magnitude of the dipole transition matrix elements $a_{\ell,\nu}$ and products of the Legendre polynomials $P_\ell(\theta)P_{\nu}(\theta)$, as well as a phase




FIG. 5. Angle-dependent phase retrieved from the S_{14} group in the RABBITT scans for IR peak intensities of (a) I_1 , (b) I_2 , and (c) I_3 .

$\phi_{\ell,\ell'}$. Here, ℓ and ℓ' span across the possible orbital angular momentum quantum numbers reached in the absorption and emission processes, respectively. Referring to Eqs. (3), we use the term “same-channel interference” [Eqs. (3b) and (3c)] to indicate interference occurring between two partial waves with the same angular momentum ($\ell = \ell'$) and “cross-channel interference” [Eq. (3d)] to indicate interference between two partial waves with different angular momenta ($\ell \neq \ell'$).

The oscillation phase $\phi_{\ell,\ell'}$ of each interference term in Eq. (3) includes the XUV chirp, the Wigner phases, and phases arising from continuum-continuum transitions. While the XUV chirp and Wigner phases remain constant across all $\phi_{\ell,\ell'}$, the contribution of the continuum-continuum coupling phase (ϕ_{cc}) varies based on the angular momentum of the states involved in the transition pathways. However, the variability in ϕ_{cc} across distinct angular momenta is typically small, usually less than $\pi/10$ for kinetic energies exceeding 5 eV. Moreover, this variability diminishes with increasing kinetic energy [15]. Consequently, differences among the oscillation phases $\phi_{\ell,\ell'}$ across various interference terms are generally small, decreasing further with increasing kinetic energies.

Due to the distinct oscillation phase $\phi_{\ell,\ell'}$ in each of the interfering terms and the variation in the associated oscillation amplitude, which contains the Legendre polynomials, with the angle of electron emission, the overall retrieved oscillation phase $\phi_R(\theta)$ is angle dependent [29,30]. A significant change in $\phi_R(\theta)$ may occur when the electron emission angle gets close to the angle where the cross-channel interference amplitude equals the combined amplitude of other interference terms. In the cross-channel interference expressions within Eqs. (3), the product $P_1(\theta)P_3(\theta)$ changes sign upon passing through a node in one of the Legendre polynomials. This sign change is effectively equivalent to adding π in the oscillation phase ϕ_{13} and ϕ_{31} . This phase addition results in a sudden change in the overall RABBITT phase ϕ_R around the angle where the cross-channel amplitude surpasses that of the same-channel interference terms. The drop in the RABBITT phase $\phi_R(\theta)$ in the central sideband in Fig. 4(a) above 40° suggests that the angle-dependent amplitude of the cross-channel (p - f) interference is larger than that of the same-channel (p - p and f - f) interferences. Since the angle-dependent behavior of the RABBITT phase for the central sideband remains consistent across all three applied probe intensities, the aforementioned arguments apply to all three cases.

In the subsequent sideband group S_{14} , the discrepancy in the oscillation phase $\phi_{\ell,\ell'}$ among various interfering terms reduces due to the further decreased variation in ϕ_{cc} between different angular momenta. This leads to an almost negligible angular dependence of the RABBITT phase until the emission angle approaches a critical point where the dominance of cross-channel interference over other interfering terms triggers an abrupt π phase shift. While this is not seen in the angular regime for which we have experimental data, it is very noticeable in the theoretical predictions for the larger angles (cf. Fig. 6 below).

We next discuss the angular variation of the RABBITT phase in the lower and higher sidebands of S_{12} for the case of

the lowest applied intensity, as shown in Fig. 4(a). In both the lower and higher sidebands, a four-photon transition populates s , d , and g states, while a two-photon process populates s and d states (cf. Fig. 1).

Based on the propensity rule for continuum-continuum transitions [13–15], the creation of a g electron in the lower sideband via emission of three IR photons is least likely. Consequently, despite the occurrence of a π jump in the oscillation phases of s - g and d - g cross-channel interferences near 30° , this does not result in a significant change in the RABBITT phase due to the reduced oscillation amplitude associated with cross-channel interferences involving a g electron. A variation becomes noticeable in the lower sideband starting around 50° when the d wave approaches its node position at 57° . However, this variation does not lead to a substantial change in the observed angular range. This suggests that none of the cross-channel interferences outweighs the other interference terms in this case.

Moving to the higher sideband, in accordance with the propensity rule, three-photon absorption in the continuum creates a g electron with notable probability. Therefore, it is reasonable to expect that the angle dependence of the higher sideband will start to vary early, following the node position of $P_4(\theta)$ around 30° . Clearly, the significant variation in the RABBITT phase around 50° in the higher sideband suggests that cross-channel interferences involving d and/or g electrons start to become comparable and outweigh the remaining interference terms above 50° . However, our current technique does not allow us to definitively determine which interference term is the most significant.

In the next sideband group, S_{14} , we observe that the higher sideband exhibits less angular variation within the observed angular range compared to the S_{12} group. Additionally, the transition-amplitude ratios to different angular momentum states change with the energy, potentially shifting the angle where cross-channel interference becomes dominant or, in some cases, nearly eliminating cross-channel interference altogether. Unfortunately, the error bars in the $S_{14,l}$ numbers are too large to draw conclusions with high confidence.

As the probe intensity increases, higher-order transition terms become increasingly significant. In such cases, the transition pathways depicted in Fig. 1 may no longer provide an adequate description, necessitating the inclusion of higher-order terms. Furthermore, we note that the angle dependence of the RABBITT phase changes slowly with varying probe intensity for most sidebands. However, a notable exception is seen in the lower sideband of S_{12} , where there is a substantial variation in the angle dependence of the RABBITT phase with changing probe intensity. This behavior can be attributed to the increasing impact of six-photon transitions involving the under-threshold harmonic (H_9) and Rydberg states. These transitions can be intensified by resonances and may also introduce rapid, energy-dependent resonance phases, resulting in a pronounced angle dependence of the RABBITT phase.

Figure 6 depicts a comparison between the experimental data and predictions from the SAE and RMT models for the S_{12} and S_{14} groups. We only show the case for the peak intensity of $I_2 = 7 \times 10^{11}$ W/cm², since the findings are similar for the other two. Starting with the S_{12} group, we see fairly




FIG. 6. SAE and RMT angle-dependent phase for the S_{12} and S_{14} groups, compared with the experimental data for an IR peak intensity of $I_2 = 7 \times 10^{11}$ W/cm².

good agreement between both sets of theoretical predictions and experiment for the center sideband $S_{12,c}$ and the higher sideband $S_{12,h}$, whereas there are quantitative differences for $S_{12,l}$. These include the relative position at small angles: experiment has the phase of $S_{12,l}$ above that of $S_{12,c}$ and $S_{12,h}$, while the SAE model in particular predicts $S_{12,l}$ to start significantly below. A possible reason is that higher-order transitions involving Rydberg states are affecting the phase of $S_{12,l}$, and it shifts gradually above that of $S_{12,c}$ as the probe intensity is increased. Nevertheless, there is agreement in the drop of the RABBITT phases beyond $\approx 40^\circ$, with $S_{12,h}$ dropping the fastest with increasing angle.

On the other hand, there is substantial disagreement between experiment and theory for the S_{14} group. While SAE and RMT in general agree well with each other, the experimental data suggest a much stronger angular dependence of all RABBITT phases, while the theoretical predictions are nearly flat. A notable exception to the otherwise good agreement between the SAE and RMT predictions is the rapid increase in the SAE RABBITT phase of $S_{14,h}$ at angles beyond $\approx 50^\circ$ and

$S_{14,c}$ beyond $\approx 65^\circ$. We note that the signal strength drops fast with increasing angle. Hence, the predictions become very sensitive to the details of the model, in this case the treatment (or the lack thereof) of exchange effects and correlations in the description of the ground state. While RMT is likely to be superior to SAE in this regard, neither model reproduces the early decreases in the measured phases of $S_{14,c}$ and particularly $S_{14,h}$. Instead, both theories predict the most rapid changes in the angular range where no experimental data are available. RMT, for example, predicts S_h to drop first around 65° , followed by S_c around 70° , and finally S_l around 75° .

VI. CONCLUSIONS AND OUTLOOK

In this joint experimental and theoretical study, we extended our previous work on argon and carried out a 3-SB experiment on helium. This target was chosen for several reasons: (1) the XUV step is much simpler in helium compared to argon, since only one orbital angular momentum is generated; (2) we expected a better chance for theory to handle helium rather than the more complex argon target; and (3) there are no known autoionizing resonances in the range of ejected-electron energies studied in the present work.

Indeed, we found that relatively simple numerical models were able to qualitatively, and in some cases also quantitatively, reproduce most of the experimental findings. Exceptions include the angle-integrated phase extracted for the threshold sideband, which is heavily affected by IR-induced transitions involving Rydberg states, and the angle dependence of the RABBITT phase for one of the two sideband groups studied in this work.

We hope that the experimental data produced in this study will serve as motivation for future work. This includes a detailed study of the threshold sideband, as well as measurements with improved statistics at lower intensity to reduce higher-order effects. In addition, we will attempt to increase the cutoff of the higher-order harmonics in order to be able to compare more sideband groups. We will also perform a thorough investigation of the sensitivity of theoretical predictions on the details of the model. We are currently extending the RMT calculations to include more coupled states to further improve the bound-state description and even to account for coupling to the ionization continuum in the spirit of the R -matrix with pseudostates (RMPS) approach [31]. Since many delays have to be scanned through, such calculations are computationally expensive and have to be planned with great care in light of the available resources.

ACKNOWLEDGMENTS

The experimental part of this work was supported by the DFG-QUTIF program under Project No. HA 8399/2-1 and IMPRS-QD. A.T.B., S.S., K.R.H., and K.B. acknowledge funding from the NSF through Grant No. PHY-2110023 as well as the Frontera Pathways allocation of Grant No. PHY-20028. A.T.B. is grateful for funding through NSERC. The calculations were performed on Stampede-2 and Frontera at the Texas Advanced Computing Center in Austin (Texas).

- [1] P. M. Paul, E. S. Toma, P. Breger, G. Mullot, F. Augé, P. Balcou, H. G. Muller, and P. Agostini, *Science* **292**, 1689 (2001).
- [2] H. G. Muller, *Appl. Phys. B* **74**, s17 (2002).
- [3] K. Klünder, J. M. Dahlström, M. Gisselbrecht, T. Fordell, M. Swoboda, D. Guénot, P. Johnsson, J. Caillat, J. Mauritsson, A. Maquet, R. Taïeb, and A. L'Huillier, *Phys. Rev. Lett.* **106**, 143002 (2011).
- [4] J. M. Dahlström, A. L'Huillier, and A. Maquet, *J. Phys. B: At. Mol. Opt. Phys.* **45**, 183001 (2012).
- [5] J. M. Dahlström and E. Lindroth, *J. Phys. B: At. Mol. Opt. Phys.* **47**, 124012 (2014).
- [6] R. Pazourek, S. Nagele, and J. Burgdörfer, *Rev. Mod. Phys.* **87**, 765 (2015).
- [7] A. Harth, N. Douguet, K. Bartschat, R. Moshhammer, and T. Pfeifer, *Phys. Rev. A* **99**, 023410 (2019).
- [8] D. Bharti, D. Atri-Schuller, G. Menning, K. R. Hamilton, R. Moshhammer, T. Pfeifer, N. Douguet, K. Bartschat, and A. Harth, *Phys. Rev. A* **103**, 022834 (2021).
- [9] D. Bharti, H. Srinivas, F. Shobeiry, K. R. Hamilton, R. Moshhammer, T. Pfeifer, K. Bartschat, and A. Harth, *Phys. Rev. A* **107**, 022801 (2023).
- [10] A. McPherson, G. Gibson, H. Jara, U. Johann, T. S. Luk, I. A. McIntyre, K. Boyer, and C. K. Rhodes, *J. Opt. Soc. Am. B* **4**, 595 (1987).
- [11] M. Ferray, A. L'Huillier, X. F. Li, L. A. Lompre, G. Mainfray, and C. Manus, *J. Phys. B: At. Mol. Opt. Phys.* **21**, L31 (1988).
- [12] J. Ullrich, R. Moshhammer, A. Dorn, R. Dörner, L. P. H. Schmidt, and H. Schmidt-Böcking, *Rep. Prog. Phys.* **66**, 1463 (2003).
- [13] D. Busto, J. Vinbladh, S. Zhong, M. Isinger, S. Nandi, S. Maclot, P. Johnsson, M. Gisselbrecht, A. L'Huillier, E. Lindroth, and J. M. Dahlström, *Phys. Rev. Lett.* **123**, 133201 (2019).
- [14] M. Bertolino, D. Busto, F. Zapata, and J. M. Dahlström, *J. Phys. B: At. Mol. Opt. Phys.* **53**, 144002 (2020).
- [15] J. Peschel, D. Busto, M. Plach, M. Bertolino, M. Hofflund, S. Maclot, J. Vinbladh, H. Wikmark, F. Zapata, E. Lindroth, M. Gisselbrecht, J. M. Dahlström, A. L'Huillier, and P. Eng-Johnsson, *Nat. Commun.* **13**, 5205 (2022).
- [16] M. Swoboda, T. Fordell, K. Klünder, J. M. Dahlström, M. Miranda, C. Buth, K. J. Schafer, J. Mauritsson, A. L'Huillier, and M. Gisselbrecht, *Phys. Rev. Lett.* **104**, 103003 (2010).
- [17] A. Jiménez-Galán, L. Argenti, and F. Martín, *Phys. Rev. Lett.* **113**, 263001 (2014).
- [18] D. M. Villeneuve, P. Hockett, M. J. J. Vrakking, and H. Niikura, *Science* **356**, 1150 (2017).
- [19] A. Kheifets, *Atoms* **9**, 66 (2021).
- [20] A. Autuori, D. Platzer, M. Lejman, G. Gallician, L. Maëder, A. Covolo, L. Bosse, M. Dalui, D. Bresteau, J.-F. Hergott, O. Tcherbakoff, H. J. B. Marroux, V. Loriot, F. Lépine, L. Poisson, R. Taïeb, J. Caillat, and P. Salières, *Sci. Adv.* **8**, eabl7594 (2022).
- [21] L. Drescher, T. Witting, O. Kornilov, and M. J. J. Vrakking, *Phys. Rev. A* **105**, L011101 (2022).
- [22] L. Neoričić, D. Busto, H. Laurell, R. Weissenbilder, M. Ammitzböll, S. Luo, J. Peschel, H. Wikmark, J. Lahl, S. Maclot, R. J. Squibb, S. Zhong, P. Eng-Johnsson, C. L. Arnold, R. Feifel, M. Gisselbrecht, E. Lindroth, and A. L'Huillier, *Front. Phys.* **10**, 964586 (2022).
- [23] H. Srinivas, F. Shobeiry, D. Bharti, T. Pfeifer, R. Moshhammer, and A. Harth, *Opt. Express* **30**, 13630 (2022).
- [24] P. Birk, V. Stooß, M. Hartmann, G. D. Borisova, A. Blättermann, T. Heldt, K. Bartschat, C. Ott, and T. Pfeifer, *J. Phys. B: At. Mol. Opt. Phys.* **53**, 124002 (2020).
- [25] S. Meister, A. Bondy, K. Schnorr, S. Augustin, H. Lindenblatt, F. Trost, X. Xie, M. Braune, R. Treusch, B. Manschwetus, N. Schirmel, H. Redlin, N. Douguet, T. Pfeifer, K. Bartschat, and R. Moshhammer, *Phys. Rev. A* **102**, 062809 (2020).
- [26] A. Kramida, Yu. Ralchenko, J. Reader, and NIST ASD Team, NIST Atomic Spectra Database (ver. 5.10), NIST, Gaithersburg, MD, 2023, <http://physics.nist.gov/asd>.
- [27] N. Douguet, A. N. Grum-Grzhimailo, E. V. Gryzlova, E. I. Staroselskaya, J. Venzke, and K. Bartschat, *Phys. Rev. A* **93**, 033402 (2016).
- [28] A. C. Brown, G. S. Armstrong, J. Benda, D. D. Clarke, J. Wragg, K. R. Hamilton, Z. Mašín, J. D. Gorfinkiel, and H. W. van der Hart, *Comput. Phys. Commun.* **250**, 107062 (2020).
- [29] S. Heuser, A. Jiménez Galán, C. Cirelli, C. Marante, M. Sabbar, R. Boge, M. Lucchini, L. Gallmann, I. Ivanov, A. S. Kheifets, J. M. Dahlström, E. Lindroth, L. Argenti, F. Martín, and U. Keller, *Phys. Rev. A* **94**, 063409 (2016).
- [30] J. Fuchs, N. Douguet, S. Donsa, F. Martin, J. Burgdörfer, L. Argenti, L. Cattaneo, and U. Keller, *Optica* **7**, 154 (2020).
- [31] K. Bartschat, E. T. Hudson, M. P. Scott, P. G. Burke, and V. M. Burke, *J. Phys. B: At. Mol. Opt. Phys.* **29**, 115 (1996).