

Entanglement classification via a single entanglement measure

Adam Burchardt^{1,*}, Gonalo M. Quinta,² and Rui Andr e³

¹*QuSoft, CWI and University of Amsterdam, Science Park 123, 1098 XG Amsterdam, The Netherlands*

²*Instituto de Telecomunikaoes, Physics of Informations and Quantum Technologies Group, 1049-001 Lisbon, Portugal*

³*Centro de Astrof sica e Gravitaao (CENTRA), Departamento de F sica, Instituto Superior T cnico (IST), Universidade de Lisboa (UL), Av. Rovisco Pais 1, 1049-001 Lisbon, Portugal*



(Received 9 June 2021; revised 18 April 2023; accepted 5 February 2024; published 22 March 2024)

We show that a single polynomial entanglement measure is enough to verify equivalence between generic n -qubit states under stochastic local operations with classical communication (SLOCC). SLOCC operations may be represented geometrically by M bius transformations on the roots of the entanglement measure on the Bloch sphere. Moreover, we show how the roots of the three-tangle measure classify four-qubit generic states, and propose a method to obtain the normal form of a four-qubit state, which bypasses the possibly infinite iterative procedure.

DOI: [10.1103/PhysRevA.109.032424](https://doi.org/10.1103/PhysRevA.109.032424)

I. INTRODUCTION

Quantum entanglement is one of the key manifestations of quantum mechanics and the main resource for technologies founded on quantum information science. In particular, quantum states with nonequivalent entanglement represent distinct resources, which may be useful for different protocols. The idea of clustering states into classes exhibiting different qualities under quantum information processing tasks resulted in their classification under stochastic local operations assisted by classical communication (SLOCC). Such a classification was successfully presented for two, three, and four qubits [1–4]. However, the full classification of larger systems is completely unknown. Even the much simpler problem of detecting if two n -qubit states ($n > 4$) are SLOCC equivalent is, in general, quite demanding [5–8].

Among several approaches to the entanglement quantification and classification problem, a particularly useful one is via SL-invariant polynomial (SLIP) measures. Well-known examples are concurrence and three-tangle, which measure the two-body and three-body quantum correlations of the system [9,10]. SLIP measures provide not only a convenient method for entanglement classification but also its practical detection. Indeed, it was shown that almost all SLOCC equivalence classes can be distinguished by ratios of such measures [3]. Any given two n -qubit states are then SLOCC equivalent if a complete set of SLIP measures has the same values for both of them [5]. For more than four qubits, however, the size of such a set grows exponentially, making it intractable to use this approach to discriminate SLOCC-equivalent states with more than four qubits [11].

In this paper, we show that, contrary to intuition, a single SLIP measure is enough to verify the SLOCC equivalence between any two generic pure n -qubit states. By generic state,

we mean the set of all pure states except of the measure zero subset with respect to the Haar measure. We achieve this by using a mathematical trick, in particular, we use a SLIP measure defined for n -qubit systems to verify equivalence of systems of $n + 1$ qubits. In essence, we look at how the roots of the SLIP measure for those states behave under SLOCC. We show that if the states are SLOCC equivalent, then the roots of the SLIP measure for each state must be related by a M bius transformation, which is straightforward to verify. In particular, we use this procedure to show that the three-tangle measure is enough to discriminate between generic four-qubit states. Finally, we show how one may use the roots of a SLIP entanglement measure to obtain the normal form of a four-qubit state, which bypasses the possibly infinite iterative standard procedure.

II. POLYNOMIAL INVARIANT MEASURES

An entanglement measure is a function $E(|\psi\rangle)$ defined for pure states of n qubits, which vanishes on the set of separable states. One of the desired features of entanglement measures is invariance under SLOCC operations. Mathematically, a SLOCC operation might be uniquely determined by the action of local invertible operators $L \in \text{SL}(2, \mathbb{C})^{\otimes n}$ [1]. An entanglement measure E defined for all pure states of n qubits is called a SL-invariant polynomial of homogeneous degree h if it is a homogeneous polynomial of degree h in the state coefficients, and it is invariant under any local operation $\mathcal{O} = \mathcal{O}_1 \otimes \dots \otimes \mathcal{O}_n$, where $\mathcal{O}_i \in \text{SL}(2, \mathbb{C})$. It is easy to see that those two conditions are equivalent to the fact that E satisfies

$$E(\kappa \mathcal{O}|\psi\rangle) = \kappa^h E(|\psi\rangle)$$

for each real constant $\kappa > 0$ and invertible linear operator \mathcal{O} [1,3,12]. Such polynomial will be denoted by SLIP_n^h , where the top index indicates the degree of the polynomial and the bottom index is related to the number of qubits.

*adam.burchardt.uam@gmail.com

Well-known examples are concurrence and three-tangle [9,10]. Both of those measures, concurrence and three-tangle, can be written as the absolute value of the antilinear expectation value of simple operators. Indeed, for a two-qubit pure state $|\psi\rangle = c_{00}|00\rangle + c_{01}|01\rangle + c_{10}|10\rangle + c_{11}|11\rangle$ its concurrence reads as $C(|\psi\rangle) = |c_{00}c_{11} - c_{01}c_{10}| = |\langle\psi|\sigma_y \otimes \sigma_y|\bar{\psi}\rangle|$. Furthermore, the three-tangle $\tau^{(3)}$ defined for three-qubit states $|\psi\rangle \in \mathcal{H}_2^3$ takes relatively simple form:

$$\tau^{(3)}(|\psi\rangle) = \left| \sum_{j=\text{Id},x,y,z} \eta_j (\langle\psi|\sigma_j \otimes \sigma_y \otimes \sigma_y|\bar{\psi}\rangle)^2 \right|, \quad (1)$$

with notation $(\eta_{\text{Id}}, \eta_x, \eta_y, \eta_z) = (-1, 0, 1, 1)$. Moreover, the degree-four polynomial invariants for four qubits described by Luque and Thibon in Ref. [13] can be also written as similar expressions [14]. This simple idea of exploring the antilinear expectation value of the tensor product of Pauli operators was further used for constructing invariants of an arbitrary number of qubits [12,14,15]. For example, the following formulas:

$$E^{(2k+1)}(|\psi\rangle) = \left| \sum_{j=\text{Id},x,y,z} \eta_j (\langle\psi|\sigma_j \otimes \underbrace{\sigma_y \otimes \dots \otimes \sigma_y}_{2k}|\bar{\psi}\rangle)^2 \right|, \quad (2)$$

$$E^{(2k)}(|\psi\rangle) = \left| \sum_{j,i=\text{Id},x,y,z} \eta_j v_i (\langle\psi|\sigma_j \otimes \sigma_i \otimes \underbrace{\sigma_y \otimes \dots \otimes \sigma_y}_{2k}|\bar{\psi}\rangle)^2 \right|, \quad (3)$$

with $(\eta_{\text{Id}}, \eta_x, \eta_y, \eta_z) = (v_{\text{Id}}, v_x, v_y, v_z) = (-1, 0, 1, 1)$ are a degree-four SLIP measure for odd and even number of qubits, respectively [14].

A general method to construct various SLIP measures based on this simple idea of exploring the antilinear expectation value of the tensor product of Pauli operators was further invented in Refs. [12,14,15]. In particular any SLIP measure E can also be extended to mixed states by determining the largest convex function on the set of mixed states, which coincides with E on the set of pure states [16]. Despite its simple definition, the evaluation of a convex roof extension requires nonlinear minimization procedure, and for a general density matrix is a challenging task [17–20]. An attempt to address this challenging task was carried out by introducing the so-called zero polytope, the convex hull of pure states with vanishing E measure [21–24]. In the particular case of rank-two density matrices ρ , the zero polytope can be represented inside a Bloch sphere, spanned by the roots of E [22,25]. We adapt this approach focusing only on the roots of polynomial invariants, equivalently the vertices of the zero polytope.

III. SYSTEM OF ROOTS

Consider a $(n+1)$ -partite qubit state $|\psi\rangle$. The state $|\psi\rangle$ can be uniquely written as

$$|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle, \quad (4)$$

providing the canonical decomposition of the reduced density matrix $\rho = |\psi_0\rangle\langle\psi_0| + |\psi_1\rangle\langle\psi_1|$ obtained by tracing out the first qubit. Note that the states $|\psi_0\rangle$ and $|\psi_1\rangle$ are in general




FIG. 1. The stereographic projection relating the family of states $|\psi_z\rangle$ on the extended complex plane with the associated family of states $|\tilde{\psi}_z\rangle$ on the Bloch sphere. The spherical coordinates (θ, ϕ) and the complex coordinate z are related via the stereographic projection $z = \text{ctg}(\theta/2) e^{-i\phi}$.

neither normalized nor orthogonal. Consider now the family of states

$$|\psi_z\rangle = z|\psi_0\rangle + |\psi_1\rangle, \quad (5)$$

where z is taken from the extended complex plane $\hat{\mathbb{C}}$, i.e., complex numbers plus infinity. We denote this the extended plane representation. In addition, consider any SLIP_n^h measure E defined on the set of n -partite pure qubit states. Since E is polynomial in the coefficients of $|\psi_z\rangle$, it is also polynomial in the complex variable z [21]. Therefore, the polynomial $E(z|\psi_0\rangle + |\psi_1\rangle)$ has exactly h roots: ζ_1, \dots, ζ_h (which may be degenerated and/or at infinity), related to the degree of E . By using the complex number z , the states $|\psi_z\rangle$ can be mapped to the surface of a sphere via the standard stereographic projection $(\theta, \phi) := (\arctan 1/|z|, -\arg z)$ written in spherical coordinates. This way, a point on the unit two-sphere (θ, ϕ) can be associated with the quantum state

$$|\tilde{\psi}_z\rangle := \cos\frac{\theta}{2}|\psi_0\rangle + \sin\frac{\theta}{2}e^{i\phi}|\psi_1\rangle \quad (6)$$

with $z = \text{ctg}(\theta/2) e^{-i\phi}$, such that $|\psi_0\rangle$ lies in the north pole and $|\psi_1\rangle$ lies in the south pole, see Fig. 1. We denote this the Bloch sphere representation. Note that $|\tilde{\psi}_z\rangle \propto |\psi_z\rangle$ and that neither of these states is normalized, since $|\psi_0\rangle$ and $|\psi_1\rangle$ are not normalized in general either.

Local operations on the system of roots

To each linear invertible operator $\mathcal{O} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, one may associate a Möbius transformation $z \mapsto z' := \frac{az+b}{cz+d}$, mapping the extended complex plane $\hat{\mathbb{C}}$ into itself [26,27]. The composition of such transformations represents the multiplication of the associated operators. In particular, $z \mapsto z' := \frac{dz-b}{-cz+a}$ is an inverse Möbius transformation related with $\mathcal{O}^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$. Note that although Möbius transformations are typically represented on the extended complex plane, one may represent them as transformations on the Bloch sphere via the stereographic projection. The correspondence between invertible operators and Möbius transformations represented on the Bloch sphere was already successfully used for SLOCC classification of permutation-symmetric states [28–30].

To study the effect of SLOCC operations on the system of roots we begin by acting on the first qubit of a state $|\psi\rangle$ written in the form of Eq. (4) with an invertible linear operator \mathcal{O} . In terms of the family of states $|\psi_z\rangle$ in Eq. (5), this operation induces the map

$$|\psi_z\rangle \mapsto |\psi_{z'}\rangle = \frac{az+b}{cz+d}|\psi_0\rangle + |\psi_1\rangle, \quad (7)$$

i.e., the index is mapped via the Möbius transformation $z \mapsto z' := \frac{az+b}{cz+d}$, see Appendix A. In addition, since $|\tilde{\psi}_z\rangle \propto |\psi_z\rangle$, we have that the family of states $|\tilde{\psi}_z\rangle$ also transforms according to Eq. (7). This reflects the fact that the states $|\psi_z\rangle$ and $|\tilde{\psi}_z\rangle$ associated to the extended complex plane and the Bloch sphere are related by a stereographic projection of the variable z . Using Eq. (7) and the defining equation $E(\zeta_i|\psi_0) + |\psi_1\rangle = 0$ for the roots ζ_i of the polynomial E , one concludes that the roots transform according to the inverse Möbius transformation associated to the operator \mathcal{O} , i.e., $\zeta_i \mapsto \frac{d\zeta_i-b}{-c\zeta_i+a}$. Finally, although the system of roots changes with local operations acting on the qubit that is being traced out in Eq. (4), it is invariant under local operations acting on any other qubit since the polynomial E is $\text{SL}(2, \mathbb{C})^{\otimes n}$ invariant. We summarize these results in the theorem below, see Appendix A for a detailed proof.

Theorem 1. Consider an $(n+1)$ -partite pure quantum state $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$. The roots ζ_i of any SLIP_n^h entanglement measure associated to the partial trace of the first qubit:

- (i) are invariant under invertible operators, i.e., invariant under $\mathbf{1} \otimes \mathcal{O}_{\bar{n}} \in \text{SL}(2, \mathbb{C})^{\otimes n}$ operators;
- (ii) transform via an inverse Möbius transformation $\zeta'_i = \frac{d\zeta_i-b}{-c\zeta_i+a}$ with respect to the $\mathcal{O} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \otimes \mathbf{1}^n$ operator.

It is crucial to emphasize that normalizing the states $|\psi_0\rangle$ and $|\psi_1\rangle$ after the action of the operator \mathcal{O} , as is the case in existing related works [21,22,24,25,31], would spoil the mapping of Eq. (7). As a consequence, the action of SLOCC operators on the states $|\psi_z\rangle$ would no longer be given by the corresponding Möbius transformation, and the statements in Theorem 1 would no longer hold.

The decomposition (4) can be performed with respect to any other subsystem, each with its own system of roots. Any local operator $\mathcal{O}_k = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ acting on the k th qubit will influence independently the corresponding k th system of roots via the Möbius transformation $\zeta_i \mapsto \frac{d\zeta_i-b}{-c\zeta_i+a}$. On the other hand, if acting globally with a local operator $\mathcal{O}_1 \otimes \dots \otimes \mathcal{O}_{n+1}$, all roots (and thus all zero polytopes) will be affected. Since a Möbius transformation is a bijective mapping on the Bloch sphere, the total number of roots will always be preserved [31]. Moreover, the existence of a local transformation between two given states becomes straightforward to verify since Möbius transformations are fully classified.

IV. VERIFICATION OF SLOCC EQUIVALENCE

Theorem 1 provides a solution for the problem of discriminating quantum states up to SLOCC equivalence. To verify if two pure states are SLOCC equivalent, one can use the following procedure, which takes a single SLIP_n^h measure and two pure $(n+1)$ -qubit states as an input. In the generic situation, it returns a set of at most $3! \binom{h}{3}^{n+1}$ SLOCC operators as an

output. The input states are SLOCC equivalent if and only if (iff) they are interconnected by one of the output operators.

Procedure 1. Choose any SLIP_n^h entanglement measure of degree $h \geq 3$ and two $(n+1)$ -qubit states.

(i) Calculate the roots of a chosen measure for each subsystem for both states. If for any subsystem the number of roots of both states is not equal, such states are not SLOCC equivalent. Otherwise, denote by h_i the number of roots of both states calculated for i th subsystem. If $h_i \geq 3$ for all $1 \leq i \leq n+1$, the procedure will be conclusive. Note that for a generic state $h_i = h \geq 3$ for each subsystem i .

(ii) Focus on one subsystem i , $1 \leq i \leq n+1$, and determine all Möbius transformations, which transform the roots of the first state into roots of the second state. For example, choose three of the h_i roots from each state and write the unique Möbius transformations between the two triplets of roots. Repeat this for all $3! \binom{h_i}{3}$ possibilities of choosing three out of h_i roots for the second state. Derive the local operators \mathcal{O}_i associated to Möbius transformations.

(iii) Repeat step (ii) for all other subsystems and then consider the tensor products $\mathcal{O}_1 \otimes \dots \otimes \mathcal{O}_{n+1}$ of all the local operators obtained.

(iv) If the two given $(n+1)$ -qubit states are SLOCC equivalent, at least one of these operators must transform one state into the other (up to the normalization). Otherwise, they are not SLOCC equivalent.

Proposition 1. Consider any SLIP_n^h measure and two $(n+1)$ -qubit states. If both states have at least three roots with respect to each subsystem, they are SLOCC equivalent iff they are interconnected by one of the operator obtained as an outcome of Procedure 1.

We refer to Appendix B for a proof of this statement. Intuitively, a generic pure quantum state will have exactly h distinct roots for each subsystem, where h is the degree of the SLIP_n^h measure. Thus, any such measure of degree $h \leq 3$ should be sufficient to verify SLOCC equivalence between generic quantum states. We confirmed this intuition by looking at several examples of degree-four measures. In particular, Eqs. 3 and 2 provides a family of SLIP measures for arbitrary number of qubits. We examined those measures, and for each number $4 \leq n \leq 20$ of qubits, we generated a sample of a thousand $(n+1)$ -qubit states and each state had, indeed, four distinct roots with respect to any subsystem [32]. Furthermore, we confirm this numerical result analytically and proved that a generic pure quantum state has, indeed, four distinct roots for each subsystem for measures defined via Eqs. 3 and 2. By generic state, we mean the set of all pure states except of the measure zero subset with respect to the Haar measure, see Appendix C for more details. This result applies for an arbitrary number of qubits, thus proving the following proposition.

Proposition 2. A single SLIP measure is enough to provide necessary and sufficient conditions for any two generic pure n -qubit states to be SLOCC equivalent. We refer to Appendix D for a technical proof of this statement.

V. NORMAL SYSTEM OF ROOTS

In the previous section we showed that in principle any SLIP measure of degree $h \geq 3$ can be used to verify SLOCC




FIG. 2. A normal system of roots $z, 1/z, -z, -1/z$ together with the conjugate points $\bar{z}, 1/\bar{z}, -\bar{z}, -1/\bar{z}$ span the cuboid whose faces are parallel to the $XZ, XY,$ and YZ planes. There are 24 rotations of the Bloch sphere, which preserve this property, composing the elements of the group \mathcal{G}_{24} . Two of them, namely the rotation by a $\pi/2$ angle around X and Y axes are presented. The system of roots transforms according to Eqs. (F1)–(F3), giving $z \mapsto z' := \frac{z-i}{-iz+1}$ and $z \mapsto z'' := \frac{z-1}{z+1}$ for the two rotations.

equivalence. In this section, we consider the special case when such a measure has degree $h = 4$.

First, recall that any three distinct points on the sphere can be transformed onto any other three distinct points via a unique Möbius transformation, see Appendix E. While this is not the case for four points, it is possible to take any four complex points z_1, z_2, z_3, z_4 and associate a so-called cross-ratio

$$\lambda(z_1, z_2, z_3, z_4) := \frac{z_3 - z_1}{z_3 - z_2} \frac{z_4 - z_2}{z_4 - z_1}, \quad (8)$$

which is preserved under Möbius transformations [26,29]. Systems of four distinct points are related via Möbius transformations if their cross-ratios are related in the same way. The cross-ratio is not invariant under permutations of points, however, and depending on the ordering taken for the four points, it takes six values: $\lambda, \frac{1}{\lambda}, 1 - \lambda, \frac{1}{1-\lambda}, \frac{\lambda-1}{\lambda}, \frac{\lambda}{\lambda-1}$ [29]. A particularly interesting set of four points is one of the form $z, 1/z, -z, -1/z$, which we call a normal system, see Fig. 2. Any set of four points may be mapped into a normal system, for which $z, 1/z, -z, -1/z$ will be the solutions of the fourth-degree equation $\lambda = 4z^2/(1+z^2)^2$, where λ is the corresponding cross-ratio from Eq. (8). Such a map is unique up to symmetries of the cube, i.e., the group of 24 rotations generated by $\pi/2$ rotations along the X, Y, Z axis, denoted by \mathcal{G}_{24} . Appendix F contains the exact description of the group \mathcal{G}_{24} . In Appendix G we formally show the following mathematical result.

Proposition 3. Each nondegenerated four points z_1, z_2, z_3, z_4 on the Bloch sphere can be transformed onto the normal form $z, \frac{1}{z}, -z, -\frac{1}{z}$ via a Möbius transformation T . The latter is uniquely defined up to 24 rotations in the group \mathcal{G}_{24} .

This mathematical result has interesting implications for the SLOCC verification and classification problems. Indeed, assume that E is SLIP_n^4 measure of degree $h = 4$, and the state $|\psi\rangle \in \mathcal{H}_2^{\otimes(n+1)}$ has exactly four distinct roots for each subsystem. By Proposition 3, one can find a Möbius transformation T_i that transforms roots for each system i into the normal form. By Theorem 1, the local operators \mathcal{O}_i corresponding to transformations T_i transform state $|\psi\rangle$ into the form with roots for i th system being in normal form, hence

$$|\psi'\rangle := \mathcal{O}_1 \otimes \cdots \otimes \mathcal{O}_{n+1} |\psi\rangle \quad (9)$$

is a state for which roots of measure E for each subsystem are in normal form. We call $|\psi'\rangle$ an E -normal form of a state $|\psi\rangle$ with respect to measure E . Note that the E -normal form of a state, if it exists, is defined up to the group \mathcal{G}_{24}^{n+1} of local rotations. We illustrate this procedure on a simple example of broadly discussed four-partite state $\frac{1}{\sqrt{2}}(|0\rangle|GHZ\rangle + |1\rangle|W\rangle)$ [22,24]. We calculate its roots for $\tau^{(3)}$ measure (1) and find the Möbius transformation transforming them into a normal system, see Appendix H for detailed calculations. As a result, we obtain SLOCC operator that transforms the aforementioned state into its normal form with respect to $\tau^{(3)}$ measure, see Fig. 3.

In this way, for the chosen SLIP_n^4 measure E , we defined the E -normal form of any state that has exactly four roots with respect to measure E for each system. As we have shown in Proposition 2, choosing, for example, measure E defined on Eqs. (3) and (2), provides E -normal form of the generic state of any number of qubits.

This allows us to address more ambitious problems, when, for example, one needs to verify the pairwise SLOCC equivalence of a larger number of states. Indeed, one can reduce each of them to E -normal form and then compare them using only a finite (and relatively small) group \mathcal{G}_{24}^{n+1} of local rotations. Furthermore, such E -normal form might be possibly used for an even more challenging task of SLOCC classification of




FIG. 3. The system of four roots (represented as blue dots) related to the three-tangle polynomial measure $\tau^{(3)}$ evaluated on the first subsystem of the state $1/\sqrt{2}(|0\rangle|GHZ\rangle + |1\rangle|W\rangle)$. This system of four points can be mapped into a normal system (i.e., symmetrically related points $z, -z, 1/z, -1/z$) by a Möbius transformation. Similar local transformations can be performed with respect to other subsystems, transforming the states into a state in the normal form.

pure states. Indeed, as we will demonstrate in the next section, the $\tau^{(3)}$ -normal form for measure (1) can be successfully used to classify generic four qubit states. Recall that the problem of classifying $n \leq 5$ states still remains open [5–8].

We will finish this section by pointing out the intriguing connection between E -normal form of a state and, so-called, normal form of a state [33]. Recall that a state is in a normal form if reduced density matrices to one subsystem are all maximally mixed, i.e., proportional to the identity. Normal form, if it exists, is defined up to the local unitary operations. The process of determining the normal form of a state, if it exists, may turn out to be an infinite iterative minimization process [33]. On the other hand, the process of obtaining E -normal form of a state is straightforward and consists of a finite number of steps. As we observed, for four-qubit states the normal form of a state coincides with its $\tau^{(3)}$ -normal form, see Appendix H. For systems with a larger number of qubits $n > 4$ (especially for $n = 5, 6$), we unsuccessfully searched for measures E for which E -normal form would coincide with normal form of a state. Such measures, if found, would lead to a simple procedure for obtaining a normal form of a state for arbitrary number of qubits.

We illustrate this procedure by transforming the widely discussed four-partite state $\frac{1}{\sqrt{2}}(|0\rangle|GHZ\rangle + |1\rangle|W\rangle)$ [22,24] into its normal form, see Fig. 3. Without this technique, the standard way of obtaining the normal form would indeed result in an infinite iterative procedure.

VI. STATE CLASSIFICATION

We show that for small numbers of qubits $n = 3, 4$ our approach might be successfully used for the more demanding problem of entanglement classification. Focusing first on the three-qubit case, genuinely entangled pure states are SLOCC equivalent to either $|GHZ\rangle = \frac{1}{\sqrt{2}}(|000\rangle + |111\rangle)$ or $|W\rangle = \frac{1}{\sqrt{3}}(|001\rangle + |010\rangle + |100\rangle)$ [34]. Using the two-tangle $\tau^{(2)}$ [9] as the entanglement measure, one may use the roots to distinguish between the two classes. Indeed, all rank-two reduced density matrices of the $|W\rangle$ state have a single root, while there are always two distinct roots for the $|GHZ\rangle$ state [25].

Contrary to the three-qubit case, there are infinitely many SLOCC classes of four-qubit states [34]. Although four-qubit states were divided into nine families [2,35,36], we will focus on generic four-qubit states, i.e., four-qubit states with random coefficients belonging to the so-called G_{abcd} family—the four-qubit SLOCC family with the most degrees of freedom. The representative state is of the form $|G_{abcd}\rangle = \frac{a+d}{2}(|0000\rangle + |1111\rangle) + \frac{a-d}{2}(|0011\rangle + |1100\rangle) + \frac{b+c}{2}(|0101\rangle + |1010\rangle) + \frac{b-c}{2}(|0110\rangle + |1001\rangle)$, where $a^2 \neq b^2 \neq c^2 \neq d^2$ are pairwise different. Choosing the three-tangle $\tau^{(3)}$ (1) as the entanglement measure, the states $|G_{abcd}\rangle$ have four nondegenerate roots already in the normal form, see Appendix I. Since the normal form of roots is unique up to the group \mathcal{G}_{24} , the problem of SLOCC equivalence of states $|G_{abcd}\rangle$ becomes solvable, with a discrete amount of solutions. Indeed, it can quickly be confirmed if two states in the G_{abcd} class are SLOCC equivalent by checking if one can be obtained from the other

by the action of an element of the finite class of operators $\mathcal{O} \in \mathcal{G}_{24}^{\otimes 4}$. We thus find that exactly 192 states of the form $|G_{abcd}\rangle$ are SLOCC equivalent.

Proposition 4. Two states $|G_{abcd}\rangle$ and $|G_{a'b'c'd'}\rangle$ are SLOCC equivalent iff their coefficients are related by the following three operations: multiplication by a phase factor $(a', b', c', d') = e^{i\phi}(a, b, c, d)$, permutation of coefficients $(a', b', c', d') = \sigma(a, b, c, d)$, and change of sign in front of two coefficients from a, b, c, d .

See Appendix I for a detailed proof of the above statement. We finish this section by pointing out some intriguing connections between this result and other related problems. Note that the symmetry in Proposition 4 is given by the Weyl group of Cartan type D_4 and it has already been observed that the generators of four-qubit polynomial invariants exhibit this type of symmetry [13,35]. As a consequence, this result constitutes a relation between four-qubit invariants and the convex roof extension of three-tangle $\tau^{(3)}$, which may shed some light on the problem of generalizing the CKW inequality [10] for four-qubit states [12,19,20,37,38], and beyond [37–39].

VII. CONCLUSIONS

In this paper, we showed how a single SLIP_n^h entanglement measure is enough to verify if two generic $(n+1)$ -qubit states to be SLOCC equivalent. Our result is applicable for any number of qubits. This was possible by showing that the roots of any SLIP_n^h measure transform via Möbius transformations under the SLOCC operations performed on the subsystems. In that way, SLOCC equivalence between two states is implied by the easily verifiable existence of a Möbius transformation relating aforementioned roots for each subsystem. Moreover, we define the E -normal form of the state with respect to the given $\text{SLIP}_n^{h=4}$ measure E , for which roots are symmetrically distributed on the Bloch sphere. Such form is simple to determine and exists for generic states. In comparison, so-called, normal form of state requires possibly infinite iterative minimization procedure. Furthermore, we demonstrated our approach on four-qubit states, and showed that the roots of the three-tangle measure $\tau^{(3)}$ are enough to fully classify four-qubit states from the most generic G_{abcd} family. Lastly, as we observed that for four-qubit states the $\tau^{(3)}$ -normal form coincides with the normal form of a state, which gives a procedure to determine the normal form of state that circumvents the possibility of an infinite iterative process of the standard procedure.

ACKNOWLEDGMENTS

The authors thank Andreas Osterloh, Karol Życzkowski, Rui Perdigo, and Yasser Omar for fruitful discussions and correspondence. The authors are especially grateful to Jens Siewert, Andreas Osterloh, Barbara Kraus, and Karol Życzkowski for stimulating conversations, which led to formal proof of Proposition 2 while staying at Centro de Ciencias de Benasque Pedro Pascual. A.B. acknowledges support from the National Science Center under Grant No. DEC-2015/18/A ST2/00274 and by NWO Vidi grant (Project No. VI.Vidi.192.109). G.Q. thanks the support from Fundação para a Ciência e a Tecnologia (Portugal), namely

through projects CEECIND/02474/2018 and Project No. UIDB/50008/2020 and IT project QuantSat-PT. R.A. acknowledges support from the Doctoral Programme in the Physics and Mathematics of Information (DP-PMI) and the Fundação para a Ciência e Tecnologia (FCT) through Grant No. PD/BD/135011/2017. This work is co-funded by the European Union (ERC, ASC-Q,101040624). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

APPENDIXES

In the Appendixes, we present detailed proofs of the statements presented in the main body of the paper, as well as a summary of the related results concerning Möbius transformations, SLIP entanglement measures, Haar measure on the set of pure quantum states, and SLOCC classification.

Appendixes A and B contain proofs of Theorem 1 and Proposition 1, respectively. Appendix C summarizes basic results regarding the Haar measure on the set of pure quantum states and discusses the notion of generic quantum states. Appendix D proves the main result of a paper, namely Proposition 2. The proof is based on two technical lemmas related to the systems of odd and even number of qubits, see Lemma 1 and Lemma 2, respectively. Appendix E presents a well-known form of a unique Möbius transformation transforming two given tuples of three points one onto another. Appendixes F and G discuss details regarding an E -normal form of a state with respect to a given measure E . Furthermore, Appendix H provides an explicit example of transformation of a state into its E -normal form. Appendix I presents proof of Proposition 4, which is reproducing the SLOCC-classification results for the generic systems of four qubits. The E -normal form proves to be useful for that purpose. Lastly, in Appendix J, we apply the presented method to verify the SLOCC equivalence between certain five-qubit states.

APPENDIX A: PROOF OF THEOREM 1

We present a proof for Theorem 1. Any $(n+1)$ -partite qubit state $|\psi\rangle \in \mathcal{H}_2^{\otimes(n+1)}$ might be written as

$$|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle. \quad (\text{A1})$$

Such a form provides the canonical decomposition of the reduced density matrix $\rho = |\psi_0\rangle\langle\psi_0| + |\psi_1\rangle\langle\psi_1|$ over the non-normalized states $|\psi_0\rangle, |\psi_1\rangle \in \mathcal{H}_2^{\otimes n}$, obtained by tracing out the first qubit. Consider now a reversible operator $\mathcal{O} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{C})$ acting on the first qubit. Under the action of this operator, the state $|\psi\rangle$ is transformed into

$$\begin{aligned} |\psi'\rangle &:= \mathcal{O}|\psi\rangle = (a|0\rangle + b|1\rangle)|\psi_0\rangle + (c|0\rangle + d|1\rangle)|\psi_1\rangle \\ &= |0\rangle|\psi'_0\rangle + |1\rangle|\psi'_1\rangle, \end{aligned}$$

where

$$|\psi'_0\rangle := a|\psi_0\rangle + c|\psi_1\rangle, \quad (\text{A2})$$

$$|\psi'_1\rangle := b|\psi_0\rangle + d|\psi_1\rangle. \quad (\text{A3})$$

Consider now any superposition of states $|\psi'_0\rangle$ and $|\psi'_1\rangle$. Observe that

$$\begin{aligned} |\psi'_z\rangle &:= z|\psi'_0\rangle + |\psi'_1\rangle = z(a|\psi_0\rangle + c|\psi_1\rangle) + b|\psi_0\rangle + d|\psi_1\rangle \\ &= (az + b)|\psi_0\rangle + (cz + d)|\psi_1\rangle \\ &\propto \frac{az + b}{cz + d}|\psi_0\rangle + |\psi_1\rangle, \end{aligned}$$

where the complex number $cz + d$ was factored out in order for the transformation to map states from the extended plane representation to the extended plane representation. In other words, we have

$$\mathcal{O}|\psi_z\rangle = |\psi'_z\rangle, \quad z' = \frac{az + b}{cz + d}, \quad (\text{A4})$$

i.e., the operator \mathcal{O} transforms states in the extended plane representation by applying a Möbius transformation on the index z . Suppose now that ζ_i is a zero of a h -degree polynomial function E , i.e., $E(\zeta_i|\psi_0\rangle + |\psi_1\rangle) = 0$. Acting on the first qubit with \mathcal{O} , the density matrix after tracing out the first qubit becomes $|\psi'_0\rangle\langle\psi'_0| + |\psi'_1\rangle\langle\psi'_1|$, so the entanglement measure E will be zero for some new roots ζ'_i , such that $E(\zeta'_i|\psi'_0\rangle + |\psi'_1\rangle) = 0$. Using Eqs. (A2)–(A3), the latter equation can be transformed into

$$E\left((c\zeta'_i + d)\left(\frac{a\zeta'_i + b}{c\zeta'_i + d}|\psi_0\rangle + |\psi_1\rangle\right)\right) = 0, \quad (\text{A5})$$

where the factor $(c\zeta'_i + d)$ is irrelevant since any root multiplied by it will still be a valid root. Comparing with the equation for the roots before the action of \mathcal{O} , we reach the conclusion that the roots transform according to the inverse Möbius transformation as

$$\zeta'_i = \frac{d\zeta_i - b}{-c\zeta_i + a}, \quad (\text{A6})$$

under the action of the operator \mathcal{O} . As a consequence, the roots of the zero polytope transform with respect to the inverse Möbius transformation associated to the operator $\mathcal{O} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Analyze now the case when \mathcal{O} is a unitary operator $\mathcal{O} = \mathcal{U}$. Since any unitary operator \mathcal{U} can be represented as a rotation $\mathcal{R} = \begin{pmatrix} \cos\alpha & \sin\alpha e^{i\phi} \\ -\sin\alpha e^{-i\phi} & \cos\alpha \end{pmatrix}$ (up to an irrelevant global phase), it will simply rotate the Bloch ball, together with the zero polytope.

Consider now multilocal operators $\mathcal{O}_{\bar{n}} = \mathcal{O}_1 \otimes \dots \otimes \mathcal{O}_n$ acting on the remaining qubits of the state $|\psi\rangle$ from Eq. (A1). The state $|\psi\rangle$ will transform accordingly as

$$|\psi'\rangle := \mathcal{O}_{\bar{n}}|\psi\rangle = |0\rangle \underbrace{\mathcal{O}_{\bar{n}}|\psi_0\rangle}_{:=|\psi'_0\rangle} + |1\rangle \underbrace{\mathcal{O}_{\bar{n}}|\psi_1\rangle}_{:=|\psi'_1\rangle}. \quad (\text{A7})$$

After the action of $\mathcal{O}_{\bar{n}}$, a state in the extended plane representation will have a value of entanglement measure E equal to

$$E(z|\psi'_0\rangle + |\psi'_1\rangle) = E(\mathcal{O}_{\bar{n}}(z|\psi_0\rangle + |\psi_1\rangle)).$$

However, since E is $\text{SL}(2, \mathbb{C})^{\otimes n}$ invariant, we have that $E(z|\psi_0\rangle + |\psi_1\rangle) = 0$ iff $E(z|\psi'_0\rangle + |\psi'_1\rangle) = 0$, and so the roots of both polynomial equations are the same. As a consequence, the roots of the zero polytope will remain

unchanged under the action of $\mathcal{O}_{\bar{n}}$. This concludes the proof of Theorem 1.

APPENDIX B: PROOF OF PROPOSITION 1

In order to prove Proposition 1, we begin with the following simple observation:

Observation 1. For two sets $\Lambda = \{z_1, \dots, z_h\}$ and $\Lambda' = \{w_1, \dots, w_h\}$ of h distinct elements of the extended complex plane $z_i, w_i \in \hat{\mathbb{C}}$, $h > 2$, there exists at most $h(h-1)(h-2) = 3! \binom{h}{3}$ Möbius transformations, which map Λ into Λ' .

Proof. Suppose that T is a Möbius transformation that transforms Λ into Λ' . Thus,

$$T(z_1) = w_{i_1}, \quad T(z_2) = w_{i_2}, \quad T(z_3) = w_{i_3},$$

for distinct indices $i_1, i_2, i_3 = 1, \dots, h$. Observe that there are exactly $h(h-1)(h-2)$ possibilities for choice of those indices. For each such possibility, there is a unique Möbius transformation, which maps $z_1 \mapsto w_{i_1}$, $z_2 \mapsto w_{i_2}$, $z_3 \mapsto w_{i_3}$, see Appendix D [40]. Therefore there are at most $h(h-1)(h-2)$ Möbius transformations that map Λ into Λ' . ■

Remark 1. Usually, there are more efficient methods to determine all Möbius transformations that map a given set Λ into the other Λ' . For example, if $|\Lambda| = |\Lambda'| = 4$, one may calculate the values of cross-ratios. If they differ for any choice of orderings in Λ and Λ' , those sets are not related by any Möbius transformation.

Proof of Proposition 1. Suppose that two $(n+1)$ -qubit states $|\psi\rangle, |\phi\rangle \in \mathcal{H}_2^{\otimes(n+1)}$ have at least three roots calculated with respect to a given SLIP $_n^h$ measure E for each subsystem. We will show that both states are SLOCC equivalent iff in the outcome of Procedure 1 there is an operator \mathcal{O} providing such equivalence.

First, notice that if in the outcome of Procedure 1 there is an operator providing SLOCC equivalence, both states are SLOCC equivalent. Hence, described condition is a sufficient condition for SLOCC equivalence.

Second, we will see that the described condition is a necessary condition for SLOCC equivalence. Indeed, suppose that states $|\psi\rangle, |\phi\rangle$ are SLOCC equivalent, i.e., there exists an operator

$$\mathcal{O} = \mathcal{O}_1 \otimes \dots \otimes \mathcal{O}_{n+1} \quad (\text{B1})$$

such that $\mathcal{O}|\psi\rangle \propto |\phi\rangle$ for some operators $\mathcal{O}_i \in \text{SL}(2, \mathbb{C})$. Denote by T_i the Möbius transformation corresponding to \mathcal{O}_i . Denote by $\Lambda_i = \{z_1^i, \dots, z_h^i\}$ the system of roots of the state $|\psi\rangle$ calculated with respect to i th subsystem. According to Theorem 1, the i th system of roots of a state $|\phi\rangle \propto \mathcal{O}|\psi\rangle$ is simply $\Lambda'_i = \{T_i(z_1^i), \dots, T_i(z_h^i)\}$. In particular T_i maps Λ_i into Λ'_i , hence will be found in Step 2 of Procedure 1. Note, that there might be other operators transforming Λ_i into Λ'_i , hence in Step 2, one may obtain multiple local operators. Similarly for any other subsystem, the transformation T_i (and the corresponding local operator \mathcal{O}_i) will be found. Hence among all local operators in the outcome of Procedure 1, there will be operator \mathcal{O} from Eq. (B1). ■

APPENDIX C: FUBINI-STUDY MEASURE AND GENERIC STATES

A Fubini-Study measure of the set of pure quantum states (known also as a Haar measure) is a unique probability distribution on \mathcal{H}_{d+1} , which is invariant under any unitary operation acting on the whole \mathcal{H}_{d+1} . It can be explicitly written as a probability density distribution

$$\Omega(v_1, \dots, v_d, \theta_1, \dots, \theta_d) = \frac{d!}{\pi^d} \prod_{i=1}^d \cos \theta_i \sin \theta_i^{2i-1}, \quad (\text{C1})$$

where $\theta_i \in [0, \frac{\pi}{2}]$, $v_i \in [0, 2\pi]$, on the set of pure quantum states parametrized as

$$|\psi\rangle = \sum_{i=0}^d \left(e^{iv_i} \cos \theta_i \prod_{\ell=i+1}^d \sin \theta_\ell \right) |i\rangle \in \mathcal{H}_{d+1}, \quad (\text{C2})$$

with the convention $\theta_0 = v_0 = 0$ [26]. Notice that states in Eq. (C2) are normalized and

$$\int_0^{2\pi} \dots \int_0^{2\pi} \int_0^{\frac{\pi}{2}} \dots \int_0^{\frac{\pi}{2}} \Omega(v_1, \dots, v_d, \theta_1, \dots, \theta_d) dv_1 \dots dv_d d\theta_1 \dots d\theta_d = 1$$

hence Ω is, indeed, a probability distribution. Notice that all random variables θ_i, v_i in Eq. (C1) are chosen independently according to the uniform distribution $\Theta(v_i) = \frac{1}{2\pi} v_i$ on $[0, 2\pi]$ for v_i variables, and according to

$$\Omega_i(\theta_i) = 2 i \cos \theta_i \sin \theta_i^{2i-1} \quad (\text{C3})$$

distribution on $[0, \frac{\pi}{2}]$.

Since $\mathcal{H}_2^{\otimes n} \cong \mathcal{H}_{2^n}$, the Fubini-Study measure is also well defined on the set of pure quantum states of n qubits $\mathcal{H}_2^{\otimes n}$. We will notice some properties of such a distribution, while expressing the state in the form $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$. For simplicity, we introduce the index sets $\mathcal{J} = \{0, 1\}^n$, $\mathcal{J}_0 = \mathcal{J} \setminus \{0, \dots, 0\}$, and for any index $I = (i_1, \dots, i_n) \in \mathcal{J}$, we denote by $|I|$ the decimal representation of the binomial string I , i.e., $|I| = \sum_{j=1}^n 2^{j-1} i_j$. In order to determine the Fubini-Study measure on $\mathcal{H}_2^{\otimes n+1}$, we will use the following isomorphism between $\mathcal{H}_2^{\otimes n+1} \cong \mathcal{H}_{2^{n+1}}$ defined on the basis vectors as $|iI\rangle \mapsto |\bar{i}2^n + |I|\rangle$ for $I \in \mathcal{J}$, and where $\bar{i} = i + 1 \pmod{2}$. In such a way, the Fubini-Study measure on $\mathcal{H}_2^{\otimes n+1}$ is given by the probability density distribution independent for each random variable and written jointly as

$$\Omega(v_I, \theta_I, v_{I'}, \theta_{I'} : I \in \mathcal{J}, I' \in \mathcal{J}_I) = \frac{(2^{n+1} - 1)!}{\pi^{2^{n+1} - 1}} \cdot \prod_{I \in \mathcal{J}} \cos \theta_I \sin \theta_I^{2^{n+1} + 2|I| - 1} \prod_{I' \in \mathcal{J}_I} \cos \theta_{I'} \sin \theta_{I'}^{2|I'| - 1}, \quad (\text{C4})$$

where $\theta_I, \theta_{I'} \in [0, \frac{\pi}{2}]$, and $v_I, v_{I'} \in [0, 2\pi]$, on the set of pure quantum states parametrized as

$$|\psi\rangle = |0\rangle|\psi_0\rangle + \left(\prod_{I \in \mathcal{J}} \sin \theta_I \right) |1\rangle|\psi_1\rangle, \quad (\text{C5})$$

where

$$\begin{aligned} |\psi_0\rangle &= \sum_{I \in \mathcal{J}} d_I |I\rangle, & d_I &= e^{iv_I} \cos \theta_I \prod_{\substack{J \in \mathcal{J}: \\ |J| > |I|}} \sin \theta_J, \\ |\psi'_1\rangle &= \sum_{I \in \mathcal{J}} c_I |I\rangle, & c_I &= e^{iv'_I} \cos \theta'_I \prod_{\substack{J \in \mathcal{J}: \\ |J| > |I|}} \sin \theta'_J, \end{aligned} \quad (\text{C6})$$

with the convention $\theta'_{0\dots 0} = v'_{0\dots 0} = 0$. Notice that the state coefficients of $|\psi_0\rangle$ depends only on θ_I, v_I variables, while state coefficients of $|\psi'_1\rangle$ depend only on θ'_I, v'_I variables. Furthermore, state coefficient d_I of $|\psi_0\rangle$ depends only on the variables v_I and θ_J for $J \in \mathcal{J}$ such that $|J| \geq |I|$. In particular, $d_{0\dots 0}$ is the only coefficient in $|\psi_0\rangle$ depending on $\theta_{0\dots 0}$, while $d_{0\dots 0}, d_{0\dots 01}$ are the only coefficients in $|\psi_0\rangle$ depending on $\theta_{0\dots 01}$, etc.

By a generic pure quantum state $|\psi\rangle \in \mathcal{H}_2^{\otimes n}$ we understand a random state chosen with respect to the Fubini-Study measure on the set of quantum states [26]. In that sense, we say that some property occurs with probability zero for generic state iff it occurs only on the set of measure zero among all pure quantum states with respect to the Fubini-Study measure. Notice that such a notion of generic states [41–43] is not the only one that appears in literature. Some authors refers to generic states as those whose stabilizers meet certain symmetry conditions [44–46]. It is worth mentioning that those two notions of generic states among pure quantum states agrees up to the measure zero subset on the set of pure quantum states with respect to the Fubini-Study measure.

APPENDIX D: PROOF OF PROPOSITION 2

In this section, we will prove that a generic pure quantum state will have exactly four distinct roots for each subsystem, for measures defined in Eqs. (2) and (3). We begin with the following lemma.

Lemma 1 (even case). Consider a generic pure quantum state $|\psi\rangle \in \mathcal{H}_2^{\otimes n+1}$ of even number of qubits written in the form $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$ where $|\psi_0\rangle, |\psi_1\rangle \in \mathcal{H}_2^{\otimes n}$, and a SLIP measure $E^{(n)}$ presented in Eq. (2). The conditional probability that the polynomial equation $E^{(n)}(|\psi_0\rangle + z|\psi_1\rangle) = 0$ has no multiple root at $z = 0$ under the condition that it has at least a single root at $z = 0$ is zero.

Proof. Consider an $(n+1)$ -qubit state $|\psi\rangle \in \mathcal{H}_2^{\otimes n+1}$ written in the form $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$, and suppose that $n = 2k + 1$ is an odd number. We will evaluate the entanglement measure presented in Eq. (2) on the following state $|\psi_0\rangle + z|\psi_1\rangle$, i.e.,

$$\begin{aligned} &E^{(2k+1)}(|\psi_0\rangle + z|\psi_1\rangle) \\ &= \left| \sum_{j=\text{ld},x,y,z} \eta_j (\langle\psi_0|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle \right. \\ &\quad + 2z\langle\psi_1|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle \\ &\quad \left. + z^2\langle\psi_1|\sigma_j \otimes \underbrace{\sigma_y \otimes \dots \otimes \sigma_y}_{2k}|\bar{\psi}_1\rangle) \right|^2. \end{aligned}$$

Furthermore, we expand the equation above as an absolute value of a degree-four polynomial in z variable, namely

$$E^{(2k+1)}(|\psi_0\rangle + z|\psi_1\rangle) = |C_0 + C_1z + C_2z^2 + C_3z^3 + C_4z^4|,$$

where the coefficients C_i are of the following form:

$$\begin{aligned} C_0 &= \sum_{j=\text{ld},x,y,z} \eta_j \langle\psi_0|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle \\ &\quad \cdot \langle\psi_0|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle = E^{(2k+1)}(|\psi_0\rangle), \\ C_1 &= 2 \sum_{j=\text{ld},x,y,z} \eta_j \langle\psi_1|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle \\ &\quad \cdot \langle\psi_0|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle. \\ &\dots \end{aligned}$$

Notice that the equation $E^{(n)}(|\psi_0\rangle + z|\psi_1\rangle) = 0$ has root at $z = 0$ iff the coefficient $C_0 = 0$, and has a multiple root at $z = 0$ iff coefficients $C_0, C_1 = 0$. Therefore we will prove that the conditional probability

$$\mathbb{P}(C_1 = 0 | C_0 = 0) = 0 \quad (\text{D1})$$

that $C_1 = 0$ under the condition $C_0 = 0$ is zero.

We can rewrite the coefficients C_0, C_1 in the following way:

$$\begin{aligned} C_0 &= \langle\psi_0|A(\psi_0)\rangle, \\ C_1 &= \langle\psi_1|A(\psi_0)\rangle, \end{aligned}$$

where $|A(\psi_0)\rangle$ is the following vector:

$$|A(\psi_0)\rangle = A(\psi_0)|\bar{\psi}_0\rangle,$$

defined by operator

$$\begin{aligned} A(\psi_0) &= \sum_{j=\text{ld},x,y,z} (\eta_j \langle\psi_0|\sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle) \cdot \\ &\quad \cdot \sigma_j \otimes \sigma_y \otimes \dots \otimes \sigma_y. \end{aligned}$$

Notice that $|A(\psi_0)\rangle$ is a function of state coefficients of $|\psi_0\rangle$ only. State coefficients of $|\psi_1\rangle$ are not related to the state coefficients of $|\psi_0\rangle$ in anyway by a norm, see Eqs. (C5) and (C6), hence the space of solutions for the equation

$$C_1 = \langle\psi_1|A(\psi_0)\rangle = 0 \quad (\text{D2})$$

is of zero measure iff $|A(\psi_0)\rangle \neq 0$. Therefore, condition Eq. (D1) can be rewritten as the conditional probability

$$\mathbb{P}(|A(\psi_0)\rangle \equiv 0 | \langle\psi_0|A(\psi_0)\rangle = 0) = 0 \quad (\text{D3})$$

that $|A(\psi_0)\rangle \equiv 0$ under the condition $\langle\psi_0|A(\psi_0)\rangle = 0$ is zero.

For simplicity, we will introduce the following notation:

$$\beta_k = \langle\psi_0|\sigma_k \otimes \sigma_y \otimes \dots \otimes \sigma_y|\bar{\psi}_0\rangle,$$

for $i = 0, 2, 3$. It is easy to see that $\langle\psi_0|A(\psi_0)\rangle = -\beta_0^2 + \beta_2^2 + \beta_3^2$, hence the condition $\langle\psi_1|A(\psi_0)\rangle = 0$ can be rewritten as

$$\sum_k \eta_k \beta_k^2 = -\beta_0^2 + \beta_2^2 + \beta_3^2 = 0, \quad (\text{D4})$$

Furthermore, the operator $A(\psi_0)$ takes the following form:

$$A(\psi_0) = \left(\sum_{\mu} \eta_{\mu} \beta_{\mu} \sigma_{\mu} \right) \otimes \sigma_y \otimes \dots \otimes \sigma_y.$$

Observe that condition $|A(\psi_0)\rangle = A(\psi_0)|\bar{\psi}_0\rangle \equiv 0$ is equivalent to requiring that $|\bar{\psi}_0\rangle$ belongs to the null space $\mathcal{N}[A(\psi_0)]$ of the matrix $A(\psi_0)$. Therefore, we have an immediate implication: if the null space of the operator $A(\psi_0)$ is trivial, i.e., $\mathcal{N}[A(\psi_0)] = \{0\}$ then $|A(\psi_0)\rangle \neq 0$. Since the null space of σ_y is empty, so is the null space of any number of tensor products of σ_y . Hence, we will find a null space $\mathcal{N}(\sum_\mu \eta_\mu \beta_\mu \sigma_\mu)$. One simple way to do this is to calculate the eigenvectors of $\sum_\mu \eta_\mu \beta_\mu \sigma_\mu$ and look at the space spanned by the eigenvectors whose eigenvalues are 0. It is straightforward to show that the eigenvalues and eigenvectors of a general combination of Pauli matrices $\sum_\mu a_\mu \sigma_\mu$ are given by

$$v_\pm = \left\{ \left(\frac{a_3 \pm \sqrt{a_1^2 + a_2^2 + a_3^2}}{a_1 + ia_2} \right) \right\}, \quad \lambda_\pm = a_0 \pm \sqrt{a_1^2 + a_2^2 + a_3^2}.$$

Hence, for $a_\mu = \eta_\mu \beta_\mu$ under the condition $C_0 = 0$ [in form Eq. (D4)], we have the eigensystem

$$v_\pm = \left\{ \left(-i \frac{\beta_3 \pm \sqrt{\beta_0^2}}{\beta_2} \right) \right\}, \quad \lambda_\pm = \beta_0 \pm \sqrt{\beta_0^2}.$$

Therefore if $\beta_0 \neq 0$ then $|A(\psi_0)\rangle \neq 0$. Recall that the condition $C_0 = 0$ is equivalent to $\beta_0^2 = \beta_2^2 + \beta_3^2$, hence the following conditional probability:

$$\mathbb{P}(\beta_0 = 0 \mid \beta_0^2 = \beta_2^2 + \beta_3^2) = 0 \quad (\text{D5})$$

that $\beta_0 = 0$ under the condition $\beta_0^2 = \beta_2^2 + \beta_3^2$ is zero implies Eq. (D3). In the remaining part of the proof, we will show that Eq. (D5) holds true.

We begin by expanding the n -qubit state $|\psi_0\rangle$ in the computational basis

$$|\psi_0\rangle = \sum_{j_1, \dots, j_n=0}^n d_{j_1 \dots j_n} |j_1, \dots, j_n\rangle.$$

Notice that β_k takes the following form in the computational basis:

$$\beta_k = \sum_{\substack{j_1, \dots, j_n \\ i_1, \dots, i_n}} d_{j_1 \dots j_n} d_{i_1 \dots i_n} \langle j_1 | \sigma_k | i_1 \rangle \times \langle j_2, \dots, j_n | \sigma_y \otimes \dots \otimes \sigma_y | i_2, \dots, i_n \rangle.$$

Notice that because of the cancellation $\beta_2 = 0$, we can further simplify the expression for β_k and single out one state coefficient, namely $d_{0 \dots 0}$. We have

$$\begin{aligned} \beta_0 &= 2(-1)^k (D_0 + D_3), \\ \beta_2 &= 0, \\ \beta_3 &= 2(-1)^k (D_0 - D_3), \end{aligned} \quad (\text{D6})$$

where

$$\begin{aligned} D_0 &= \sum_{I \in \mathcal{I}} (-1)^{|I|} d_{0I} d_{0\bar{I}}, \\ D_1 &= \sum_{I \in \mathcal{I}} (-1)^{|I|} d_{1I} d_{1\bar{I}}, \end{aligned} \quad (\text{D7})$$

where $\mathcal{I} = \{0, 1\}^{n-1}$ is the index set introduced for consistency of the notation, and for any bit string $I = (i_1, \dots, i_{n-1})$, we define its complement \bar{I} as $\bar{I} := (\bar{i}_1, \dots, \bar{i}_{n-1})$ where

$\bar{i}_j = i_j + 1 \pmod{2}$, and $|I| = \sum_{j=1}^{n-1} i_j$. Expanding β_k according to Eq. (D6), the condition $\beta_0^2 = \beta_2^2 + \beta_3^2$ becomes the following:

$$(D_0 + D_1)^2 = (D_0 - D_1)^2 \quad (\text{D8})$$

and hence has exactly two solutions, either $D_0 = 0$ or $D_1 = 0$. Notice that under any of those two solutions the condition $\beta_0 = 0$, i.e., $D_0 + D_1 = 0$ becomes $D_0 = D_1 = 0$. Hence Eq. (D5) can be rewritten as the following conditional probability:

$$\mathbb{P}(D_0 = 0 \wedge D_1 = 0 \mid D_0 = 0 \vee D_1 = 0) = 0. \quad (\text{D9})$$

Notice that

$$\begin{aligned} \mathbb{P}(D_0 = 0 \wedge D_1 = 0 \mid D_0 = 0 \vee D_1 = 0) &\leq \\ \mathbb{P}(D_0 = 0 \mid D_1 = 0) + \mathbb{P}(D_1 = 0 \mid D_0 = 0), \end{aligned} \quad (\text{D10})$$

hence we will show that both terms on the right-hand side of the equation above vanish.

Note that equation $D_0 = 0$ imposes conditions on the state coefficients d_{0I} , $I \in \mathcal{I}$ only, while equation $D_1 = 0$ on the state coefficients d_{1I} , $I \in \mathcal{I}$, see Eq. (D7). Since equations $D_1 = 0$ and $D_0 = 0$ impose conditions on different state coefficients, intuitively condition $D_0 = 0$ should not enforce $D_1 = 0$ being satisfied (and vice versa). This can be rigorously shown by the properties of Fubini-Study measure. Recall that state coefficients d_{1I} in $|\psi_0\rangle$ depend on θ_{1I} , and v_{1I} where $I \in \{0, 1\}^{n-1}$ only, see Eqs. (C5) and (C6). Therefore, the condition $D_1 = 0$ can be rewritten in θ_{1I} and v_{1I} variables as

$$f(\theta_{1I}, v_{1I} : I \in \mathcal{I}) = 0, \quad (\text{D11})$$

where f is an elementary function in $\theta_{1I}, v_{1I} : I \in \mathcal{I}$ variables. The exact form of f can be traced back from Eq. (D7) and Eq. (C6). In particular f , as an elementary function, is continuous on its domain. Similarly, the condition $D_0 = 0$ can be rewritten in θ_{0I} and v_{0I} variables as

$$g(\theta_{0I}, v_{0I} : I \in \mathcal{I}) = 0, \quad (\text{D12})$$

where g is an elementary, hence continuous, function. Therefore, $\mathbb{P}(D_1 = 0 \mid D_0 = 0)$ can be rewritten as the following conditional probability:

$$\mathbb{P}(f(\theta_{1I}, v_{1I} : I \in \mathcal{I}) = 0 \mid g(\theta_{0I}, v_{0I} : I \in \mathcal{I}) = 0) = 0, \quad (\text{D13})$$

where f, g are elementary (in particular continuous) functions on its domains defined on different variables. Recall that the random variables $\theta_I, v_I : I \in \{0, 1\}^n$ were chosen independently with continuous probability density distributions described in Eq. (C4), i.e., v_I are chosen independently according to the uniform distribution $\Theta(v_I) = \frac{1}{2\pi} v_I$ on $[0, 2\pi]$, and θ_I are chosen according to

$$\Omega_I(\theta_I) = (2^{n+1} + 2|I|) \cos \theta_I \sin \theta_I^{2^{n+1} + 2|I| - 1} \quad (\text{D14})$$

distribution on $[0, \frac{\pi}{2}]$. Therefore, a random quantum state $|\psi_0\rangle$, which satisfies condition $g(\theta_{0I}, v_{0I} : I \in \mathcal{I}) = 0$ is given by randomly chosen values of $\theta_I, v_I : I \in \{0, 1\}^n$ with respect to the probability distributions in Eq. (D14) for which $g(\theta_{0I}, v_{0I} : I \in \mathcal{I}) = 0$. It does not impose, however, any further constrains on $v_{1I} \theta_{1I}$ coefficients. Hence the value

of $f(\theta_{1I}, \nu_{1I} : I \in \mathcal{I})$ under condition $g(\theta_{0I}, \nu_{0I} : I \in \mathcal{I}) = 0$ does not vanish except of measure zero subspace for such induced probability distribution, hence $\mathbb{P}(D_1 = 0 | D_0 = 0) = 0$. An analogous argument shows that $\mathbb{P}(D_0 = 0 | D_1 = 0) = 0$, and hence that Eq. (D9) holds true, which finishes the proof of Lemma 1. ■

Below, we prove the result analogous to Lemma 1, for any odd number of qubits.

Lemma 2 (odd case). Consider a generic pure quantum state $|\psi\rangle \in \mathcal{H}_2^{\otimes n+1}$ of odd number of qubits written in the form $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$ where $|\psi_0\rangle, |\psi_1\rangle \in \mathcal{H}_2^{\otimes n}$, and a SLIP measure $E^{(n)}$ presented in Eq. (3). The conditional probability that the polynomial equation $E^{(n)}(|\psi_0\rangle + z|\psi_1\rangle) = 0$ has no multiple root at $z = 0$ under the condition that it has at least single root at $z = 0$ is zero.

Proof. Consider an n -qubit state $|\psi\rangle \in \mathcal{H}_2^{\otimes n}$ written in the form $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$, and suppose that $n = 2k$ is an even number. We will evaluate the value of a measure $E^{(2k)}$ presented in Eq. (3) on the following state $|\psi_0\rangle + z|\psi_1\rangle$. Thus, we have

$$\begin{aligned} E^{(2k)}(|\psi_0\rangle + z|\psi_1\rangle) &= \left| \sum_{j,i=\text{Id},x,y,z} \eta_j \nu_i (\langle \psi_0 | \sigma_j \otimes \sigma_i \otimes \sigma_y \otimes \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle \right. \\ &\quad + 2z \langle \psi_1 | \sigma_j \otimes \sigma_i \otimes \sigma_y \otimes \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle \\ &\quad \left. + z^2 \langle \psi_1 | \sigma_j \otimes \sigma_i \otimes \underbrace{\sigma_y \otimes \cdots \otimes \sigma_y}_{2k} | \bar{\psi}_1 \rangle \right|^2. \end{aligned} \quad (\text{D15})$$

Observe that by expanding it further, it is an absolute value of a degree-four polynomial in the z variable:

$$E^{(2k)}(|\psi_0\rangle + z|\psi_1\rangle) = |C_0 + C_1 z + C_2 z^2 + C_3 z^3 + C_4 z^4|,$$

where the coefficients are of the following form:

$$\begin{aligned} C_0 &= \sum_{j,i=\text{Id},x,y,z} \eta_j \nu_i \langle \psi_0 | \sigma_j \otimes \sigma_i \otimes \sigma_y \otimes \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle \\ &\quad \cdot \langle \psi_0 | \sigma_j \otimes \sigma_i \otimes \sigma_y \otimes \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle = E^{(2k)}(|\psi_0\rangle), \\ C_1 &= 2 \sum_{j,i=\text{Id},x,y,z} \eta_j \nu_i \langle \psi_1 | \sigma_j \otimes \sigma_i \otimes \sigma_y \otimes \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle \\ &\quad \cdot \langle \psi_0 | \sigma_j \otimes \sigma_i \otimes \sigma_y \otimes \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle, \\ &\dots \end{aligned}$$

with $(\eta_{\text{Id}}, \eta_x, \eta_y, \eta_z) = (\nu_{\text{Id}}, \nu_x, \nu_y, \nu_z) = (-1, 0, 1, 1)$. Notice that the equation $E^{(n)}(|\psi_0\rangle + z|\psi_1\rangle) = 0$ has a root at $z = 0$ iff the coefficient $C_0 = 0$, and has a multiple root at $z = 0$ iff coefficients $C_0, C_1 = 0$. Therefore we will prove the conditional probability

$$\mathbb{P}(C_1 = 0 | C_0 = 0) = 0 \quad (\text{D16})$$

that $C_1 = 0$ under the condition $C_0 = 0$ is zero.

We can rewrite the coefficients C_0, C_1 in the following way:

$$\begin{aligned} C_0 &= \langle \psi_0 | A(\psi_0) \rangle, \\ C_1 &= \langle \psi_1 | A(\psi_0) \rangle, \end{aligned}$$

where $|A(\psi_0)\rangle$ is the following vector:

$$|A(\psi_0)\rangle = A(\psi_0) |\bar{\psi}_0\rangle, \quad (\text{D17})$$

defined by operator

$$\begin{aligned} A(\psi_0) &= \sum_{i,j=\text{Id},x,y,z} (\eta_j \nu_i \langle \psi_0 | \sigma_i \otimes \sigma_j \otimes \sigma_y \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle) \\ &\quad \cdot \sigma_i \otimes \sigma_j \otimes \sigma_y \cdots \otimes \sigma_y. \end{aligned}$$

Notice that $|A(\psi_0)\rangle$ is a function of state coefficients of $|\psi_0\rangle$ only. State coefficients of $|\psi_1\rangle$ are not related to the state coefficients of $|\psi_0\rangle$ in anyway by a norm, see Eqs. (C5) and (C6), hence the space of solutions for the equation

$$C_1 = \langle \psi_1 | A(\psi_0) \rangle = 0 \quad (\text{D18})$$

is of zero measure iff $|A(\psi_0)\rangle \neq 0$. Therefore, condition Eq. (D16) can be rewritten as the conditional probability

$$\mathbb{P}(|A(\psi_0)\rangle \equiv 0 | \langle \psi_0 | A(\psi_0) \rangle = 0) = 0 \quad (\text{D19})$$

that $|A(\psi_0)\rangle \equiv 0$ under the condition $\langle \psi_0 | A(\psi_0) \rangle = 0$ is zero. Observe that in order to show that $|A(\psi_0)\rangle = A(\psi_0) |\bar{\psi}_0\rangle \neq 0$ it is enough to show that the first coefficient of $|A(\psi_0)\rangle$ is non-vanishing, i.e., $\langle 0 \dots 0 | A(\psi_0) \rangle \neq 0$. Therefore the following conditional probability:

$$\mathbb{P}(\langle 0 \dots 0 | A(\psi_0) \rangle = 0 | \langle \psi_0 | A(\psi_0) \rangle = 0) = 0 \quad (\text{D20})$$

that $\langle 0 \dots 0 | A(\psi_0) \rangle = 0$ under the condition $\langle \psi_0 | A(\psi_0) \rangle = 0$ is zero implies Eq. (D19). In the remaining part of the proof, we will show that Eq. (D20) holds true.

For simplicity, we will introduce the following notation:

$$\beta_{kp} = \langle \psi_0 | \sigma_k \otimes \sigma_p \otimes \sigma_y \cdots \otimes \sigma_y | \bar{\psi}_0 \rangle,$$

for $i, j = 0, 2, 3$. It is easy to see that the condition $\langle \psi_0 | A(\psi_0) \rangle = 0$ can be rewritten as

$$\begin{aligned} 0 &= \sum_{k,p} \eta_k \eta_p \beta_{kp}^2 \\ &= \beta_{00}^2 - \beta_{02}^2 - \beta_{03}^2 - \beta_{20}^2 + \beta_{22}^2 + \beta_{23}^2 - \beta_{30}^2 + \beta_{32}^2 + \beta_{33}^2. \end{aligned} \quad (\text{D21})$$

Furthermore, the operator $A(\psi_0)$ takes the following form:

$$\begin{aligned} A(\psi_0) &= \left(\sum_{k,p} \eta_k \eta_p \beta_{kp} \sigma_k \otimes \sigma_p \right) \otimes \sigma_y \otimes \cdots \otimes \sigma_y \\ &= B \otimes \sigma_y \otimes \cdots \otimes \sigma_y, \end{aligned} \quad (\text{D22})$$

where matrix B is presented on next Equation.

$$B = \begin{pmatrix} \beta_{00} - \beta_{03} - \beta_{30} + \beta_{33} & i(\beta_{02} - \beta_{32}) & i(\beta_{20} - \beta_{23}) & -\beta_{22} \\ -i(\beta_{02} - \beta_{32}) & \beta_{00} + \beta_{03} - \beta_{30} - \beta_{33} & \beta_{22} & i(\beta_{20} + \beta_{23}) \\ -i(\beta_{20} - \beta_{23}) & \beta_{22} & \beta_{00} - \beta_{03} + \beta_{30} - \beta_{33} & i(\beta_{02} + \beta_{32}) \\ -\beta_{22} & -i(\beta_{20} + \beta_{23}) & -i(\beta_{02} + \beta_{32}) & \beta_{00} + \beta_{03} + \beta_{30} + \beta_{33} \end{pmatrix}.$$

We will expand an n -qubit state $|\psi_0\rangle$ in the computational basis

$$|\psi_0\rangle = \sum_{j_1, \dots, j_n=0}^n d_{j_1 \dots j_n} |j_1, \dots, j_n\rangle.$$

Notice that β_{kp} takes the following form in the computational basis:

$$\beta_{kp} = \sum_{\substack{j_1, \dots, j_n \\ i_1, \dots, i_n}} d_{j_1 \dots j_n} d_{i_1 \dots i_n} \langle j_1 | \sigma_k | i_1 \rangle \langle j_2 | \sigma_p | i_2 \rangle \cdot \langle j_3, \dots, j_n | \sigma_y \otimes \dots \otimes \sigma_y | i_3, \dots, i_n \rangle.$$

We can further simplify the expression for β_{kp} :

$$\begin{aligned} \beta_{00} &= (-1)^{k-1} (D_{00} + D_{01} + D_{10} + D_{11}), \\ \beta_{03} &= (-1)^{k-1} (D_{00} - D_{01} + D_{10} - D_{11}), \\ \beta_{30} &= (-1)^{k-1} (D_{00} + D_{01} - D_{10} - D_{11}), \\ \beta_{33} &= (-1)^{k-1} (D_{00} - D_{01} - D_{10} + D_{11}), \\ \beta_{22} &= 2(-1)^k (B_0 - B_1), \\ \beta_{02} &= \beta_{32} = \beta_{20} = \beta_{23} = 0, \end{aligned} \quad (\text{D23})$$

where we define the following quantities:

$$\begin{aligned} D_{ij} &= \sum_{I \in \mathcal{I}} (-1)^{|I|} d_{ijI} d_{ij\bar{I}}, \\ B_0 &= \sum_{I \in \mathcal{I}} (-1)^{|I|} d_{00I} d_{11\bar{I}}, \\ B_1 &= \sum_{I \in \mathcal{I}} (-1)^{|I|} d_{01I} d_{10\bar{I}}, \end{aligned}$$

and $\mathcal{I} = \{0, 1\}^{n-2}$ is the index set introduced for consistency of the notation, and for $I = (i_1, \dots, i_{n-2})$, we define $\bar{I} = (\bar{i}_1, \dots, \bar{i}_{n-2})$ where $\bar{i}_j = i_j + 1 \pmod{2}$, and $|I| = \sum_{j=1}^{n-2} i_j$. With above notation at hand, Eq. (D21) becomes the following:

$$8(D_{01}D_{10} + D_{11}D_{00}) - 2(B_0 - B_1) = 0. \quad (\text{D24})$$

We will single out $d_{0\dots 0}$ state coefficient from the equation above. Notice that the state coefficient $d_{0\dots 0}$ appears in D_{00} and B_0 terms only. Denote by

$$\begin{aligned} \bar{D}_{00} &= D_{00} - 2d_{0\dots 0}d_{001\dots 1}, \\ \hat{D}_{00} &= \bar{D}_{00} + 2d_{0\dots 01}d_{001\dots 10}, \\ \bar{B}_0 &= B_0 - 2d_{0\dots 0}d_{1\dots 1}, \\ \hat{B}_0 &= \bar{B}_0 + 2d_{0\dots 01}d_{1\dots 10}. \end{aligned} \quad (\text{D25})$$

Observe that $d_{0\dots 0}$ appears in D_{00} and B_0 , but does not appear in any of the following terms $\bar{D}_{00}, D_{01}, D_{10}, D_{11}, \bar{B}_0, \hat{B}_1$. Hence Eq. (D24) might be solved with respect to $d_{0\dots 0}$ variable, and become

$$d_{0\dots 0} = \frac{4(D_{01}D_{10} + D_{11}\bar{D}_{00}) - (\bar{B}_0 - B_1)}{2(d_{1\dots 1} - 4D_{11}d_{001\dots 1})}. \quad (\text{D26})$$

In such a way, we expressed the condition $\langle \psi_0 | A(\psi_0) \rangle = 0$, which appears in Eq. (D20) as an equation satisfied by the

$d_{0\dots 0}$ state coefficient. As a next step, we will investigate the following equation:

$$\langle 0 \dots 0 | A(\psi_0) \rangle = 0. \quad (\text{D27})$$

From (D22), (D17), and (D23), we have

$$\langle A(\psi_0) | 0 \dots 0 \rangle = 4D_{11}d_{001\dots 1} + 2(B_0 - B_1)d_{1\dots 1}.$$

Notice that in the equation above the $d_{0\dots 0}$ coefficient appears only in B_0 term. By expressing B_0 according to (D25), and substituting $d_{0\dots 0}$ according to (D26), Eq. (D27) becomes

$$\begin{aligned} 0 &= 4d_{001\dots 1}D_{11} + 8d_{1\dots 1} \frac{d_{1\dots 1} - 4d_{001\dots 1}D_{11}}{d_{1\dots 1} - 4d_{001\dots 1}D_{11}} \\ &\cdot (d_{1\dots 1}D_{01}D_{10} + D_{11}(d_{001\dots 1}(B_1 - \bar{B}_0) + d_{1\dots 1}\hat{D}_{00})). \end{aligned} \quad (\text{D28})$$

Notice that in the equation above, the coefficient $d_{0\dots 01}$ appears only in \hat{D}_{00} and \bar{B}_0 terms. With the notation (D25), we can single out the $d_{0\dots 01}$ coefficient and solve Eq. (D28) with respect to $d_{0\dots 01}$ variable:

$$\begin{aligned} d_{0\dots 01} &= \frac{2d_{1\dots 1}^2 D_{01}D_{10} + D_{11}}{4d_{1\dots 1}D_{11}(d_{001\dots 10}d_{1\dots 1} - d_{001\dots 1}d_{1\dots 10})} \\ &\cdot ((d_{001\dots 1}d_{1\dots 1}(2B_1 + 1) \\ &- 2d_{001\dots 1}(d_{1\dots 1}\hat{B}_0 + 2d_{001\dots 1}D_{11}) + 2d_{1\dots 1}^2\hat{D}_{00})). \end{aligned} \quad (\text{D29})$$

Notice that with Eqs. (D26) and (D29) in hand, Eq. (D5) can be rewritten as the following conditional probability:

$$\mathbb{P}(\text{Eq. (D29) holds} | \text{Eq. (D26) holds}) = 0 \quad (\text{D30})$$

that the probability that Eq. (D29) holds under the condition that Eq. (D26) holds is zero.

Note that Eq. (D26) expresses state coefficient $d_{0\dots 0}$ as a function of other state coefficients, and it defines a subspace in the state space in which it holds. Intuitively, there should be no restriction on other state coefficients, as Eq. (D29), in such a subspace. Hence Eq. (D30) should be satisfied. This intuition can be rigorously shown by properties of the Fubini-Study measure. In the remaining part of the proof, we show that Eq. (D30) holds, indeed, true. Notice that $d_{0\dots 0}$ is on the left-hand side of Eq. (D26) and does not appear on the right-hand side of Eq. (D26). We will recall now some properties of Fubini-Study distribution on the set $\mathcal{H}_2^{\otimes n+1}$ presented in Eqs. (C5) and (C6). Recall that, $d_{0\dots 0}$ is the only coefficient in $|\psi_0\rangle$ depending on $\theta_{0\dots 0}$. Therefore, by multiplying Eq. (D29) by $\prod_{I \in \mathcal{J}_0} \frac{1}{\sin \theta_I}$, where we used notation $\mathcal{J}_0 = \{0, 1\}^n \setminus \{0, \dots, 0\}$, Eq. (D29) takes the following form:

$$e^{i\theta_{0\dots 0}} \cos \theta_{0\dots 0} = f(\theta_I, \nu_I : I \in \mathcal{J}_0), \quad (\text{D31})$$

where f is an elementary function in $\theta_I, \nu_I : I \in \mathcal{J}_0$ variables, where exact form of f can be traced back from Eq. (D29) by substituting successively Eq. (D7) and Eq. (C6). In particular f , as an elementary function, is continuous on its domain. Similarly, the coefficient $d_{0\dots 01}$ is on the left-hand side of Eq. (D26) and does not appear on the right-hand side of Eq. (D26), moreover, there is no $d_{0\dots 0}$ coefficient in both sides of Eq. (D26). As we noticed in Eq. (C6), while $d_{0\dots 0}, d_{0\dots 01}$ are the only coefficients in $|\psi_0\rangle$ depending on $\theta_{0\dots 01}$, therefore, by

multiplying Eq. (D26) by $\prod_{I \in \mathcal{J}_1} \frac{1}{\sin \theta_I}$, where we used notation $\mathcal{J}_1 = \mathcal{J}_0 \setminus \{0, \dots, 0, 1\}$, Eq. (D26) takes the following form:

$$e^{i v_{0\dots 01}} \cos \theta_{0\dots 01} = g(\theta_I, v_I : I \in \mathcal{J}_1), \quad (\text{D32})$$

where g is an elementary function in $\theta_I, v_I : I \in \mathcal{J}_1$ variables, where exact form of g can be traced back from Eq. (D26) by substituting successively Eq. (D7) and Eq. (C6). In particular g , as an elementary function, is continuous on its domain. In summary, Eq. (D30) can be rewritten as the following conditional probability:

$$\mathbb{P}(e^{i v_{0\dots 01}} \cos \theta_{0\dots 01} = g(\theta_I, v_I : I \in \mathcal{J}_1) \mid e^{i v_{0\dots 0}} \cos \theta_{0\dots 0} = f(\theta_I, v_I : I \in \mathcal{J}_0)) = 0 \quad (\text{D33})$$

that the probability that $e^{i v_{0\dots 01}} \cos \theta_{0\dots 01} = g(\theta_I, v_I : I \in \mathcal{J}_1)$ under the condition that $e^{i v_{0\dots 0}} \cos \theta_{0\dots 0} = f(\theta_I, v_I : I \in \mathcal{J}_0)$ is zero, where f, g are elementary (in particular continuous) functions on its domains in $\theta_I, v_I : I \in \mathcal{J}_0$ and $\theta_I, v_I : I \in \mathcal{J}_1$ variables, respectively. Recall that the random variables $\theta_I, v_I : I \in \mathcal{J}$ were chosen independently with continuous probability density distributions described in Eq. (C4), i.e., v_I are chosen independently according to the uniform distribution $\Theta(v_I) = \frac{1}{2\pi} v_I$ on $[0, 2\pi]$, and θ_I are chosen according to

$$\Omega_I(\theta_I) = (2^{n+1} + 2|I|) \cos \theta_I \sin \theta_I^{2^{n+1} + 2|I| - 1} \quad (\text{D34})$$

distribution on $[0, \frac{\pi}{2}]$. Therefore, a random quantum state $|\psi_0\rangle$, which satisfies condition $e^{i v_{0\dots 0}} \cos \theta_{0\dots 0} = f(\theta_I, v_I : I \in \mathcal{J}_0)$ is given by randomly chosen values of $\theta_I, v_I : I \in \mathcal{I}$ with respect to the probability distributions in Eq. (D34) for which the norm of $f(\theta_I, v_I : I \in \mathcal{J}_0)$ is smaller than one, and coefficients $v_{0\dots 0}, \theta_{0\dots 0}$ uniquely determined by the value of function f . It does not impose, however, any further constrains on $v_{0\dots 01} \theta_{0\dots 01}$ coefficient in terms of $\theta_I, v_I : I \in \mathcal{J}_1$ coefficients (except of the norm of f being sufficiently small). Hence the following equation $e^{i v_{0\dots 01}} \cos \theta_{0\dots 01} = g(\theta_I, v_I : I \in \mathcal{J}_1)$ will be not satisfied except of measure zero subspace for such induced probability distribution. This shows that Eq. (D10) holds true, and hence finishes the proof of Lemma 2. ■

Proposition 5. Consider a generic pure quantum state $|\psi\rangle \in \mathcal{H}_2^{\otimes n+1}$ written in the form $|\psi\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$ where $|\psi_0\rangle, |\psi_1\rangle \in \mathcal{H}_2^{\otimes n}$. Furthermore, consider the polynomial equation

$$E^{(n)}(|\psi_0\rangle + z|\psi_1\rangle) = 0 \quad (\text{D35})$$

in z variable, where $E^{(n)}$ is a SLIP measure presented in Eq. (2) for an even number of qubits $n = 2(k+2)$ and Eq. (3) for an odd number of qubits $n = 2k+1$. For a generic state $|\psi\rangle \in \mathcal{H}_2^{\otimes n+1}$ the above polynomial equation has exactly four distinct roots with probability one. In other words, the set of states for which Eq. (D35) has less than four distinct roots is of measure zero with respect to the Fubini-Study measure.

Proof of Proposition 5. We will see that the statement of Proposition 5 easily follows from Lemma 1 and Lemma 2 and properties of Fubini-Study measure. Recall that Fubini-Study probability distribution on $\mathcal{H}_2^{\otimes n+1}$ is invariant under any unitary operation acting on the whole space $\mathcal{H}_2^{\otimes n+1}$. Hence the probability that Eq. (D35) has multiple roots is bigger or equal to the probability that Eq. (D35) has multiple root at $z = 0$

under the condition that it has at least single root at $z = 0$ is zero. Hence Lemma 1 and Lemma 2 justify the statement. ■

Observe that while using a SLIP measure defined in Eqs. (2) and (3) (depending on the parity of the number of qubits n) in Procedure 1 (described in the main body of the paper) a generic pure n -qubit state has exactly four roots with probability one. Therefore, together with Proposition 1 from the main body of the paper (proven in Appendix B), it shows that a single SLIP measure is enough to provide necessary and sufficient conditions for any two generic pure n -qubit states to be SLOCC equivalent. Hence, we have the following corollary.

Corollary 1. A single SLIP measure, Eq. (2) for an even number of qubits $n = 2(k+2)$ and Eq. (3) for an odd number of qubits $n = 2k+1$, is enough to provide necessary and sufficient conditions for any two generic pure n -qubit states to be SLOCC equivalent. Corollary 1 can be written shortly as Proposition 2 from the main body of the paper.

APPENDIX E: UNIQUE MÖBIUS TRANSFORMATION

It is well known that for a given tuple of three distinct points z_1, z_2, z_3 and any second tuple of such points w_1, w_2, w_3 on the extended complex plane $\hat{\mathbb{C}}$, there is a unique Möbius transformation T , which transforms one tuple into the other, i.e.,

$$T(z_1) = w_1, \quad T(z_2) = w_2, \quad T(z_3) = w_3.$$

There are several ways to determine the form of T . An explicit form can be found by evaluating the determinant [40]:

$$T(z) := \det \begin{pmatrix} zw & z & w & 1 \\ z_1 w_1 & z_1 & w_1 & 1 \\ z_2 w_2 & z_2 & w_2 & 1 \\ z_3 w_3 & z_3 & w_3 & 1 \end{pmatrix}. \quad (\text{E1})$$

This results into the following form of T :

$$T(z) = \frac{az + b}{cz + d}, \quad (\text{E2})$$

where

$$a = \det \begin{pmatrix} z_1 w_1 & w_1 & 1 \\ z_2 w_2 & w_2 & 1 \\ z_3 w_3 & w_3 & 1 \end{pmatrix}, \quad b = \det \begin{pmatrix} z_1 w_1 & z_1 & w_1 \\ z_2 w_2 & z_2 & w_2 \\ z_3 w_3 & z_3 & w_3 \end{pmatrix},$$

$$c = \det \begin{pmatrix} z_1 & w_1 & 1 \\ z_2 & w_2 & 1 \\ z_3 & w_3 & 1 \end{pmatrix}, \quad d = \det \begin{pmatrix} z_1 w_1 & z_1 & 1 \\ z_2 w_2 & z_2 & 1 \\ z_3 w_3 & z_3 & 1 \end{pmatrix}.$$

Note that the corresponding $\text{SL}(2, \mathbb{C})$ operator is of the form $\mathcal{O} = \frac{1}{N} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with the normalization constant

$$N = \sqrt{\det \begin{pmatrix} a & b \\ c & d \end{pmatrix}} = \sqrt{(z_1 - z_2)(z_1 - z_3)(z_2 - z_3)} \cdot \sqrt{(w_1 - w_2)(w_1 - w_3)(w_2 - w_3)}.$$

APPENDIX F: ROTATION GROUP \mathcal{G}_{24}

Consider the set of four symmetrically related points $\Phi = \{z, \frac{1}{z}, -z, -\frac{1}{z}\}$. It is very convenient to associate with them the

cuboid spanned by eight points:

$$\Phi \cup \bar{\Phi} = \left\{ z, \frac{1}{z}, -z, -\frac{1}{z}, \bar{z}, \frac{1}{\bar{z}}, -\bar{z}, -\frac{1}{\bar{z}} \right\},$$

as it is presented on Fig. 2. Observe that all six faces of the cuboid are parallel to one of the planes: XZ , XY , or YZ . In fact, this property is equivalent to the initial assumption that the set of points Φ is in normal form. Clearly, all rotations of the Bloch ball preserve the form of the cuboid. Nevertheless, only a special subgroup of all rotations preserve faces of the cuboid being parallel to XZ , XY , or YZ . This special subgroup \mathcal{G}_{24} contains 24 elements spanned by three rotations of $\pi/2$ around X , Y , and Z axis, given by:

$$\mathbf{R}_x(\pi/2) = \begin{pmatrix} \cos \pi/4 & -i \sin \pi/4 \\ -i \sin \pi/4 & \cos \pi/4 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix}, \quad (\text{F1})$$

$$\mathbf{R}_y(\pi/2) = \begin{pmatrix} \cos \pi/4 & -\sin \pi/4 \\ \sin \pi/4 & \cos \pi/4 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \quad (\text{F2})$$

$$\mathbf{R}_z(\pi/2) = \begin{pmatrix} e^{-i\pi/4} & 0 \\ 0 & e^{i\pi/4} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1-i & 0 \\ 0 & 1+i \end{pmatrix}. \quad (\text{F3})$$

In fact, this is a group of rotations preserving the regular cube (the group of orientable cube symmetries). Clearly, all rotations in the \mathcal{G}_{24} group preserve the normal-form structure of Φ . On the other hand, the normal form is uniquely determined up to 24 rotations in the \mathcal{G}_{24} group.

APPENDIX G: PROOF OF PROPOSITION 3

We present a proof of Proposition 3. For each complex number λ there exists another complex number z , such that the cross-ratio of the four points is equal to λ , i.e.,

$$\left(z, \frac{1}{z}; -z, -\frac{1}{z} \right) = \lambda. \quad (\text{G1})$$

Indeed, the cross-ratio on the left side equals $4z^2/(1+z^2)^2$, and the equation $4z^2/(1+z^2)^2 = \lambda$ has exactly four solutions

$$z_0 = \frac{4 - 2\lambda + \sqrt{1-\lambda}}{2\lambda}, \quad \frac{1}{z_0}, \quad -z_0, \quad -\frac{1}{z_0}. \quad (\text{G2})$$

Therefore, for a given value λ there exists a unique λ -normal system, such that the cross-ratio of its vertices is given by $(z_0, \frac{1}{z_0}; -z_0, -\frac{1}{z_0}) = \lambda$. Replacing the vertex z_0 by any other vertex $z_0, 1/z_0, -z_0$, or $-1/z_0$ does not change the value of the cross-ratio $(z_0, \frac{1}{z_0}; -z_0, -\frac{1}{z_0}) = \lambda$. Note that there exists a unique Möbius transformation T , which maps z_1, z_2, z_3 onto $z_0, 1/z_0, -z_0$, with the remaining z_4 mapped onto $-1/z_0$. Observe as well that the value of z_0 is unique up to its inverse, opposite, and opposite inverse elements, according to Eq. (G2), with the corresponding Möbius transformations associated to the matrices $T, \sigma_x T, \sigma_y T$, and $\sigma_z T$. Each of those transformations maps the set of points $\{z_1, z_2, z_3, z_4\}$ onto the same set of points $\{z_0, 1/z_0, -z_0, -1/z_0\}$, although the exact bijection between those two sets of roots is different for each transformation.

Depending on the order of four points $\{z_1, z_2, z_3, z_4\}$, the corresponding cross-ratio takes six values: $\lambda, \frac{1}{\lambda}, 1 -$

$\lambda, \frac{1}{1-\lambda}, \frac{\lambda-1}{\lambda}$, and $\frac{\lambda}{\lambda-1}$. For each of these, there is a corresponding set of solutions of the form $\{z_0, 1/z_0, -z_0, -1/z_0\}$ via Eq. (G2) with four related Möbius transformations. Therefore, there are in total 24 Möbius transformations that map any four nondegenerated points onto a normal system, each of them related by an element of the group \mathcal{G}_{24} , which has exactly 24 elements.

APPENDIX H: TRANSFORMATION OF A STATE INTO ITS NORMAL FORM

We illustrate the procedure to determine the normal form of four-qubit states on the following example:

$$\begin{aligned} |\psi\rangle &\propto |0\rangle|\text{GHZ}\rangle + |1\rangle|\text{W}\rangle \\ &= |0000\rangle + |0111\rangle + |1100\rangle + |1010\rangle + |1001\rangle \\ &= |0\rangle \underbrace{|000\rangle}_{|\psi_0\rangle} + |1\rangle \underbrace{|100\rangle + |010\rangle + |001\rangle}_{|\psi_1\rangle} \end{aligned} \quad (\text{H1})$$

of the widely discussed four-partite state [22,24]. We will focus attention on the first subsystem. Corresponding states $|\psi_0\rangle$ and $|\psi_1\rangle$ are indicated on Eq. (H1). We will use the three-tangle measure. The following polynomial $\tau(3)(|\psi_0\rangle + z|\psi_1\rangle)$ is of a degree four in variable z , and has exactly four distinct roots:

$$\begin{aligned} z_1 &= 0, \\ z_2 &= -\sqrt[3]{4}, \\ z_3 &= -\sqrt[3]{4} e^{2\pi i/3}, \\ z_4 &= -\sqrt[3]{4} e^{4\pi i/3}. \end{aligned}$$

The corresponding value of the cross-ratio is equal to $\lambda = 1 + e^{4\pi i/3}$. As it is shown in Proposition 3, there is a unique Möbius transformation T , which maps z_1, z_2, z_3, z_4 onto the system $z_0, \frac{1}{z_0}, -z_0, -\frac{1}{z_0}$ where z_0 is a root of a polynomial $4z^2/(1+z^2)^2 = \lambda$. Choose one of its roots, e.g., $z_0 = 1 + \sqrt{2}$. Appendix C presents one way of obtaining the exact transformation T , giving

$$T(z) = \frac{z - \frac{1+\sqrt{3}}{\sqrt[3]{4}}}{\frac{(1+\sqrt{3})(1+i)}{2}z + \frac{1+i}{\sqrt[3]{4}}}, \quad (\text{H2})$$

which performs the mapping

$$T(z_1) = z_0 = 1 + \sqrt{2}, \quad T(z_2) = \frac{1}{z_0}, \quad T(z_3) = -z_0.$$

According to Proposition 3, $T(z_4) = -\frac{1}{z_0}$ and hence T maps z_1, z_2, z_3, z_4 into the normal system of roots. Note that the corresponding $\text{SL}(2, \mathbb{C})$ operator is of the form

$$\mathcal{O}_1 = \frac{\sqrt[3]{2}}{(1+i)(3+\sqrt{3})} \begin{pmatrix} 1 & -\frac{1+\sqrt{3}}{\sqrt[3]{4}} \\ \frac{(1+\sqrt{3})(1+i)}{2} & \frac{1+i}{\sqrt[3]{4}} \end{pmatrix}, \quad (\text{H3})$$

which is presented on Fig. 3. Similar calculations for three remaining subsystems lead to the following SLOCC operator:

$$\mathcal{O} = \mathcal{O}_1 \otimes \mathcal{O}_2 \otimes \mathcal{O}_3 \otimes \mathcal{O}_4, \quad (\text{H4})$$

where \mathcal{O}_1 is presented on Eq. (H3) and $\mathcal{O}_2 = \mathcal{O}_3 = \mathcal{O}_4 := \mathcal{O}_1 \sigma_x$, which transforms state (H1) into the $|G_{abcd}\rangle$ form

$$\begin{aligned} |G_{abcd}\rangle &= \frac{a+d}{2}(|0000\rangle + |1111\rangle) + \frac{a-d}{2}(|0011\rangle + |1100\rangle) \\ &+ \frac{b+c}{2}(|0101\rangle + |1010\rangle) \\ &+ \frac{b-c}{2}(|0110\rangle + |1001\rangle) \end{aligned}$$

with parameters

$$\begin{aligned} a &= (-8 + 4i) - \frac{(12 - 8i)}{\sqrt{3}}, \\ b &= \frac{8}{3}i(3 + 2\sqrt{3}), \\ c &= 0, \\ d &= -\frac{4}{3}((6 + 3i) + (3 + 2i)\sqrt{3}). \end{aligned}$$

APPENDIX I: PROOF OF PROPOSITION 4

Consider the state $|G_{abcd}\rangle$ and its decomposition with respect to the first subsystem $|G_{abcd}\rangle = |0\rangle|\psi_0\rangle + |1\rangle|\psi_1\rangle$, where

$$\begin{aligned} |\psi_0\rangle &= \frac{a+d}{2}|000\rangle + \frac{a-d}{2}|011\rangle + \frac{b+c}{2}|101\rangle + \frac{b-c}{2}|110\rangle, \\ |\psi_1\rangle &= \frac{a+d}{2}|111\rangle + \frac{a-d}{2}|100\rangle + \frac{b+c}{2}|010\rangle + \frac{b-c}{2}|001\rangle. \end{aligned}$$

Suppose that $\tau^{(3)}(\zeta|\psi_0\rangle + |\psi_1\rangle) = 0$. Since $\tau^{(3)}$ is a $\text{SL}(2, \mathbb{C})^{\otimes 3}$ invariant, for any local operators $\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3$ we have

$$\tau^{(3)}((\mathcal{O}_1 \otimes \mathcal{O}_2 \otimes \mathcal{O}_3)(\zeta|\psi_0\rangle + |\psi_1\rangle)) = 0.$$

Observe that

$$\begin{aligned} |\psi_0\rangle &= (\sigma_x \otimes \sigma_x \otimes \sigma_x)|\psi_1\rangle, \\ |\psi_1\rangle &= (\sigma_x \otimes \sigma_x \otimes \sigma_x)|\psi_0\rangle, \end{aligned}$$

where $\sigma_x, \sigma_y, \sigma_z$ are Pauli matrices. Therefore by taking all local operators $\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3$ equal to σ_x , we may conclude that

$$\begin{aligned} 0 &= \tau^{(3)}((\sigma_x \otimes \sigma_x \otimes \sigma_x)(\zeta|\psi_0\rangle + |\psi_1\rangle)) \\ &= \zeta|\psi_1\rangle + |\psi_0\rangle \propto \frac{1}{\zeta}|\psi_0\rangle + |\psi_1\rangle, \end{aligned}$$

hence $1/\zeta$ is another root of $\tau^{(3)}$. Similarly, by considering $(\sigma_y \otimes \sigma_y \otimes \sigma_y)$ and $(\sigma_z \otimes \sigma_z \otimes \sigma_z)$, one may find another two roots $-\zeta, -1/\zeta$ of $\tau^{(3)}$. This shows that the roots of $\tau^{(3)}$ evaluated on any state from the G_{abcd} family are symmetrical with respect to rotations around X, Y, Z axes by the angle π . Writing $\tau^{(3)}(z|\psi_0\rangle + |\psi_1\rangle) = 0$ explicitly, we obtain the equation

$$\tau^{(3)}(z|\psi_0\rangle + |\psi_1\rangle) = Az^4 - 2(2B + A)z^2 + A = 0,$$

where $A = (b^2 - c^2)(a^2 - d^2)$ and $B = (c^2 - d^2)(a^2 - b^2)$. The above equation is nondegenerated iff $A, B, A + 2B \neq 0$, which happens iff $a^2 \neq b^2 \neq c^2 \neq d^2$ are pairwise different.

Lemma 3. Any local operator $\mathcal{O} = \mathcal{O}_1 \otimes \mathcal{O}_2 \otimes \mathcal{O}_3 \otimes \mathcal{O}_4 \in \text{SL}(2, \mathbb{C})^{\otimes 4}$, which transforms states $|G_{a'b'c'd'}\rangle \propto \mathcal{O}|G_{abcd}\rangle$ with $a^2 \neq b^2 \neq c^2 \neq d^2$, is of the form $\mathcal{O}_i \in \mathcal{G}_{24}$.

Proof. A local operator \mathcal{O}_1 acting on the first qubit and transforming the state $|G_{abcd}\rangle$ onto $|G_{a'b'c'd'}\rangle$, also transforms their systems of roots denoted as Λ and Λ' , respectively, via the action of the corresponding Möbius transformation. Note that both systems Λ and Λ' are in the normal form, therefore, according to Proposition 3, we have that $\mathcal{O}_i \in \mathcal{G}_{24}$. A similar analysis with respect to all other qubits shows that $\mathcal{O}_2, \mathcal{O}_3, \mathcal{O}_4 \in \mathcal{G}_{24}$. ■

This way, searching for SLOCC equivalence between the states $|G_{abcd}\rangle$ and $|G_{a'b'c'd'}\rangle$ becomes restricted to the search within the finite class of operators $\mathcal{O} \in \mathcal{G}_{24}^{\otimes 4}$. Since the group \mathcal{G}_{24} has only 24 elements, one may numerically verify that there are exactly $8 \times 24 = 192$ states in the G_{abcd} family, which are SLOCC equivalent to $|G_{abcd}\rangle$ by $\mathcal{O} \in \mathcal{G}_{24}^{\otimes 4}$. For example, the following operation:

$$\mathbf{R}_x\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_x\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_x\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_x\left(\frac{\pi}{2}\right) \quad (11)$$

transforms state $|G_{abcd}\rangle$ into $|G_{-b-acd}\rangle$. This might be simply written as a transformation of a tuples of indices: the tuple (a, b, c, d) is transformed into the tuple $(-b, -a, c, b)$. Similarly, the operators showed on the following right-hand sides provide the corresponding transformations of the tuple (a, b, c, d) on the left-hand side:

$$\begin{aligned} \mathbf{R}_y\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_y\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_y\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_y\left(\frac{\pi}{2}\right) &\longleftrightarrow (a, d, c, b), \\ \mathbf{R}_z\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_z\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_z\left(\frac{\pi}{2}\right) \otimes \mathbf{R}_z\left(\frac{\pi}{2}\right) &\longleftrightarrow (-d, b, c, -a), \\ \mathbf{R}_y(\pi) \otimes \mathbf{R}_y(\pi) \otimes \mathbf{1} \otimes \mathbf{1} &\longleftrightarrow (a, -b, -c, d), \\ \mathbf{R}_x(\pi) \otimes \mathbf{R}_x(\pi) \otimes \mathbf{1} \otimes \mathbf{1} &\longleftrightarrow (a, b, -c, -d), \\ \mathbf{R}_y(\pi) \otimes \mathbf{1} \otimes \mathbf{R}_y(\pi) \otimes \mathbf{1} &\longleftrightarrow (d, c, b, a), \\ \mathbf{R}_x(\pi) \otimes \mathbf{1} \otimes \mathbf{R}_x(\pi) \otimes \mathbf{1} &\longleftrightarrow (b, a, d, c). \end{aligned}$$

Additionally, the tuples (a, b, c, d) and $(-a, -b, -c, -d)$ represent the same state. Note that any composition of the

above operations also provides SLOCC equivalences between $|G_{abcd}\rangle$ states. The eight aforementioned transformations of

tuples generate all permutations of the a, b, c, d indices, together with the change of a sign of any two or all four indices. There are exactly 24 permutations and for each permutation the signs can be matched in exactly $1 + \binom{4}{2} + 1 = 8$ ways. This gives in total 192 tuples representing SLOCC-equivalent states, which perfectly matches the numerical result.

Finally, another trivial manipulation with indices a, b, c, d comes from multiplying by a global phase, which is an irrelevant operation due to the fact that quantum states are elements of a projective space. This operation transforms the indices as

$$(e^{i\theta} a, e^{i\theta} b, e^{i\theta} c, e^{i\theta} d) \sim (a, b, c, d),$$

resulting in the same quantum state for any real number $\theta \in [0, 2\pi)$. In particular, for $\theta = \pi$, we observe that system of opposite indices determines the same state as the initial one, i.e., $(-a, -b, -c, -d) \sim (a, b, c, d)$.

APPENDIX J: EXACT SCENARIOS

In order to exemplify the viability of the results in this work, we present another nontrivial scenario where our method is useful. We show how a single four-qubit entanglement SLIP_4^4 measure might be used to verify whether five-qubit states are SLOCC equivalent. Among several SLIP_4^h measures defined for system of four qubits, the so-called M measure has degree 4 [13,35] and is defined as the determinant

$$M(|\psi\rangle) := \det \begin{pmatrix} c_{0000} & c_{1000} & c_{0010} & c_{1010} \\ c_{0001} & c_{1001} & c_{0011} & c_{1011} \\ c_{0100} & c_{1100} & c_{0110} & c_{1110} \\ c_{0101} & c_{1101} & c_{0111} & c_{1111} \end{pmatrix}, \quad (\text{J1})$$

where c_{i_1, i_2, i_3, i_4} are state coefficients, i.e.,

$$|\psi\rangle = \sum_{i_1, i_2, i_3, i_4=0}^1 c_{i_1, i_2, i_3, i_4} |i_1, i_2, i_3, i_4\rangle.$$

We demonstrate now how the M measure can be used to verify whether two given five-qubit states are SLOCC equivalent. As

TABLE I. Cross-ratio of roots of M measure calculated for each subsystem of $|\psi\rangle$ state rounded to the fourth decimal place.

Number of subsystem	Value of cross-ratio
1	$0.8656 + 0.5008i$
2	$0.5 + 0.5906i$
3	$0.5 - 0.4844i$
4	$0.5 + 0.5397i$
5	$0.9231 + 0.3845i$

an example, consider the following five-qubit state:

$$\begin{aligned} |\psi\rangle = & |00001\rangle + |00111\rangle + |01001\rangle + |01010\rangle + |01101\rangle \\ & + |10000\rangle + |10010\rangle + |10101\rangle + |10110\rangle + |11011\rangle \\ & + \frac{1}{2}|11100\rangle. \end{aligned}$$

By permuting the five subsystems of the state $|\psi\rangle$, we obtain in total $5! = 120$ five-qubit states. In the following, we will show that such states are not pairwise SLOCC equivalent. Indeed, for each subsystem of a state $|\psi\rangle$, we may calculate the roots of the M measure according to Procedure 1. For any subsystem, there are four distinct roots. The values of the corresponding cross-ratios are presented in Table I. It is straightforward to show that no two values of cross-ratios corresponding to different subsystems (see Table I) are related by $\lambda, \frac{1}{\lambda}, 1 - \lambda, \frac{1}{1-\lambda}, \frac{\lambda-1}{\lambda}$, or $\frac{\lambda}{\lambda-1}$. As a direct consequence of Theorem 1, one concludes that those states are not SLOCC equivalent. Indeed, the states obtained from $|\psi\rangle$ by nonidentical permutation of subsystems will have corresponding cross ratios simultaneously permuted. Therefore, for some subsystems, the values of cross-ratios corresponding to any two such states are not related via $\lambda, \frac{1}{\lambda}, 1 - \lambda, \frac{1}{1-\lambda}, \frac{\lambda-1}{\lambda}$, or $\frac{\lambda}{\lambda-1}$. As a consequence, there is no Möbius transformation relating the roots of any subsystem (see Remark 1), and hence there is no SLOCC operator, which might relate those two states. A similar reasoning holds true for any pair of states obtained from $|\psi\rangle$ by permuting its subsystems by two distinct permutations. Finally, we note that these results would otherwise be highly computationally demanding if standard procedures were applied.

[1] W. Dür, G. Vidal, and J. I. Cirac, *Phys. Rev. A* **62**, 062314 (2000).
 [2] F. Verstraete, J. Dehaene, B. De Moor, and H. Verschelde, *Phys. Rev. A* **65**, 052112 (2002).
 [3] G. Gour and N. R. Wallach, *Phys. Rev. Lett.* **111**, 060502 (2013).
 [4] D. Li, X. Li, H. Huang, and X. Li, *Quantum Inf. Comput.* **9**, 778 (2009).
 [5] O. Viehmann, C. Eltschka, and J. Siewert, *Phys. Rev. A* **83**, 052330 (2011).
 [6] T. Zhang, M.-J. Zhao, and X. Huang, *J. Phys. A: Math. Theor.* **49**, 405301 (2016).
 [7] G. Gour and N. Wallach, *New J. Phys.* **13**, 073013 (2011).
 [8] A. Burchardt and Z. Raissi, *Phys. Rev. A* **102**, 022413 (2020).

[9] W. K. Wootters, *Phys. Rev. Lett.* **80**, 2245 (1998).
 [10] V. Coffman, J. Kundu, and W. K. Wootters, *Phys. Rev. A* **61**, 052306 (2000).
 [11] P. Love, A. Brink, A. Smirnov, M. Amin, M. Grajcar, E. Il'ichev, A. Izmailkov, and A. Zagoskin, *Quantum Inf. Proc.* **6**, 187 (2007).
 [12] C. Eltschka and J. Siewert, *J. Phys. A: Math. Theor.* **47**, 424005 (2014).
 [13] J.-G. Luque and J.-Y. Thibon, *Phys. Rev. A* **67**, 042303 (2003).
 [14] C. Eltschka, T. Bastin, A. Osterloh, and J. Siewert, *Phys. Rev. A* **85**, 022301 (2012).
 [15] A. Osterloh and J. Siewert, *Phys. Rev. A* **72**, 012337 (2005).
 [16] A. Uhlmann, *Open Syst. Inf. Dyn.* **5**, 209 (1998).
 [17] G. Tóth, T. Moroder, and O. Gühne, *Phys. Rev. Lett.* **114**, 160501 (2015).

- [18] T. J. Osborne and F. Verstraete, *Phys. Rev. Lett.* **96**, 220503 (2006).
- [19] B. Regula, S. Di Martino, S. Lee, and G. Adesso, *Phys. Rev. Lett.* **113**, 110501 (2014).
- [20] B. Regula, A. Osterloh, and G. Adesso, *Phys. Rev. A* **93**, 052338 (2016).
- [21] A. Osterloh, J. Siewert, and A. Uhlmann, *Phys. Rev. A* **77**, 032310 (2008).
- [22] R. Lohmayer, A. Osterloh, J. Siewert, and A. Uhlmann, *Phys. Rev. Lett.* **97**, 260502 (2006).
- [23] A. Osterloh, *Phys. Rev. A* **94**, 062333 (2016).
- [24] S. Gartzke and A. Osterloh, *Phys. Rev. A* **98**, 052307 (2018).
- [25] B. Regula and G. Adesso, *Phys. Rev. Lett.* **116**, 070504 (2016).
- [26] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement*, 2nd ed. (Cambridge University Press, Cambridge, 2017).
- [27] M. Aulbach, *Int. J. Quantum. Inform.* **10**, 1230004 (2012).
- [28] T. Bastin, S. Krins, P. Mathonet, M. Godefroid, L. Lamata, and E. Solano, *Phys. Rev. Lett.* **103**, 070503 (2009).
- [29] P. Ribeiro and R. Mosseri, *Phys. Rev. Lett.* **106**, 180502 (2011).
- [30] C.-F. Kam and R.-B. Liu, *Phys. Rev. A* **101**, 032318 (2020).
- [31] B. Regula and G. Adesso, *Phys. Rev. A* **94**, 022324 (2016).
- [32] The coefficients of states were independently randomly chosen complex numbers with normal distribution $N(0, 1)$ for imaginary and real part. A sampling of $(n + 1)$ -qubit states and calculating their entanglement measure was swift on a PC computer for $n < 20$.
- [33] F. Verstraete, J. Dehaene, and B. De Moor, *Phys. Rev. A* **68**, 012103 (2003).
- [34] A. Sudbery, *J. Phys. A: Math. Gen.* **34**, 643 (2001).
- [35] O. Chterental and D. Z. Djokovic, *Linear Algebra Research Advances* (Nova Science Publishers, New York, 2007), Chap. 4, pp. 133–167.
- [36] C. Spee, J. I. de Vicente, and B. Kraus, *J. Math. Phys.* **57**, 052201 (2016).
- [37] G. Gour and N. Wallach, *J. Math. Phys.* **51**, 112201 (2010).
- [38] C. Eltschka, A. Osterloh, and J. Siewert, *Phys. Rev. A* **80**, 032313 (2009).
- [39] C. Eltschka and J. Siewert, *Phys. Rev. Lett.* **114**, 140402 (2015).
- [40] G. Toth, *Finite Möbius Groups, Minimal Immersions of Spheres, and Moduli* (Springer, Berlin, 2002).
- [41] P. Hayden, D. W. Leung, and A. Winter, *Commun. Math. Phys.* **265**, 95 (2006).
- [42] G. Aubrun and C. Lancie, *Quantum Inf. Comput.* **15**, 513 (2015).
- [43] Z. Puchała, L. Pawela, and K. Życzkowski, *Phys. Rev. A* **93**, 062112 (2016).
- [44] G. Gour, B. Kraus, and N. R. Wallach, *J. Math. Phys.* **58**, 092204 (2017).
- [45] C. Spee, J. I. de Vicente, D. Sauerwein, and B. Kraus, *Phys. Rev. Lett.* **118**, 040503 (2017).
- [46] M. Hebenstreit, C. Spee, N. K. H. Li, B. Kraus, and J. I. de Vicente, *Phys. Rev. A* **105**, 032458 (2022).