

**Completely entangled subspaces of entanglement depth  $k$** Maciej Demianowicz<sup>1</sup>, Kajetan Vogtt<sup>2,3</sup> and Remigiusz Augusiak<sup>2</sup><sup>1</sup>*Institute of Physics and Applied Computer Science, Faculty of Applied Physics and Mathematics, Gdańsk University of Technology, Narutowicza 11/12, 80-233 Gdańsk, Poland*<sup>2</sup>*Center for Theoretical Physics, Polish Academy of Sciences, Aleja Lotników 32/46, 02-668 Warsaw, Poland*<sup>3</sup>*Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland*

(Received 20 December 2023; accepted 7 June 2024; published 1 July 2024)

We introduce a class of entangled subspaces: completely entangled subspaces of entanglement depth  $k$  ( $k$ -CESs). These are subspaces of multipartite Hilbert spaces containing only pure states with an entanglement depth of at least  $k$ . We present an efficient construction of  $k$ -CESs of any achievable dimensionality in any multipartite scenario. Further, we discuss the relation between these subspaces and unextendible product bases (UPBs). In particular, we establish that there is a nontrivial bound on the cardinality of a UPB whose orthocomplement is a  $k$ -CES. Further, we discuss the existence of such UPBs for qubit systems.

DOI: [10.1103/PhysRevA.110.012403](https://doi.org/10.1103/PhysRevA.110.012403)**I. INTRODUCTION**

Entanglement has long been known to be an enabling resource for a variety of information processing protocols such as quantum communication [1] and quantum metrology [2,3]. The ongoing rapid development of quantum network technologies requires a thorough theoretical characterization of its multipartite facet. It is a very challenging problem because the description of quantum systems with multiple nodes is far more complex than that of systems with only two subsystems [4,5]. The topic has attracted much attention in recent years, and many profound results related to its different aspects have been reported in the literature (see, e.g., [6–11]). It is noteworthy that the theoretical advancements have largely been made in parallel with remarkable progress in the experimental domain (see, e.g., [12–16]). Still, despite all the efforts, many aspects of many-body entanglement remain insufficiently explored, and further research to better understand it is highly necessary. This work is in line with this important trend.

A particular useful tool for the characterization of entanglement in multipartite systems is the entanglement depth [17], which indicates how many particles in a given ensemble are genuinely entangled. The concept has found applications, for example, in the domain of cold gases [18–20]. The notion of entanglement can also be meaningfully considered for subspaces in addition to individual states. The most general notion is that of completely entangled subspaces (CESs), which are those subspaces that contain only pure entangled states [21–24]. More specific classes of CESs considered in the literature are, e.g., subspaces containing only states with bounded tensor rank ( $r$ -entangled subspaces) [25,26] or genuinely entangled subspaces [22,27], which are particularly interesting in the multipartite scenario because they contain only genuinely entangled states—the most resourceful states in this framework. Interesting examples of genuinely entangled subspaces are those spanned by the stabilizer error-correction codes, including the five-qubit [28] and toric [29] codes.

Entangled subspaces play an important role in quantum information science, making them objects of intrinsic interest (see, e.g., [30–32] for some recent results). First of all, they constitute an invaluable tool in the theory of entanglement because they can be readily utilized for the construction of mixed entangled states owing to the simple fact that any state supported on an entangled subspace is necessarily entangled. An important class of such constructed states comprises those built from unextendible product bases [33] and having positive partial transpose (PPT). These states share the appealing property of being bound entangled; i.e., their entanglement cannot be distilled into singlets with local operations and classical communication. In fact, the construction from Ref. [33] was the first general construction of multipartite bound entangled states harnessing the notion of completely entangled subspaces. Second, it was shown that almost all subspaces are completely entangled, provided they are not too large, i.e., their dimension does not exceed the maximal permissible one for CESs [24]. This result was utilized to prove that almost all sets of states (separable or entangled) are locally unambiguously distinguishable if the number of elements in the set is not larger than the difference between the dimension of the whole system and the maximal dimension of a CES in the given setup [24,34]. Moreover, entangled subspaces, such as the antisymmetric one and that introduced in Ref. [22], have been exploited to provide counterexamples to the additivity of the minimum output Rényi entropy of quantum channels [35,36]. Their practical applicability further adds to their significance. It has been recognized that they are relevant in the dynamically growing field of quantum error correction [37,38]. A recent development in the research exploring this connection revealed that there exists a certain trade-off relating the larger capability of a code to correct errors to the higher entanglement of the code space [39]. Furthermore, a link between entangled subspaces and self-testing was established [40–43], pointing to their use, e.g., in cryptography.

Our current contribution is to marry the notions of the entanglement depth and entangled subspaces. Precisely, we

introduce completely entangled subspaces of entanglement depth  $k$  ( $k$ -CESs) as those subspaces containing only pure states whose entanglement depth is at least  $k$ . We find the maximal dimensions of such subspaces and present their universal construction from sets of nonorthogonal product vectors in any multipartite setup. Further, we discuss the relation between  $k$ -CESs and unextendible product bases (UPBs). This includes the derivation of a nontrivial bound on the cardinalities of UPBs leading to  $k$ -CESs and a discussion with a positive conclusion about their existence.

## II. PRELIMINARIES

In this section we introduce the relevant notions and terminology.

We consider quantum states defined on  $n$ -partite ( $n \geq 3$ ) Hilbert spaces

$$\mathcal{H}_{d_1 \dots d_n} := \bigotimes_{i=1}^n \mathcal{H}_i = \mathbb{C}^{d_1} \otimes \mathbb{C}^{d_2} \otimes \dots \otimes \mathbb{C}^{d_n}. \quad (1)$$

We will mainly focus on the scenario with equal local dimensions  $d_i = d$ , in which case the space will be denoted as  $\mathcal{H}_{d^n}$ . Single-partite subsystems (parties) are denoted  $A_1, A_2, \dots, A_n := \mathbf{A}$ . A division of the set of parties into  $K$  nonoverlapping nonempty sets  $\mathcal{S}_i$  such that  $\bigcup_i \mathcal{S}_i = \mathbf{A}$  is called a  $K$ -partition and is denoted  $\mathcal{S}_1 | \dots | \mathcal{S}_K$ .

An  $n$ -partite state  $|\Phi\rangle \in \mathcal{H}_{d_1 \dots d_n}$  is called  $k$ -*producible* if it can be written as

$$|\Phi\rangle = |\phi_1\rangle \otimes \dots \otimes |\phi_M\rangle, \quad (2)$$

where  $|\phi_i\rangle$ 's are at most  $k$ -partite states. If a  $k$ -producible state is not  $(k-1)$ -producible at the same time, it is said to have *entanglement depth*  $k$  [17,44]. States with entanglement depth  $k=1$  are named *fully product* because they are products of pure states on individual subsystems  $A_i$ . All other states with  $k \geq 2$  are said to be entangled, and those with entanglement depth equal to  $n$ , in particular, are called *genuinely multipartite entangled* (GME). These are the states which cannot be written as a product for any bipartition of the parties. Excellent examples of GME states are the Greenberger-Horne-Zeilinger (GHZ) states [45],

$$|\text{GHZ}_n\rangle = \frac{1}{\sqrt{2}}(|0\rangle^{\otimes n} + |1\rangle^{\otimes n}), \quad (3)$$

which belong to a general class of GME states called graph states [46].

Let us now recall the concept of entangled subspaces. A subspace of a multipartite Hilbert space  $\mathcal{H}_{d_1 \dots d_n}$  is called *completely entangled* if all pure states belonging to it are entangled [21,22]. Importantly, entanglement of these states can be of any kind, including only bipartite. In the extreme case of all states from a subspace being GME one deals with genuinely entangled subspaces (GESs) [27]. This is the case, for instance, for the antisymmetric subspace.

Entangled subspaces are closely related to the so-called unextendible product bases, and we will extensively exploit this connection in later parts of the paper. Below we recall the basic necessary facts about the discussed notions.

Consider a set of fully product mutually orthogonal  $n$ -partite vectors from  $\mathcal{H}_{d_1 \dots d_n}$ ,

$$\mathcal{B} := \left\{ |\psi^{(j)}\rangle = \bigotimes_{i=1}^n |\varphi_i^{(j)}\rangle \right\}_{j=1}^m, \quad (4)$$

where  $|\varphi_i^{(j)}\rangle \in \mathbb{C}^{d_i}$ .  $\mathcal{B}$  is called an *unextendible product basis* iff it does not span the total Hilbert space  $\mathcal{H}_{d_1 \dots d_n}$  and a fully product vector in the subspace complementary to  $\text{span } \mathcal{B}$  does not exist [33,47]. We will refer to the number of elements in  $\mathcal{B}$  as the size or cardinality of it.

Clearly, by the very definition of a UPB, its complementary subspace is a CES. To illustrate this general link with a simple example let us consider the following set of three-qubit fully product vectors introduced in Ref. [33]:

$$\mathcal{S} := \{|000\rangle, |1+-\rangle, |-1+\rangle, |+ -1\rangle\}, \quad (5)$$

where  $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$ . One can easily verify that  $\mathcal{S}$  is, indeed, a UPB and only entangled vectors belong to  $(\text{span } \mathcal{S})^\perp$ , which itself is thus a CES. Notably, the latter subspace is not a GES because it contains biproduct vectors, e.g.,  $|1\rangle \otimes |\phi\rangle$ , where  $|\phi\rangle$  is an entangled two-qubit vector orthogonal to  $|+-\rangle, |1+\rangle$ , and  $|-1\rangle$ .

Concluding this section, we give two results regarding the unextendibility of a set of product vectors that will be our main tools in later parts. The first is the following observation.

*Fact 1* [33,47]. Consider a set of  $n$ -partite product vectors  $\mathcal{B}$  defined in Eq. (4). A fully product vector orthogonal to  $\mathcal{B}$  exists if and only if a partition of  $\mathcal{B}$  into  $n$  disjoint subsets exists:  $\mathcal{B} = \mathcal{B}_1 \cup \dots \cup \mathcal{B}_n$ , such that for all subsets  $\mathcal{B}_i$  ( $i = 1, 2, \dots, n$ ), the local vectors  $\{|\varphi_i^{(j)}\rangle : |\psi^{(j)}\rangle \in \mathcal{B}_i\}$  do not span the corresponding local Hilbert spaces  $\mathcal{H}_i = \mathbb{C}^{d_i}$ .

This elegant result often serves as a basis for the constructions of UPBs, and it will be used by us for a direct check of whether one of the considered bases possesses a certain property of unextendibility. Its immediate consequence is a lower bound on the cardinalities of unextendible sets:

$$m \geq \sum_{i=1}^n (d_i - 1) + 1. \quad (6)$$

A stronger version of Fact 1 is given in the following corollary.

*Corollary 1.* Consider a set of product vectors  $\mathcal{B}$  defined in Eq. (4) satisfying the condition (6). If for every  $i$  any  $d_i$ -tuple of local vectors  $\{|\varphi_i^{(j)}\rangle\}_j$  spans the corresponding local Hilbert space  $\mathcal{H}_i = \mathbb{C}^{d_i}$ , then the set  $\mathcal{B}$  is unextendible with product vectors.

In fact, for sets with the minimal cardinality  $\sum_i (d_i - 1) + 1$  the constituent vectors must necessarily have the property of full local spanning on subsystems given in the corollary above.

Importantly, both results related to unextendibility, Fact 1 and Corollary 1, are also applicable to sets of nonorthogonal vectors.

### III. COMPLETELY ENTANGLED SUBSPACES OF ENTANGLEMENT DEPTH $k$

Let us move to the main body of the paper. We introduce a class of entangled subspaces defined through the entanglement depth of vectors belonging to them. We propose the following.

*Definition 1.* A subspace  $S \subset \mathcal{H}_{d_1 \dots d_n}$  is called a completely entangled subspace of entanglement depth  $k$  if the entanglement depth of any vector belonging to  $S$  is at least  $k$ . Equivalently, a  $k$ -CES is void of  $(k-1)$ -producible states.

Let us remark that for  $k=2$  the above definition recovers the definition of completely entangled subspaces, whereas for  $k=n$  it recovers the definition of genuinely entangled subspaces.

Before delving into a characterization of  $k$ -CESs, let us give a few simple examples of them, which will allow for a quick grasp of the notion.

As the first example let us take the above-discussed CES complementary to the UPB given in Eq. (5). Within our terminology we can now say that it is a 2-CES (CES), but it is not a 3-CES (GES) because, as discussed, the complementary subspace contains a three-qubit vector with an entanglement depth of 2.

For the second example, we turn our attention to quantum error correction. Consider the celebrated nine-qubit Shor's code [48] described by the vectors

$$|\psi_0\rangle = |\text{GHZ}_3\rangle \otimes |\text{GHZ}_3\rangle \otimes |\text{GHZ}_3\rangle, \quad (7)$$

$$|\psi_1\rangle = |\overline{\text{GHZ}}_3\rangle \otimes |\overline{\text{GHZ}}_3\rangle \otimes |\overline{\text{GHZ}}_3\rangle, \quad (8)$$

where  $|\overline{\text{GHZ}}_3\rangle = (1/\sqrt{2})(|000\rangle - |111\rangle)$  and  $|\text{GHZ}_3\rangle$  is defined in Eq. (3). Now, the subspace  $\text{span}\{|\psi_0\rangle, |\psi_1\rangle\}$  is a 3-CES because the entanglement depth of both  $|\psi_0\rangle$  and  $|\psi_1\rangle$  is 3, and at the same time, any linear combination of these vectors is GME, i.e., it has an entanglement depth equal to 9.

As the final example we give an elementary general construction. With this aim we take a GES of a  $k$ -partite Hilbert space. Then, we take a tensor product of all the vectors from this subspace with an arbitrary, possibly different for each vector, fully product  $(n-k)$ -partite vector. This clearly results in a  $k$ -CES of an  $n$ -partite Hilbert space. This rather trivial method does not lead to  $k$ -CESs of the maximal dimensions for  $k < n$ , as can be seen from dimension considerations (see Appendix B), and in Sec. IV we will provide a more elaborate construction which does this job. However, it is noteworthy that if we choose the otherwise arbitrary vectors to be the same for each of the vectors from a GES, the resulting subspace will be a  $k$ -CES with all vectors belonging to it having an entanglement depth equal exactly to  $k$ .

One of the central problems of the theory of entangled subspaces is the problem of verifying whether a given subspace is entangled and, further, determining to which class it belongs. It is worth noting that some tools have been developed in the literature so far which could also be applied for the case of  $k$ -CESs. Specifically, the simple criterion based on the entanglement of basis vectors from Ref. [49] is directly applicable, and the hierarchical method from Ref. [31] can be readily adapted. Another vital problem in the area is the quantification of subspace entanglement (see [50] for recent results). Here,

also, we find certain methods which can be used in the current case, e.g., the semidefinite programming bounds of Ref. [51].

Let us now investigate how big—in terms of their dimensions— $k$ -CESs can be. The theory of completely entangled subspaces helps us address this question successfully. For simplicity, we concentrate on the case of equal local dimensions in our derivation, but a properly adjusted argument is applicable in the general case too. We have the following proposition.

*Proposition 1.* The maximal attainable dimension of a  $k$ -CES of  $\mathcal{H}_{d^n}$  equals

$$D_{k\text{-CES}}^{\max} = d^n - (td^{k-1} + d^{n-(k-1)t} - t), \quad (9)$$

where

$$t = \left\lfloor \frac{n}{k-1} \right\rfloor. \quad (10)$$

*Proof.* By definition, a  $k$ -CES  $S \subset \mathcal{H}_{d^n}$  does not contain  $(k-1)$ -producible pure states. This implies that  $S$  is a CES in any  $r$ -partite Hilbert space

$$(\mathbb{C}^d)^{\otimes n_1} \otimes (\mathbb{C}^d)^{\otimes n_2} \otimes \dots \otimes (\mathbb{C}^d)^{\otimes n_r}, \quad (11)$$

where  $n_i \leq k-1$  and  $i = 1, 2, \dots, r$ , corresponding to an  $r$ -partition  $S_1|S_2|\dots|S_r$  of the parties, where the size of each subset  $S_i$  is  $|S_i| = n_i$ .

References [21,22] showed that the maximal attainable dimension of a CES of  $\mathcal{H}_{d_1 \dots d_n}$  is given by  $\prod_{i=1}^n d_i - [\sum_{i=1}^n (d_i - 1) + 1]$  [note that the term in the square brackets happens to match the right-hand side of Eq. (6)]. Applying this result to (11), we obtain

$$d^n - \left[ \sum_{i=1}^r (d^{n_i} - 1) + 1 \right], \quad (12)$$

which must further be minimized (equivalently, the term in the square brackets must be maximized) over  $n_i$ 's to obtain an upper bound on the maximal dimension of a  $k$ -CES. This is achieved for  $n_i = k-1$ ,  $i = 1, 2, \dots, t$ , and  $n_{t+1} = n - t(k-1)$ , where  $t = \lfloor n/(k-1) \rfloor$  (see Appendix A). The minimal value of (12) is thus

$$d^n - (td^{k-1} + d^{n-(k-1)t} - t). \quad (13)$$

The choice above corresponds to the partitions of the parties as follows (we assume that for  $k-1|n$  we have  $t$ -partitions):

$$(\mathbb{C}^{d^{k-1}})^{\otimes t} \otimes \mathbb{C}^{d^{n-(k-1)t}}. \quad (14)$$

Further, generic subspaces are known to saturate bounds on the dimensions of entangled subspaces (cf. [25]), meaning that a random subspace of a proper dimension will be a  $k$ -CES. ■

The maximal dimension (9) is a strictly decreasing function of  $k$  for given  $d$  and  $n$  (see Appendix B). For  $k=2$  and  $k=n$ , it obviously recovers the known maximal dimensions of CESs and GESs, respectively, which are explicitly given by

$$D_{2\text{-CES}}^{\max} = d^n - n(d-1) - 1 \quad (15)$$

and

$$D_{n\text{-CES}}^{\max} = (d^{n-1} - 1)(d-1). \quad (16)$$

#### IV. UNIVERSAL CONSTRUCTION OF $k$ -CESs

While random subspaces do have significance in quantum information theory (see, e.g., Ref. [52]), it is the explicit constructions which often help one analyze these objects in more detail, for example, to compute entanglement measures of states supported on them. In this section, we provide a universal analytical construction of  $k$ -CESs; that is, we introduce a method for building  $k$ -CESs of any dimension for any combination of the numbers  $k$ ,  $n$ , and  $d$ . For this aim we utilize the approach put forward recently in [53], which was originally designed for the construction of GESs. It turns out that it is versatile enough to be translated directly to the current case.

The core idea of the construction is to use a set of product vectors for which the full local spanning of Corollary 1 is apparent for properly chosen  $r$ -partitions of the parties [see Eq. (11)]. Consequently, the subspace complementary to the set will be automatically inferred to be void of  $(k-1)$ -producible states, as needed. The desired properties are held by the Vandermonde vectors:

$$\begin{aligned} |v_{d^n}(x)\rangle_{\mathbf{A}} &:= (1, x, x^2, \dots, x^{d^n-1})_{\mathbf{A}}^T \\ &= \bigotimes_{m=1}^n \left( \sum_{s_m=0}^{d-1} x^{q_{m,s_m}} |s_m\rangle \right)_{A_m}, \end{aligned} \quad (17)$$

where

$$q_{m,s_m} = s_m d^{n-m}. \quad (18)$$

Crucially, the Vandermonde matrices, that is, matrices with rows being Vandermonde vectors,

$$V_{K,L} = \sum_{k=0}^{K-1} |k\rangle \langle v_L(x_k)|, \quad (19)$$

with arbitrary  $K$  and  $L$ , are totally positive (i.e., all their minors are strictly positive) whenever the nodes  $\{x_i\}_i$  obey the relation

$$0 < x_0 < x_1 < \dots < x_{K-1}. \quad (20)$$

Let us now consider the following set of linearly independent vectors:

$$\{|v_{d^n}(x_i)\rangle_{\mathbf{A}}\}_{i=0}^{K-1}, \quad (21)$$

where the number  $K$  satisfies the inequality

$$K \geq t d^{k-1} + d^{n-t(k-1)} - t, \quad (22)$$

which follows from Eq. (9), and the nodes  $x_i \in \mathbb{R}$  are ordered as in Eq. (20).

The claim is that the vectors in Eq. (21) are not extendible with vectors of the entanglement depth less than or equal to  $k-1$ , i.e., the subspace orthogonal to their span is a  $k$ -CES. The proof is almost immediate. We write

$$|v_{d^n}(x_i)\rangle_{\mathbf{A}} = |\varphi_1^{(i)}\rangle_{S_1} \otimes |\varphi_2^{(i)}\rangle_{S_2} \otimes \dots \otimes |\varphi_r^{(i)}\rangle_{S_r}, \quad (23)$$

for an  $r$ -partition  $S_1|S_2|\dots|S_r$  of the parties corresponding to (11), in particular (14). We consider matrices with rows being

local vectors on  $S'_j$ s,

$$\mathfrak{S}'_j = \sum_{k=0}^{K-1} |k\rangle \langle \varphi_j^{(k)}|, \quad j = 1, 2, \dots, r. \quad (24)$$

From the total positivity of Vandermonde matrices with positive nodes (see above) it follows that  $\mathfrak{S}'_j$ s are also totally positive. This further means that for every  $j$  any set of  $d^{n_j}$  vectors  $\{|\varphi_j^{(i)}\rangle\}_i$  spans the Hilbert space  $(\mathbb{C}^d)^{\otimes n_j}$  of  $S_j$ . This is the case for any  $r$ -partition with  $n_i \leq k-1$ . By Corollary 1, we conclude that, indeed, there does not exist a  $(k-1)$ -producible state in the orthocomplement of the span of vectors (21), i.e., the resulting subspace is a  $k$ -CES. Its dimension is  $d^n - K$ , which for the smallest  $K$  corresponds to the maximal dimension given in Eq. (9). Explicit (nonorthogonal) bases for such constructed subspaces were given in [53].

Similar to Ref. [53], one could also use the discrete Fourier transform matrices in the construction. The method presented here can be viewed as a construction of “nonorthogonal” UPB $_{k-1}$ ’s (see Sec. V), whose elements are not mutually orthogonal.

We observe that the construction is flexible in that for a given  $k$ -CES we can always increase  $k$  by adding some number of Vandermonde vectors with properly chosen nodes to the construction set (21). For example, consider an 11-dimensional 2-CES, or, simply, a CES in the standard terminology, in the system of four qubits. Its construction requires five Vandermonde vectors. If we now add two more vectors to (21), we obtain a nine-dimensional 3-CES (in Appendix C we give an example of such a subspace). If we again add two vectors, we obtain a seven-dimensional 4-CES, i.e., a GES, of the maximal permissible dimension. Further enlargement of the set of Vandermonde vectors leads to GESs of smaller dimensions.

#### V. $k$ -CESs AND UNEXTENDIBLE PRODUCT BASES

In this section, we explore the link between  $k$ -CESs and UPBs. This line of study is motivated by the important fact that UPBs lead directly to a general construction of multipartite states with undistillable entanglement [33].

We are interested in a certain type of UPBs, namely, those bases which are unextendible with vectors of the entanglement depth less than  $k$  that thus lead to  $k$ -CESs. We thus propose the following.

*Definition 2.* A UPB which is unextendible with  $(k-1)$ -producible vectors, i.e., states with an entanglement depth less than or equal to  $k-1$ , is called a  $(k-1)$ -unextendible product basis, denoted UPB $_{k-1}$ .

For  $k=2$  we have the standard UPBs (see Sec. II), while for  $k=n$  we have genuinely unextendible product bases (GUPBs), that is, UPBs unextendible even with biproduct vectors [27,54], whose existence is currently unknown (see Refs. [54–56] for recent results related to this topic).

By definition, the orthocomplement of a UPB $_{k-1}$  is a  $k$ -CES. Importantly, the reverse statement is not necessarily true even in the cases when the dimension of a  $k$ -CES is such that its complement may admit a UPB $_{k-1}$ . In fact, the orthocomplement of a  $k$ -CES may also be a  $k$ -CES itself.

We observe that the theoretical minimal cardinality of a  $\text{UPB}_{k-1}$  is [see Eq. (10) for the definition of  $t$ ]

$$m_{\min}^{\text{theor}} = \begin{cases} d^n & d = 2 \text{ and } k = n, \\ t(d^{k-1} - 1) + d^{n-(k-1)t} & \text{odd } d \text{ or } [\text{even } d \text{ and } \{(k-1|n \text{ and odd } t) \text{ or } (k-1 \nmid n \text{ and even } t)\}], \\ t(d^{k-1} - 1) + d^{n-(k-1)t} + 1 & \text{otherwise.} \end{cases} \quad (25)$$

These numbers correspond to the minimal cardinalities of UPBs [57–59] for systems defined on Hilbert spaces as in Eq. (14). We will refer to these bounds as *trivial* and discuss their improvement in the next section.

### A. Bound on cardinalities of $\text{UPB}_{k-1}$

The trivial bounds from Eq. (25) can, in many cases, be strengthened if the internal product structure of partitions is appropriately taken into account. The improvement can be achieved by a generalization of the technique recently introduced in Ref. [54] to lower bound permissible cardinalities of  $\text{UPB}_{n-1}$ 's, i.e., GUPBs. There, the pigeonhole principle was the key resource; here, its generalization naturally comes in handy. We have the following.

*Proposition 2.* The number of states  $m$  in a  $\text{UPB}_{k-1}$  is bounded as follows:

$$m \geq d^{k-1} + (n - k + 1) \left( \left\lfloor \frac{d^{k-1} - 2}{k - 1} \right\rfloor + 1 \right). \quad (26)$$

*Proof.* Consider a set of mutually orthogonal fully product vectors

$$\mathcal{K} = \{|v_i\rangle_{\mathbf{A}}\}_{i=1}^m, \quad |v_i\rangle_{\mathbf{A}} = \bigotimes_{j=1}^n |u_j^{(i)}\rangle_{A_j}. \quad (27)$$

We will show that if the number of elements in  $\mathcal{K}$  is strictly smaller than the right-hand side of (26), then  $\mathcal{K}$  is extendible with vectors of the entanglement depth  $k - 1$  or smaller, i.e., it is not a  $\text{UPB}_{k-1}$ .

Any two vectors from  $\mathcal{K}$ ,  $|v_p\rangle$  and  $|v_q\rangle$ , are orthogonal because their local vectors on at least one site  $A_k$  are orthogonal, i.e.,  $\langle u_k^{(p)} | u_k^{(q)} \rangle = 0$  for at least one value of index  $k$ . By the generalized pigeonhole principle [60] (see Appendix D), we conclude that for any vector  $|v_i\rangle$  there are  $w$  sites such that the total number of vectors orthogonal to  $|v_i\rangle$  on these sites is at least

$$s := w \left\lfloor \frac{m-1}{n} \right\rfloor + \min \left( w, m-1 - n \left\lfloor \frac{m-1}{n} \right\rfloor \right). \quad (28)$$

Let us focus on one of the vectors from  $\mathcal{K}$ , e.g.,  $|v_1\rangle$ . Assume that said  $s$  vectors which are orthogonal to  $|v_1\rangle$  are  $\mathcal{K}_1 = \{|v_2\rangle, |v_3\rangle, \dots, |v_{s+1}\rangle\}$ , with orthogonality holding on sites from  $\mathbf{A}_w := A_1, A_2, \dots, A_w$ . Since a vector orthogonal to  $\mathcal{K}_1$  exists on one of the sites from  $\mathbf{A}_w$ , we immediately infer that the elements of  $\mathcal{K}_1$  do not span the whole Hilbert space corresponding to  $\mathbf{A}_w$ , that is,

$$\dim \text{span} \left\{ \bigotimes_{j=1}^w |u_j^{(i)}\rangle_{A_j} \right\}_{i=2}^{s+1} < \dim \mathcal{H}_{\mathbf{A}_w}. \quad (29)$$

Now, if the remaining vectors  $\mathcal{K}_2 := \mathcal{K} \setminus \mathcal{K}_1 = \{|v_1\rangle, |v_{s+2}\rangle, \dots, |v_m\rangle\}$  do not span  $\mathbf{A} \setminus \mathbf{A}_w$ , the corresponding

Hilbert space, i.e.,

$$\dim \text{span } \mathcal{K}_2^{\mathbf{A} \setminus \mathbf{A}_w} < \dim \mathcal{H}_{\mathbf{A} \setminus \mathbf{A}_w}, \quad (30)$$

where

$$\mathcal{K}_2^{\mathbf{A} \setminus \mathbf{A}_w} = \left\{ \bigotimes_{j=w+1}^n |u_j^{(i)}\rangle_{A_j} \right\}_{i=1, s+2, s+3, \dots, m}, \quad (31)$$

then we will find a vector orthogonal to  $\mathcal{K}$  in the form

$$\left( \bigotimes_{j=1}^w |u_j^{(1)}\rangle_{A_j} \right) \otimes |\xi\rangle_{\mathbf{A} \setminus \mathbf{A}_w}, \quad (32)$$

with an arbitrary  $(n - w)$ -partite vector  $|\xi\rangle$  orthogonal to  $\mathcal{K}_2^{\mathbf{A} \setminus \mathbf{A}_w}$ . Clearly, the vector  $|\xi\rangle$  would have, in this case, an entanglement depth of at most  $(n - w)$ , and as a consequence,  $\mathcal{K}$  would not be a  $\text{UPB}_{n-w}$ .

We are interested in the permissible cardinalities of  $\text{UPB}_{k-1}$ 's, in which case we need to consider

$$w = n - k + 1 \quad (33)$$

and verify when the condition (30) holds true. This will provide us with the forbidden cardinalities of the bases, and by negating the obtained bound we will arrive at (26). A sufficient condition for (30) is simply that the number of states in  $\mathcal{K}_2^{\mathbf{A} \setminus \mathbf{A}_w}$  is smaller than the dimension of  $\mathcal{H}_{\mathbf{A} \setminus \mathbf{A}_w}$ , that is,

$$m - s \leq d^{n-w} - 1 := r, \quad (34)$$

which needs to be solved for  $m$ .

Let us define the following function corresponding to the left-hand side of Eq. (34):

$$f_w(m) = m - s = m - w \left\lfloor \frac{m-1}{n} \right\rfloor - \min \left( w, m-1 - n \left\lfloor \frac{m-1}{n} \right\rfloor \right). \quad (35)$$

Using the identities  $\min(x, y) = x + y - \max(x, y)$  and  $\max(x, y) = \max(x - z, y - z) + z$ , we can write this function as

$$f_w(m) = (n - w) \left\lfloor \frac{m-1}{n} \right\rfloor + \max \left( m - w - n \left\lfloor \frac{m-1}{n} \right\rfloor, 1 \right). \quad (36)$$

It is straightforward to show that this function has the following property (see Appendix E):

$$f_w(m+1) = \begin{cases} f_w(m) + 1 & \text{if } m - n \left\lfloor \frac{m}{n} \right\rfloor \geq w + 1 \text{ or } n \mid m, \\ f_w(m) & \text{if } m - n \left\lfloor \frac{m}{n} \right\rfloor \leq w \text{ and } n \nmid m. \end{cases} \quad (37)$$

It then follows that for a fixed  $w$  function  $f_w(m)$  is nondecreasing in  $m$  and takes all integer values. This allows us to look for the largest solution of  $f_w(m) = r$ , that is,

$$(n-w) \left\lfloor \frac{m-1}{n} \right\rfloor + \max \left( m - w - n \left\lfloor \frac{m-1}{n} \right\rfloor, 1 \right) = r. \quad (38)$$

We now observe that we can replace the maximum with the function inside of it, as for any  $m$  for which this function is less than 1, we can find another argument  $\tilde{m} > m$  that reaches 1 without changing the value of the whole function under scrutiny; precisely, we can take  $\tilde{m} = n \left\lfloor \frac{m-1}{n} \right\rfloor + w + 1$  for which  $f_w(m) = f_w(\tilde{m})$ . In turn, we can consider

$$(n-w) \left\lfloor \frac{m-1}{n} \right\rfloor + m - w - n \left\lfloor \frac{m-1}{n} \right\rfloor = r \quad (39)$$

with the assumption that

$$\left\lfloor \frac{m-1}{n} \right\rfloor \leq \frac{m-w-1}{n}. \quad (40)$$

Equation (39) simplifies to

$$m - w - q = r + w, \quad q := \left\lfloor \frac{m-1}{n} \right\rfloor. \quad (41)$$

From Eqs. (40) and (41) we have

$$q \leq \frac{r + qw - 1}{n}, \quad (42)$$

which after solving for  $q$  gives

$$q \leq \frac{r-1}{n-w}, \quad (43)$$

implying further that the largest solution to Eq. (41) is

$$m = r + w + w \left\lfloor \frac{r-1}{n-w} \right\rfloor. \quad (44)$$

Plugging  $r = d^{n-w} - 1$  back in and substituting  $w = n - k + 1$  [(33)], we finally obtain the largest excluded cardinality:

$$m = d^{k-1} + n - k + (n - k + 1) \left\lfloor \frac{d^{k-1} - 2}{k-1} \right\rfloor, \quad (45)$$

meaning that the minimal permissible size of a  $\text{UPB}_{k-1}$  is

$$d^{k-1} + (n - k + 1) \left( \left\lfloor \frac{d^{k-1} - 2}{k-1} \right\rfloor + 1 \right), \quad (46)$$

as claimed in Eq. (26).  $\blacksquare$

We have proved the statement for equal local dimensions, but a similar reasoning can be applied to the general case. We omit the derivation here.

We now show when the obtained bound is nontrivial, i.e., when it gives a lower bound strictly larger than the one

provided by Eq. (25). We have the following chain of inequalities:

$$\begin{aligned} & d^{k-1} + (n - k + 1) \left( \left\lfloor \frac{d^{k-1} - 2}{k-1} \right\rfloor + 1 \right) \\ & \geq d^{k-1} + (n - k + 1) \frac{d^{k-1} - 1}{k-1} \end{aligned} \quad (47)$$

$$\begin{aligned} & = t(d^{k-1} - 1) + 1 + (d^{k-1} - 1) \left( \frac{n}{k-1} - t \right) \\ & \geq t(d^{k-1} - 1) + d^{(k-1)(\frac{n}{k-1} - t)} \end{aligned} \quad (48)$$

$$= t(d^{k-1} - 1) + d^{n - (k-1)t}, \quad (49)$$

where  $t = \lfloor \frac{n}{k-1} \rfloor$ . The first inequality follows from the fact that for any pair of integers  $x$  and  $y > 0$  the following inequality holds true:

$$\left\lfloor \frac{x}{y} \right\rfloor = \left\lceil \frac{x+1}{y} \right\rceil - 1 \geq \frac{x+1}{y} - 1, \quad (50)$$

and the equality in (47) holds iff  $k-1$  divides  $d^{k-1} - 1$ . The second inequality, Eq. (48), follows from Bernoulli's inequality, stating that  $(1+x)^r \leq 1+xr$ , with  $x > -1$  and  $0 \leq r \leq 1$ , here applied with  $x = d^{k-1} - 1$  and  $r = n/(k-1) - t$ . The equality in this case holds iff  $k-1$  divides  $n$  (then  $r = 0$ ;  $r = 1$  is never the case).

Confronting the above conditions for equalities with Eq. (25) provides us with the cases when Eq. (26) is certainly nontrivial. For example, this is the case for odd  $d$  if it additionally holds that  $k-1 \nmid d^{k-1} - 1$  or  $k-1 \nmid n$ . On the other hand, there are clearly cases when it is trivial. This happens, for example, in the case of 3-CESs in systems of four qubits ( $n = 4, d = 2, k = 3$ ). Then, our bound gives 8 as the minimal cardinality, and the same number is obtained from the theory of bipartite UPBs applied to a  $4 \otimes 4$  system.

Note that for  $k = n$  (GUPBs) the bound reproduces, as it should, the one from [54], in which case it is always nontrivial. This bound was recently improved in [56]. At this point, it is not clear whether our current bound on the permissible cardinalities of  $\text{UPB}_{k-1}$ 's can be improved using the graph-theory techniques of [56].

## B. Construction of $k$ -CESs from UPBs

While the bound of Proposition 2 puts limitations on the sizes of  $\text{UPB}_{k-1}$ 's, it does not say anything about the possibility of the actual constructions of bases with permissible cardinalities. This is particularly important in view of the unknown status of this problem for GUPBs ( $k = n$ ). It turns out that  $\text{UPB}_{k-1}$ 's do exist for  $k < n$ , and in fact, there are already examples in the literature for systems of four qubits [61].

We will now reconstruct one of the four-qubit bases from Ref. [61] and show the existence of a  $\text{UPB}_3$  in the case of

five qubits. In both situations, we are interested in the case of  $k = n - 1$ . For this aim we use the results of Ref. [62], where a recursive procedure was given for a construction of UPBs of size  $2^{n-1}$  for any  $n$ .

Let us begin with  $n = 4$ . The UPB from Ref. [62] is given by

$$\mathcal{K}^{(4)} = \{|0000\rangle, |01fe\rangle, |1e1e\rangle, |1fe0\rangle, |e001\rangle, |e1ff\rangle, |fe1f\rangle, |ffel\rangle\}, \quad (51)$$

where  $\{|e\rangle, |f\rangle\}$  is any orthonormal basis different from  $\{|0\rangle, |1\rangle\}$ , e.g.,  $\{|+\rangle, |-\rangle\}$ . It is easy to see that a vector with an entanglement depth equal to 2 orthogonal to  $\mathcal{K}^{(4)}$  exists, i.e., this set is not a  $\text{UPB}_2$ . For example, such a vector is given by  $|\varphi\rangle_{A_1A_4} \otimes |\psi\rangle_{A_2A_3}$ , where  $|\varphi\rangle \perp \{|10\rangle, |f1\rangle\}$  and  $|\psi\rangle \perp \{|00\rangle, |1f\rangle, |e1\rangle\}$ . We can see that this stems from the fact that there are pairs of vectors in  $\mathcal{K}^{(4)}$  which are identical in two subsystems. Here, there are three such pairs, but for sets with a cardinality of 8 this would already be an obstacle if there were only two of them. However, the following amendment can be made to  $\mathcal{K}^{(4)}$  to avoid this problem: in the pairs of vectors with  $|1\rangle$  and  $|e\rangle$  on the first site ( $A_1$ ), we swap vectors on the third site ( $A_3$ ). This leads to the following set:

$$\overline{\mathcal{K}^{(4)}} = \{|0000\rangle, |01fe\rangle, |1eee\rangle, |1f10\rangle, |e0f1\rangle, |e10f\rangle, |fe1f\rangle, |ffel\rangle\}, \quad (52)$$

which recovers one of the UPBs given in Ref. [61], which showed that it is unextendible for any of the two vs two party bipartitions, i.e., that  $\overline{\mathcal{K}^{(4)}}$  is a  $\text{UPB}_2$ . As noted in Sec. V A, its cardinality is the minimal possible.

In the  $n = 5$  case the UPB from Ref. [62] is as follows:

$$\mathcal{K}^{(5)} = \{|00000\rangle, |001fe\rangle, |e001f\rangle, |fe001\rangle, |1fe00\rangle, |01fe0\rangle, |01e1e\rangle, |e01e1\rangle, |1e01e\rangle, |e1e01\rangle, |1e1e0\rangle, |1fffe\rangle, |e1fff\rangle, |fe1ff\rangle, |ffe1f\rangle, |fffel\rangle\}.$$

The set has a visibly cyclic structure, and when proving the (non)existence of a vector with an entanglement depth of 3 orthogonal to all its elements, one needs to consider only two bipartitions because the rest are equivalent to one of them. These bipartitions are  $A_1A_2|A_3A_4A_5$  and  $A_1A_3|A_2A_4A_5$ . With an exhaustive search based on Fact 1 we have verified that  $\mathcal{K}^{(5)}$  is already a  $\text{UPB}_3$  and no modifications of it are necessary.

Since the UPBs of Ref. [62] were constructed recursively, it is plausible that they could be used, directly or after some modifications similar to those shown above in the four-qubit case, to obtain  $\text{UPB}_{n-2}$ 's for any  $n$ .

Let us conclude with an observation that in general, the case  $k - 1|n$  is somewhat special because then we deal with systems with  $t$  subsystems of equal local dimensions  $d^{k-1}$  (recall our assumption of dimension  $d$  of each subsystem  $A_i$ ) in the ‘‘coarse-grained’’ Hilbert space [Eq. (14)]. Constructions of UPBs for homogeneous systems (i.e., with equal local dimensions) are far better explored than those for heterogeneous systems (i.e., different local dimensions). A possible route to find  $\text{UPB}_{k-1}$ 's could then be to build UPBs first for (14) and then look for their proper ‘‘fine-grained’’ versions in the original space.

## VI. CONCLUSIONS AND OUTLOOK

We proposed to consider the notion of the entanglement depth in the context of entangled subspaces by introducing the notion of completely entangled subspaces of entanglement depth  $k$  ( $k$ -CESs), that is, subspaces composed solely of pure states with an entanglement depth of at least  $k$ . We presented a universal construction of such subspaces that works for any multipartite system. We also considered the relationship between unextendible product bases (UPBs) and  $k$ -CESs. In particular, we provided a nontrivial bound on the cardinalities of UPBs whose orthocomplements are  $k$ -CESs. Further, we discussed the problem of constructing such UPBs and provided some examples in systems of several qubits.

From the general point of view, it is natural to expect that the introduced notion may turn out to be relevant for analyses of many-body systems, where the entanglement depth is a figure of merit. The already established connection between entangled subspaces and quantum error correction may suggest that  $k$ -CESs could be of some use also in this area, and their analysis could help us to understand this relationship better. Moreover, it would be interesting to see whether the  $k$ -CESs we have provided could be useful in further studies of counterexamples to the additivity of the minimum output Rényi entropy of quantum channels.

There are also several concrete open problems that naturally emerge from our study. For example, it would be desirable to see whether the obtained bound on the cardinalities of UPBs leading to  $k$ -CESs can be further improved, in particular, using graph-theoretic methods, and to research the possibility of a general practical construction of such UPBs, as this would automatically entail a construction of PPT states. Finally, there is constant demand for practical methods of subspace entanglement certification and quantification tailored to specific types of entangled subspaces, so this line of research is also worth exploration.

## ACKNOWLEDGMENTS

K.V. and R.A. acknowledge the support of the National Science Center (Poland) through the SONATA BIS project (Grant No. 2019/34/E/ST2/00369). Enlightening discussions with M. Wieśniak are acknowledged.

## APPENDIX A: MINIMAL VALUE OF (12)

Let us start with two auxiliary results.

*Lemma 1.* Let  $n$  and  $k \leq n$  be positive integers. Further, let

$$\mathbf{x} = (x_1, x_2, \dots, x_n) \quad (A1)$$

be a nonincreasing sequence of non-negative integers, such that

$$x_{i+1} \leq x_i \leq k - 1, \quad \sum_i x_i = n, \quad (A2)$$

and

$$\mathbf{y} = (\underbrace{k - 1, \dots, k - 1}_{t \text{ times}}, n - t(k - 1), 0, \dots, 0), \quad (A3)$$

where

$$t = \left\lfloor \frac{n}{k - 1} \right\rfloor. \quad (A4)$$

Then, for any  $p \leq n$  it holds that

$$\sum_{i=1}^p x_i \leq \sum_{i=1}^p y_i, \quad (\text{A5})$$

i.e.,  $\mathbf{x}$  is majorized by  $\mathbf{y}$ , written as  $\mathbf{x} \prec \mathbf{y}$ .

*Proof.* The result is obvious.

*Lemma 2.* (See, e.g., [63]). Let  $g(\cdot)$  be a convex function and  $\mathbf{x} \prec \mathbf{y}$ . It then holds that

$$\sum_i g(x_i) \leq \sum_i g(y_i). \quad (\text{A6})$$

Furthermore, if  $g(\cdot)$  is strictly convex and  $\mathbf{x}$  is not a permutation of  $\mathbf{y}$ , then the inequality in Eq. (A6) is strict.

We can now prove the following.

*Fact 2.* The maximal value of

$$\sum_{i=1}^n (d^{n_i} - 1) + 1 \quad (\text{A7})$$

over all  $\mathbf{n} = (n_1, n_2, \dots, n_n)$  such that  $\sum_i n_i = n$  and  $0 \leq n_i \leq k - 1$  is

$$t d^{k-1} - t + d^{n-t(k-1)}, \quad (\text{A8})$$

where  $t = \lfloor \frac{n}{k-1} \rfloor$ .

*Proof.* From Lemma 1 and Lemma 2 with  $g(x) = d^x - 1$ , which is strictly convex, it follows immediately that the maximal value of  $\sum_i (d^{n_i} - 1)$  over all permissible  $\mathbf{n}$ 's corresponds to the choice  $\mathbf{n} = \mathbf{y}$ , with  $\mathbf{y}$  defined in Eq. (A3). Equation (A8) then follows from the direct substitution of the optimal  $n_i$ 's.

The minimal value of the expression in Eq. (12) is obtained by subtracting the obtained value (A8) from  $d^n$ .

## APPENDIX B: PROPERTIES OF $D_{k\text{-CES}}^{\max}$

In this Appendix we prove some basic properties of the maximal dimension of a  $k$ -CES. In what follows we denote

$$D(d, k, n) := D_{k\text{-CES}}^{\max} = d^n - (t_{n,k} d^{k-1} + d^{T_{n,k}} - t_{n,k}), \quad (\text{B1})$$

Since  $2 \leq k \leq n$ , we can easily verify that  $D(d, k, n+1) > D(d, k, n)$ . ■

Case 3 supports the claim made in Sec. III that the construction of  $(n-1)$ -CESs in  $n$ -partite systems from GESs in  $(n-1)$ -partite ones does not lead to subspaces of the maximal dimension.

## APPENDIX C: MAXIMAL 3-CES: FOUR-QUBIT CASE

In this Appendix, we give a nine-dimensional, i.e., maximal, 3-CES in the case of four qubits ( $n=4$ ,  $d=2$ ,  $k=3$ ) obtained with the construction in Sec. IV.

With the choice of nodes  $x_i = i+1$ ,  $i=0, 1, \dots, 6$ , the Vandermonde vectors (21) are now of the form (we omit the

with

$$t_{n,k} := t = \left\lfloor \frac{n}{k-1} \right\rfloor, \quad T_{n,k} := n - (k-1)t_{n,k}. \quad (\text{B2})$$

*Fact 3.* The maximal dimension of a  $k$ -CES is strictly (1) increasing in  $d$  for fixed  $k$  and  $n$ ,  $D(d+1, k, n) > D(d, k, n)$ , (2) decreasing in  $k$  for fixed  $d$  and  $n$ ,  $D(d, k+1, n) < D(d, k, n)$ , and (3) increasing in  $n$  for fixed  $k$  and  $d$ ,  $D(d, k, n+1) > D(d, k, n)$ .

*Proof.* *Case 1.* Assume now  $d$  is a continuous parameter such that  $d \in [2, \infty)$ . It holds that

$$\begin{aligned} \frac{\partial D(d, k, n)}{\partial d} &= \frac{1}{d} \{ n d^n - [t_{n,k}(k-1)d^{k-1} + T_{n,k}d^{T_{n,k}}] \} \\ &> \frac{n}{d} \{ d^n - [t_{n,k}d^{k-1} + \text{sgn}(T_{n,k})d^{T_{n,k}}] \}. \end{aligned} \quad (\text{B3})$$

The term in the curly brackets is, in fact, the difference of the product and the sum of numbers, which are all larger than or equal to 2, and as such, it is larger than or equal to zero, implying that  $\partial D / \partial d > 0$ . This, in turn, means that  $D(d, k, n)$  is strictly increasing in  $d$ . ■

*Case 2.* Recall from Fact 2 in Appendix A that the maximal dimensions  $D(d, k, n)$  and  $D(d, k+1, n)$  correspond in (A7) to, respectively,  $\mathbf{y} = (k-1, \dots, k-1, T_{n,k})$  with  $(k-1)$  appearing  $t_{n,k}$  times [see Eq. (A3)] and

$$\tilde{\mathbf{y}} = (\underbrace{k, \dots, k}_{t_{n,k+1} \text{ times}}, T_{n,k+1}, 0, \dots, 0). \quad (\text{B4})$$

(Note that  $t_{n,k} \geq t_{n,k+1}$ , and we may need to pad zeros in  $\tilde{\mathbf{y}}$ , so that the vectors match in length.) Clearly,  $\mathbf{y} \prec \tilde{\mathbf{y}}$ , and by Lemma 2 it holds that  $D(d, k+1, n) < D(d, k, n)$ . ■

*Case 3.* Taking into account that

$$t_{n+1,k} = \begin{cases} t_{n,k} + 1 & \text{if } k-1 \mid n+1, \\ t_{n,k} & \text{if } k-1 \nmid n+1, \end{cases} \quad (\text{B5})$$

we have

$$D(d, k, n+1) - D(d, k, n) = \begin{cases} d^{n+1} - d^n - d^{k-1} + 1 + d^{T_{n,k}-k+2}(d^{k-2} - 1) & \text{if } k-1 \mid n+1, \\ (d^n - d^{T_{n,k}})(d-1) & \text{if } k-1 \nmid n+1. \end{cases} \quad (\text{B6})$$

transpose and shift the index)

$$\begin{aligned} &(1, i, i^2, \dots, i^{15})_{\mathbf{A}} \\ &= (1, i^8)_{\mathbf{A}_1} \otimes (1, i^4)_{\mathbf{A}_2} \otimes (1, i^2)_{\mathbf{A}_3} \otimes (1, i)_{\mathbf{A}_4}, \end{aligned} \quad (\text{C1})$$

with  $i = 1, 2, \dots, 7$ . The matrix built from these vectors is totally positive. This implies that matrices constructed from bipartite vectors on any pair  $A_m A_n$  [see Eq. (24)] are always full rank, meaning, in particular, that any four ( $d^2$  with  $d=2$ ) such vectors span the whole four-dimensional Hilbert space. Corollary 1 tells us then that a vector of the form  $|\phi\rangle_{A_m A_n} \otimes |\varphi\rangle_{A \setminus A_m A_n}$ , i.e., a 2-producible vector, which is orthogonal to all the vectors (C1) does not exist. Consequently, the orthocomplement of the span of the vectors (C1) is a 3-CES.

We present a nonorthogonal basis for this 3-CES as a matrix whose rows are the basis vectors:



0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

(C2) and

$$f_w(m + 1) = m - w + 1 - w \left\lfloor \frac{m}{n} \right\rfloor = f_w(m) + 1. \tag{E10}$$

**APPENDIX D: GENERALIZED PIGEONHOLE PRINCIPLE**

For convenience, we recall here the statement of the generalized pigeonhole principle [60].

*Fact 4.* If  $pq + r$  objects are put into  $q$  boxes, then for each  $0 \leq s \leq q$ ,  $s$  boxes with the total number of objects in them being at least  $ps + \min(r, s)$  exist.

**APPENDIX E: PROOF OF EQUATION (37)**

Let us start by recalling the definition of the function in question:

$$f_w(m) = (n - w) \left\lfloor \frac{m - 1}{n} \right\rfloor + \max \left( m - w - n \left\lfloor \frac{m - 1}{n} \right\rfloor, 1 \right), \tag{E1}$$

with  $w = n - k + 1$ .

*Case 1.  $n|m$ .* We begin with the case of  $n$  being a divisor of  $m$ . We then have

$$\left\lfloor \frac{m}{n} \right\rfloor = \frac{m}{n}, \quad \left\lfloor \frac{m - 1}{n} \right\rfloor = \frac{m}{n} - 1. \tag{E2}$$

It follows that

$$\begin{aligned} f_w(m) &= (n - w) \left( \frac{m}{n} - 1 \right) + \max(n - w, 1) \\ &= (n - w) \frac{m}{n} \end{aligned} \tag{E3}$$

since  $n - w = k - 1 \geq 1$ . Further,

$$f_w(m + 1) = (n - w) \left\lfloor \frac{m}{n} \right\rfloor + \max \left( m + 1 - w - n \left\lfloor \frac{m}{n} \right\rfloor, 1 \right) \tag{E4}$$

$$= (n - w) \frac{m}{n} + \max(1 - w, 1) \tag{E5}$$

$$= (n - w) \frac{m}{n} + 1 = f_w(m) + 1 \tag{E6}$$

because  $w = n - k + 1 \geq 1$ .

*Case 2.  $n \nmid m$ .* We now consider the opposite case of  $n$  not being a divisor of  $m$ . It now holds that

$$\left\lfloor \frac{m - 1}{n} \right\rfloor = \left\lfloor \frac{m}{n} \right\rfloor. \tag{E7}$$

Within the current case we analyze two subcases related to the value of the function under the maximum in  $f_w(m)$ .

*Subcase 2a.  $m - n \lfloor \frac{m}{n} \rfloor - w \geq 1$ .* We just note that this condition can never be satisfied simultaneously with case 1 and is valid only for  $n \nmid m$ . We have

$$m - n \left\lfloor \frac{m - 1}{n} \right\rfloor - w = m - n \left\lfloor \frac{m}{n} \right\rfloor - w \geq 1 \tag{E8}$$

in view of Eq. (E7). As a consequence, in both  $f_w(m)$  and  $f_w(m + 1)$  we can take the function under the maximum as its value. We then have, after trivial simplifications,

$$f_w(m) = m - w - w \left\lfloor \frac{m - 1}{n} \right\rfloor \tag{E9}$$

*Subcase 2b.*  $m - n\lfloor \frac{m}{n} \rfloor - w \leq 0$ . Now, the maximum is easily seen to be equal to 1 for both  $f_w(m)$  and  $f_w(m+1)$ ,

which, taking into account Eq. (E7), results in this subcase in  $f_w(m) = f_w(m+1)$ .

This concludes the proof of Eq. (37).

- 
- [1] R. Ursin, F. Tiefenbacher, T. Schmitt-Manderbach, H. Weier, T. Scheidl, M. Lindenthal, B. Blauensteiner, T. Jennewein, J. Perdigues, P. Trojek, B. Ömer, M. Fürst, M. Meyenburg, J. Rarity, Z. Sodnik, C. Barbieri, H. Weinfurter, and A. Zeilinger, Entanglement-based quantum communication over 144 km, *Nat. Phys.* **3**, 481 (2007).
- [2] V. Giovannetti, S. Lloyd, and L. Maccone, Advances in quantum metrology, *Nat. Photonics* **5**, 222 (2011).
- [3] G. Tóth and I. Apellaniz, Quantum metrology from a quantum information science perspective, *J. Phys. A: Math. Theor.* **47**, 424006 (2014).
- [4] W. Dür, G. Vidal, and J. I. Cirac, Three qubits can be entangled in two inequivalent ways, *Phys. Rev. A* **62**, 062314 (2000).
- [5] F. Verstraete, J. Dehaene, B. De Moor, and H. Verschelde, Four qubits can be entangled in nine different ways, *Phys. Rev. A* **65**, 052112 (2002).
- [6] W. Dür and J. I. Cirac, Activating bound entanglement in multipartite systems, *Phys. Rev. A* **62**, 022302 (2000).
- [7] M. Horodecki, J. Oppenheim, and A. Winter, Partial quantum information, *Nature (London)* **436**, 673 (2005).
- [8] G. Adesso and I. Fuentes-Schuller, Correlation loss and multipartite entanglement across a black hole horizon, *Quantum Inf. Comput.* **9**, 0657 (2009).
- [9] L. Chen, E. Chitambar, R. Duan, Z. Ji, and A. Winter, Tensor rank and stochastic entanglement catalysis for multipartite pure states, *Phys. Rev. Lett.* **105**, 200501 (2010).
- [10] R. Chaves, D. Cavalcanti, and L. Aolita, Causal hierarchy of multipartite Bell nonlocality, *Quantum* **1**, 23 (2017).
- [11] H. Yamasaki, S. Morelli, M. Miethlinger, J. Bavaresco, N. Friis, and M. Huber, Activation of genuine multipartite entanglement: Beyond the single-copy paradigm of entanglement characterisation, *Quantum* **6**, 695 (2022).
- [12] J.-W. Pan, Z.-B. Chen, C.-Y. Lu, H. Weinfurter, A. Zeilinger, and M. Żukowski, Multiphoton entanglement and interferometry, *Rev. Mod. Phys.* **84**, 777 (2012).
- [13] W. McCutcheon, A. Pappa, B. A. Bell, A. McMillan, A. Chailloux, T. Lawson, M. Mafu, D. Markham, E. Diamanti, I. Kerenidis, J. G. Rarity, and M. S. Tame, Experimental verification of multipartite entanglement in quantum networks, *Nat. Commun.* **7**, 13251 (2016).
- [14] R. Schmied, J.-D. Bancal, B. Allard, M. Fadel, V. Scarani, P. Treutlein, and N. Sangouard, Bell correlations in a Bose-Einstein condensate, *Science* **352**, 441 (2016).
- [15] N. J. Engelsens, R. Krishnakumar, O. Hosten, and M. A. Kasevich, Bell correlations in spin-squeezed states of 500 000 atoms, *Phys. Rev. Lett.* **118**, 140401 (2017).
- [16] F. Fröwis, P. Strassmann, A. Tiranov, C. Gut, J. Lavoie, N. Brunner, F. Bussières, M. Afzelius, and N. Gisin, Experimental certification of millions of genuinely entangled atoms in a solid, *Nat. Commun.* **8**, 907 (2017).
- [17] A. S. Sørensen and K. Mølmer, Entanglement and extreme spin squeezing, *Phys. Rev. Lett.* **86**, 4431 (2001).
- [18] C. Gross, T. Zibold, E. Nicklas, J. Estève, and M. K. Oberthaler, Nonlinear atom interferometer surpasses classical precision limit, *Nature (London)* **464**, 1165 (2010).
- [19] L.-M. Duan, Entanglement detection in the vicinity of arbitrary Dicke states, *Phys. Rev. Lett.* **107**, 180502 (2011).
- [20] B. Lücke, J. Peise, G. Vitagliano, J. Arlt, L. Santos, G. Tóth, and C. Klempt, Detecting multiparticle entanglement of Dicke states, *Phys. Rev. Lett.* **112**, 155304 (2014).
- [21] N. R. Wallach, An unentangled Gleason's theorem, *Contemp. Math.* **305**, 291 (2002).
- [22] K. Parthasarathy, On the maximal dimension of a completely entangled subspace for finite level quantum systems, *Proc. Math. Sci.* **114**, 365 (2004).
- [23] B. V. Rajarama Bhat, A completely entangled subspace of maximal dimension, *Int. J. Quantum Inf.* **04**, 325 (2006).
- [24] J. Walgate and A. J. Scott, Generic local distinguishability and completely entangled subspaces, *J. Phys. A: Math. Theor.* **41**, 375305 (2008).
- [25] T. Cubitt, A. Montanaro, and A. Winter, On the dimension of subspaces with bounded Schmidt rank, *J. Math. Phys.* **49**, 022107 (2008).
- [26] B. Lovitz and N. Johnston, Entangled subspaces and generic local state discrimination with pre-shared entanglement, *Quantum* **6**, 760 (2022).
- [27] M. Demianowicz and R. Augusiak, From unextendible product bases to genuinely entangled subspaces, *Phys. Rev. A* **98**, 012313 (2018).
- [28] R. Laflamme, C. Miquel, J. P. Paz, and W. H. Żurek, Perfect quantum error correcting code, *Phys. Rev. Lett.* **77**, 198 (1996).
- [29] A. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys. (NY)* **321**, 2 (2006).
- [30] O. Makuta, B. Kuzaka, and R. Augusiak, Fully non-positive-partial-transpose genuinely entangled subspaces, *Quantum* **7**, 915 (2023).
- [31] N. Johnston, B. Lovitz, and A. Vijayaraghavan, Complete hierarchy of linear systems for certifying quantum entanglement of subspaces, *Phys. Rev. A* **106**, 062443 (2022).
- [32] N. Johnston, B. Lovitz, and A. Vijayaraghavan, Computing linear sections of varieties: Quantum entanglement, tensor decompositions and beyond, in *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)* (IEEE Computer Society, Los Alamitos, CA, USA, 2023), pp. 1316–1336.
- [33] C. H. Bennett, D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. M. Terhal, Unextendible product bases and bound entanglement, *Phys. Rev. Lett.* **82**, 5385 (1999).
- [34] A. Chefles, Condition for unambiguous state discrimination using local operations and classical communication, *Phys. Rev. A* **69**, 050307(R) (2004).
- [35] A. Grudka, M. Horodecki, and Ł. Pankowski, Constructive counterexamples to the additivity of the minimum output Rényi entropy of quantum channels for all  $p > 2$ , *J. Phys. A: Math. Theor.* **43**, 425304 (2010).
- [36] K. Szczygielski and M. Studziński, New constructive counterexamples to additivity of minimum output Rényi entropy of quantum channels, [arXiv:2301.07428](https://arxiv.org/abs/2301.07428).
- [37] G. Gour and N. R. Wallach, Entanglement of subspaces and error-correcting codes, *Phys. Rev. A* **76**, 042309 (2007).

- [38] A. J. Scott, Multipartite entanglement, quantum-error-correcting codes, and entangling power of quantum evolutions, *Phys. Rev. A* **69**, 052330 (2004).
- [39] F. Huber and M. Grassl, Quantum codes of maximal distance and highly entangled subspaces, *Quantum* **4**, 284 (2020).
- [40] A. H. Shenoy and R. Srikanth, Maximally nonlocal subspaces, *J. Phys. A: Math. Theor.* **52**, 095302 (2019).
- [41] F. Baccari, R. Augusiak, I. Šupić, and A. Acín, Device-independent certification of genuinely entangled subspaces, *Phys. Rev. Lett.* **125**, 260507 (2020).
- [42] O. Makuta and R. Augusiak, Self-testing maximally-dimensional genuinely entangled subspaces within the stabilizer formalism, *New J. Phys.* **23**, 043042 (2021).
- [43] I. Frérot and A. Acín, Coarse-grained self-testing, *Phys. Rev. Lett.* **127**, 240401 (2021).
- [44] O. Gühne, G. Tóth, and H. J. Briegel, Multipartite entanglement in spin chains, *New J. Phys.* **7**, 229 (2005).
- [45] D. M. Greenberger, M. A. Horne, and A. Zeilinger, Going beyond Bell's theorem, in *Bell's Theorem, Quantum Theory and Conceptions of the Universe*, edited by M. Kafatos (Springer, New York, 2004), pp. 69–72.
- [46] M. Hein, W. Dür, J. Eisert, R. Raussendorf, M. Van den Nest, and H.-J. Briegel, Entanglement in graph states and its applications, in *Quantum Computers, Algorithms and Chaos, Proceedings of the International School of Physics "Enrico Fermi"* Vol. 162 (IOS Press, Amsterdam, 2006), pp. 115–218.
- [47] D. P. DiVincenzo, T. Mor, P. W. Shor, J. A. Smolin, and B. M. Terhal, Unextendible product bases, uncompletable product bases and bound entanglement, *Commun. Math. Phys.* **238**, 379 (2003).
- [48] P. W. Shor, Scheme for reducing decoherence in quantum computer memory, *Phys. Rev. A* **52**, R2493(R) (1995).
- [49] M. Demianowicz, G. Rajchel-Mieldzioc, and R. Augusiak, Simple sufficient condition for subspace to be completely or genuinely entangled, *New J. Phys.* **23**, 103016 (2021).
- [50] X. Zhu, Ch. Zhang, and B. Zeng, Quantifying subspace entanglement with geometric measures, [arXiv:2311.10353](https://arxiv.org/abs/2311.10353).
- [51] M. Demianowicz and R. Augusiak, Entanglement of genuinely entangled subspaces and states: Exact, approximate, and numerical results, *Phys. Rev. A* **100**, 062318 (2019).
- [52] P. Hayden, Entanglement in random subspaces, *AIP Conf. Proc.* **734**, 226 (2004).
- [53] M. Demianowicz, Universal construction of genuinely entangled subspaces of any size, *Quantum* **6**, 854 (2022).
- [54] M. Demianowicz, Negative result about the construction of genuinely entangled subspaces from unextendible product bases, *Phys. Rev. A* **106**, 012442 (2022).
- [55] F. Shi, M.-S. Li, X. Zhang, and Q. Zhao, Unextendible and uncompletable product bases in every bipartition, *New J. Phys.* **24**, 113025 (2022).
- [56] F. Shi, G. Bai, X. Zhang, Q. Zhao, and G. Chiribella, Graph-theoretic characterization of unextendible product bases, *Phys. Rev. Res.* **5**, 033144 (2023).
- [57] N. Alon and L. Lovász, Unextendible product bases, *J. Comb. Theory, Ser. A* **95**, 169 (2001).
- [58] K. Feng, Unextendible product bases and 1-factorization of complete graphs, *Discrete Appl. Math.* **154**, 942 (2006).
- [59] J. Chen and N. Johnston, The minimum size of unextendible product bases in the bipartite case (and some multipartite cases), *Commun. Math. Phys.* **333**, 351 (2015).
- [60] Ch. W. Wu, On graphs whose Laplacian matrix's multipartite separability is invariant under graph isomorphism, *Discrete Math.* **310**, 2811 (2010).
- [61] K. Wang, L. Chen, L. Zhao, and Y. Guo,  $4 \times 4$  unextendible product basis and genuinely entangled space, *Quantum Inf. Process.* **18**, 202 (2019).
- [62] R. Augusiak, T. Fritz, M. Kotowski, M. Kotowski, M. Pawłowski, M. Lewenstein, and A. Acín, Tight Bell inequalities with no quantum violation from qubit unextendible product bases, *Phys. Rev. A* **85**, 042113 (2012).
- [63] A. W. Marshall, I. Olkin, and B. C. Arnold, *Inequalities: Theory of Majorization and Its Applications*, Springer Series in Statistics (Springer, New York, NY, 2010).