


Generation of multiple bound states in the continuum through doubly degenerate quasiguided modes

Tetsuyuki Ochiai *Research Center for Electronic and Optical Materials, National Institute for Materials Science, Tsukuba 305-0044, Japan*

(Received 2 May 2024; revised 5 July 2024; accepted 10 July 2024; published 2 August 2024)

We present a detailed theoretical analysis of a peculiar generation of multiple bound states in the continuum (BICs) in two-dimensional periodic arrays of dielectric spheres. They emerge in high-symmetry lattices with the C_{6v} and C_{4v} point groups and involve doubly degenerate quasiguided modes at the Γ point that can couple to external radiation. By tuning a system parameter, the doubly degenerate modes can exhibit accidental BICs at a critical parameter. In the vicinity of the critical parameter, the two bands originating from the degenerate mode exhibit multiple off- Γ BICs. They move and annihilate across the two bands by changing the parameter around the critical one. A ring-like high- Q channel pinned with multiple BICs emerges particularly for the C_{6v} case. Across the critical coupling, the total vorticity of the multiple BICs is conserved. The $\mathbf{k} \cdot \mathbf{p}$ perturbation theory explains some features of the phenomena reasonably well.

DOI: [10.1103/PhysRevA.110.023504](https://doi.org/10.1103/PhysRevA.110.023504)

I. INTRODUCTION

Optical bound states in the continuum (BIC) are localized eigenstates embedded in the radiation continuum [1]. They have infinite quality factors or, in other words, vanishing decay rates, although they are inside the light cone. Strong light confinement via the infinite quality factor enables us to investigate various applications such as lasing [2], sensing [3], nonlinear optics [4], and so on, via BICs.

The BICs are found typically in monolayers of spheres and photonic crystal slabs at the Γ point [5–8]. Off- Γ BICs are also found at generic \mathbf{k} points [9,10] via the formation of polarization vortices [11]. The former BICs are symmetry-protected and irrelevant to physical parameters unless the relevant symmetry is unchanged. The latter ones are topologically protected, moving in momentum space. Therefore, a parameter scan is necessary to find the latter BICs at a given \mathbf{k} point other than Γ . In this sense, the off- Γ BICs are sometimes called accidental.

Recently, merging of an at- Γ BIC and off- Γ BICs in a nondegenerate isolated band attracts much interest as the so-called super-BIC [12,13]. This BIC is obtained by tuning system parameters such that the off- Γ BICs move toward the Γ point, where the symmetry-protected BIC exists. The super BIC has superb properties as it involves an extreme suppression of the decay rate in a broad region of the momentum space around Γ . The decay rate behaves as k^6 (k^8 for higher-order charges [14]), in striking contrast to the k^2 (k^4 for higher-order charges) behavior of the symmetry-protected BICs. Merging among BICs is available also off the Γ point [15], resulting in a $(k - k_{\text{BIC}})^4$ behavior.

Another type of super-BICs was found recently in monolayers of spheres [16]. From now on, we call it the next-to-super-BIC. It is obtained at the Γ point by tuning system parameters and exhibits a k^4 scaling of the decay rate at a critical parameter. In contrast to the ordinary super-BIC,

it involves doubly degenerate eigenmodes at the Γ point. Moreover, off-critical parameters result in a ring-like high- Q channel in the triangular lattice system. This high- Q channel is either P - or S -polarization-like and can be switched by changing the parameter across the critical value.

Once we have a high- Q channel, we have a continuous distribution of optical modes with strongly enhanced light-matter interactions. The channel can be selectively excited by the polarization of the incident light. As a result, we can steer the coherent radiation continuously along the ring. The output light has nearly the same frequency, but the angle changes continuously. Such a beam steering paves the way for various applications.

In this paper, we further investigate the next-to-super-BICs and show that multiple off- Γ BICs are generated at off-critical parameters nearby. We show more complex behaviors than in Ref. [16] can be observed. In addition to the ring-like channel, discrete multiple BICs can be generated through the next-to-super-BICs. In this case, the ring is formed asymmetrically in the parameter space. If we change the parameter across the critical value, the ring disappears, and multiple discrete BICs are again generated. Across the critical parameter, the total vorticity of the BICs is conserved. In addition, the appearance of the ring depends on the lattice structure.

One of the points behind the above phenomena is spatial symmetry. We show that the $\mathbf{k} \cdot \mathbf{p}$ perturbation theory based on spatial symmetry explains some properties of the phenomena reasonably well.

The ring-like high- Q channel can be explained in terms of the multipolar lattice [17,18]. However, the lattice-structure dependence needs complementary approaches to understand the phenomena. The $\mathbf{k} \cdot \mathbf{p}$ perturbation provides one reasonable scenario as shown in this paper.

The multiple off- Γ BICs are also produced by breaking spatial symmetries of photonic crystal (PhC) slabs [19–21]. There, degenerate or nondegenerate symmetry-protected

BICs at Γ turn into multiple off- Γ BICs. Since the original symmetry-protected BICs exist irrespective of system parameters, the resulting multiple BICs are deterministic and parameter tuning is unnecessary.

This paper is organized as follows. In Secs. II and III, we present numerical examples of the accidental at- Γ BICs and resulting multiple off- Γ BICs in the triangular and square lattices of dielectric spheres, respectively. In Sec. IV, we analyze these phenomena via the $\mathbf{k} \cdot \mathbf{p}$ perturbation theory of the Maxwell equation. Finally, in Sec. V, we summarize the results.

II. TRIANGULAR LATTICE

Let us first consider the next-to-super-BICs in a monolayer of dielectric spheres arranged in a triangular lattice. The system has the C_{6v} point group symmetry [22]. Therefore, the eigenmodes are classified according to the irreducible representations of C_{6v} . At the Γ point, the eigenmodes with the E_1 representation of C_{6v} can couple to external radiation. The other eigenmodes of $A_1, A_2, B_1, B_2,$ and E_2 representations are uncoupled, provided their eigenfrequencies are below the diffraction threshold. Thus, the former eigenmodes generally have finite lifetimes, whereas the latter eigenmodes have infinite lifetimes and are symmetry-protected BICs.

However, tuning system parameters can cause the former eigenmodes to have infinite lifetimes. In this way, accidental at- Γ BICs can occur for the E_1 modes.

Figure 1 shows the resonance angular frequency ω_k and width γ_k of the E_1 modes at the Γ point ($k = 0$). They are evaluated with the photonic Korringa-Kohn-Rostoker (KKR) method [23–25] together with the curve fitting to the Breit-Wigner formula [26] of the scattering phase shift δ as a function of angular frequency ω :

$$e^{2i\delta} = e^{2i\delta_k} \left(\frac{\omega - \omega_k - i\gamma_k}{\omega - \omega_k + i\gamma_k} \right)^n, \quad (1)$$

where δ_k is the background (frequency-independent) phase shift, and n is the degree of degeneracy of the resonant mode concerned. The scattering phase shift is derived from the S matrix of the monolayer [27]. At several radii, the eigenmodes exhibit infinite quality factor accidentally.

Such an accidental at- Γ BIC can be easily found theoretically and experimentally by monitoring the change of the resonance signal in the transmission spectrum of the normal incidence. For instance, Fig. 2 shows the change of the transmission spectrum with the sphere radius around the critical one of BIC1 in Fig. 1. Since the modes other than E_1 are symmetry-protected, they do not affect the transmission spectrum. Solely the modes of the E_1 representation emerge as an asymmetric resonance signal of the Fano shape [28] in the spectrum. The spectrum changes with the radius, and we can find a narrowing of the resonance width toward the critical radius. The vanishing resonance width corresponds to the accidental at- Γ BIC.

The above property of the next-to-super-BIC presents a marked contrast to the super BIC. In the latter case, the BIC consists of a symmetry-protected BIC and topologically protected BICs, so it is impossible to observe the super BIC




FIG. 1. The resonance (a) angular frequencies ω_0 and (b) widths γ_0 of the E_1 modes at the Γ point in the triangular-lattice monolayer of dielectric spheres with the C_{6v} point group symmetry, as a function of sphere radius r_s . The modes are classified according to the parity in the z direction, assuming the centers of the spheres are on the $z = 0$ plane. The dielectric constant of the spheres is taken to be 12, and that of the background material is 1. The lattice constant is denoted by a . Accidental at- Γ BICs (next-to-super BICs) occur around the modes indicated by solid circles. The inset in (a) stands for the geometry of the monolayer.

via the transmission spectrum of the normal incidence. It is available only through a detailed analysis of the momentum and parameter dependence of the spectrum.

Figure 3 shows the real and imaginary photonic band structure around the critical radius of BIC1. Here, the real photonic band structure is referred to as the resonance angular frequency ω_k , and the imaginary one is to the resonance width γ_k , as a function of Bloch momentum k . Although the real band structure does not change so much in its shape, the imaginary band structure exhibits a clear contrast between the two bands of opposite parities and among the three values of on- and off-critical radii. Below the critical radius [Figs. 3(a) and 3(b)], the odd-parity band exhibits the off- Γ BIC at about $ka/2\pi = 0.1$. At the critical radius [Figs. 3(c) and 3(d)], both the bands exhibit a flat region of nearly zero values in their imaginary parts. This suppression of the decay rate resembles that in the super-BIC. However, now γ_k behaves as k^4 instead of the k^6 behavior of the super-BIC. Above the critical radius [Figs. 3(e) and 3(f)], the even-parity band exhibits the off- Γ BIC.

For comparison, the photonic bands originating from the symmetry-protected BICs at Γ exhibit a quick blow-up of the imaginary parts with the k^2 scaling as shown in Fig. 4. Near the Γ point, we do not observe the trend of decreasing the imaginary part toward a minimum. We should point out that the imaginary part is much larger than that from the E_1 mode, whose imaginary part is significantly suppressed.




FIG. 2. Specular transmission spectra of the normal incidence in the triangular lattice monolayer of dielectric spheres. The sphere radius varies from $0.36a$ to $0.41a$ with a $0.005a$ step. The other parameters are the same as in Fig. 1. The vanishing resonance signal of the red curve indicates the accidental at- Γ BIC of BIC1 in Fig. 1.

By scanning the Bloch momentum in all the directions around the Γ point, the real band structure consists of two surfaces touched quadratically at the Γ point, as shown in Fig. 5. The upper and lower bands are S - and P -polarization-like, respectively.

Figure 6 shows the Q -value map of the photonic band modes at the off-critical radii. It is remarkable that the ring-like high- Q channels found in Ref. [16] are formed. The channels lie in the S -polarization-like upper band at $r_s < r_c$ and in the P -polarization-like lower band for $r_s > r_c$. The other bands (the lower band at $r_s < r_c$ and the upper band at $r_s > r_c$) do not have such channels. These channels indicate that the minimum decay rate γ_k at the off-critical radii found in Fig. 3 emerges at nearly the same distance from the Γ point regardless of momentum orientation. As we move the radius r_s toward r_c , we can show the ring shrinks to the Γ point and across $r_s = r_c$, the ring moves from the upper band to the lower band or vice versa.

Looking closely at the rings, we find 12 off- Γ BICs: six on the equivalent ΓM axes and six on the equivalent ΓK axes. These axes are the mirror axes of the triangular-lattice Brillouin zone. The polarization vortices of the BICs are very elongated along the rings. The vortex charge diagram is shown in Fig. 7. The vortex charges are opposite between ΓM and ΓK . Off the mirror axes, the Q values in the ring are still very high, of order 10^{-8} . Thus, the ring-like high Q channel is of quasi-BICs pinned with the true BICs on the mirror axes.

In the above argument, the polarization ellipse in two-dimensional (2D) momentum space is for the complex 2D




FIG. 3. Real and imaginary photonic band structures of the z -even parity originating from the E_1 mode around the critical radius r_c of BIC1 in Fig. 1. The momentum is taken along the ΓM and ΓK directions. The band structure is classified according to the in-plane parity concerning the ΓM and ΓK axes. The sphere radius is fixed as (a,b) $r_s = 0.38a$, (c,d) $r_s = 0.385a$ ($\simeq r_c$), and (e,f) $r_s = 0.388a$. The other parameters are the same as in Fig. 1. The inset in (b) represents the first Brillouin zone of the triangular lattice.

polarization vector \mathbf{e}^+ defined by

$$\mathbf{e}^+ = t_{p0}^+ \hat{\mathbf{k}} + t_{s0}^+ \hat{\mathbf{k}}_\perp, \quad (2)$$

$$\mathbf{E}^\pm(\mathbf{x}) = \sum_{\mathbf{g}} (t_{pg}^\pm \mathbf{P}_g^\pm + t_{sg}^\pm \mathbf{S}_g) e^{i\mathbf{K}_g^\pm \cdot \mathbf{x}}, \quad (3)$$

$$\mathbf{P}_g^\pm = \pm \frac{\Gamma_g}{q_0} \hat{\mathbf{k}}_g - \frac{|\mathbf{k}_g|}{q_0} \hat{\mathbf{z}}, \quad \mathbf{S}_g = (\hat{\mathbf{k}}_g)_\perp, \quad (4)$$

$$\mathbf{K}_g^\pm = \mathbf{k}_g \pm \Gamma_g \hat{\mathbf{z}}, \quad \Gamma_g = \sqrt{q_0^2 - (\mathbf{k}_g)^2}, \quad (5)$$

$$\mathbf{k}_g = \mathbf{k} + \mathbf{g}, \quad q_0 = \frac{\omega}{c}, \quad (6)$$

where $\mathbf{E}^\pm(\mathbf{x})$ is the electric field of the photonic band mode above (superscript $+$) and below ($-$) the monolayer (see Appendix B), \mathbf{k} is a 2D Bloch momentum, and \mathbf{g} is a 2D reciprocal lattice. In the far field, solely the $\mathbf{g} = 0$ component survives, so that the far-field polarization is determined by t_{p0}^+ and t_{s0}^+ . In addition, the vortex charge q is defined by

$$q = \frac{1}{2\pi} \oint d\theta_k, \quad (7)$$




FIG. 4. Real and imaginary photonic band structure originating from symmetry-protected BICs at the Γ point. The z parity is odd. The momentum is taken along the ΓK direction. The sphere radius is fixed as $r_s = 0.38a$. The other parameters are the same as in Fig. 1.

where θ_k is the argument of the long axis of the polarization ellipse, and the integration contour orbits the vortex core.

Around the BIC2 in Fig. 1, we have similar behavior in the ring-like quasi-BICs pinned with multiple true BICs on the mirror axis. However, a slightly different behavior than in Fig. 6 is observed. Figure 8 shows the evolution of the multiple BICs through the critical radius of BIC2. Across the critical radius, the ring-like BIC moves from the upper band to the lower band. At $r_s = 0.452a (> r_c)$, there are extra BICs on the ΓM axes other than the ring-like BICs of the accidental




FIG. 5. Real band structure originating from the E_1 mode at the Γ point near the accidental BIC of BIC1 in Fig. 1. The sphere radius is $0.38a$. The band structure with $|ka/2\pi| \leq 0.15$ is plotted. The other parameters are the same as in Fig. 1.




FIG. 6. Q -value map of the two bands originating from the E_1 mode at the Γ point near the accidental BIC of BIC1 in Fig. 1. The (a) upper and (c) lower bands (in the real band structure) at $r_s = 0.38a (< r_c)$. The real band structure at this parameter is shown in Fig. 5. The (b) upper and (d) lower bands at $r_s = 0.388a (> r_c)$. Insets show the close-up views overlaid by the polarization ellipse map of the photonic band with off- Γ BICs. The vortex charge is denoted by q .

at- Γ BIC origin. The latter ring is deformed from a circular shape to a hexagonal shape by a substantial perturbation. If we further increase the radius, the extra BICs merge with the true BICs on ΓM in the ring, and the ring is further deformed.

The BIC3 in Fig. 1 is found at $r_s = 0.495a$, near the close-packing condition of the triangular lattice. Within the photonic KKR method employed in this paper, there is not




FIG. 7. Schematic illustration of the vortex charge diagram in the ring. The charge distribution is common between the two rings of Fig. 6.




FIG. 8. Evolution of multiple off- Γ BICs around the critical radius $r_c \simeq 0.45a$ of BIC2 in Fig. 1. The upper two rows show the Q value map of the upper and lower bands originating from the E_1 mode at the Γ point. The lower row shows the vortex charge diagram of the off- Γ BICs. The inset shows the real band structure around the critical radius.

enough parameter space to enlarge the sphere radius beyond the close-packing condition. We do not consider this BIC here.

The BIC4 has a relatively high resonance frequency, giving rise to a more complex multiple off- Γ BIC distribution around the critical radius. Figure 9 shows the evolution of the multiple off- Γ BICs through the critical radius of BIC4. Now the ring is formed for the lower band at $r_s = 0.45a (> r_c)$. The upper band exhibits additional BICs on ΓM . This ring pinned with true BICs on ΓM and ΓK , and the additional BICs shrink to the Γ point as $r_s \rightarrow r_c$. Then, at $r_s = 0.44a (< r_c)$, the ring disappears in the lower band, and discrete BICs are generated from the Γ point of the upper band. The total vorticity is conserved across the critical radius.

The BIC5 also have a high resonance frequency, resulting in a complex behavior of multiple off- Γ BICs affected by a nearby band of a higher-order vortex charge at the Γ point. We do not consider this BIC here.

III. SQUARE LATTICE

A similar design of the accidental at- Γ BIC and subsequent multiple off- Γ BIC generation is available for the square-lattice systems with the C_{4v} point group. A fine-tuning of a system parameter results in the accidental at- Γ BIC of a doubly degenerate E mode at the Γ point. Above and below the critical parameter, multiple off- Γ BICs emerge in the two bands originating from the E mode at Γ . The eigenmodes of A_1 , A_2 , B_1 , and B_2 representations of C_{4v} at the Γ point are symmetry-protected BICs provided that there are no open diffraction channels other than the specular one.




FIG. 9. Evolution of multiple off- Γ BICs around the critical radius $r_c \simeq 0.445a$ of BIC4 in Fig. 1. The upper two rows show the Q value map of the upper and lower bands originating from the E_1 mode at the Γ point. The lower row shows the vortex charge diagram of the off- Γ BICs. The inset shows the real band structure around the critical radius.

Figure 10 shows the design of the accidental at- Γ BIC in the square lattice of identical spheres. By changing the sphere radius, we can find three critical radii of the accidental at- Γ BICs at $r_s \simeq 0.415a$, $0.485a$, and $0.49a$ for the z -even, z -odd, and z -even parities, respectively.

Considering the parameter regions around the critical radius r_c , we have multiple off- Γ BICs. Figure 11 shows the real and imaginary photonic band structures of the z -even parity around the critical radius of BIC1 in Fig. 10. As in Fig. 3, the real band structure does not change so much, while the imaginary band structure changes remarkably by changing the sphere radius. Concerning the ΓM direction, a similar trend as in Fig. 3 is observed. Namely, above and below the critical radius, the band that exhibits the off- Γ BIC at a nonzero momentum is interchanged. Below the critical radius [Figs. 11(a) and 11(b)], the odd-parity band exhibits the BIC while above the critical radius [Figs. 11(e) and 11(f)], the even-parity band exhibits the BIC in ΓM . The suppression of the decay rate at about the critical radius is weaker in ΓM but is stronger in ΓX . Such an anisotropy is manifest also in the real band structure. As for the ΓX direction, both the even- and odd-parity bands exhibit the off- Γ BICs below the critical radius, while the BICs are absent above the critical radius.

By scanning all the directions around the Γ point, the real band structure is shown in Fig. 12. The anisotropy is evident in the band structure. The upper band is S -like, and the lower band is P -like.




FIG. 10. Resonance (a) angular frequencies ω_0 and (b) widths γ_0 of the E modes at the Γ point in the square-lattice monolayer of dielectric spheres with the C_{4v} point group symmetry, as a function of sphere radius r_s . The modes are classified according to the parity in the z direction, assuming the centers of the spheres are on the $z = 0$ plane. The dielectric constant of the spheres is 12, and that of the background material is 1. The lattice constant is denoted by a . Accidental at- Γ BICs occur around the modes indicated by the solid circles. The inset in (a) stands for the geometry of the monolayer.

Figure 13 shows the Q -value maps of the two photonic bands originating from the E mode at the off-critical radii of Fig. 11. Now, the ring-like BICs found in the triangular-lattice system are absent. A similar pattern as in Fig. 13(a) was observed in Ref. [16]. At $r_s = 0.41a$, the total vorticity vanishes in the upper band. However, the lower band sustains the additional BICs on the ΓX axes. The net vorticity of the upper and lower bands is thus nonzero. All the BICs are discrete and move toward the Γ point as $r_s \rightarrow r_c$. At $r_s = r_c$, the BICs are collapsed there. Across the critical coupling, multiple off- Γ BICs are again generated now on the ΓM axes. The total vorticity is conserved across the critical coupling.

As for the BIC2 of Fig. 10, the evolution of the multiple off- Γ BICs across the critical radius is shown in Fig. 14. Since the two bands intersect in ΓM , the Q -value becomes singular there. At $r_s = 0.48a$, the upper band exhibits the eight off- Γ BICs. They are elongated, and their polarizations are perpendicular and parallel to the ΓX and ΓM axes, respectively. The lower band holds four off- Γ BICs on the ΓM axes. All the BICs moves toward the Γ point as $r_s \rightarrow r_c$. After the collapse, the four off- Γ BICs are generated as in BIC1.

The BIC3 exists in a close vicinity of another E_1 mode, resulting in mixed four bands. We do not consider this BIC here.

IV. $k \cdot p$ PERTURBATION THEORY

Let us consider the phenomena obtained in Secs. II and III in the $k \cdot p$ perturbation. The eigenmodes in the monolayer are




FIG. 11. Real and imaginary photonic band structure of the z -even parity, originating from the E mode around the critical radius r_c of BIC1 in Fig. 10. The momentum is scanned in the ΓX and ΓM directions. The photonic bands are further classified according to the parity in these directions. The sphere radius is (a,b) $0.41a$, (c,d) $0.415a (\approx r_c)$, (e,f) and $0.42a$. The other parameters are the same as in Fig. 10. The inset in (b) represents the first Brillouin zone of the square lattice.

determined by

$$\nabla \times \left(\frac{1}{\varepsilon(\mathbf{x})} \nabla \times \mathbf{H}(\mathbf{x}) \right) = \frac{\omega^2}{c^2} \mathbf{H}(\mathbf{x}). \quad (8)$$

Here \mathbf{H} is the time-harmonic magnetic field with angular frequency ω , and $\varepsilon(\mathbf{x})$ is the dielectric function that is periodic in the in-plane coordinate \mathbf{x}_{\parallel} . The inverse dielectric function can be expanded as

$$\frac{1}{\varepsilon(\mathbf{x})} = \sum_g e^{ig \cdot \mathbf{x}_{\parallel}} \eta_g(z). \quad (9)$$

In addition, the radiation field is also expanded by the plane waves via Bloch's theorem as

$$\mathbf{H}(\mathbf{x}) = \sum_g e^{i(k+g) \cdot \mathbf{x}_{\parallel}} \mathbf{h}_g(z). \quad (10)$$

Accordingly, the equation to be solved becomes

$$\begin{aligned} & \sum_{g'} (i(\mathbf{k} + \mathbf{g}) + \hat{z}\partial_z) \{ \eta_{g-g'}(z) [i(\mathbf{k} + \mathbf{g}') + \hat{z}\partial_z] \times \mathbf{h}_{g'}(z) \} \\ & = \frac{\omega^2}{c^2} \mathbf{h}_g(z), \end{aligned} \quad (11)$$




FIG. 12. Real band structure originating from the E mode at the Γ point near the accidental BIC of BIC1 in Fig. 10. The sphere radius is $0.41a$. The z parity is even. The band structure with $|k_a/2\pi| \leq 0.2$ is plotted. The other parameters are the same as in Fig. 10.

which is symbolically expressed as

$$\mathcal{H}|\psi\rangle = \mathcal{E}|\psi\rangle, \quad \mathcal{E} = \frac{\omega^2}{c^2}. \quad (12)$$

Suppose we have a nearly accidental BIC of a doubly degenerate eigenmode at the Γ point. We perform the $\mathbf{k} \cdot \mathbf{p}$ perturbation starting from these states as the zeroth-order




FIG. 13. Q -value map of the two bands originating from the E mode at the Γ point near accidental BIC of BIC1 in Fig. 10. The (a) upper and (c) lower bands at $r_s = 0.41a (< r_c)$. The (b) upper and (d) lower bands at $r_s = 0.42a (> r_c)$. The insets show the close-up view of the Q -value map overlaid with the polarization ellipse map.




FIG. 14. Evolution of multiple off- Γ BICs around the critical radius $r_c \simeq 0.485a$ of BIC2 in Fig. 10. The upper two rows show the Q -value map of the upper and lower bands originating from the E mode at the Γ point. The lower row shows the vortex charge diagram of the off- Γ BICs. The inset shows the real band structure (at $r_s = 0.495a$).

ones. The effective Hamiltonian to be diagonalized is given by [29]

$$\begin{aligned} \mathcal{H}_{ab}^{\text{eff}} &= \langle \phi_{0a} | \mathcal{H}^{(2)} | \phi_{0b} \rangle \\ &+ \sum_{n \neq 0} \frac{1}{\epsilon_0 - \epsilon_n} \langle \phi_{0a} | \mathcal{H}^{(1)} | \phi_n \rangle \langle \phi_n | \mathcal{H}^{(1)} | \phi_{0b} \rangle, \end{aligned} \quad (13)$$

$$\mathcal{H}^{(0)} | \phi_{0a} \rangle = \epsilon_0 | \phi_{0a} \rangle, \quad \mathcal{H}^{(0)} | \phi_n \rangle = \epsilon_n | \phi_n \rangle, \quad (14)$$

where $\mathcal{H}^{(i)}$ is the operator in the left-hand side of Eq. (11) expanded by Bloch momentum \mathbf{k} ($i = 0, 1$, and 2 imply $\mathbf{k} = 0$, \mathbf{k} linear, and \mathbf{k} quadratic, respectively). The state $| \phi_{0a} \rangle$ ($a = 1, 2$) represents the doubly degenerate nearly accidental BIC mode at Γ . Here we assume the degeneracy is not lifted in the first-order perturbation. That is,

$$\langle \phi_{0a} | \mathcal{H}^{(1)} | \phi_{0b} \rangle = 0. \quad (15)$$

This property is verified explicitly by the C_{6v} or C_{4v} symmetry. More generally, the matrix element $\langle \phi_m | \mathcal{H}^{(1)} | \phi_n \rangle$ is nonzero only if (ϕ_m, ϕ_n) is attributed to (E_1, E_2) , (E_1, A_1) , (E_1, A_2) , (E_2, B_1) , (E_2, B_2) , and vice versa for C_{6v} , and to (E, A_1) , (E, A_2) , (E, B_1) , (E, B_2) , and vice versa for C_{4v} (see Appendix A).

The sum in the second term of Eq. (13) generally includes the continuous radiation modes and discrete quasiguided

modes other than the mode concerned, as the intermediate states. The continuous radiation modes give the decay rate to the original eigenmode via Fermi's golden rule [30,31]. However, if we start from the mode of the nearly accidental at- Γ BIC, the decay rate is strongly suppressed.

The continuous radiation modes belong to the same irreducible representation as the original doubly degenerate mode below the diffraction threshold. Therefore, the matrix elements vanish between the nearly accidental at- Γ BIC and radiation modes. That is why the decay rate is strongly suppressed.

In the C_{6v} case, the effective Hamiltonian is expressed as

$$\mathcal{H}^{\text{eff}} = \alpha \mathbf{k}^2 1 + \beta [(k_x^2 - k_y^2)\sigma_3 + 2k_x k_y \sigma_1], \quad (16)$$

$$\alpha = \alpha_{E_1 E_1}^{(2)} + \sum_{n \in E_2} \frac{|\alpha_{E_1 E_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{E_1 A_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{E_1 A_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}, \quad (17)$$

$$\beta = \beta_{E_1 E_1}^{(2)} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{E_1 A_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} - \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{E_1 A_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}, \quad (18)$$

where $\sigma_i (i = 1, 2, 3)$ is the Pauli matrix and parameters $\alpha_{E_1 E_1}^{(2)}$ and so on are given in Appendix A. The resulting effective parameters α and β are complex with $\text{Im}[\alpha] \geq 0$. Their imaginary parts come solely from the one in the eigenfrequency of the E_1 mode, namely, $\epsilon_0 = (\omega_0 - i\gamma_0)^2/c^2$, provided that the intermediate modes are lossless ($\text{Im}[\epsilon_n] = 0$). Therefore, if we start with the E_1 mode of the perfect accidental at- Γ BIC ($\gamma_0 = 0$), $\text{Im}[\alpha] = \text{Im}[\beta] = 0$ so that we still have a vanishing decay rate even at finite \mathbf{k} , within the second-order $\mathbf{k} \cdot \mathbf{p}$ perturbation. This is the case of the extreme suppression of the decay rate around the Γ point of the accidental (next-to-super) BIC. Since the third-order terms vanish by the inversion symmetry, the imaginary frequencies of the two bands behave k^4 as pointed out in Ref. [16].

By diagonalizing the effective Hamiltonian, the eigenfrequencies of the two bands that stem from the E_1 mode at Γ are given by

$$\frac{\omega^2}{c^2} = \frac{(\omega_0 - i\gamma_0)^2}{c^2} + (\alpha \pm \beta) \mathbf{k}^2, \quad (19)$$

$$\mathbf{H}_k^+(\mathbf{x}) \propto k_x \mathbf{H}_{E_1}^{(01)}(\mathbf{x}) + k_y \mathbf{H}_{E_1}^{(02)}(\mathbf{x}), \quad (20)$$

$$\mathbf{H}_k^-(\mathbf{x}) \propto -k_y \mathbf{H}_{E_1}^{(01)}(\mathbf{x}) + k_x \mathbf{H}_{E_1}^{(02)}(\mathbf{x}), \quad (21)$$

where $(\mathbf{H}_{E_1}^{(01)}, \mathbf{H}_{E_1}^{(02)})$ forms the E_1 representation at Γ . If $\gamma_0 \ll \omega_0$, the decay rate γ_k is evaluated as

$$\gamma_k \simeq \gamma_0 - \frac{c^2}{2\omega_0} \text{Im}[\alpha \pm \beta] \mathbf{k}^2. \quad (22)$$

The radiation field of $\mathbf{H}_k^{+(-)}$ is P -like (S -like). The property of $\text{Im}[\alpha] > 0$ promises a monotonic decreasing of the decay rate γ_k with increasing $|\mathbf{k}|$ for at least one band because either $\text{Im}[\alpha + \beta]$ or $\text{Im}[\alpha - \beta]$ is promised to be positive. The behavior of the other band depends on the relative magnitude of $\text{Im}[\alpha]$ and $\text{Im}[\beta]$. Across $\text{Im}[\beta] = 0$, the band with fast




FIG. 15. Comparison in the complex band structures of Figs. 3(a) and 3(b) between the KKR result and the $\mathbf{k} \cdot \mathbf{p}$ perturbation of Eq. (19) with a numerical fitting. Fitted values are $\alpha = 0.111 + i9.56 \times 10^{-5}$ and $\beta = -0.132 + i1.20 \times 10^{-3}$.

decreasing decay rate with $|\mathbf{k}|$ is interchanged. This condition is satisfied at the critical coupling. There, $\gamma_0 = 0$ and thus $\text{Im}[\beta] = 0$. Moreover, since the dispersion relation is isotropic, we can have ring-like BICs at the minimum of γ_k . These results explain Fig. 6 reasonably well.

More direct and quantitative comparison between the KKR calculation and the $\mathbf{k} \cdot \mathbf{p}$ perturbation needs special care. The perturbation requires a regularization of the nonnormalizable (due to the radiation loss) quasiguided modes of the E_1 representation. Therefore, the parameters $\alpha_{E_1 A_1}^{(1)}$ and so on become regularization dependent. To avoid such an artifact, we consider the fitting of the real and imaginary band structures around the Γ point by complex parameters of α and β . The result is shown in Fig. 15. Since the $\mathbf{k} \cdot \mathbf{p}$ perturbation is valid only for small $|\mathbf{k}|$, it can describe the trend there reasonably well. In particular, the fitting of the real band structure is perfect. However, the fitting of the imaginary band structure is not good enough. In the second-order $\mathbf{k} \cdot \mathbf{p}$ perturbation, the lower bound of γ_k ($\gamma_k \geq 0$) is ignored, so that the fitted γ_k can go down below the lower bound. This is a drawback of the present approach.

As for Figs. 8 and 9, the spatial anisotropy is very strong, so that the $\mathbf{k} \cdot \mathbf{p}$ perturbation that predicts the isotropic dispersion is not so efficient.

In the C_{4v} case, the effective Hamiltonian becomes

$$\mathcal{H}^{\text{eff}} = \alpha \mathbf{k}^2 1 + \beta_1 (k_x^2 - k_y^2) \sigma_3 + 2\beta_2 k_x k_y \sigma_1, \quad (23)$$

$$\alpha = \alpha_{E E}^{(2)} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{E A_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in B_1} \frac{|\alpha_{E B_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{E A_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in B_2} \frac{|\alpha_{E B_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}, \quad (24)$$

$$\beta_1 = \beta_{1:EE}^{(2)} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{EA_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in B_1} \frac{|\alpha_{EB_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} - \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{EA_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n} - \frac{1}{2} \sum_{n \in B_2} \frac{|\alpha_{EB_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}, \quad (25)$$

$$\beta_2 = \beta_{2:EE}^{(2)} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{EA_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} - \frac{1}{2} \sum_{n \in B_1} \frac{|\alpha_{EB_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} - \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{EA_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in B_2} \frac{|\alpha_{EB_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}. \quad (26)$$

By diagonalizing the effective Hamiltonian, the eigenfrequency becomes

$$\frac{\omega^2}{c^2} = \frac{(\omega_0 - i\gamma_0)^2}{c^2} + \alpha k^2 \pm \sqrt{\beta_1^2 (k_x^2 - k_y^2)^2 + 4\beta_2^2 k_x^2 k_y^2}. \quad (27)$$

Again, $\text{Im}[\alpha] > 0$, promising a monotonic decreasing of the imaginary part in Eq. (27) with increasing $|k|$ for at least one band branch. Now, Eq. (27) is not isotropic. Thus, the ring-like high- Q channel is not formed. This trend is entirely consistent with the numerical results obtained in Sec. III.

On the ΓX axis ($k_y = 0$), the two eigenstates become

$$\frac{\omega^2}{c^2} = \frac{(\omega_0 - i\gamma_0)^2}{c^2} + (\alpha \pm \beta_1) k_x^2, \quad (28)$$

$$\mathbf{H}_k^+(\mathbf{x}) = \mathbf{H}_E^{(01)}(\mathbf{x}), \quad \mathbf{H}_k^-(\mathbf{x}) = \mathbf{H}_E^{(02)}(\mathbf{x}), \quad (29)$$

where $(\mathbf{H}_E^{(01)}, \mathbf{H}_E^{(02)})$ forms the E representation at Γ . Since the E representation behaves as (x, y) , \mathbf{H}_k^+ is P polarized and \mathbf{H}_k^- is S polarized. One of the eigenmodes is promised to have a decreasing decay rate with $|k_x|$, and the two eigenmodes are interchanged at $\text{Im}[\beta_1] = 0$ via the critical coupling condition $\text{Im}[\gamma_0] = 0$.

On the ΓM axis ($k_x = k_y$), the eigenstates become

$$\frac{\omega^2}{c^2} = \frac{(\omega_0 - i\gamma_0)^2}{c^2} + 2(\alpha \pm \beta_2) k_x^2, \quad (30)$$

$$\mathbf{H}_k^\pm(\mathbf{x}) \propto \mathbf{H}_E^{(01)}(\mathbf{x}) \pm \mathbf{H}_E^{(02)}(\mathbf{x}). \quad (31)$$

The eigenmode of superscript “+” is P polarized, and “-” is S polarized. One of the two eigenmodes is promised to have a decreasing decay rate with $|k_x|$, and the two eigenmodes are interchanged at $\text{Im}[\beta_2] = 0$ via the critical coupling condition $\text{Im}[\gamma_0] = 0$. Again, these features are consistent with Figs. 13 and 14.

The symmetry-protected BICs are also available for C_{3v} point group systems. There, the eigenmodes of the A_1 and A_2 representations at the Γ point are symmetry-protected, provided there are no open diffraction channels other than the specular one. In this case, it is possible to have accidental at- Γ BICs for the eigenmodes of the E representation by tuning system parameters.

The effective Hamiltonian around the nearly accidental at- Γ BIC mode is given by

$$\mathcal{H}^{\text{eff}} = \alpha k^2 1 + \beta [(k_x^2 - k_y^2)\sigma_3 + 2k_x k_y \sigma_1], \quad (32)$$

$$\alpha = \alpha_{EE}^{(2)} + \sum'_{n \in E} \frac{|\alpha_{EE}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{EA_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} + \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{EA_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}, \quad (33)$$

$$\beta = \beta_{EE}^{(2)} + \frac{1}{2} \sum_{n \in A_1} \frac{|\alpha_{EA_1}^{(1)}|^2}{\epsilon_0 - \epsilon_n} - \frac{1}{2} \sum_{n \in A_2} \frac{|\alpha_{EA_2}^{(1)}|^2}{\epsilon_0 - \epsilon_n}, \quad (34)$$

where the prime in the second term of Eq. (33) means the unperturbed E mode is excluded in the sum. In this case, a significant contribution to the decay rate emerges from the continuous radiation modes of the E representation via Fermi's golden rule. Therefore, suppressing of the decay rate at nonzero $|k|$ near the accidental BIC (at Γ) does not occur. Consequently, the multiple off- Γ BICs are absent in the $\mathbf{k} \cdot \mathbf{p}$ perturbation theory.

In the above arguments, we do not rely on the monolayer of spheres presented in Secs. II and III, but on the spatial symmetry of C_{6v} , C_{4v} , and C_{3v} . Accordingly, the multiple BIC generation with the present scenario is available in other photonic membranes with C_{6v} or C_{4v} . We can show PhC slabs with the triangular or square lattice of circular air holes exhibit multiple BIC generations. The ring-like high- Q channel is also formed in the C_{6v} case.

V. SUMMARY

In summary, we presented a detailed theoretical analysis of the multiple BIC generation in the monolayers of spheres with C_{6v} or C_{4v} point group. A tuning of system parameters results in an accidental at Γ BIC of doubly degenerate eigenmodes at the Γ point. This BIC, obtained at a critical parameter, is the next-to-super-BIC with extremely suppressed decay rates around the Γ point. Off-critical parameters yield multiple off- Γ BICs that are generated from the next-to-super-BIC at the Γ point.

In the C_{6v} point group system, a ring-like high- Q channel pinned with multiple off- Γ BICs on the mirror axes can be formed in the two bands originating from the doubly degenerate E_1 mode at the Γ point. The spatial anisotropy destroys the ring in the C_{4v} point group system, but multiple BICs are certainly formed. Across the critical parameters, these BICs move from the upper band to the lower band or vice versa, conserving the total vorticity of the BICs.

We also show that these phenomena are owing to the spatial symmetry of C_{6v} or C_{4v} and the $\mathbf{k} \cdot \mathbf{p}$ perturbation explains the phenomena reasonably well. In addition, the C_{3v} system does not support the multiple BIC generation through a similar design.

ACKNOWLEDGMENT

This work was supported by JSPS KAKENHI Grant No. 22K03488.

APPENDIX A: MATRIX ELEMENTS IN THE $\mathbf{k} \cdot \mathbf{p}$ PERTURBATION

We summarize various matrix elements relevant to the effective Hamiltonian. The symmetry relation under

point-group operations is crucial. It is given by

$$\mathcal{H}_{R_1 R_2}^{(i)}(\mathbf{k}) = D_{R_1}^\dagger(A) \mathcal{H}_{R_1 R_2}^{(i)}(A\mathbf{k}) D_{R_2}(A), \quad (\text{A1})$$

where the matrix element is the one between the modes of irreducible representations R_1 and R_2 , and $D_R(A)$ is the representation matrix of group element A in the irreducible representation R . Equation (A1) constrains the possible form of the matrix elements as follows.

In the C_{6v} case, The nonzero matrix elements in the first-order $\mathbf{k} \cdot \mathbf{p}$ perturbation are given by

$$\mathcal{H}_{E_1 E_2}^{(1)}(\mathbf{k}) = \alpha_{E_1 E_2}^{(1)}(k_x \sigma_1 + k_y \sigma_3), \quad (\text{A2})$$

$$\mathcal{H}_{E_1 A_1}^{(1)}(\mathbf{k}) = \alpha_{E_1 A_1}^{(1)} \begin{pmatrix} k_x \\ k_y \end{pmatrix}, \quad (\text{A3})$$

$$\mathcal{H}_{E_1 A_2}^{(1)}(\mathbf{k}) = \alpha_{E_1 A_2}^{(1)} \begin{pmatrix} -k_y \\ k_x \end{pmatrix}, \quad (\text{A4})$$

$$\mathcal{H}_{E_2 B_1}^{(1)}(\mathbf{k}) = \alpha_{E_2 B_1}^{(1)} \begin{pmatrix} -k_y \\ k_x \end{pmatrix}, \quad (\text{A5})$$

$$\mathcal{H}_{E_2 B_2}^{(1)}(\mathbf{k}) = \alpha_{E_2 B_2}^{(1)} \begin{pmatrix} k_x \\ k_y \end{pmatrix}, \quad (\text{A6})$$

where $\alpha_{E_1 E_2}^{(1)}$ and so on are complex. The vanishing matrix elements between E_1 and $B_1(B_2)$ are crucial for the k^4 scaling of the decay rate in the symmetry-protected BICs of the B_1 and B_2 representations, which have a higher-order vortex charge of $q = -2$.

In the second order, the relevant matrix element becomes

$$\mathcal{H}_{E_1 E_1}^{(2)}(\mathbf{k}) = \alpha_{E_1 E_1}^{(2)} \mathbf{k}^2 + \beta_{E_1 E_1}^{(2)} ((k_x^2 - k_y^2) \sigma_3 + 2k_x k_y \sigma_1), \quad (\text{A7})$$

where $\alpha_{E_1 E_1}^{(2)}$ and $\beta_{E_1 E_1}^{(2)}$ are real.

In the C_{4v} case, the nonzero matrix elements in the first-order $\mathbf{k} \cdot \mathbf{p}$ perturbation are given by

$$\mathcal{H}_{EA_1}^{(1)}(\mathbf{k}) = \alpha_{EA_1}^{(1)} \begin{pmatrix} k_x \\ k_y \end{pmatrix}, \quad (\text{A8})$$

$$\mathcal{H}_{EA_2}^{(1)}(\mathbf{k}) = \alpha_{EA_2}^{(1)} \begin{pmatrix} -k_y \\ k_x \end{pmatrix}, \quad (\text{A9})$$

$$\mathcal{H}_{EB_1}^{(1)}(\mathbf{k}) = \alpha_{EB_1}^{(1)} \begin{pmatrix} k_x \\ -k_y \end{pmatrix}, \quad (\text{A10})$$

$$\mathcal{H}_{EB_2}^{(1)}(\mathbf{k}) = \alpha_{EB_2}^{(1)} \begin{pmatrix} k_y \\ k_x \end{pmatrix}, \quad (\text{A11})$$

where $\alpha_{EA_1}^{(1)}$ and so on are complex. In the second order, the relevant matrix element becomes

$$\mathcal{H}_{EE}^{(2)}(\mathbf{k}) = \alpha_{EE}^{(2)} \mathbf{k}^2 + \beta_{1:EE}^{(2)} (k_x^2 - k_y^2) \sigma_3 + \beta_{2:EE}^{(2)} 2k_x k_y \sigma_1, \quad (\text{A12})$$

where $\alpha_{E_1 E_1}^{(2)}$, $\beta_{1:E_1 E_1}^{(2)}$, $\beta_{2:E_1 E_1}^{(2)}$ are real.

In the C_{3v} case, the nonzero matrix elements in the first-order $\mathbf{k} \cdot \mathbf{p}$ perturbation are given by

$$\mathcal{H}_{EE}^{(1)}(\mathbf{k}) = \alpha_{EE}^{(1)}(k_x \sigma_1 + k_y \sigma_3), \quad (\text{A13})$$

$$\mathcal{H}_{EA_1}^{(1)}(\mathbf{k}) = \alpha_{EA_1}^{(1)} \begin{pmatrix} k_x \\ k_y \end{pmatrix}, \quad (\text{A14})$$

$$\mathcal{H}_{EA_2}^{(1)}(\mathbf{k}) = \alpha_{EA_2}^{(1)} \begin{pmatrix} -k_y \\ k_x \end{pmatrix}, \quad (\text{A15})$$

where $\alpha_{EE}^{(1)}$ and so on are complex. The matrix element of Eq. (A13) vanishes by the time-reversal symmetry if the two E modes that form the matrix element are identical. In the second order, the relevant matrix element becomes

$$\mathcal{H}_{EE}^{(2)}(\mathbf{k}) = \alpha_{EE}^{(2)} \mathbf{k}^2 + \beta_{EE}^{(2)} [(k_x^2 - k_y^2) \sigma_3 + 2k_x k_y \sigma_1], \quad (\text{A16})$$

where $\alpha_{EE}^{(2)}$ and $\beta_{EE}^{(2)}$ are real.

The parameters $\alpha_{R_1 R_2}^{(1,2)}$ and $\beta_{R_1 R_2}^{(2)}$ are not determined solely from the symmetry relation [Eq. (A1)].

APPENDIX B: ELECTRIC FIELDS OF QUASIGUIDED MODES

We employ the following method for evaluating the polarization ellipse map. At a true BIC point, the kernel of a secular matrix gives the eigenmode and its determinant vanishes on the real frequency axis.

In the monolayer of spheres, the secular matrix \mathcal{S} is given by [23,32]

$$\mathcal{S}_{(L\beta)(L'\beta')} = \delta_{LL'} \delta_{\beta\beta'} - t_l^\beta G_{LL'}^{\beta\beta'}, \quad (\text{B1})$$

$$G_{LL'}^{\beta\beta'} = \frac{1}{l(l+1)} \sum_{L_1 L_2} [(\mathbf{P}^\beta)^\dagger]_{LL_1} G_{L_1 L_2} [\mathbf{P}^{\beta'}]_{L_2 L'}, \quad (\text{B2})$$

$$G_{LL'} = 4\pi \sum_{L''} i^{l-l''} \langle L|L''|L' \rangle S_{L''}, \quad (\text{B3})$$

$$S_L = \sum_{X \neq 0} h_l^{(1)}(q_0 |X|) Y_L^*(\hat{X}) e^{i\mathbf{k} \cdot X}, \quad (\text{B4})$$

where $L = (l, m)$ ($|m| \leq l$) is the angular momentum index, $\beta (= M, N)$ is the index to classify two transverse vector spherical waves, t_l^β is the so-called t matrix of the sphere, \mathbf{P}^β is the transformation matrix from scalar spherical waves to vector spherical waves, $\langle L|L''|L' \rangle$ is the Clebsh-Gordan coefficient, $h_l^{(1)}$ is the spherical hankel function of the first kind, Y_L is the spherical harmonics, and X is the 2D real-lattice vector. The structure constant S_L is efficiently calculated via the Ewald method [33]. Through $\psi_L^\beta = \text{Ker}[\mathcal{S}]$, the electric field outside the monolayer is expressed as

$$\mathbf{E}(\mathbf{x}) = \sum_{LX} h_l^{(1)}(q_0 |\mathbf{x} - X|) Y_L(\widehat{\mathbf{x} - X}) \mathbf{V}_L e^{i\mathbf{k} \cdot X}, \quad (\text{B5})$$

$$\mathbf{V}_L = \sum_{L'\beta} \mathbf{P}_{LL'}^\beta \psi_{L'}^\beta. \quad (\text{B6})$$

Its plane-wave-expansion coefficient $t_{\sigma g}^\pm$ ($\sigma = P, S$) becomes

$$t_{\sigma g}^\pm = \frac{2\pi}{q_0 \Gamma_g A_{UC}} \sum_L (-i)^l Y_L(\hat{\mathbf{k}}_g^\pm) \sigma_g^\pm \cdot \mathbf{V}_L, \quad (\text{B7})$$

where A_{UC} is the area of the unit cell.

Off the BIC point, the determinant is nonzero on the real frequency axis but can be zero in the complex frequency plane. This zero should be the resonance pole of the S matrix, namely, $\Omega_k = \omega_k - i\gamma_k$. We assume that the right eigenstate

of \mathcal{S} with the least absolute eigenvalue can be approximated as the kernel on the real frequency axis near the zero. Then, we can obtain the plane-wave coefficients $t_{\sigma g}^{\pm}(\omega)$ through Eq. (B7). We further assume that it can be Taylor expanded around $\omega = \Omega_k$, namely,

$$t_{\sigma g}^{\pm}(\omega) \simeq t_{\sigma g}^{\pm}(\Omega_k) + (\omega - \Omega_k)C_{\sigma g}^{\pm} \quad (\sigma = P, S). \quad (\text{B8})$$

Then, from the two points ω_a and ω_b on the real frequency axis, we can estimate $t_{\sigma g}^{\pm}(\Omega_k)$ as

$$t_{\sigma g}^{\pm}(\Omega_k) = \frac{(\omega_a - \Omega_k)t_{\sigma g}^{\pm}(\omega_b) - (\omega_b - \Omega_k)t_{\sigma g}^{\pm}(\omega_a)}{\omega_a - \omega_b}. \quad (\text{B9})$$

We confirm that the resulting $t_{\sigma 0}^{\pm}(\Omega_k)$ vanishes at the BIC points.

-
- [1] C. W. Hsu, B. Zhen, A. D. Stone, J. D. Joannopoulos, M. Soljacic, and M. Soljačić, *Nat. Rev. Mater.* **1**, 16048 (2016).
- [2] A. Kodigala, T. Lepetit, Q. Gu, B. Bahari, Y. Fainman, and B. Kanté, *Nature (London)* **541**, 196 (2017).
- [3] S. Romano, G. Zito, S. Torino, G. Calafiore, E. Penzo, G. Coppola, S. Cabrini, I. Rendina, and V. Mocella, *Photonics Res.* **6**, 726 (2018).
- [4] K. Koshelev, S. Kruk, E. Melik-Gaykazyan, J. H. Choi, A. Bogdanov, H. G. Park, and Y. Kivshar, *Science* **367**, 288 (2020).
- [5] H. Miyazaki and K. Ohtaka, *Phys. Rev. B* **58**, 6920 (1998).
- [6] P. Paddon and J. F. Young, *Phys. Rev. B* **61**, 2090 (2000).
- [7] T. Ochiai and K. Sakoda, *Phys. Rev. B* **63**, 125107 (2001).
- [8] S. H. Fan and J. D. Joannopoulos, *Phys. Rev. B* **65**, 235112 (2002).
- [9] C. W. Hsu, B. Zhen, J. Lee, S. L. Chua, S. G. Johnson, J. D. Joannopoulos, and M. Soljačić, *Nature (London)* **499**, 188 (2013).
- [10] Q. Jiang, P. Hu, J. Wang, D. Han, and J. Zi, *Phys. Rev. Lett.* **131**, 013801 (2023).
- [11] B. Zhen, C. W. Hsu, L. Lu, A. D. Stone, and M. Soljačić, *Phys. Rev. Lett.* **113**, 257401 (2014).
- [12] J. Jin, X. Yin, L. Ni, M. Soljačić, B. Zhen, and C. Peng, *Nature (London)* **574**, 501 (2019).
- [13] M.-S. S. Hwang, H.-C. C. Lee, K.-H. H. Kim, K.-Y. Y. Jeong, S.-H. H. Kwon, K. Koshelev, Y. Kivshar, and H.-G. G. Park, *Nat. Commun.* **12**, 4135 (2021).
- [14] M. Kang, L. Mao, S. Zhang, M. Xiao, H. Xu, and C. T. Chan, *Light Sci. Appl.* **11**, 228 (2022).
- [15] M. Kang, S. Zhang, M. Xiao, and H. Xu, *Phys. Rev. Lett.* **126**, 117402 (2021).
- [16] A. S. Kostyukov, V. S. Gerasimov, A. E. Ershov, and E. N. Bulgakov, *Phys. Rev. B* **105**, 075404 (2022).
- [17] Z. Sadrieva, K. Frizyuk, M. Petrov, Y. Kivshar, and A. Bogdanov, *Phys. Rev. B* **100**, 115303 (2019).
- [18] S. Gladyshev, A. Shalev, K. Frizyuk, K. Ladutenko, and A. Bogdanov, *Phys. Rev. B* **105**, L241301 (2022).
- [19] T. Yoda and M. Notomi, *Phys. Rev. Lett.* **125**, 053902 (2020).
- [20] C. F. Doiron, I. Brener, and A. Cerjan, *Nat. Commun.* **13**, 7534 (2022).
- [21] X. Wang, J. Wang, X. Zhao, L. Shi, and J. Zi, *ACS Photonics* **10**, 2316 (2023).
- [22] T. Inui, Y. Tanabe, and Y. Onodera, *Group Theory and its Applications in Physics* (Springer, New York, 1996).
- [23] K. Ohtaka, *J. Phys. C: Solid State Phys.* **13**, 667 (1980).
- [24] A. Modinos, *Physica A* **141**, 575 (1987).
- [25] N. Stefanou, V. Karathanos, and A. Modinos, *J. Phys.: Condens. Matter* **4**, 7389 (1992).
- [26] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics: Non-Relativistic Theory* (Pergamon, Oxford, 1977).
- [27] K. Ohtaka, J. I. Inoue, and S. Yamaguti, *Phys. Rev. B* **70**, 035109 (2004).
- [28] U. Fano, *Phys. Rev.* **124**, 1866 (1961).
- [29] T. Ochiai, *Phys. Rev. B* **86**, 075152 (2012).
- [30] T. Ochiai and K. Sakoda, *Phys. Rev. B* **64**, 045108 (2001).
- [31] L. C. Andreani and D. Gerace, *Phys. Rev. B* **73**, 235114 (2006).
- [32] K. Ohtaka, *Phys. Rev. B* **19**, 5057 (1979).
- [33] K. Kambe, *Z. Naturforsch., A: Phys. Sci.* **22**, 322 (1967).