
COMMENTS AND ADDENDA

The *Comments and Addenda* section is for short communications which are not of such urgency as to justify publication in *Physical Review Letters* and are not appropriate for regular *Articles*. It includes only the following types of communications: (1) comments on papers previously published in *The Physical Review* or *Physical Review Letters*; (2) addenda to papers previously published in *The Physical Review* or *Physical Review Letters*, in which the additional information can be presented without the need for writing a complete article. Manuscripts intended for this section may be accompanied by a brief abstract for information-retrieval purposes. Accepted manuscripts will follow the same publication schedule as articles in this journal, and galley proofs will be sent to authors.

Comment on the Characteristic Time of Spontaneous Decay in Jaynes's Semiclassical Radiation Theory

Darryl Leiter*

Physics Department, Boston College, Chestnut Hill, Massachusetts 02167

(Received 1 October 1969)

We examine the implication in the fact that, in a semiclassical theory of atomic structure proposed by Jaynes and Crisp, the initial state of an atom plays a critical role in determining its average lifetime in an excited state.

In a recent publication,¹ Jaynes and Crisp have studied the behavior of atoms within the framework of a semiclassical theory, which includes the effects upon the atom of the fields created by the atomic currents. They state that, in the absence of an applied field, an atom will spontaneously decay from an excited state with a characteristic time which is equal to the reciprocal of the Einstein A coefficient for the transition. The purpose of this note is to point out that statement is not entirely precise. To see this we note that the solution to the nonlinear density-matrix equation, for the diagonal matrix elements in a two-level decay, in a spontaneous transition with no applied fields can be written

$$\rho_{11}(t) = 1 / [\exp(-A_{21}(t - t_m)) + 1], \quad (1)$$

$$\rho_{22}(t) = [\exp(A_{21}(t - t_m)) + 1]^{-1}, \quad (2)$$

where $\rho_{11}(t) + \rho_{22}(t) = 1$

$$\text{and } t_m = A_{21}^{-1} [\ln(\rho_{22}(0)/\rho_{11}(0))] \quad (3)$$

is related to the initial state of the atom at $t=0$. If we temporarily neglect that part of the self-field which yields only a small frequency shift, the solution for the off-diagonal elements are given by

$$\begin{aligned} \rho_{12}(t) &= \rho_{21}^*(t) \\ &= \left(\frac{\rho_{12}(0)}{\rho_{22}(0)} \right) \frac{\exp(-i\Omega_{12}t + A_{21}t/2)}{\exp[A_{21}(t - t_m)] + 1}, \end{aligned} \quad (4)$$

where $\rho_{nm}(0) \equiv C_n(0)C_m^*(0)$ and $C_n(0)$ are the initial values of the coefficients ($n=1, 2,$) in the wavefunction of the atom which describes the transition process. We first note, that the value of t_m determines the point at which the maximum atomic-dipole moment occurs [the effective-dipole moment of the transition is proportional to $(\rho_{12} + \rho_{21})$]. In Jaynes's semiclassical theory, it is assumed that the expectation value of the dipole moment of the atom is responsible for the radiation process. Hence, an excited atom radiates slowly until its dipole moment grows to an appreciable value, and then begins to radiate its energy away very rapidly. While this characteristic behavior was duly noted by Jaynes and Crisp in their article, the role that the value of t_m (the point in time where the maximum dipole radiation occurs) plays in determining the average lifetime of the atom was not clarified. While it is certainly true that Eqs. (1), (2), and (4) imply that most of the transition energy is radiated during a time interval which is proportional to the reciprocal of the associated Einstein A coefficient A_{21} , this "characteristic energy transfer time" is *not* equal to the average lifetime of the atom. In particular, if we define the average lifetime of the atom as that time required for $\rho_{22}(0) \simeq 1$, at $t=0$, to decay to $1/e$ th of its initial value, we find from Eq. (2) that

$$\tau_2 \sim (A_{21}^{-1}) \ln[(1.718)\rho_{22}(0)/\rho_{11}(0)](\text{sec}). \quad (5)$$

Hence, we see that the initial values of $\rho_{22}(0)$

and $\rho_{11}(0)$ [where $\rho_{11}(0) + \rho_{22}(0) = 1$] play a critical role in determining the atoms average lifetime, while the "characteristic energy transfer time" (occurring in some interval² of time Δt about t_m) is essentially independent of the atoms initial condition at $t = 0$. The implication here is that there is a need to include an additional element into the theory, one which can account for the experimental

fact that the average lifetime of an atom is essentially *independent* of its preparation. One possibility is that the presence of a "semiclassically described" vacuum state³ might produce the required behavior when properly included in the associated nonlinear density-matrix equations of the theory.

*Present address: Physics Department, University of Windsor, Windsor, 11, Ontario.

¹E. T. Jaynes and M. D. Crisp, Phys. Rev. **179**, 1253 (1969).

²Since the maximum-dipole moment occurs at $t = t_m$ in Eq. (4), the "time halfwidth" of the associated dipole radiation pulse about the maximum is given by

$$\Delta t \sim 2A_{21}^{-1} \ln \left(\frac{2 + \sqrt{3}}{2 - \sqrt{3}} \right).$$

Hence the associated frequency halfwidth is $\Delta\omega \sim (\Delta t)^{-1} < A_{21}$, which is smaller than that predicted quantum electrodynamics.

³E. A. Uehling, Phys. Rev. **48**, 55 (1935).

Reply to Leiter's Comment

E. T. Jaynes

Arthur Holly Compton Laboratory of Physics, Washington University, St. Louis, Missouri 63130

(Received 1 January 1970)

Leiter¹ raises an important point which illustrates the need for more refined experiments before we could claim to understand the dynamics of spontaneous emission and other radiation processes. The same point was raised by Schawlow at the 1966 Rochester Coherence Conference, and answered in the ensuing discussion. We welcome the opportunity to clarify matters to a wider audience.

The semiclassical or "neoclassical" theory (NCT) in question was developed by the writer and his colleagues²⁻⁷ with the following motivation. Our present quantum electrodynamics (QED) has not achieved any satisfactory final form; it contains many important "elements of truth," but is mixed up with clear "elements of nonsense." The divergence and other difficulties indicate, that at least one of its underlying principles must be modified; but for forty years we have lacked experimental clues suggesting where and how this should be done, and nobody has seen how to disentangle the truth from the nonsense.

A possible way out of this impasse is to try to construct alternative theories in which various objectionable features of QED are eliminated by *fiat*, and see whether they suggest new experi-

ments capable of deciding among them. If some alternative theory could be shown to contain just one grain of truth that is not contained in present QED, then we would have the missing clue showing how QED must be modified.

NCT automatically removes all divergences arising from field quantization and infinite vacuum fluctuations, but retains the conventional Schrödinger equation to describe the behavior of matter. Although energy exchanges between field and matter then take place continuously, there is a strong tendency for this to occur in units of $\hbar\omega$, explained by NCT in a completely mechanistic and causal manner as a consequence of the equations of motion for matter – just as Planck and Schrödinger always believed must be true.

To the best of our knowledge, NCT agrees with existing experiments in every case where accurate calculations have been completed.⁸ But the predictions always differ from those of QED in finer details on which we have as yet no experimental evidence. The case of spontaneous emission discussed by Leiter is one example of this. Considering for simplicity only two levels, when an atom is excited (for example, by electron impact) we have to expect that, in general, it will not be left in ex-