

TABLE A-V-E. Corrected partial wave contributions to SEPna A_2 elastic and absorption cross sections (a_0^2) at 30–100 eV for e-Ne scattering.

E (eV)	l	Elastic	Absorption
30	0	5.57	5.50(-2)
	1	6.28	1.43(-1)
	2	1.47	3.67(-1)
	3	8.03(-2)	1.35(-1)
	4	1.21(-2)	4.12(-2)
	≥ 5	8.69(-3)	1.38(-2)
50	0	2.98	8.82(-2)
	1	5.42	2.63(-1)
	2	1.97	7.49(-1)
	3	1.42(-1)	4.38(-1)
	4	1.79(-2)	1.99(-1)
	≥ 5	8.21(-3)	1.14(-1)
100	0	7.78(-1)	8.55(-2)
	1	3.55	2.71(-1)
	2	2.15	6.89(-1)
	3	3.15(-1)	7.12(-1)
	4	5.98(-2)	5.21(-1)
	5	1.33(-2)	3.19(-1)
	6	3.24(-3)	1.72(-1)
	≥ 7	2.16(-3)	1.50(-1)

Erratum: Antiparticle excitation of atomic inner shells
 [Phys. Rev. A 27, 578 (1983)]

Werner Brandt and George Basbas

Figure 3 is mislabeled: The high- Z_2 target is tin ($Z_2=50$), not silicon ($Z_2=14$). The associated text and figure caption require no changes. A factor of 9 should be inserted in Eq. (5) to multiply the integral; the right-hand side is correct as printed.

Erratum: Amplitude combinations in the critical binary fluid nitrobenzene and *n*-hexane
 [Phys. Rev. A 28, 440 (1983)]

G. Zalczer, A. Bourgou, and D. Beysens

There is an error in the caption of Table IV, where the formula $\Delta\rho = \dots$ should read

$$\Delta\rho/\rho = T_c [(1/\rho)(\partial\rho/\partial T)_{p,M}^{\text{reg}}] t + R_\rho t^{0.89} .$$