

### Nonvariational calculation of the sticking probability and fusion rate for the $\mu dt$ molecular ion

R. Krivec

*Physics Department, The Hebrew University, Jerusalem 91904, Israel  
and J. Stefan Institute, Jamova 39, P.O. Box 100, 61111 Ljubljana, Slovenia*

V. B. Mandelzweig

*Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel*

(Received 13 December 1994)

The sticking probability for the  $\mu dt$  molecular ion is estimated with the help of directly calculated nonvariational wave functions obtained by the correlation function hyperspherical harmonic method. The method generates an accurate, locally correct  $\mu dt$  wave function which leads to precise estimates of the partial-wave sticking coefficients and the fusion rate. Our calculations are compared with other precision computations.

PACS number(s): 36.10.Gv, 31.15.Ja

In the past few years the muon-catalyzed fusion reaction

$$dt\mu \begin{cases} \xrightarrow{\omega_s^{(0)}} (\alpha\mu + n + \Delta) \\ \xrightarrow{1-\omega_s^{(0)}} (\alpha + \mu + n + \Delta) \end{cases} \quad (1)$$

where  $\Delta = 17.6$  MeV is the energy release, has attracted considerable attention. In (1) the sticking coefficient  $\omega_s^{(0)}$  is the probability of the first reaction, when the muon in the final state is bound to the  $\alpha$  particle. The value of  $\omega_s^{(0)}$  plays an essential role in the number of fusion reactions a muon can produce before it decays. This quantity limits the number of synthesis reactions that a muon can catalyze during its lifetime, and thus determines the possibility of using this reaction in energetics.

Calculation of the muon- $\alpha$ -particle sticking coefficient is, therefore, of special importance in muon-catalysis problems. Also, there is an uncertainty in the experimental estimation of this coefficient that makes an accurate theoretical determination of this quantity to be of great significance.

The measured quantity in experiments is not  $\omega_s^{(0)}$ , but the effective sticking coefficient  $\omega_s = (1-R)\omega_s^{(0)}$ , where  $R$  contains the effect of the stripping, i.e., the reactivation of muons by collisions. Some experimental values [1-3] of  $\omega_s$  (in %) are  $0.45 \pm 0.05$ ,  $0.35 \pm 0.07$ , and  $0.59 \pm 0.07$ ;  $R = 0.35 \pm 0.05$  [1,2]. The corresponding estimates of  $\omega_s^{(0)}$  (in %) are  $0.7 \pm 0.13$ ,  $0.54 \pm 0.15$ , and  $0.91 \pm 0.18$ , respectively. For recent results, see Ref. [4].

In recent years, many different calculations [4-15, 17, 18, 34] of  $\omega_s^{(0)}$  were performed that give appreciably larger values of the sticking coefficient. Some values from the literature [4-9] (in %) are 0.8824, 0.886, 0.897, 0.895, 0.845, and 0.858. These theoretical estimates are based on taking into account the Coulomb in-

teraction between the constituents and do not account for corrections due to nuclear interaction in the  $dt$  subsystem. These corrections are of the same order of magnitude [4] as the discrepancies in the literature.

The sticking coefficient  $\omega_s^{(0)}$  was calculated by many different methods such as adiabatic [17,18], variational [4-6,9-14], and statistical [7] (Green-function Monte Carlo). However, a direct nonvariational approach to the problem using three-body techniques was not yet employed. The purpose of this work is to apply the correlation function hyperspherical harmonic (CFHH) method [19-31] of solving the Schrödinger equation, which is geared to provide not only precise estimates of energy but also wave functions accurate in the whole range of interparticle distances. Use of these wave functions leads in turn to precise estimates of the expectation values of the Hamiltonian and of different operators, including the sticking coefficient.

In the CFHH method [19-31], which is described in Ref. [22], one writes the wave function as a product of two factors,

$$\Psi = \chi \phi, \quad (2)$$

where  $\chi$  is the "correlation factor" and  $\phi$  is expanded in the usual hyperspherical harmonic (HH) functions. If the correlation factor  $\chi$  is chosen to describe the singular features of  $\Psi$  (like cusps) as well as its asymptotic behavior, the function  $\phi$  will be compact, that is, little expanded in space and the HH expansion for it should be rapid. The solution for  $\phi$  proceeds as in the usual HH method, except that the potential  $V$  is replaced by an effective velocity-dependent potential  $V'$ ,

$$V' = V - \frac{1}{2} \frac{\nabla^2 \chi}{\chi} - (\nabla \ln \chi) \nabla, \quad (3)$$

where  $\nabla$  is the six-dimensional gradient operator and  $V$  is the sum of the interparticle potentials.

In the first CFHH calculation of the  $\mu dt$  system [28], the correlation factor  $\chi = \exp(f)$  with a simple linear correlation function

$$f = \sum_{i=1}^3 b_i r_i \quad (4)$$

was employed. Here  $b_i$  are adjustable parameters, chosen to describe the cusp singularities,

$$b_i = \frac{m_j m_k}{m_j + m_k} Z_j Z_k, \quad (5)$$

where  $m_j$  and  $Z_j$  are the mass and the charge of the particle  $j$ , and  $r_i$  are the interparticle distances in the odd-man-out notation, with  $\{d, t, \mu\}$  corresponding to particles  $\{1, 2, 3\}$ , respectively. However, such a choice of the parameter  $b_3$ , which is positive, leads to wrong asymptotic behavior of the correlation factor. For systems with large  $m_1 m_2$ , this prevents the CFHH method from converging. Therefore, the linear form (4) forces one to use a different value of this parameter. This corrects the asymptotic behavior of the wave function but destroys the proper description of the  $dt$  cusp. The recent more sophisticated  $\mu dt$  calculation [31] employs a nonlinear correlation function

$$f = \sum_{i=1}^3 [a_i + (b_i - a_i) \exp(-c_i r_i)] r_i \quad (6)$$

geared to reproduce both all the cusps and the asymptotic behavior of the wave function. This guarantees smoothness of the factor  $\phi$  in (2) and correspondingly a fast convergence of the hyperspherical expansion both at small and large interparticle distances  $r_i$ . Indeed, at  $r_i \rightarrow 0$   $f$  has the form

$$f = \sum_{i=1}^3 [b_i r_i + \mathcal{O}(r_i^2)] \quad (7)$$

so the parameters  $b_i$  have to describe the cusp singularities, while at  $r_i \rightarrow \infty$   $f$  has the form

$$f \approx \sum_{i=1}^3 a_i r_i \quad (8)$$

and the parameters  $a_i$  can be used to provide a proper asymptotic description. Usually  $a_i$  are defined by constraints obtained from known cluster asymptotics. Parameters  $c_i$  determine the start of the asymptotic region. Use of such a correlation function allows a precise estimate of the wave function for all interparticle distances including the  $dt$  coalescence point where the knowledge of the wave function is essential for an accurate calculation of the fusion rate and of the muon sticking probability. In addition, calculation shows [31] that the use of the nonlinear correlation function (6) improves the overall convergence patterns of the expectation values of different operators, including the Hamiltonian, decreasing, for example, the error in energy by an order of magnitude, if the same number of hyperspherical harmonic functions are used. The advantage of the nonlinear correlation function (6) *vis a vis* a linear one is especially obvious from the comparison of Figs. 1 and 2, where values of  $H\Psi/E\Psi - 1$  (which are identically zero for the exact wave function and whose values at different points characterize, therefore, the extent of the local deviation




FIG. 1. Values of  $H\Psi/E\Psi - 1$ , in percent, for the linear correlation function  $f$  (parametrization A) and  $K_m = 40$ . The coordinate  $r_3$  is the distance between  $d$  and  $t$ , and  $s_3$  is the distance between  $\mu$  and the center of mass of  $d$  and  $t$ . The angle between the corresponding Jacobi vectors is  $175^\circ$  so  $\mu$  is close to  $d$  if  $r_3$  and  $s_3$  have a fixed ratio.

from the precise wave function) are presented. Note that the scales of Figs. 1 and 2 differ by an order of magnitude.

For the present calculation of the sticking probabilities and fusion rates we, therefore, use the wave functions with the nonlinear correlation function  $f$  obtained in our work of Ref. [31]. In this work, the following values of the deuteron, triton, and muon masses and of the Rydberg constant ( $\mathcal{R}$ ) most commonly employed in muonic molecules binding energy calculations were used,

$$\begin{aligned} m_\mu &= 206.769 m_e, & m_d &= 3670.481 m_e, \\ m_t &= 5496.918 m_e, & \mathcal{R} &= 13.605\,8041 \text{ eV}. \end{aligned} \quad (9)$$

The choice of parameters  $a_i$ ,  $b_i$ , and  $c_i$  is discussed in Ref. [31]. The parameters  $b_i$  are given by Eq. (5) while the parameters  $c_i$  equal 0.0990, 0.0943, and 0.519




FIG. 2. Same as in Fig. 1, but for the nonlinear correlation function (parametrization C). The vertical scale is different from that of Fig. 1.

TABLE I. Correlation function parameters  $a_i$  (in d.a.u., deuteron atomic units, where mass of the deuteron is set to unity) for different parametrizations  $A$ ,  $B$ ,  $C$ , and  $D$ . Each parametrization may be used with different values of  $c_i$ .  $A$  refers to linear  $f$ .  $B$  is defined by the constraint that the asymptotic conditions are imposed on  $\exp(f)$ .  $C$  is such that the asymptotic conditions are imposed self-consistently on the full wave function  $\Psi$  at  $K_m=0$ . (This is a simple nonlinear problem that can be solved before the actual calculation.)  $D$  refers to choices without constraints.

	$a_1$	$a_2$	$a_3$
$A^a$	-0.054 290 772 4	-0.053 328 776 8	-0.281 664 72
$B$	-0.017 485 511 192 784	-0.012 356 005 126 688	-0.05
$C^b$	-0.054 290 772 402 005	-0.053 328 776 812 877	-0.29
$D$	0.0	0.0	0.0

<sup>a</sup>Listed are the values of  $b_i$ , cf. Eq. (4);  $b_3$  differs from the cusp value, 0.599 615 878, as required for the CFHH method to converge.

<sup>b</sup>The value of  $a_3$  is taken approximately equal to the  $b_3$  of  $A$  (first entry). In both  $A$  and  $C$  parametrizations this parameter is defined by the same constraint.

$\mu$ .a.u.<sup>-1</sup> (muonic atomic units where mass of the muon is set to unity) for  $i = 1, 2$ , and  $3$ , respectively. (Only  $c_3$  appears in the parametrization  $C$ .) The parameters  $a_i$  are presented in Table I.

The sticking probability  $\omega_s^{(0)}$  is the sum of the partial sticking probabilities  $\omega_{nl}$  defined by the equation [7,11]

$$\omega_{nl} = 4\pi(2l+1) \left| \int_0^\infty dr r^2 R_{nl}(r) j_l(Qr) \psi(r) \right|^2 \quad (10)$$

in which averaging over the direction of the outgoing neutron and summation over magnetic quantum numbers is performed.

Here  $\psi(r)$  is the three-body wave function  $\Psi$  at the point where  $d$  and  $t$  are located on top of each other, normalized so that the integral of  $|\psi(r)|^2$  in the muonic coor-

dinate equals unity,  $R_{nl}(r)$  is the radial function of muonic helium,  $j_l(Qr)$  is the spherical Bessel function, and  $Q$  equals 5.844 in the  $\mu$ .a.u. (although sometimes a slightly different value of  $Q = 5.846$ , obtained from relativistic kinematics, is used [10,11]).

The results of our calculations of the partial ground-state sticking probabilities and their comparison with previous precision computations are presented in Table II. One can see that the values of the sticking probability are very stable and change very little with the increase of  $K_m$ .  $K_m$  is the maximum global angular momentum and the number of the HH used is  $\frac{1}{2}(K_m/2+1)(K_m/2+2)$ . The dependence on the parametrization of  $f$  is also negligible and is shown in Fig. 3. From comparison of the values for  $K_m = 40$  and  $48$  one concludes that the uncer-

TABLE II. Ground-state sticking probabilities  $\omega_{nl}$  for  $Q = 5.844$  calculated with the nonlinear correlation function  $f$ . Prescriptions  $B$ ,  $C$ , and  $D$  give the same values of  $\omega_{nl}$  to the number of quoted digits. The second line for  $K_m = 40$  is for  $Q = 5.846$ .

$K_m$	1s	2s	3s	4s	2p	3p	4p	3d	4d
32	0.6822	0.0978	0.0297	0.0127	0.0238	0.0086	0.0039	0.0002	0.0001
40	0.6820	0.0978	0.0297	0.0127	0.0238	0.0086	0.0038	0.0002	0.0001
	0.6807	0.0976	0.0297	0.0126	0.0237	0.0085			
48	0.6819	0.0978	0.0297	0.0126	0.0238	0.0086	0.0038	0.0002	0.0001
Ref. [6]	0.6932	0.0992	0.0302	0.0128	0.0241	0.0087	0.0039		
Ref. [7]	0.689	0.099	0.030	0.013	0.024	0.009			
Ref. [10] <sup>a</sup>	0.6846								
	0.6826	0.0979	0.0297	0.0127	0.0238	0.0086	0.0039	0.0002	
Ref. [11] <sup>b</sup>	0.6817								
	0.6800	0.0975	0.0296	0.0126	0.0237	0.0085	0.0038	0.0002	
Ref. [12] <sup>c</sup>	0.6842								
	0.6825	0.0978	0.0297	0.0127	0.0238	0.0086	0.0039	0.0002	
Ref. [4] <sup>d</sup>	0.6802	0.0975	0.0296	0.0126	0.0237	0.0086	0.0039	0.0002	

<sup>a</sup> $Q = 5.8460$  (second line). Authors state that the change of the total sticking fraction if  $Q$  is increased to 5.847 is about  $-0.001$ . For our value (5.844) the change would be  $+0.002$  (this extrapolation is shown in the first line, giving even larger disagreement with our value).

<sup>b</sup>Authors quote values for  $Q = 5.846$  (second line). They also compare with values for  $Q = 5.844$ ; the latter values are said to be larger by 0.0017 (in our calculation this difference is 0.0013). The first line was obtained by adding 0.0017.

<sup>c</sup> $Q = 5.846$  (second line); the first line was obtained by adding 0.0017.

<sup>d</sup> $Q = 5.844$ . Quoted are the purely Coulombic results corresponding to an  $R$ -matrix formulation imposing the nuclear boundary condition.




FIG. 3. Ground-state sticking coefficient  $\omega_{10}$  (in percent) for different parametrizations and its comparison with variational calculations. Upper values correspond to  $Q=5.844$  and lower values to  $Q=5.846$ . Pairs of points on the right are crosses (nonextrapolated variational values) and diamonds (extrapolated variational values). The variational results in sequence from left to right on the graph correspond to Refs. [10, 11, and 12], respectively.

tainty of our  $\omega_{10}=0.6819$  value (for  $Q=5.844$ ), for example, is 0.0001 and is less than the differences in the literature for a given value of  $Q$ . In particular, the values of Ref. [10] adjusted for  $Q$  deviate more from our results than the published values. The recent results of Ref. [4] also deviate appreciably, in the opposite direction.

Table III displays the dependence of the ground-state probability  $\rho_0$  (see Ref. [16]) of the deuteron and triton being together at the same point on  $K_m$  (in units of  $10^{-13} \text{ fm}^{-3}$ ). This probability determines the fusion rate [33]:  $\lambda_f = A_s \rho_0$ . We present  $\rho_0$  rather than  $\lambda_f$  because the convention and value of  $A_s$  vary in the literature. The value is [13,32]  $A_s = 1.3 \times 10^{-14} \text{ cm}^3 \text{ sec}^{-1}$ ; Ref. [14] uses  $\lambda_f = A_r \rho_0$ , where  $A_r = \frac{3}{2} 1.361 \times 10^{-14} \text{ cm}^3 \text{ sec}^{-1}$ ; Ref. [34] used likewise  $\frac{3}{2} A_s = \frac{3}{2} 1.36 \times 10^{-14} \text{ cm}^3 \text{ sec}^{-1}$ , the factor  $\frac{3}{2}$  coming from spin symmetry.

The dependence of  $\rho_0$  on  $K_m$  is also shown graphically on Fig. 4. Again, one can see that different parametrizations B, C, and D for the nonlinear correlation function converge to the same value with the increase of  $K_m$  and that the uncertainty of our  $\rho_0=0.5280 \pm 0.0011$  value for  $K_m=48$  is much less than the discrepancies in the literature.

TABLE III. Probability of  $d$  and  $t$  being at the same point,  $\rho_0 = \langle \delta(\mathbf{r}_{dt}) \rangle$  (in units of  $10^{-13} \text{ fm}^{-3}$ ), for the ground state of  $\mu dt$  for different parametrizations B, C, and D, and its comparison with variational calculations.

$K_m$	C	B	D
32	0.5204	0.5250	0.5285
40	0.5250	0.5273	0.5290
48	0.5269	0.5280	0.5287
Ref. [13]		0.5296	
Ref. [14] <sup>a</sup>		0.5502	
		0.5294	
Ref. [15] <sup>b</sup>		0.594	
Ref. [32] <sup>c</sup>		0.6	

<sup>a</sup>Given are fusion rates. We calculated  $\rho_0$  by dividing by the quoted constant,  $A_r = 2.042 \times 10^{-14} \text{ cm}^3/\text{sec}$ . The first line is for the adiabatic case, the second line for the nonadiabatic case.

<sup>b</sup>Misquoted in Ref. [13] as  $7.47 \times 10^{-13}$  (should be divided by  $4\pi$ ).

<sup>c</sup>Extrapolation to nuclear interaction radius zero from Ref. [32], using the quoted coefficient  $A_{L=0} = 1.3 \times 10^{-14} \text{ cm}^3/\text{sec}$ . Reference [32] takes into account the nuclear interaction to estimate  $\rho_0$ .




FIG. 4. Expectation values of the  $\delta$ -function operator of the distance between the deuteron and the triton, proportional to the fusion rate according to the Jackson formula [33], in units of  $10^{-13} \text{ fm}^{-3}$ , for the linear (parametrization A) and nonlinear (parametrizations B, C, and D) correlation functions. Separate points on the right display the results of variational calculations. The variational results in sequence from left to right on the graph correspond to Refs. [13 and 14], respectively, with the upper (lower) points corresponding to the adiabatic (nonadiabatic) values.




FIG. 5. Ratio of the CFHH wave function  $\Psi$  with the nonlinear correlation function (parametrization  $C$ ,  $K_m = 40$ ) at the  $dt$  coalescence point (where  $\Psi = \psi$ ) and the Green-function Monte-Carlo wave function  $\Psi_{MC}$  of Ceperley and Alder [7].  $r_{\mu-dt}$  is the distance between  $\mu$  and the center of mass of  $d$  and  $t$ .

It is interesting to compare our wave function with the Green-function Monte-Carlo wave function of Ceperley and Alder [7], which was also used in a nonvariational calculation of the sticking probabilities. The result of the comparison is presented in Figs. 5 and 6, where the ratio of both wave functions and the divided differences of the logarithms of both wave functions at the  $dt$  coalescent point are calculated. Figure 5 shows unexpected quasi-periodic deviations from unity. Therefore, we estimated the derivative (Fig. 6) and found that the deviations are caused by the roughness of the Ceperley-Adler wave function.

We have calculated the partial wave sticking probabilities  $\omega_{nl}^{(0)}$  and the probability of  $d$  and  $t$  being at the same point  $\rho_0$  and compared them to values in the literature.

There are significant discrepancies in the literature on the sticking probability and  $\rho_0$  (more so for fusion rates which also depend on the constant  $A_s$  adopted), from the point of view of the reliability of various precision computation methods.

Using the CFHH method with a nonlinear correlation function, we have obtained results that have converged much better than discrepancies between different calculations in the literature, both older and recent. Since the




FIG. 6. Divided differences (approximate derivative) of the logarithm of the wave function at the  $dt$  coalescence point. The differences correspond to intervals of 0.5 for  $r_{\mu-dt} \leq 5$  and to intervals of 1 for  $r_{\mu-dt} > 5$  (in  $\mu.a.u.$ ). The solid curve is the CFHH wave function calculated with the nonlinear correlation function (parametrization  $C$ ,  $K_m = 40$ ) while the dashed curve represents the Green-function Monte-Carlo wave function of Ceperley and Alder [7]. The Born-Oppenheimer wave function (an exponential) would give a constant value.

expectation values in the CFHH method converge to the correct limits, uncertainty of the results can be deduced from the convergence patterns (Figs. 3 and 4).

No modifications of the method were necessary to get precise wave function at coalescence points; accuracy of the various expectation values is uniform. In variational calculations, it is usually necessary to modify the basis to better describe certain expectation values; frequently, quadruple precision arithmetic has to be used.

The work also confirmed that different parametrizations of the correlation function  $f$  give insignificantly different results, as in the case for other expectation values [31]. Also, it is sufficient to use the one parametrization ( $B$ ) for which the correct cluster asymptotics is imposed on  $\exp(f)$ . This is the most straightforward of the various prescriptions for choosing the parameters of  $f$  (see Table I).

This work has been partially supported by the Ministry of Science and Technology of Slovenia, Grant No. P1-5056-0106. The calculations have been done on the Convex C3860 of the J. Stefan Institute.

- [1] W. H. Breunlich *et al.*, Phys. Rev. Lett. **58**, 329 (1987).
- [2] S. E. Jones *et al.*, Phys. Rev. Lett. **56**, 588 (1986).
- [3] C. Petitjean *et al.*, Muon-Catalyzed Fusion **5**, 261 (1990); K. Nagamine *et al.*, Muon-Catalyzed Fusion **5**, 289 (1990).
- [4] C.-Y. Hu, G. M. Hale, and J. S. Cohen, Phys. Rev. A **49**, 4481 (1994).
- [5] S. E. Haywood, H. J. Monkhorst, and K. Szalewicz, Phys. Rev. A **39**, 1634 (1989).
- [6] C.-Y. Hu, Phys. Rev. A **34**, 2536 (1986).
- [7] D. Ceperley and B. J. Alder, Phys. Rev. A **31**, 1999 (1985).
- [8] L. N. Bogdanova *et al.*, Nucl. Phys. A **454**, 653 (1986).
- [9] C.-Y. Hu, Phys. Rev. A **36**, 4135 (1987).
- [10] S. E. Haywood *et al.*, Phys. Rev. A **37**, 3393 (1988).
- [11] B. P. Carter and C.-Y. Hu, Phys. Rev. A **41**, 6509 (1990).
- [12] M. Kamimura, in *Muon Catalyzed Fusion*, Proceedings of the Conference on Muon-Catalyzed Fusion, Sanibel Island, Florida, 1988, edited by S. E. Jones, J. Rafelski, and H. J. Monkhorst, AIP Conf. Proc. No. 181 (AIP, New York, 1989), p. 330.
- [13] S. A. Alexander *et al.*, Phys. Rev. A **41**, 2854 (1990).
- [14] K. Szalewicz *et al.*, Phys. Rev. A **42**, 3768 (1990).
- [15] L. N. Bogdanova *et al.*, Zh. Eksp. Teor. Phys. **81**, 829 (1981) [Sov. Phys. JETP **54**, 174 (1981)].
- [16] J. P. Friar *et al.*, Phys. Rev. Lett. **66**, 1827 (1991).
- [17] S. Gerstein *et al.*, Zh. Eksp. Teor. Phys. **80**, 1690 (1981) [Sov. Phys. JETP **53**, 872 (1981)].
- [18] L. Bracci and G. Fiorentini, Nucl. Phys. A **364**, 383 (1981).
- [19] V. B. Mandelzweig, Nucl. Phys. A **508**, 63c (1990).
- [20] M. I. Haftel and V. B. Mandelzweig, Phys. Lett. A **120**, 232 (1987).
- [21] M. I. Haftel and V. B. Mandelzweig, Phys. Rev. A **38**, 5995 (1988).
- [22] M. I. Haftel and V. B. Mandelzweig, Ann. Phys. **189**, 29 (1989); **195**, 420 (1989).
- [23] M. I. Haftel and V. B. Mandelzweig, Phys. Rev. A **39**, 2813 (1989).
- [24] M. I. Haftel and V. B. Mandelzweig, Phys. Rev. A **41**, 2339 (1990).
- [25] M. I. Haftel and V. B. Mandelzweig, Phys. Rev. A **42**, 6324 (1990).
- [26] R. Krivec, M. I. Haftel, and V. B. Mandelzweig, Phys. Rev. A **44**, 7158 (1991).
- [27] M. I. Haftel and V. B. Mandelzweig, Phys. Rev. A **46**, 142 (1992).
- [28] R. Krivec, M. I. Haftel, and V. B. Mandelzweig, Phys. Rev. A **46**, 6903 (1992).
- [29] R. Krivec, M. I. Haftel, and V. B. Mandelzweig, Phys. Rev. A **47**, 911 (1993).
- [30] M. I. Haftel and V. B. Mandelzweig, Phys. Rev. A **49**, 3344 (1994).
- [31] R. Krivec, M. I. Haftel, and V. B. Mandelzweig, Few-Body Systems **17**, 229 (1994).
- [32] K. Langanke and D. Lukas, Few-Body Systems **13**, 87 (1992).
- [33] J. D. Jackson, Phys. Rev. **106**, 330 (1957).
- [34] P. Froelich *et al.*, Phys. Rev. Lett. **70**, 1595 (1993).