

Quantum tunneling in a dissipative system

Li Hua Yu

National Synchrotron Light Source, Brookhaven National Laboratory, New York 11973

(Received 13 May 1996)

Applying a technique developed recently [L. H. Yu and C. P. Sun, Phys. Rev. A **49**, 592 (1994); L. H. Yu, Phys. Lett. **202**, 167 (1995)] for a harmonic oscillator coupled to a bath of harmonic oscillators, we present an exact solution for the tunneling problem in an Ohmic dissipative system with inverted harmonic potential. The result shows that while the dissipation tends to suppress the tunneling, the Brownian motion tends to enhance the tunneling. Whether the tunneling rate increases or not would then depend on the initial conditions. We give a specific formula to calculate the tunneling probability determined by various parameters and the initial conditions. [S1050-2947(96)00511-2]

PACS number(s): 03.65.Bz

I. INTRODUCTION

Quantum tunneling with dissipation has been studied by many people after the work by Caldeira and Leggett [1-4]. These studies use different approximations to calculate the tunneling probability. Among these works, there is a widely discussed question about whether the dissipation suppresses or enhances the quantum tunneling [5]. In this paper we shall show that for a special potential barrier, the inverted harmonic potential well, the tunneling problem can be solved exactly, thus we shall answer this question precisely. The technique we used for this solution has been developed recently to solve the wave function evolution for another dissipative system problem: a harmonic oscillator coupled to an environment [6,7]. In this paper we shall show that this technique is equally applied to the inverted harmonic potential.

The simplest example of a dissipative system, a harmonic oscillator coupled to an environment of a bath of harmonic oscillators, has been the subject of extensive studies (see [6] for references). In a recent paper [6] we obtained a simple and exact solution for the wave function of the system plus the bath, in the Ohmic case. It is described by the direct product in two independent Hilbert spaces. One of them is described by the Caldirola-Kanai (CK) Hamiltonian, the other represents the effect of the bath, i.e., the Brownian motion. In a second paper [7], using this wave function solution, we derived a simple formula for the probability distribution at finite temperature, expressed in terms of the wave function solution for the CK Hamiltonian only.

In this paper we shall apply these results to a different system, the inverted harmonic potential well, to study the quantum tunneling problem of dissipative system. We shall use the same notations as Refs. [6,7] and quote formulas from there. The Hamiltonian of this system is

$$H = \frac{p^2}{2M} - \frac{1}{2}M(\omega_0^2 + \Delta\omega^2)q^2 + q \sum_j c_j x_j + \sum_j \left(\frac{p_j^2}{2m_j} + \frac{1}{2}m_j\omega_j^2 x_j^2 \right). \quad (1)$$

The only difference of this Hamiltonian from that in Ref. [6] is the sign of the second term of the right hand side, i.e., the

potential is now an inverted harmonic potential well. Assuming the Ohmic bath oscillator density to be the same as given in [6],

$$\rho(\omega_j) = \frac{2\eta M}{\pi} \frac{m_j \omega_j^2}{c_j^2}. \quad (2)$$

Following exactly the same procedure as [6], we derive the Langevin equation of motion for the system

$$\ddot{q} + \eta \dot{q} - \omega_0^2 q = f(t), \quad (3)$$

with the Brownian motion driving force

$$f(t) = - \sum_j \frac{c_j}{M} \left(x_{j0} \cos \omega_j t + \dot{x}_{j0} \frac{\sin \omega_j t}{\omega_j} \right). \quad (4)$$

This equation is easily solved to give

$$q(t) = a_1(t)q_0 + a_2(t)\dot{q}_0 + \sum_j [b_{j1}(t)x_{j0} + b_{j2}(t)\dot{x}_{j0}], \quad (5)$$

where $q_0, \dot{q}_0, x_{j0}, \dot{x}_{j0}$ are the initial position and velocity operators of the system and bath, respectively, and

$$a_1 = e^{-(\eta/2)t} \left(\cosh \omega t + \frac{\eta}{2\omega} \sinh \omega t \right), \quad a_2 = e^{-(\eta/2)t} \frac{\sinh \omega t}{\omega}, \quad (6)$$

$$b_{j1}(t) = - \frac{c_j}{M} \int_0^t a_2(t') \cos \omega_j(t-t') dt',$$

$$b_{j2}(t) = - \frac{c_j}{M} \int_0^t a_1(t') \sin \omega_j(t-t') dt', \quad (7)$$

with $\omega = (\omega_0^2 + (\eta^2/4))^{1/2}$.

Then, based on the same arguments as [6], this solution can be used to show that the wave function of the system plus the bath can be written as

$$\Psi(q, \{\xi_j\}, t) = \psi\left(q - \sum_j \xi_j, t\right) \prod_{j=1}^N \chi_j(\xi_j, t), \quad (8)$$

where the wave function $\psi(Q, t)$ is a solution of Schrödinger equation with the time dependent CK Hamiltonian

$$H_Q = e^{-\eta t} \frac{P^2}{2M} - \frac{1}{2} M \omega_0^2 e^{\eta t} Q^2, \quad (9)$$

and the commutation $[Q, P] = i\hbar$, while $\xi_j = b_{j1}(t)x_{j0} + b_{j2}(t)\dot{x}_{j0}$ is the contribution of the j th bath oscillator to the Brownian motion. The function $\chi_j(\xi_j, t)$ is given by $\chi_j(\xi_j, t) = \langle \theta_{\xi_j} | \chi_{j0} \rangle$, where $\chi_{j0}(x_{j0})$ is the initial state of the bath oscillator, and $|\theta_{\xi_j}\rangle$ is an eigenstate of the operator ξ_j with eigenvalue ξ_j , as given by Eq. (4) of [7]:

$$\theta_{\xi_j}(x_{j0}, t) = \left(\frac{m_j}{2\pi\hbar b_{j2}}\right)^{1/2} \exp\left[-\frac{im_j}{2\hbar b_{j2}}(b_{j1}x_{j0}^2 - 2x_{j0}\xi_j)\right]. \quad (10)$$

According the analysis of [7], assuming the bath is at temperature T , averaging over the Boltzmann distribution of the wave function $\chi_{j0}(x_{j0})$ of the bath oscillators, and using Eq. (8), we find the probability distribution the same as derived in [7]:

$$\rho(q, t) = \frac{1}{\sqrt{2\pi}\sigma_\xi} \int |\psi(q - \xi, t)|^2 e^{-\xi^2/2\sigma_\xi^2} d\xi, \quad (11)$$

where σ_ξ is the Brownian motion width:

$$\sigma_\xi^2(t) = \int_0^{\omega_c} \frac{\hbar}{2m_j\omega_j} [|b_{j1}(t)|^2 + \omega_j^2 |b_{j2}(t)|^2] \times \coth\left(\frac{\hbar\omega_j}{2kT}\right) \rho(\omega_j) d\omega_j. \quad (12)$$

This result is apparently the same as that given in [7], but the expressions for $b_{j1}(t)$ and $b_{j2}(t)$ are different. There are also subtleties associated with a logarithmic divergence of the integration over ω_j , which is removed by introducing a cutoff frequency ω_c . This width is zero initially, but then increases exponentially to infinity as time evolves. In the following, for simplicity, we consider only the low temperature limit, then the width approaches in a time range of about $1/\omega$ to the asymptotic value

$$\sigma_\xi^2 = e^{(2\omega - \eta)t} \sigma_0^2 \frac{\eta}{2\omega} \frac{\omega_0}{\omega} B, \quad (13)$$

with $\sigma_0^2 = \hbar/(2M\omega_0)$ (the width of the ground state of a harmonic oscillator with frequency ω_0) and

$$B \equiv \frac{1}{\pi} \ln \left[1 + \left(\frac{\omega_c}{\omega - \eta/2} \right)^2 \right]. \quad (14)$$

II. THE WAVE FUNCTION EVOLUTION

To calculate the tunneling probability, we assume that initially the wave function is a Gaussian wave packet cen-

tered at the right of the peak of the potential by z_0 with a velocity $v_0 = -\hbar k/M$ towards the potential barrier peak:

$$\psi_0(q) \equiv \psi(q; t=0) = (2\pi\sigma^2)^{-1/4} e^{-(q-z_0)^2/4\sigma^2 + ikq}. \quad (15)$$

First, following exactly the same method as in [6], we derive the Green function of the time dependent Hamiltonian Eq. (9):

$$G(q, q_0; t, 0) = \left(\frac{M}{2\pi i\hbar a_2}\right)^{1/2} \times \exp\left[\frac{iM}{2\hbar a_2}(a_1 q_0^2 + \dot{a}_2 e^{\eta t} q^2 - 2q_0 q)\right]. \quad (16)$$

Then we calculate the wave function $\psi(q, t)$:

$$\begin{aligned} \psi(q, t) &= \int G(q, q_0; t, 0) \psi_0(q_0) dq_0 \\ &= (2\pi\sigma^2)^{-1/4} (a_1 + i\omega_0 a_2 r^2) \\ &\quad \times e^{-(q-z_0)^2/4\sigma^2 + i(c_2 q^2 + c_1 q + c_0)}. \end{aligned} \quad (17)$$

This result represents a Gaussian distribution centered at $q_c = a_1 z_0 - a_2 v_0$ (the classical trajectory of a particle initially at z_0 with velocity $-v_0$) and with a width of

$$\sigma_\theta^2 \equiv \sigma^2 (a_1^2 + r^4 \omega_0^2 a_2^2), \quad (19)$$

where $r \equiv \sigma_0/\sigma$. The coefficients in the phase factor exponent are

$$c_2 = \frac{M e^{\eta t}}{4\hbar} \frac{d}{dt} (\ln \sigma_\theta^2), \quad (20)$$

$$c_1 = \frac{M e^{\eta t}}{4\hbar} \left(-2q_c \frac{d}{dt} (\ln \sigma_\theta^2) + 4\dot{q}_c \right), \quad (21)$$

$$c_0 = \frac{ka_2}{4} e^{\eta t} \left(q_c \frac{d}{dt} (\ln \sigma_\theta^2) - 2\dot{q}_c \right) + \frac{q_c z_0}{4\sigma_\theta^2} \omega_0 a_2 r^2. \quad (22)$$

This phase factor is related to the current density, as will be explained later. It is straightforward to verify that Eq. (18) indeed satisfies the Schrödinger equation with the time dependent Hamiltonian Eq. (9). With these provisions, we are ready to calculate the tunneling probability.

III. THE TUNNELING PROBABILITY AND CURRENT DENSITY

Using the density Eq. (11) and wave function Eq. (18), we find a very simple expression for the probability density:

$$\rho(q, t) = \frac{1}{\sqrt{2\pi}\sigma_t} e^{-(q-q_c)^2/2\sigma_t^2}, \quad (23)$$

where

$$\sigma_t^2 \equiv \sigma_\theta^2 + \sigma_\xi^2 \quad (24)$$

is the total width including the Brownian motion width.

The probability for the particle to pass to the left of position q at t is then

$$P(q,t) = \int_{-\infty}^q \rho(q',t) dt' = F\left(\frac{q-q_c}{\sqrt{2}\sigma_t}\right), \quad (25)$$

where the function F is

$$F(W) \equiv \frac{1}{\sqrt{\pi}} \int_{-\infty}^W e^{-u^2} du. \quad (26)$$

Hence the current density is found to be

$$I = -\frac{\partial P(q,t)}{\partial t} = |\psi|^2 \left(\frac{1}{2}(q-q_c) \frac{d}{dt} (\ln \sigma_t^2) + \dot{q}_c \right). \quad (27)$$

On the other hand, if the Brownian motion can be ignored (replacing σ_t by σ_θ), the solution Eq. (18) of the time dependent Schrödinger equation should also satisfy the expression for the current density:

$$I = \frac{\hbar}{2Me^{\eta t}} \left(\psi \frac{\partial \psi^*}{\partial q} - \psi^* \frac{\partial \psi}{\partial q} \right) = \frac{\hbar}{Me^{\eta t}} |\psi|^2 (2c_2 q + c_1). \quad (28)$$

The expressions for c_1 and c_2 , Eqs. (21) and (20), indeed satisfy Eq. (27), hence they acquire the following clear physical meaning. If we denote the velocity of the ‘‘fluid’’ distribution $|\psi|^2$ as v , the current density is then $I = |\psi|^2 v$. Therefore a comparison of Eq. (27) with Eq. (28) leads to

$$v = \frac{\hbar}{Me^{\eta t}} (2c_2 q + c_1) = \frac{1}{2}(q-q_c) \frac{d}{dt} (\ln \sigma_\theta^2) + \dot{q}_c. \quad (29)$$

Thus those particles with larger q acquire larger velocity falling from the apex of the potential. Since the width σ_θ^2 is proportional to $\exp(2\omega - \eta)t$, as time evolves, most particles move rapidly away from the apex of the potential. In the meantime, for any fixed $q \neq 0$, the wave number $2c_2 q + c_1$ also increases exponentially by a factor $e^{\eta t}$ according to Eqs. (21) and (20) because the equivalent mass is $Me^{\eta t}$ instead of M . Hence the wavelength decays exponentially, as pointed out by Ref. [7]. When $t \gg 1/\eta$ the system approaches the classical limit.

IV. TUNNELING PROBABILITY AS TIME APPROACHES INFINITY

We define the tunneling probability P as the probability for the particle to be at the left of the peak. Then, applying Eqs. (25), (24), and (19), we find $P = F(-W)$, with

$$W = \frac{q_c}{\sqrt{2}\sigma_t} = \frac{a_1 z_0 - a_2 v_0}{\sqrt{2[\sigma^2(a_1^2 + r^4 \omega_0^2 a_2^2) + \sigma_\xi^2]}}. \quad (30)$$

Because a_1, a_2 , and σ_ξ are all proportional to $e^{(\omega - \eta/2)t}$ as time approaches infinity, W approaches a finite limit, which determines the final tunneling probability. To simplify the expression for this limit, we use the following scaled variables:

$$Z \equiv \frac{z_0}{\sigma}, \quad V \equiv \frac{v_0}{\omega_0 z_0}, \quad \epsilon \equiv \frac{\eta}{2\omega}, \quad r \equiv \frac{\sigma_0}{\sigma},$$

$$B \equiv \frac{1}{\pi} \ln \left[1 + \left(\frac{\omega_c}{\omega - \eta/2} \right)^2 \right]. \quad (31)$$

Then, using Eq. (6) for a_1, a_2 , and Eq. (13) for the Brownian motion width, we have

$$W = \frac{Z}{\sqrt{2}} \frac{1 - V \sqrt{(1-\epsilon)/(1+\epsilon)}}{\sqrt{1 + (1-\epsilon)/(1+\epsilon)r^4 + 4B\epsilon \sqrt{1-\epsilon^2}r^2}}. \quad (32)$$

Notice that by the definition of ω following Eq. (7) and the definition of ϵ in Eq. (31), we always have $\epsilon \leq 1$. Thus W is always a real number, as required by the physical meaning of this variable.

To understand the meaning of this expression, we remark that $Z \gg 1$ represents a case where the wave packet width is much smaller than its distance from the origin. We also remark that if $\eta = 0$, as time approaches infinity, $q_c = e^{\omega_0 t} (z_0 - v_0/\omega_0)/2$. Hence $V < 1$ means that a classical particle with initial velocity v_0 and position z_0 does not have enough kinetic energy to pass the potential barrier if there is no dissipation. Thus $Z \gg 1$ and $V < 1$ represent a case relevant to the quantum tunneling problem.

The term $4B\epsilon \sqrt{1-\epsilon^2}r^2$ comes from the Brownian motion. If $\epsilon \ll 1$, i.e., if the dissipation is small, this term increases as ϵ increases, and reduces W , which in turn increases $F(-W)$, hence enhances the tunneling. This effect is reduced if $r \ll 1$, or, in other words, if the initial wave packet width is much larger than σ_0 , the increase due to the Brownian motion is insignificant. If the initial velocity is not zero, the second term in the numerator increases W as ϵ increases, thus the damping suppresses the tunneling. Intuitively this can be explained as that the damping makes the classical particle unable to move to the barrier peak as close as if there is no damping, and hence increases the barrier height.

To get an idea about the effect of dissipation on the tunneling probability, in Fig. 1 we plot P as a function of V and ϵ for $Z = 3$, $r = 0.3$, and $B = 3$, which corresponds to a cutoff frequency about 100 time larger than ω . Because of the logarithmic dependence of B on ω_c , the result is very insensitive to the cutoff frequency. The plot clearly shows that when V is large, the tunneling is suppressed by the dissipation, while if initial velocity is zero, the dissipation enhances the tunneling.

V. TUNNELING PROBABILITY AND UNCERTAINTY PRINCIPLE

Finally, it is interesting to examine the relation between the initial momentum and position distribution and the tunneling probability. For simplicity, we consider here only the case without dissipation. After Fourier transform of initial wave function Eq. (15), the wave function in momentum space is

$$\psi_0(k) = \left(\frac{2\sigma^2}{\pi} \right)^{1/4} e^{-(k+k_0)^2 - i(k+k_0)z_0}. \quad (33)$$




FIG. 1. The tunneling probability as a function of the scaled dissipation coefficient ϵ and the scaled initial velocity V , assuming the scaled initial position $Z=3$, the scaled inverse wave packet size $r=0.3$, and the scaled cutoff frequency $B=3$.

Thus from the classical point of view, the probability for the particle to pass the potential barrier is the probability of the initial velocity $v < -\omega_0 z_0$, or $k < -z_0/2\sigma_0^2$ (notice that $v = \hbar k/M$):

$$P = \int_{-\infty}^{-z_0/2\sigma_0^2} |\psi_0(k)|^2 dk = F\left(-\frac{Z(1-V)}{\sqrt{2}r^2}\right). \quad (34)$$

A comparison with Eq. (32) shows that if $r \gg 1$, i.e., if the initial width σ is much smaller than the minimum wave packet width σ_0 , or, in other words, if the initial momentum spread is very large, this crude estimate is correct.

Another extreme is when $r \ll 1$, i.e., when we can consider the initial momentum spread is very small, and all the particles have velocity v_0 , but initial position has large spread. Then from the classical point of view, the probability for a particle to pass the potential barrier is its initial $q < v_0/\omega_0$, i.e.,

$$P = \int_{-\infty}^{v_0/\omega_0} |\psi_0(q)|^2 dq = F\left(-\frac{Z(1-V)}{\sqrt{2}}\right). \quad (35)$$

Again, a comparison with Eq. (32) shows that if $r \ll 1$, i.e., if the initial width σ is much larger than the minimum wave packet width σ_0 , this crude estimate is also correct. These estimates indicate that the tunneling probability has a very simple relation with the uncertainty principle, and we have a very simple way to estimate the tunneling probability.

ACKNOWLEDGMENTS

The author thanks Professor C.N. Yang for many sessions of stimulating discussions. The author also likes to thank C.P. Sun for interesting discussions. The work is performed under the auspices of the U.S. Department of Energy under Contract No. DE-AC02-76CH00016.

- [1] A.O. Caldeira and A.J. Leggett, Phys. Rev. Lett. **46**, 211 (1981); Ann. Phys. (N.Y.) **149**, 374 (1983).
- [2] A.J. Bray and M.A. Moore, Phys. Rev. Lett. **46**, 1545 (1982); S. Chakravarty, *ibid.* **49**, 681 (1982).
- [3] G. Schön and A.D. Zaikin, Phys. Rep. **198**, 237 (1990), and references therein.
- [4] S. Chakravarty and A.J. Leggett, Phys. Rev. Lett. **52**, 5 (1984); A.J. Leggett, S. Chakravarty, A.T. Dorsey, M.P.A. Fisher, A. Garg, and W. Zwegler, Rev. Mod. Phys. **59**, 1 (1987).
- [5] About the discussion on whether the dissipation suppresses or

enhances the quantum tunneling, see, for example, A. J. Leggett and Satoshi Iso, in Proceedings of the 4th International Symposium: Foundations of Quantum Mechanics in the Light of New Technology, edited by M. Tsukata, S. Kobayashi, S. Kurihara, and S. Nomura [Jpn. J. Appl. Phys. (1992)]; K. Fujikawa, S. Iso, M. Sasaki, and H. Suzuki, Phys. Rev. Lett. **68**, 1093 (1992).

- [6] L.H. Yu and C.P. Sun, Phys. Rev. A **49**, 592 (1994).
- [7] L.H. Yu, Phys. Rev. Lett. A **202**, 167 (1995).