

Activating bound entanglement in multiparticle systems

W. Dür and J. I. Cirac

Institut für Theoretische Physik, Universität Innsbruck, A-6020 Innsbruck, Austria

(Received 9 February 2000; published 10 July 2000)

We analyze the existence of activable bound entangled states in multiparticle systems. We first give a series of examples that illustrate some different ways in which bound entangled states can be activated by letting some of the parties share maximally entangled states. Then, we derive necessary conditions for a state to be distillable as well as to be activable. These conditions turn out to be also sufficient for a certain family of multiqubit states. We use these results to explicitly construct states displaying properties related to bound entanglement and its activation.

PACS number(s): 03.67.Hk, 03.65.Bz, 03.65.Ca, 03.67.-a

I. INTRODUCTION

The existence of bound entanglement (BE) [1] has been one of the most intriguing surprises in quantum information during the last few years. A state is BE if (despite being entangled) one cannot distill [2] maximally entangled states (MES)¹ out of it by using local operations and classical communication (LOCC). From this definition it follows that with bound entangled states (BES) one cannot perform reliable teleportation [3,4], quantum communication [5], etc., i.e., they seem not to be useful for quantum information purposes. However, it has been shown that under certain conditions BE can be “activated” [6]. That is, with the help of some other entangled states they enable to carry out certain tasks that could not be performed by using these other entangled states and LOCC alone.

The first examples of BE arose in the context of *two systems*. In particular, the Horodecki showed that a necessary condition for a state of two systems to be distillable is that the corresponding density operator had nonpositive partial transposition [7,8]. Thus, all entangled states with a positive partial transposition [9,10] cannot be distilled, and therefore they are examples of BES [11]. Very recently, we showed that a different kind of BE can also appear in *multiparticle systems* [12].² In particular, we considered states of three systems A , B , and C , spatially separated, that cannot be prepared locally and that have the following properties: (i) if we would be allowed to join systems A and C in one place, the state could be prepared locally; (ii) if we would be allowed to join systems B and C in one place, the state could be prepared locally; (iii) but if we would be allowed to join systems A and B in one place, the state could not be prepared locally. The property (i) immediately implies that one cannot distill a MES between the systems A and B (or B and C) by using LOCC, since even if we would allow nonlocal opera-

tions between A and C we would not be able to do it. This is due to the fact that (a) nonseparability is a necessary condition for distillation, and (b) manipulations that are impossible when joining two systems [here (AC)] are clearly also impossible when the systems remain separated. Analogously, the property (ii) implies that one cannot distill a MES between the systems A and C . Thus, no MES can be distilled between any two of the systems as long as they remain separated from each other. The property (iii), however, indicates that the state is entangled, since even if we would allow nonlocal operations between A and B we could not prepare it locally. Thus, despite the fact that the state is entangled, one cannot distill any MES out of it, and therefore it is a BES. Nevertheless, the state presented in Ref. [12] can be activated. This follows from the fact that if we allow A and B to share some maximally entangled states (which is equivalent to allow A and B to join), then one can indeed distill a MES shared between A , B and C . Another example of activable BE has been given by Smolin in the context of unlockable entanglement for the case of four systems [13]. In that case, the state has the extra property that the entanglement between A and B can be activated using a single copy of the state only and by letting C and D share only one MES. These examples show that a new kind of BE can arise when we split some free entanglement among more than two parties. They also show that the possibility of activating BE is sometimes counterintuitive and may have some applications related to the process of secret sharing [14].

In this paper we construct multiparticle states that display properties related to BE. They illustrate the richness of multiparticle as compared to two-particle entangled states regarding the possibility of activating BE. This paper is organized as follows. In Sec. II, we state the problem of activation of BE in multiparticle systems and present various intriguing examples of activable BE. The rest of the paper provides the tools needed to construct states having the properties given in all those examples, as well as the guidelines to construct other states with different properties. In Sec. III, we review both the concept of bipartite splittings [15] and the family of N qubit states introduced in Ref. [12]. These states will play a central role throughout the remaining paper; as we will show, they give rise to a vast variety of activable BES. In Sec. IV, we consider the distillability properties of a multiparticle state when we allow the particles to join into

¹A MES of n parties is given by $|\Psi\rangle = (1/\sqrt{2})(|0^{\otimes n}\rangle + |1^{\otimes n}\rangle)$.

²BES in multipartite systems are states with the following properties: (i) they cannot be prepared by means of local actions by the parties (and are thus entangled) (ii) from this state no MES shared between any two of the (separated) parties can be created by means of LOCC.

two groups (i.e., according to a bipartite splitting of the particles). We will show that it is possible to find BES that can be activated if and only if the particles are joint according to certain bipartite splittings. Moreover, we will show that one can choose the bipartite splittings for which the BE can be activated without any restriction, and that our family of states covers all possible examples of this kind. In Sec. V, we consider a more general framework where the particles are allowed to join into more than two groups. We derive necessary conditions for distillability and activation based on the distillability properties of bipartite splittings. We show that these conditions are also sufficient for our family of states, which implies that they are an important ingredient to construct generic examples of activable BES. In particular, they allow us to construct BES fulfilling the properties introduced in the examples of Sec. II. We finally summarize our results in Sec. VI.

II. ACTIVATION OF BOUND ENTANGLEMENT

Let us consider N parties, A_1, \dots, A_N , at different locations, each of them possessing M qubits. We will assume that the state of the qubits is described by a density operator of the form $\rho^{\otimes M}$, where ρ denotes an entangled state of exactly N qubits, each of them belonging to a different party. Thus, the parties have M copies of the same state, where M can be as large as we wish. This ensures that the parties can use distillation protocols [2] in order to obtain MES between some of them. In that case we will say that the state ρ is distillable (with respect to the specific parties that can obtain a MES). A state ρ is BE if it is not distillable when the parties remain separated from each other. We say that a BES can be activated if it becomes distillable once some of the parties join and form groups to act together. Note that instead of allowing some parties to join we could have allowed them to share some MES. In that case we would have the same situation given the fact that separated parties sharing MES can perform any arbitrary joint operation by simply teleporting back and forth the states of their particles.

A. Examples

In this subsection we introduce some relevant examples of BES, by specifying their properties with respect to activation. In the following sections we will explicitly construct density operators that fulfill the properties given here for each of the examples. The goal of this subsection is to display the underlying properties of activable BES.

Example I. The state ρ_I becomes distillable if and only if the parties form two groups with exactly j and $N-j$ members, respectively. Furthermore, it does not matter which of the parties join in each group, but only the number of members. For example, if $N=8$ and $j=3$ [see Fig. 1(a)], we have that ρ_I is distillable if exactly 3 and 5 parties join, but remains undistillable when the parties form two groups with 1-7, 2-6, 4-4 members, or if they form more than two groups. In particular, ρ_I is not distillable if the parties remain separated from each other, which corresponds to having 8 groups.

Example II. The state ρ_{II} is distillable if and only if the parties form exactly two groups, each of them containing say




FIG. 1. (a) $N=8$ and the state ρ_I is distillable if and only if three and five parties form a group, and ρ_I is not distillable otherwise. (b) $N=25$ and ρ_{II} is distillable if and only if two groups of approximately the same size are formed (here 12-13), while ρ_{II} is not distillable if and only if one group contains less than 40% of the particles ($j < 10$). In both examples, it does not matter which of the parties belongs to which group.

between 40% and 60% of the parties. In particular, the state ρ_{II} remains undistillable if the parties form more than two groups or if they form two groups, with one of them containing less than 40% of the parties. Again, it does not matter whether certain parties belong to the same group; only the total number of particles within each group is important. We can view ρ_{II} as a state having “macroscopic” entanglement, but no (distillable) “microscopic” entanglement [see Fig. 1(b)]. Note that one can also have the opposite effect, i.e., a state ρ'_{II} which is distillable if and only if the parties form two groups, and one of the groups contains less than say 20% of the parties.

Example III. The state ρ_{III} becomes distillable if and only if the parties form two groups, where the first group contains a *specific* set of L parties $A = \{A_{k_1}, \dots, A_{k_L}\}$, and the second group contains the remaining parties. For all other configurations in groups ρ_{III} remain undistillable. For example, we have for $N=5$ and $A = \{A_1, A_3, A_5\}$ that ρ_{III} is distillable if and only if the parties form two groups, $(A_1 A_3 A_5) - (A_2 A_4)$, and not distillable otherwise.

Example IV. The state ρ_{IV} has the following properties: Given any two groups, each of them with a specific number j (or more) parties, they can distill (with the help of the other parties) a MES between them. In particular, if there are several groups of j or more parties, they can distill a Greenberger-Horne-Zeilinger (GHZ) –like state [16] between all the groups. When a group with less than j members is formed, it cannot distill a MES with any other group [see Fig. 2(b)]. We call this effect clustering, since the parties have to form clusters with at least j members in order to be able to create MES. In a similar way, one can also choose the state such that not only the number but also the specific parties that have to form groups in order to create a MES is given.

In the preceding examples we have that some number of parties have to join into groups in order to distill some MES with some other parties. In the following, we will give examples in which the parties that join enable the remaining ones to distill entanglement.

Example V. The state ρ_V is such that once any $(N-2)$ parties form a group, the remaining two parties can distill a




FIG. 2. (a) $N=20$, $j=3$, and ρ_{IV} is such that entanglement between any two groups that contain more or equal to $j=3$ members can be distilled (those groups can thus share a GHZ state), while groups with less than three members cannot create entanglement with any other group. (b) $N=10$ and ρ_V is such that once any $(N-2)$ parties join, the remaining two can create entanglement, while no entanglement can be shared among the remaining parties if less than $(N-2)$ parties join.

MES, but ρ_V remains undistillable if less than $(N-2)$ parties join [see Fig. 2(b)].

Example VI. ρ_{VI} is a state of $N=4$ parties for which once the parties (A_3A_4) form a group a GHZ-like state can be distilled among A_1, A_2 , and the group (A_3A_4) , whereas it is undistillable whenever any other parties but (A_3A_4) are joint [see Fig. 3(a)]. In contrast to the previous example, in this one it is not only possible to create a MES between A_1 and A_2 by joining (A_3A_4) , but also this last group can distill a MES with the remaining parties.

Example VII. ρ_{VII} is a state of $N=5$ parties such that a MES between parties A_1 and A_2 can be created if and only if either the parties (A_3A_4) or (A_3A_5) join [see Fig. 3(b)], but no entanglement can be distilled if the parties (A_4A_5) join.

In the following, we will explain how all those examples can be constructed and understood by giving necessary conditions for distillation and activation in multiparticle systems. These conditions also provide us with the tools to construct other examples of activable BES. We will also introduce a family of states that includes all examples I–VII as well as all those examples that can be constructed using




FIG. 3. (a) $N=4$ and ρ_{VI} is such that a GHZ state shared among the groups $A_1-A_2-(A_3A_4)$ can be created if and only if the parties (A_3A_4) join, and no entanglement can be distilled if any other two parties form a group. (b) $N=5$ and ρ_{VII} is such that entanglement between A_1 and A_2 can be created if and only if either party (A_3A_4) or (A_3A_5) join, however no entanglement can be distilled if (A_4A_5) joins.

the rules following from the necessary conditions for distillation and activation, which are also sufficient for this family.

III. DEFINITIONS AND NOTATION

In this section we first review the concept of bipartite splittings, and introduce some notation that will be used in the following sections. Then we review the properties of the family of N -qubit states ρ_N introduced in Ref. [12]. As mentioned above, these states give rise to a wide range of activable BES. In particular, among them one can find states corresponding to the examples of activable BES introduced in the previous section.

A. Bipartite splittings

Let us denote by \mathcal{P} the set of all possible (bipartite) splittings of N parties into two groups. For example, for three parties, \mathcal{P} contains the splittings $(A_1A_3)-(A_2)$, $(A_2A_3)-(A_1)$, and $(A_3)-(A_1A_2)$. We will denote these bipartite splittings by P_k , where $k=k_1k_2\cdots k_{N-1}$ is a chain of $N-1$ bits, such that $k_n=0,1$ if the n th party belongs to the same group as the last party or not. For example, for three parties the splittings $(A_1A_3)-(A_2)$, $(A_2A_3)-(A_1)$, and $(A_3)-(A_1A_2)$ will be denoted by P_{01}, P_{10} , and P_{11} , respectively. We will denote by A the side of the splitting to which the party N belongs and by B the other side. In general, there exist $s=2^{N-1}-1$ of such splittings. In the following, when we consider bipartite splittings, the parties in each of the groups will be allowed to act together (i.e., to perform joint operations).

B. Family of states ρ_N

Let us consider ρ_N , the family of N -qubit states introduced in Ref. [12]. We have that $\rho \in \rho_N$ if it can be written as

$$\rho = \sum_{\sigma=\pm} \lambda_0^\sigma |\Psi_0^\sigma\rangle\langle\Psi_0^\sigma| + \sum_{k \neq 0} \lambda_k (|\Psi_k^+\rangle\langle\Psi_k^+| + |\Psi_k^-\rangle\langle\Psi_k^-|), \quad (1)$$

where

$$|\Psi_k^\pm\rangle \equiv \frac{1}{\sqrt{2}} (|k_1k_2\cdots k_{N-1}0\rangle \pm |\bar{k}_1\bar{k}_2\cdots\bar{k}_{N-1}1\rangle) \quad (2)$$

are GHZ-like states with $k=k_1k_2\cdots k_{N-1}$ being a chain of $N-1$ bits, and $\bar{k}_i=0,1$ if $k_i=1,0$, respectively. We have that ρ_N is parametrized by 2^{N-1} independent real numbers. The labeling is chosen such that $\Delta \equiv \lambda_0^+ - \lambda_0^- \geq 0$. As we will see below, both the separability and distillability properties of the states belonging to this family are completely determined by the coefficients

$$s_k \equiv \begin{cases} 1 & \text{if } \lambda_k < \Delta/2 \\ 0 & \text{if } \lambda_k \geq \Delta/2. \end{cases} \quad (3)$$

Let us emphasize that the notation used for the states of this family parallels the one used to denote the partitions P_k . In

fact, as shown in Ref. [15] the separability properties of ρ_N for a given partition P_k are directly related to the coefficient s_k :

Lemma 0. [15] For any bipartite splitting $P_k \in \mathcal{P}$, and $\rho \in \rho_N$ we have $\rho^{TA} \geq 0 \Leftrightarrow s_k = 0 \Leftrightarrow \rho$ is separable with respect to this splitting.³

Thus the coefficient s_k determines whether ρ is separable or not with respect the bipartite splitting P_k . Note that there are no restrictions to the values of these coefficients; that is, for any choice of $\{s_k\}$ there always exists a state $\rho \in \rho_N$ with these values. This automatically implies that the family ρ_N provides us with all possible examples of states in which the separability properties of the bipartite splittings are completely specified. As we will see in the next section, the same is true regarding distillability with respect to bipartite splittings.

IV. ACTIVATION OF BE FOR BIPARTITE SPLITTINGS: EXAMPLES I-III

With the states introduced in the previous section, we are at the position of examining examples I–III. If we take a closer look at them, we find that they all have a common feature: the states ρ_I, ρ_{II} , and ρ_{III} are distillable with respect to certain bipartite splittings, and not distillable with respect to other bipartite splittings. In this section we will show that given a subset of all possible bipartite splittings we can always find states in ρ_N such that they are distillable if and only if the parties join according to a bipartite splitting contained in this set. This general result clearly allows to find states within our family that correspond to examples I–III.

Let us be more specific. We consider the set of all possible bipartite splittings \mathcal{P} . Let us specify for each splitting P_k whether we want that one can distill a MES or not. To do that, we will assign to each splitting a 0 if we do not want it to be possible and a 1 otherwise. That is, each possible specification corresponds to a function $f: \mathcal{P} \rightarrow \{0,1\}$. There are 2^s of such functions, which we will call specifications. Now, given a specification f we define the set of splittings $\mathcal{S}_f = \{P \in \mathcal{P} \text{ such that } f(P) = 1\}$. Thus, the problem reduces to finding a state ρ such that only for the splittings contained in \mathcal{S}_f can one distill a maximally entangled state. Note that examples I–III are just different specifications f . What we will show here is that there exist states $\rho \in \rho_N$ fulfilling any given specification.

A. Family of states ρ_N

We will show here that for the family of states ρ_N , bipartite inseparability is equivalent to distillability. This equivalence is expressed in the following Lemma:

Lemma 1. Given the bipartite splitting $P_k \in \mathcal{P}$ and $\rho \in \rho_N$, we have that ρ is distillable with respect to $P_k \Leftrightarrow s_k = 1$.

³ ρ^{TA} denotes the partial transposition with respect to the parties A . For the definition of partial transposition in multiparticle systems see Refs. [9] and [15]. The relation between subsystem A and P_k is given in Sec. III A.

Proof. Using Lemma 0 we have that if $s_k = 0$ then $\rho^{TA} \geq 0$, and therefore the state is not distillable, since nonpositive partial transposition is a necessary condition for distillation [1]. Thus, we just have to show that if $s_k = 1$ then the state is distillable. We denote by $|0\rangle_{A,B}$ and $|1\rangle_{A,B}$ the states in which all qubits in side A or B are in state 0 and 1, respectively. We have that $|\Psi_0^\pm\rangle = 1/\sqrt{2}(|0\rangle_A|0\rangle_B \pm |1\rangle_A|1\rangle_B)$ and $|\Psi_k^\pm\rangle = 1/\sqrt{2}(|0\rangle_A|1\rangle_B \pm |1\rangle_A|0\rangle_B)$. By measuring the projectors $|0\rangle_{A,B}\langle 0| + |1\rangle_{A,B}\langle 1|$, in A and B , respectively, we only get contributions from the states $|\Psi_0^\pm\rangle$ and $|\Psi_k^\pm\rangle$ and one obtains that the state after a successful measurement is

$$\rho \propto \lambda_0^+ |\Psi_0^+\rangle\langle \Psi_0^+| + \lambda_0^- |\Psi_0^-\rangle\langle \Psi_0^-| + \lambda_k (|0,1\rangle\langle 0,1| + |1,0\rangle\langle 1,0|). \quad (4)$$

This state is known to be distillable for $s_k = 1$ [2]. \square

This Lemma tells us that for a given specification we just have to find a state in ρ_N such that $s_k = f(P_k)$ for all k . Since the values s_k are not restricted in any way, we have that for any specification, we can find states $\rho \in \rho_N$ fulfilling it, and thus our family provides all different kinds of bipartite distillable and nondistillable states.

Let us now come back to examples I–III. We can take as state ρ_I [example I] one from the family ρ_N that has $s_k = f(P_k) = 1$ if and only if the number of ones in k is j or $(N-j)$ and $s_k = f(P_k) = 0$ otherwise (this means that all bipartite splittings that contain exactly j members in one group are distillable, and all others are separable). In the example II, we choose $\rho_{II} \in \rho_N$ such that $s_k = f(P_k) = 1$ if and only if P_k has between 40% and 60% of the parties in B . In example III we take $\rho_{III} \in \rho_N$ such that $s_k = f(P_k) = 1$ only for one specific P_k . We also emphasize that in a similar way one may construct many other interesting examples.

V. ACTIVATION OF BE BY EXTERNAL ACTION: EXAMPLES IV–VII

While the separability and distillability properties with respect to bipartite splittings of a state ρ were sufficient to completely understand and construct examples I–III, in the remaining examples we are faced with a slightly more complicated situation. Now we have that the parties can join into more than two groups, and it may even happen that not all of the parties are needed in order to activate the BE. Thus, the bipartite splittings P_k do no longer provide a complete description of the problem. However, we can still use them to derive *necessary* conditions regarding distillability and even activation of BE between any number of groups. And what is more important, we will show that these necessary conditions turn out to be also *sufficient* for the family of states ρ_N . On the one hand, this allows us to construct various different kinds of activable BES, such as those corresponding to examples IV–VII. On the other hand, it ensures that the family ρ_N provides examples for all possible kinds of BES that can be constructed by using the necessary conditions for distillation based on the bipartite properties of a state ρ . We wish to emphasize that due to the fact that the conditions we obtain

are only necessary and not sufficient in general, there may exist other kinds of activable BES that cannot be obtained with the methods developed here. Nevertheless, these methods also give indications about some other kinds of BES.

A. Necessary conditions for distillation

The separability properties of the bipartite splittings P_k of a multiparticle state ρ provide necessary conditions for the distillability properties of ρ . This is expressed in the following result:

Theorem 1. Let $C = \{A_{i_1}, \dots, A_{i_M}\}$ and $D = \{A_{j_1}, \dots, A_{j_L}\}$ be two disjoint groups of M and L parties, respectively, whereas the rest of the parties are separated. If a MES between C and D can be distilled then ρ has to be nonseparable with respect to all those bipartite splittings P_k in which the groups C and D are located on different sides.

Proof. Let us assume that ρ is separable with respect to one of those bipartite splittings P_k , so that $C \subset A$ and $D \subset B$. This means that even if we allow the groups C and D to join some other parties (belonging to A and B , respectively), they will not be able to distill a MES. This is due to the fact that nonseparability is a necessary condition for distillability. \square

Theorem 1 relates the distillability properties of ρ to the classification with respect to the separability properties of a multiparticle state given in Ref. [15]. We also note here that the k -separability properties with respect to k -partite splittings ($k > 2$) provide no additional information about the distillability properties of a state ρ . Theorem 1 also determines the necessary conditions for the creation of GHZ-like states, since the possibility of creating maximally entangled pairs between any two out of l parties is a necessary and sufficient condition for the creation of a GHZ-like state among those l parties. We can also change "nonseparable" in the theorem to "distillable," which provides an even stronger condition. It is not clear whether this condition is then also sufficient. One may think of the existence of bound entangled states that are distillable with respect to all possible bipartite splittings, but that are not distillable when considering the parties independently. It is not known whether such states exist; see, however, [17].

We also see from Theorem 1 that in order for a MES between two specific, separated parties, to be distillable, it is necessary that the corresponding state is inseparable with respect to at least 2^{N-2} different bipartite splittings. Thus there exist many states that are inseparable, but do not fulfill the necessary condition for distillability between any two parties. All those states are obviously BE. For example, any state that has less than 2^{N-2} (and more than one) inseparable bipartite splittings is clearly bound entangled. Naturally, the question arises under which conditions this BE can be activated.

B. Necessary conditions for activation of BES

Clearly, the situation changes when the parties are allowed to form groups and act together. In this case, the parties may be able to change the separability properties of certain bipartite splittings P_k . For example, for $N=3$ parties

A, B , and C , if the parties A and B form a group they may be able to change the separability properties of the bipartite splittings (i) $(B)-(AC)$ and (ii) $(A)-(BC)$. This is due to the fact that "joining" is equivalent to having some extra entanglement available. In this case, this extra entanglement between A and B can be used to change the separability properties of the bipartite splittings in question. Note, however, that this extra entanglement does not help to change the separability properties of the bipartite splitting (iii) $(C)-(AB)$, since (AB) were already joint in this bipartite splitting. This also allows us to understand the example of an activable BE three party state given in the Introduction [12], where we had that (i) and (ii) are separable, while (iii) is inseparable and the state is thus BE according to Theorem 1. By joining the parties A and B , one may however change the bipartite splittings (i) and (ii) from separable to inseparable, so the necessary conditions for distillation may now be fulfilled, since all bipartite splittings can now be inseparable in principle. As shown in Ref. [12], this activation can in fact be achieved, i.e., the change of the separability properties of the splittings (i) and (ii) as well as the distillation are possible. In a similar way, one can explain the example given in Ref. [13] for $N=4$. The effect of activation of BE for a state ρ can thus be viewed as a consequence of the following theorem:

Theorem 2. Consider a state ρ that is separable with respect to a given bipartite splitting $(A)-(B)$. When joining M parties $C = \{A_{i_1}, \dots, A_{i_M}\}$, necessary conditions that we can make ρ distillable with respect to the splitting $(A)-(B)$ are that (i) $C \subset A, B$ and (ii) by using an operation acting on C one can transform the state such that it is now nonseparable with respect to the bipartite splitting $(A)-(B)$.

Proof. (ii) follows trivially from Theorem 1, whereas (i) follows from (ii). \square

According to this theorem, when joining some parties into a group C , they may change the separability and distillability properties of certain bipartite splittings, namely, all those splittings $(A)-(B)$ for which $C \subset A, B$. So, it may happen that the conditions for distillability, which were not fulfilled before joining the parties, are now fulfilled, and a MES shared between some parties can now be created in principle. Theorems 1 and 2 together provide necessary conditions for the activation of multiparticle BES and provide thus the framework for the construction of generic examples of different kinds of activable bound entangled states. To achieve this, one chooses at the beginning the separability and distillability of the bipartite splittings P_k of a state ρ such that the state is not distillable if the parties remain separated from each other, but that the distillability conditions may be fulfilled when some of the parties are allowed to form groups. We also note here that due to the fact that both theorems only provide necessary conditions in general, we do not have that the distillability and activation properties of a state ρ are completely determined by the separability properties of its bipartite splittings. On the one hand, there might exist states that cannot be distilled or activated even though they fulfill the necessary conditions for distillation/activation, i.e., those states are further protected against activation. On the other

hand, we already see that there are various kinds of nonactivable bound entangled states. For example, all states that are biseparable with respect to all bipartite splittings but are inseparable with respect to any k -partite splitting ($k > 2$) (see also classification proposed in Ref. [15]) are clearly bound entangled and can be neither distilled nor activated. For $N = 3$, such an example is known [18].

Using Theorems 1 and 2, it is now straightforward to check that in example I the state ρ_I remains undistillable whenever the parties form more than two groups, since the necessary condition for distillation of a MES between any two groups cannot be fulfilled. On the other hand, it is already clear that if the parties form two groups with a different number of members than j and $N-j$, we have by construction that the state $\rho_I \in \rho_N$ is separable with respect to this bipartite splitting and thus not distillable. In a similar way, one can check that in examples II and III the states ρ_{II} and ρ_{III} are only distillable if and only if the parties join in two groups of required size (example II) or required members (example III), respectively.

C. Family of states ρ_N

In this section, we show that the necessary condition for distillation and activation expressed in Theorems 1 and 2 are also sufficient for the family of states ρ_N . We first show that if the necessary conditions for the distillation of MES between any two groups of parties, both not including a certain party A_l , are fulfilled, one can disentangle party A_l from the remaining system while keeping the necessary conditions for distillation between the two groups in question. In order to achieve this, the party A_l has to cooperate, i.e., its help is required. This is expressed in the following lemma:

Lemma 2. One can convert any N -qubit state $\rho \in \rho_N$ to a $(N-1)$ -qubit state $\tilde{\rho} \in \rho_{N-1}$ by measuring a certain party A_l , such that for all bipartite splittings P_k the following property is fulfilled: If ρ is inseparable with respect to the bipartite splittings $(A_l C)$ -(rest) and (C) -(A_l rest) $\Rightarrow \tilde{\rho}$ is inseparable with respect to the bipartite splitting (C) -(rest).

Proof. We assume without loss of generality that $A_l = A_1$. If we measure in A_1 the projector $P_+ = |+\rangle_{A_1}\langle +|$ where $|+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$, we find that the remaining $(N-1)$ particles are in a state of the form ρ_{N-1} with new coefficients $\tilde{\lambda}_{j_2 j_3 \dots j_{N-1}} = \lambda_{0 j_2 j_3 \dots j_{N-1}} + \lambda_{1 j_2 j_3 \dots j_{N-1}}$ and similar for λ_0^\pm , so that $\tilde{\Delta} = \Delta$. The property we want to show is simply that if and only if both

$$\lambda_{0 j_2 j_3 \dots j_{N-1}} < \Delta/2 \text{ and } \lambda_{1 j_2 j_3 \dots j_{N-1}} < \Delta/2, \quad (5)$$

it follows that

$$\tilde{\lambda}_{j_2 j_3 \dots j_{N-1}} < \tilde{\Delta}/2. \quad (6)$$

It may happen that although Eq. (5) is fulfilled, Eq. (6) is not. In this case, we apply the first step of the distillation procedure proposed in Ref. [15] first, where one measures certain POVM elements on M copies of the initial state and is left with a new (unnormalized) state of the form ρ_N with new

coefficients $\lambda'_k = \lambda_k^M$ and $(\Delta'/2) = (\Delta/2)^M$. For M sufficiently large, we can now always have that after applying P_+ , the remaining state is such that Eq. (6) is fulfilled, since the new $\Delta/2$ is exponentially amplified compared to any $\lambda_{k_{1,2}} < \Delta/2$ and thus $(\Delta/2)^M > \lambda_{k_1}^M + \lambda_{k_2}^M$ for M sufficiently large. \square

Theorem 3. For the family of states ρ_N , we have that the necessary condition for distillability given in Theorem 1 is also sufficient.

Proof. To show this, one just has to sequentially apply Lemma 2 to all particles $A_i \in \{C, D\}$, which leaves us with a $(M+L)$ qubit-state $\rho \in \rho_{M+L}$, which has $\rho^{Tc} \geq 0$ (which, according to Lemma 0 means that the corresponding $s_k = 1$) and thus Lemma 1 can be applied, ensuring that any state $\rho \in \rho_N$ fulfilling the necessary conditions for any kind of distillation is in fact distillable. \square

Note that the help of *all* parties is required, independent of whether they finally share a MES or not. This follows from the fact that Lemma 2 is based on the cooperation of party A_l , which is separated from the remaining system after the procedure described above. If one would just trace out party A_l (i.e., the party does not cooperate with the remaining parties), one finds that the remaining parties cannot create any entanglement at all.

Theorem 4. For $\rho \in \rho_N$, and the situation of Theorem 2, we can in fact change *all* those bipartite splittings (A) -(B) for which $C \not\subset A, B$ from separable to inseparable (which is equivalent to distillable in this case) without changing the separability properties of the remaining bipartite splittings.

Proof. We assume without loss of generality that we join the first M parties $C = \{A_1, \dots, A_M\}$. Let $j = j_1 j_2 \dots j_M$ ($l = l_1 \dots l_{N-M-1}$) be $M[(N-M-1)]$ digit binary numbers. We have to show that we can change all those bipartite splittings that do not contain all parties C on one side from separable to inseparable. This is equivalent to showing that for all λ_{j_l} with $j \neq \{0, 2^M - 1\}$, we can have $\Delta/2 > \lambda_{j_l}$. When applying the POVM element

$$\tilde{P} = \sum_{j=0}^{2^M-1} \sqrt{y_j} |j\rangle_C \langle j|,$$

we find that we obtain again an (unnormalized) state of the form ρ_N with new coefficients $\tilde{\lambda}_{j_l} = y_j \lambda_{j_l}$. Choosing $y_0 = y_{2^M-1} = 1$ and all other y_j sufficiently small, we have that $\tilde{\Delta} = \Delta$ and for $j \neq \{0, 2^M - 1\}$ we can obtain that $\tilde{\Delta}/2 > \tilde{\lambda}_{j_l}$ as required. Furthermore, all other relations $\Delta/2 \times \lambda_{j_l}$, with $\times \in \{>, \leq\}$ and thus the separability properties of all those bipartite splittings for which $C \subset A, B$ remain unchanged. \square

Theorems 3 and 4 together ensure that any BES within the family ρ_N that is activable in principle can in fact be activated, i.e., the necessary conditions for the activation of bound entangled states given in Theorem 1 and 2 are also sufficient for the family ρ_N .

D. Examples IV–VII

We are now at the position to construct and explain examples IV–VII, as well as to provide many other interesting examples of activable BES.

Let us start with example IV: In this case, we choose the state $\rho_{IV} \in \rho_N$ such that it is separable with respect to all bipartite splittings P_k where either the group A or B has less than j members. All other bipartite splittings are chosen to be distillable. It is clear that if a group with less than j members is formed, they cannot distill a MES with any other group (since the corresponding bipartite splitting is separable). However once the parties form two groups, each having more than j members, it is straightforward to check (using Theorems 4 and 3) that these two groups can in fact create a MES.

In example V, we choose the state $\rho_V \in \rho_N$ such that it is distillable with respect to all bipartite splittings P_k where either the group A or B contains exactly one particle (so that the number of ones in k is either 1 or $N-1$), and is separable with respect to all other bipartite splittings P_k . It is clear that it is sufficient to join any $N-2$ particles, since this allows—according to Theorem 4—to make inseparable (and thus distillable) all those bipartite splittings where the remaining particles are located in different groups, so that distillation of a MES between the remaining two particles becomes possible, according to Theorem 3. On the other hand, if less than $N-2$ parties join, one can easily verify that for any two of the remaining parties, there always remains at least one bipartite splitting separable that has to be inseparable in order that distillation can be possible. Thus the remaining parties cannot share entanglement if less than $N-2$ parties join.

In example VI we choose $\rho_{VI} \in \rho_4$ such that it is inseparable with respect to the bipartite splittings $(A_1A_2)-(A_3A_4)$, $(A_1)-(A_2A_3A_4)$, and $(A_2)-(A_1A_3A_4)$ and separable with respect to all other bipartite splittings. Clearly, this state is BE since at least $2^{N-2} = 2^2 = 4$ bipartite splittings have to be inseparable so that a pair between any two separated parties can be distilled. Furthermore, one can check (using Theorems 3 and 4) that ρ_{VI} remains undistillable when joining any two parties but (A_3A_4) , but one can create a GHZ state shared among $A_1-A_2-(A_3A_4)$ once one joins the parties (A_3A_4) .

Finally, in example VII we have $N=5$ and choose $\rho_{VII} \in \rho_5$ such that the state is inseparable with respect to all bipartite splittings that contain A_1 and A_2 in different groups, except the splitting $(A_1A_3)-(A_2A_4A_5)$ which is chosen to be separable as well as all other bipartite splittings. One can readily check that ρ_{VII} has the desired properties, i.e., it is BE and can be activated when joining the parties (A_3A_4) or (A_3A_5) .

If we demand however that entanglement between A_1 and A_2 should be created once *any* two of the remaining parties

join, one finds that such a state cannot be constructed using our method. In this case, we have to demand that the initial state $\rho_{VII} \in \rho_N$ has to be separable with respect to at least one of the bipartite splittings where A_1 and A_2 belong to different groups. Otherwise, a MES between A_1 and A_2 could be distilled from the beginning and the state is thus not BE. Let us assume without loss of generality that the separable splitting is either (i) $(A_1A_3)-(A_2A_4A_5)$ or (ii) $(A_1)-(A_2A_3A_4A_5)$. In both cases, the separability properties of this splitting cannot be changed when joining the parties (A_4A_5) and so no entanglement can be created between A_1 and A_2 although two of the remaining parties join (note that if we had taken any other bipartite splitting, with A_1 and A_2 belonging to different groups, to be separable, we would have found some other two of the remaining parties that can join but not change the properties of this splitting).

Due to the fact that the conditions for distillation and activation we give here are only necessary in general, they do not allow us to rule out the possibility that a state having the desired properties can exist. In this case, the activation would not be based on the change of the separability properties of the bipartite splittings, but on some other mechanism.

VI. SUMMARY

In summary, we have given rules to construct activable bound entangled states using the separability and distillability properties of a density operator with respect to bipartite splittings. This method allows us to construct examples for all possible kinds of activable BES where the parties join into exactly two groups. In particular, the family of states introduced in Ref. [12] contains examples of all these kinds of activable BES. We have also given some relevant examples of activation of BE in which the parties join into more than two groups, and where the role of some of the groups is just to help the others to create a MES.

ACKNOWLEDGMENTS

We thank G. Vidal and J. Smolin for discussions. This work was supported by the Austrian Science Foundation under the SFB “control and measurement of coherent quantum systems (Project 11),” the European Community under the TMR network ERB-FMRX-CT96-0087, the European Science Foundation, and the Institute for Quantum Information GmbH.

-
- [1] P. Horodecki, Phys. Lett. A **232**, 333 (1997).
 [2] C.H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J.A. Smolin, and W.K. Wootters, Phys. Rev. Lett. **76**, 722 (1996); C.H. Bennett, H.J. Bernstein, S. Popescu, and B. Schumacher, Phys. Rev. A **53**, 2046 (1996).
 [3] C.H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W.K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
 [4] N. Linden and S. Popescu, Fortschr. Phys. **46**, 567 (1998).

- [5] C.H. Bennett, D.P. DiVincenzo, J.A. Smolin, and W.K. Wootters, Phys. Rev. A **54**, 3824 (1996); H.-J. Briegel, W. Dür, J.I. Cirac, and P. Zoller, Phys. Rev. Lett. **81**, 5932 (1998).
 [6] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **82**, 1056 (1999).
 [7] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Rev. Lett. **80**, 5239 (1998).
 [8] There are strong indications which suggest that this condition

is not sufficient, i.e., there are states with a nonpositive partial transposition that are BE; see W. Dür, J.I. Cirac, M. Lewenstein, and D. Bruss, Phys. Rev. A **61**, 062313 (2000); D.P. DiVincenzo, P.W. Shor, J.A. Smolin, B.M. Terhal, and A.V. Thapliyal, *ibid.* **61**, 062312 (2000).

- [9] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).
- [10] M. Horodecki, P. Horodecki, and R. Horodecki, Phys. Lett. A **223**, 8 (1996).
- [11] D. Bruss and A. Peres, Phys. Rev. A **61**, 030301 (R) (2000); D.P. DiVincenzo, Tal Mor, P.W. Shor, J.A. Smolin, B.M. Terhal, e-print quant-ph/9908070 (unpublished); P. Horodecki, J.A. Smolin, B.M. Terhal, A.V. Thapliyal, e-print quant-ph/9910122 (unpublished); P. Horodecki, M. Lewenstein, e-print quant-ph/0001035 (unpublished).
- [12] W. Dür, J.I. Cirac, and R. Tarrach, Phys. Rev. Lett. **83**, 3562 (1999).
- [13] J.A. Smolin, e-print quant-ph/0001001 (unpublished).
- [14] R. Cleve, D. Gottesman, and Hoi-Kwong Lo, Phys. Rev. Lett. **83**, 648 (1998); D. Gottesman, e-print quant-ph/9910067 (unpublished); M. Hillery, V. Bužek, and A. Berthiaume, Phys. Rev. A **59**, 1829 (1999).
- [15] W. Dür, and J.I. Cirac, Phys. Rev. A **61**, 042314 (2000).
- [16] D. M. Greenberger, M. Horne, and A. Zeilinger, *Bell's Theorem, Quantum Theory, and Conceptions of the Universe*, edited by M. Kafatos (Kluwer, Dordrecht, 1989), p. 69; D. Bouwmeester *et al.*, Phys. Rev. Lett. **82**, 1345 (1999).
- [17] J. A. Smolin and A. V. Thapliyal (unpublished).
- [18] C.H. Bennett, D.P. DiVincenzo, Tal Mor, P.W. Shor, J.A. Smolin, and B.M. Terhal, Phys. Rev. Lett. **82**, 5385 (1999).