

Continuous-variable teleportation of single-photon states

Toshiki Ide,¹ Holger F. Hofmann,² Takayoshi Kobayashi,¹ and Akira Furusawa³

¹*Department of Physics, Faculty of Science, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

²*CREST, Japan Science and Technology Corporation (JST), Research Institute for Electronic Science, Hokkaido University, Sapporo 060-0812, Japan*

³*Department of Applied Physics, Faculty of Engineering, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8656, Japan*
(Received 4 April 2001; revised manuscript received 30 July 2001; published 12 December 2001)

The properties of continuous-variable teleportation of single-photon states are investigated. The output state is different from the input state due to the nonmaximal entanglement in the Einstein-Podolsky-Rosen beams. The photon statistics of the teleportation output are determined and the correlation between the field information β obtained in the teleportation process and the change in photon number is discussed. The results of the output photon statistics are applied to the transmission of a qubit encoded in the polarization of a single photon.

DOI: 10.1103/PhysRevA.65.012313

PACS number(s): 03.67.Lx, 42.50.-p

I. INTRODUCTION

Quantum teleportation is a method for Alice (sender) to transmit an unknown quantum input state to Bob (receiver) at a distant place by sending only classical information using a shared entangled state as a resource. Originally quantum teleportation was proposed for discrete variables in two-dimensional Hilbert spaces [1]. Later, it was applied to continuous variables (two components of the electromagnetic field) in infinite-dimensional Hilbert spaces [2]. However, continuous-variable teleportation ideally requires a maximally entangled state that has infinite energy as a resource. Nevertheless, it has been shown theoretically that quantum teleportation realized by using a nonmaximally entangled state may still transfer nonclassical features of quantum states [3]. Experimentally, such a continuous-variable teleportation has been realized by Furusawa *et al.* [4]. In [3], the physics of continuous-variable teleportation was described in terms of the Wigner function. Reference [5] described it in terms of discrete basis states. Reference [6] formulates the whole process of the quantum teleportation by a transfer operator that is acting on arbitrary input states.

In the experiment of [4], a coherent state was used as an input state. But any quantum state may be teleported by this method. Therefore, the transfer of nonclassical states is of interest. In the following, the transfer operator formalism derived in [6] is used for analyzing the photon statistics of the output state of a one-photon-state teleportation. It is shown that the change in photon number is strongly dependent on the field measurement result obtained in the process of teleportation. This result is then applied to the two-mode teleportation of a polarized photon, illustrating the possibility of using continuous-variable teleportation for the transfer of single photon qubits.

II. TRANSFER OPERATOR

Figure 1 shows the schematic sets of the quantum teleportation according to [4]. Alice transmits an unknown quantum state $|\psi\rangle_A$ to Bob. Alice and Bob share Einstein-Podolsky-Rosen (EPR) beams in advance. The quantum state of the EPR beams reads [5,6]

$$|q\rangle_{R,B} = \sqrt{1-q^2} \sum_{n=0}^{\infty} q^n |n\rangle_R |n\rangle_B. \quad (1)$$

where R, B are the modes for reference and Bob. q is a parameter that stands for the degree of entanglement. It varies from zero to one, with one being maximal entanglement and zero being no entanglement. The degree of entanglement depends on the squeezing achieved in the parametric amplification. In the experiment of [4], 3dB ($q=0.33$) squeezed light was used. Squeezing of up to 10dB ($q=0.82$) should be possible with the available technology.

Alice mixes her input state with the reference EPR beam by a 50% beam splitter and performs an entanglement measurement of the complex field value $\beta = x_- + iy_+$, where

$$\begin{aligned} \hat{x}_- &= \hat{x}_A - \hat{x}_R, \\ \hat{y}_+ &= \hat{y}_A + \hat{y}_R. \end{aligned} \quad (2)$$

This measurement projects A and R onto the eigenstate

$$|\beta\rangle_{A,R} = \frac{1}{\sqrt{\pi}} \sum_{n=0}^{\infty} \hat{D}_A(\beta) |n\rangle_A |n\rangle_R, \quad (3)$$




FIG. 1. Schematic representation of the quantum teleportation setup.

where $\hat{D}_A(\beta)$ is a displacement operator acting on the mode A with a displacement amplitude of β . The output state $|\psi(\beta)\rangle_B$ conditioned by the measurement process may be written as a projection of the initial product state onto the eigenstate $|\beta\rangle_{A,R}$,

$$\begin{aligned} |\psi(\beta)\rangle_B &= {}_{A,R}\langle\beta|\psi\rangle_A|q\rangle_{R,B} \\ &= \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n |n\rangle_{BA} \langle n|\hat{D}_A(-\beta)|\psi\rangle_A. \end{aligned} \quad (4)$$

$|\psi(\beta)\rangle_B$ is not normalized, since the probability of obtaining the field measurement value β is given by $P_q(\beta) = \langle\psi_{\text{out}}(\beta)|\psi_{\text{out}}(\beta)\rangle$.

After Bob gets the information of the field measurement value β from Alice, Bob applies a displacement to the output state by mixing the coherent field of a local oscillator with the output EPR beam B . The output state $|\psi_{\text{out}}(\beta)\rangle_B = \hat{D}_B(\beta)|\psi(\beta)\rangle_B$ may also be written as

$$|\psi_{\text{out}}(\beta)\rangle_B = \hat{T}_q(\beta)|\psi\rangle_A, \quad (5)$$

where $\hat{T}_q(\beta)$ is a transfer operator that represents all processes of the quantum teleportation [6]. In its diagonalized form, it reads

$$\hat{T}_q(\beta) = \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n \hat{D}(\beta)|n\rangle_{BA} \langle n|\hat{D}(-\beta). \quad (6)$$

When $q \rightarrow 1$, $\hat{T}_q(\beta)$ becomes similar to the equivalence operator $\hat{1}$ except for a change of the mode from A to B , which indicates a perfect teleportation with no modification to the input state. For general q , the transfer operator at $\beta=0$ is diagonal in the photon number states.

III. ONE-PHOTON-STATE TELEPORTATION

In the experiment of [4], a coherent state was transferred. In this case, the output state is modified but is still a coherent state [6]. In case of a one-photon input state, the output state is

$$\begin{aligned} \hat{T}_q(\beta)|1\rangle &= \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n \hat{D}(\beta)|n\rangle \langle n|\hat{D}(-\beta)|1\rangle \\ &= \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n \hat{D}(\beta)|n\rangle \langle n|\hat{D}(-\beta)\hat{a}^\dagger|0\rangle \\ &= \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n \hat{D}(\beta)|n\rangle \langle n|(\hat{a}^\dagger + \beta^*) \\ &\quad \times \hat{D}(-\beta)|0\rangle \\ &= \sqrt{\frac{1-q^2}{\pi}} e^{-(1-q^2)\frac{|\beta|^2}{2}} \hat{D}[(1-q)\beta] \\ &\quad \times [(1-q^2)\beta^*|0\rangle + q|1\rangle]. \end{aligned} \quad (7)$$




FIG. 2. The probability distribution of $P_q(\beta)$ for the field measurement value $\beta = x_- + iy_+$ for $q = 1/2$.

For simplicity, the suffixes for the modes are not explicitly given. In general, this output state is quite different from the original one-photon input state.

β represents a coherent field measurement performed on the input state. The probability distribution over field measurement values β for one-photon input states is given by

$$\begin{aligned} P_q(\beta) &= \langle\psi_{\text{out}}(\beta)|\psi_{\text{out}}(\beta)\rangle \\ &= \langle 1|\hat{T}_q^\dagger(\beta)\hat{T}_q(\beta)|1\rangle \\ &= \frac{1-q^2}{\pi} e^{-(1-q^2)|\beta|^2} [(1-q^2)^2|\beta|^2 + q^2]. \end{aligned} \quad (8)$$

Figure 2 shows the probability distribution $P_q(\beta)$ for $q = 1/2$. The circular symmetric distribution with a dip in the middle and a maximum for amplitude of $|\beta| = 1$ is characteristic of the one-photon input state.

The properties of the output state may be investigated experimentally by several kinds of detection setups. If Bob has a photon counting setup that may discriminate photon numbers, he may obtain the photon statistics of the output state. The overall photon statistics of the output are obtained by integrating over β ,

$$\begin{aligned} P_q(n) &= \int d^2\beta |\langle n|\hat{T}_q(\beta)|1\rangle|^2 \\ &= \frac{1+q}{2} \left(\frac{1-q}{2}\right)^{n+1} \left[1 + \left(\frac{1+q}{1-q}\right)^2 n\right]. \end{aligned} \quad (9)$$

$P_q(n)$ is the probability of counting n photons after the teleportation. Figure 3 shows the probability distribution over n in the case of $q = 1/2$. The probability of detecting one pho-




FIG. 3. The probability of n -photon counting $P_q(n)$ for one-photon-state teleportation in the case of $q=1/2$.

ton is maximal and the probabilities of higher-photon numbers get less likely as photon number increases.

The changes in photon number may be summarized in terms of photon loss $n=0$, successful photon transfer $n=1$, and photon gain $n \geq 2$. The q dependence of these probabilities is given by

$$P_q(0) = \frac{1}{4}(1 - q^2), \quad (10)$$

$$P_q(1) = \frac{1}{4}(1 + q + q^2 + q^3), \quad (11)$$

$$P_q(n \geq 2) = \frac{1}{4}(2 - q - q^3). \quad (12)$$

Since the input state is a one-photon state, Eq. (11) gives the fidelity of the teleportation. Figure 4 shows the q dependence of $P_q(0)$, $P_q(1)$, $P_q(n \geq 2)$. In the case of a maximally entangled state ($q \rightarrow 1$), Bob may receive nothing but a one-photon state [$P_q(1) \rightarrow 1$] and all other probabilities vanish




FIG. 4. The probability of zero photon, one photon, and more-than-one photon counting for one-photon-state teleportation.




FIG. 5. The probability of n -photon counting $P_q(n)$ for one-photon-state teleportation in the case of $q=1/2$. The thick solid line shows $P_{1/2}(\beta)$, the thin solid line shows $P_{1/2}(1, \beta)$, the dotted line shows $P_{1/2}(0, \beta)$ and the dot-bar line shows $P_{1/2}(n \geq 2, \beta)$.

[$P_q(0) \rightarrow 0, P_q(n \geq 2) \rightarrow 0$], which indicates perfect teleportation. In the case of nonmaximally entangled states ($q < 1$), the probabilities of zero-photon counting and more-than-one-photon counting appear. Note that the probability of photon gain $n \geq 2$ is always greater than that of photon loss $n=0$.

In order to investigate the change in the photon number given by Eq. (9) in more detail, we now derive the conditional probability distributions over β for the cases of zero-photon, one-photon, and more-than-one-photon counting. These probabilities may be obtained from $P_q(n, \beta) = |\langle n | \hat{T}_q(\beta) | 1 \rangle|^2$. The results read

$$P_q(0, \beta) = \frac{1 - q^2}{\pi} e^{-2(1-q)|\beta|^2} (1 - q)^2 |\beta|^2, \quad (13)$$

$$P_q(1, \beta) = \frac{1 - q^2}{\pi} e^{-2(1-q)|\beta|^2} [(1 - q^2)^2 |\beta|^2 + q^2], \quad (14)$$

$$P_q(n \geq 2, \beta) = P_q(\beta) - P_q(0, \beta) - P_q(1, \beta). \quad (15)$$

Figure 5 shows these different contributions to the total probability distribution over $|\beta|$ in case of $q=0.5$. For high values of $|\beta|$, every probability vanishes. In the $\beta=0$ case, we see that only the probability of obtaining a one-photon output is nonzero. This means that the output of the EPR beam B is already a one-photon state without any additional displacement. In general, the photon number states are the eigenstates of $\hat{T}_q(\beta=0)$, as can be seen from Eq. (6). A measurement value of $\beta=0$ thus indicates that the output beam photon number is equal to the input beam photon number. Note also that a low-field amplitude provides little information about the phase of the teleported field. Therefore, the lack of phase information in the input photon number state is preserved in the teleportation process.

In the region of $|\beta| \ll 1$, the probability of obtaining a one-photon output is nearly constant and that of zero photon

and more-than-one photon are increasing with the increase of $|\beta|$. The increase of these two probabilities gives rise to the peak of the total probability. The probability of photon loss ($n=0$) and photon gain ($n\geq 2$) are crossing at around $|\beta|=1$. Below $|\beta|=1$, the probability of photon loss ($n=0$) is greater than that of photon gain ($n\geq 2$). Above $|\beta|=1$, the probability of photon gain ($n\geq 2$) is dominant. For high-field measurement results $|\beta|\gg 1$, the teleportation process generates more photons in the output. Since β may be considered as a measurement of the coherent input field amplitude, this result is similar to the correlation of the field measurement and photon number discussed in [7].

IV. APPLICATION TO SINGLE-PHOTON POLARIZATION

Since the polarization of the light field may be used to encode information using single photons, the case of a polarization-sensitive continuous-variable teleportation is of considerable interest. The results obtained for the single-mode case may be applied to the teleportation of two polarization modes, H and V , by applying a separate transfer operator to each mode. Note that this does not imply that the teleportation H and V has to be conducted separately. For example, the two-dimensional measurement amplitude (β_H, β_V) could also be obtained by measuring the circular polarization components $(\beta_H \pm i\beta_V)$. The continuous-variable teleportation of polarized photons therefore does not require any previous knowledge of the input polarization and the results obtained for successful transfers and for polarization flips may be applied directly to the teleportation of an unknown photon polarization.

The output state of a one-photon state with polarization H may be written as a product of the one-photon teleportation given in Eq. (7) and a vacuum teleportation, which is a special case of the conventional coherent-state teleportation discussed in [6]. The result reads

$$\begin{aligned} & \hat{T}_{Hq}(\beta_H)\hat{T}_{Vq}(\beta_V)|1\rangle_H|0\rangle_V \\ &= \sqrt{\frac{1-q^2}{\pi}} \sum_{n=0}^{\infty} q^n \hat{D}_H(\beta_H)|n\rangle_{HH}\langle n|\hat{D}_H(-\beta_H)|1\rangle_H \\ & \otimes \sqrt{\frac{1-q^2}{\pi}} \sum_{m=0}^{\infty} q^m \hat{D}_V(\beta_V)|m\rangle_{VV}\langle m|\hat{D}_V(-\beta_V)|0\rangle_V \end{aligned}$$

$$\begin{aligned} P_{\text{trans}}(q) &= \underbrace{\int d^2\beta_H | \langle 1 | \hat{T}_{Hq}(\beta_H) | 1 \rangle_H |^2}_{=P_q(1)} \underbrace{\int d^2\beta_V | \langle 0 | \hat{T}_{Vq}(\beta_V) | 0 \rangle_V |^2}_{=\frac{1+q}{2}} \\ &= \left(\frac{1+q}{2}\right)^2 \frac{1+q^2}{2}. \end{aligned} \quad (17)$$

The total chance of a polarization flip while preserving the total photon number of one P_{flip} is equal to the product of the probabilities for a vacuum output in a one-photon teleportation $P_q(0)$, and the reverse situation where the vacuum input produces a one-photon output. Since $\hat{T}_q(\beta)$ is Hermitian, however, these two probabilities are equal and the result reads




FIG. 6. The q dependence of the probabilities of the polarized photon teleportation.

$$\begin{aligned} &= \sqrt{\frac{1-q^2}{\pi}} e^{-(1-q^2)(|\beta_H|^2/2)} \hat{D}[(1-q)\beta_H] \\ & \times [(1-q^2)\beta_H^*|0\rangle_H + q|1\rangle_H] \\ & \otimes \sqrt{\frac{1-q^2}{\pi}} e^{-(1-q^2)(|\beta_V|^2/2)} \hat{D}[(1-q)\beta_V]|0\rangle_V. \end{aligned} \quad (16)$$

This result describes all details of single-photon teleportation, including the information obtained from the measurement result (β_H, β_V) . In the following, however, we will concentrate on the transmission errors induced by the teleportation. As will be discussed below, these results may be expressed entirely in terms of the probabilities $P_q(n)$ given by Eq. (9) and the well-known coherent-state fidelity of $(1+q)/2$.

The fidelity of the teleportation is the total chance of successfully transmitting a photon with the correct polarization P_{trans} . It is equal to the product of the probabilities for successfully teleporting a single-photon $P_q(1)$, and the probability of successfully teleporting the vacuum $(1+q)/2$ as given by

$$\begin{aligned}
P_{\text{flip}}(q) &= \underbrace{\int d^2\beta_H |_H\langle 0|\hat{T}_{Hq}(\beta_H)|1\rangle_H|^2}_{=P_q(0)} \underbrace{\int d^2\beta_V |_V\langle 1|\hat{T}_{Vq}(\beta_V)|0\rangle_V|^2}_{=P_q(0)} \\
&= \left(\frac{1+q}{2}\right)^2 \frac{1-q^2}{2}.
\end{aligned} \tag{18}$$

Finally, there are also probabilities for changes in the total photon number. The chance of obtaining no photon P_{zero} is equal to the product of the probabilities for a vacuum output in a one-photon teleportation $P_q(0)$, and for the successful teleportation of the vacuum given by the coherent-state fidelity $(1+q)/2$,

$$\begin{aligned}
P_{\text{zero}}(q) &= \underbrace{\int d^2\beta_H |_H\langle 0|\hat{T}_{Hq}(\beta_H)|1\rangle_H|^2}_{=P_q(0)} \underbrace{\int d^2\beta_V |_V\langle 0|\hat{T}_{Vq}(\beta_V)|0\rangle_V|^2}_{=\frac{1+q}{2}} \\
&= \left(\frac{1+q}{2}\right)^2 \frac{1-q}{2}.
\end{aligned} \tag{19}$$

The total chance of obtaining more than one photon in the output may then be obtained by

$$\begin{aligned}
P_{n \geq 2}(q) &= 1 - P_{\text{flip}}(q) - P_{\text{trans}}(q) - P_{\text{zero}}(q) \\
&= 1 - \left(\frac{1+q}{2}\right)^2 \frac{5-4q+3q^2}{4}.
\end{aligned} \tag{20}$$

Figure (6) shows the q dependence of the above probabilities. The fidelity P_{trans} increases with increasing entanglement q , while the probabilities of the various error sources decrease. P_{trans} exceeds 1/2 around $q \sim 0.7$ and 2/3 around $q \sim 0.8$, illustrating that the entanglement requirements for high-fidelity single-photon transfers could be fulfilled using the best squeezing sources presently available. The dominant source of error is the chance of generating additional photons $P_{n \geq 2}$, while the probability of flipping a polarization P_{flip} is always significantly lower than all the other probabilities. Therefore, the photon loss and gain processes are a more serious problem than the flip of a polarization for the transmission of the qubit.

It is interesting to compare this type of error with the postselection problems inherent in the previously realized teleportation by entangled photon pairs [8] using coincidence counting as a trigger. In particular, the fidelity of continuous-variable teleportation may also be increased by postselecting the one-photon outputs, eliminating the multiphoton and photon-loss errors. The conditional fidelity is now given by

$F_{\text{cond}} = P_{\text{trans}} / (P_{\text{trans}} + P_{\text{flip}})$. This fidelity is already 2/3 at $q = 0$ and reaches 10/11 at $q = 1/2$. While such postselection may be difficult to realize due to the limited quantum efficiency and saturation characteristics of conventional photon detectors, it illustrates how well the polarization property itself is preserved in the continuous-variable teleportation.

V. CONCLUSION

The properties of continuous-variable teleportation of single-photon states have been investigated. The difference between the input state and the output state is due to the nonmaximal entanglement in the EPR beams shared by Alice and Bob. The field measurement conditions the output state. Nearly zero values of the field measurement tend to preserve the initial one-photon input state because the eigenstates of the field measurement are close to a photon number states. In the intermediate range of field measurement values, photon loss and photon gain processes occur during teleportation. At high values of the field measurement, the probability of photon gain is dominant, corresponding to the high amplitude observed. An application of this analysis to the teleportation of a polarized photon shows that the photon loss and gain processes are a more serious problem than the polarization flips. The results imply that unconditional continuous-variable teleportation of single-photon polarization could be considered an alternative to the postselected scheme [8] using entangled photon pairs.

- [1] C.H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres, and W.K. Wootters, Phys. Rev. Lett. **70**, 1895 (1993).
- [2] L. Vaidman, Phys. Rev. A **49**, 1473 (1994).
- [3] S.L. Braunstein and H.J. Kimble, Phys. Rev. Lett. **80**, 869 (1998).
- [4] A. Furusawa, J.L. Sørensen, S.L. Braunstein, C.A. Fuchs, H.J. Kimble, and E.J. Polzik, Science **282**, 706 (1998).

- [5] S.J. van Enk, Phys. Rev. A **60**, 5095 (1999).
- [6] H.F. Hofmann, T. Ide, T. Kobayashi, and A. Furusawa, Phys. Rev. A **62**, 062304 (2000).
- [7] H.F. Hofmann, T. Kobayashi, and A. Furusawa, Phys. Rev. A **62**, 013806 (2000).
- [8] D. Boumeester, J.W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, Nature (London) **390**, 575 (1997).