

Weak values, quantum trajectories, and the cavity-QED experiment on wave-particle correlation

H. M. Wiseman

Centre for Quantum Dynamics, School of Science, Griffith University, Brisbane, Queensland 4111, Australia

(Received 18 October 2001; published 14 February 2002)

Weak values as introduced by Aharonov, Albert, and Vaidman (AAV) are ensemble-average values for the results of weak measurements. They are interesting when the ensemble is preselected on a particular initial state and postselected on a particular final measurement result. It is shown that weak values arise naturally in quantum optics, as weak measurements occur whenever an open system is monitored (as by a photodetector). The quantum-trajectory theory is used to derive a generalization of AAV's formula to include (a) mixed initial conditions, (b) nonunitary evolution, (c) a generalized (nonprojective) final measurement, and (d) a non-back-action-evading weak measurement. This theory is applied to the recent cavity-QED experiment demonstrating wave particle duality [G. T. Foster, L. A. Orozco, H. M. Castro-Beltran, and H. J. Carmichael, *Phys. Rev. Lett.* **85**, 3149 (2000)]. It is shown that the "fractional-order" correlation function measured in that experiment can be recast as a weak value in a form as simple as that introduced by AAV.

DOI: 10.1103/PhysRevA.65.032111

PACS number(s): 03.65.Ta, 42.50.Lc, 42.50.Ct, 42.50.Ar

I. INTRODUCTION

The concept of weak values in quantum mechanics was formulated by Aharonov, Albert, and Vaidman (AAV) [1] using the earlier two-wave-function formalism of Ref. [2]. A weak value is the ensemble-average value of a weak-measurement result. A weak measurement is one that minimally disrupts the system, while consequently yielding a minimal amount of information about the observable measured. For a given initial system state, the ensemble average of weak-measurement results is the same as for strong (i.e., projective) measurement results. Where weak measurements are interesting is when a *final* as well as an initial state is specified. Here the final state is the result of a second measurement (a strong one), so that the ensemble average is taken over a *postselected* ensemble, in which the desired result for the final measurement was obtained.

Let the desired initial system state be $|\psi\rangle$, and let ψ denote the event that this is successfully prepared. Let the desired final system state be $|\phi\rangle$, and let ϕ denote the event that the appropriate result is obtained by the final projective measurement. Let the observable to be measured at an intermediate time have operator \hat{X} , and let the result of the measurement be denoted by X_n . Here the nature n of the measurement could be strong ($n=s$) or weak ($n=w$). In the former case, which corresponds to a projective measurement, X_s reproduces the statistics of \hat{X} . In the latter case, X_w does not, and indeed is not confined to the spectrum of \hat{X} . We do require, however, that in all cases X_n be an unbiased estimator in the sense that

$$\langle X_n \rangle_\psi \equiv E[X_n | \psi] = \langle \psi | \hat{X} | \psi \rangle. \quad (1.1)$$

Here E denotes expectation value and the vertical line denotes "given." The postselected value we desire is

$$\phi \langle X_n \rangle_\psi \equiv E[X_n | \psi, \phi] = \sum_x x \text{Prob}[X_n = x | \phi, \psi] \quad (1.2)$$

$$= \frac{\sum_x x \text{Prob}[\phi | X_n = x, \psi] \text{Prob}[X_n = x | \psi]}{\sum_y \text{Prob}[\phi, X_n = y | \psi]}. \quad (1.3)$$

Here $\text{Prob}[H]$ denotes the probability for event H .

For simplicity, assume for the moment that there is no evolution between measurements. Then for a strong intermediate measurement, the postselected average value of X_s is

$$\phi \langle X_s \rangle_\psi = \frac{\sum_x x \text{Prob}[\phi | X_s = x] \text{Prob}[X_s = x | \psi]}{\sum_y \text{Prob}[\phi | X_s = y] \text{Prob}[X_s = y | \psi]} \quad (1.4)$$

$$= \frac{\sum_x |\langle \phi | x \rangle|^2 x |\langle x | \psi \rangle|^2}{\sum_y |\langle \phi | y \rangle|^2 |\langle y | \psi \rangle|^2}. \quad (1.5)$$

Here $|x\rangle$ denotes an eigenstate of \hat{X} . In the final result there is no direct connection between the initial state and the final state. The intermediate strong measurement has destroyed that connection, as it enables the dropping of the conditional ψ from $\text{Prob}[\phi | X_s = x, \psi]$ in Eq. (1.3). Basically there are no surprises here.

Contrast this result with that from a weak measurement, as derived by AAV,

$$\phi \langle X_w \rangle_\psi = \text{Re} \frac{\langle \phi | \hat{X} | \psi \rangle}{\langle \phi | \psi \rangle}. \quad (1.6)$$

Aharonov and Vaidman derive this in detail in Ref. [3], and it is also derived here, in Sec. II, from a somewhat different perspective. This formula appears to have much more of a mysterious “quantum” nature than Eq. (1.5), as it involves the initial and final states linearly (rather than bilinearly) on the numerator and denominator. As a consequence, weak values can have bizarre properties, especially when the outcome ϕ used for postselection is improbable [3]. For example, the weak value $\phi\langle X_w \rangle_\psi$ may lie *outside* the range of eigenvalues for \hat{X} [1,3]. Counterfactual paradoxes [4] can also be expressed and resolved in this way [5,6].

It might be thought that weak values would be mainly of theoretical interest, because arbitrarily weak measurements would not arise naturally in experiments [7]. But in fact arbitrarily weak measurements arise all the time, whenever a system is monitored (i.e., measured continuously in time). The measurement theory for monitored systems was developed by mathematical physicists [8–11]. It was subsequently rediscovered by workers in quantum optics [12–14] when it was found necessary to describe the conditioning of systems on an observed photocurrent. In quantum optics, the theory of continuous monitoring, especially for homodyne and heterodyne measurement, has come to be known as the *quantum-trajectory theory* [14,15].

In this paper I show in Sec. II how the AAV formula can be derived using the approach of the quantum-trajectory theory. I derive a simple generalization arising from removing the restriction that the weak measurement of X be a quantum nondemolition measurement. In Sec. III, I derive a more complete generalization that allows for (a) mixed initial conditions, (b) nonunitary evolution between measurements, and (c) a nonprojective final measurement. Within this context, the AAV formula can be seen to be a special case of the correlation functions that have been used in quantum optics for a long time [16]. Finally, in Sec. IV I show that the correlation function measured in the recent cavity-QED experiment [17] can be reformulated as a weak value almost exactly as originally proposed by AAV.

II. WEAK VALUES: THE SPECIAL THEORY

Monitoring a quantum system can be treated by considering a sequence of discrete weak measurements, each taking time δt and letting $\delta t \rightarrow dt$. To obtain a sensible limit, the strength of each measurement must scale as δt . By this I mean that the postmeasurement system state should, on average, be different from the premeasurement state by an amount of order δt .

To describe weak measurements, we require a generalized (nonprojective) quantum measurement theory [18–20]. Let the measurement result be the random variable X_n as above. In the case of efficient measurements, the measurement can be described in terms of a set of *measurement operators* $\{\hat{M}_x\}_x$, each associated with a result $X_n = x$. The postmeasurement conditioned state is given by

$$\rho_x(t + \delta t) = \frac{\hat{M}_x \rho(t) \hat{M}_x^\dagger}{\text{Prob}[X_n = x]}, \quad (2.1)$$

where the probability for the result is

$$\text{Prob}[X_n = x] = \text{Tr}[\hat{M}_x \rho(t) \hat{M}_x^\dagger]. \quad (2.2)$$

By conservation of probability, the measurement operators must obey a completeness relation

$$\sum_x \hat{E}_x = \hat{1}, \quad (2.3)$$

where $\hat{E}_x = \hat{M}_x^\dagger \hat{M}_x$ is known as the *probability operator or effect* for the result x .

As above, we require the measurement of \hat{X} to be unbiased, so that $\langle X_n(t) \rangle_{\rho(t)} = \text{Tr}[\rho(t) \hat{X}]$. This is satisfied by the following measurement operator, which also has a suitably weak effect on the system [21]:

$$\hat{M}_x = (2\pi/\delta t)^{-1/4} \exp(-x^2 \delta t/2) [1 + \delta t(x\hat{c} - \hat{c}^\dagger \hat{c}/2)]. \quad (2.4)$$

Here \hat{c} is an operator such that

$$\hat{c} + \hat{c}^\dagger = \hat{X}. \quad (2.5)$$

For convenience we are measuring time in units such that $\hat{c}^\dagger \hat{c} \delta t$ is dimensionless. Note that by allowing \hat{c} to be non-Hermitian we are allowing for non-back-action-evading measurements of \hat{X} , even ignoring any evolution between measurements.

The measurement operators satisfy a continuous version of the completeness relation (2.3) to order $(\delta t)^2$,

$$\int dx \hat{M}_x^\dagger \hat{M}_x = \hat{1} + O(\delta t^2). \quad (2.6)$$

In the limit $\delta t \rightarrow dt$, the correction vanishes and it is not difficult to show [21] that the measurement result X_w has statistics given by

$$X_w(t) = \text{Tr}[\hat{X} \rho(t)] + \xi(t). \quad (2.7)$$

Here is a Gaussian white-noise term, defined by $\xi(t) = dW(t)/dt$, where

$$E[dW(t)] = 0, \quad (2.8)$$

$$dW(t)^2 = dt. \quad (2.9)$$

Moreover, the stochastic postmeasurement conditioned state is given by

$$\rho_{X_w}(t + dt) = \{1 + \mathcal{D}[\hat{c}]dt + \mathcal{H}[\hat{c}]dW(t)\} \rho(t). \quad (2.10)$$

Here for arbitrary operators \hat{A} and \hat{B}

$$\mathcal{D}[\hat{A}]\hat{B} \equiv \hat{A}\hat{B}\hat{A}^\dagger - \frac{1}{2}\{\hat{A}^\dagger \hat{A}, \hat{B}\}, \quad (2.11)$$

$$\mathcal{H}[\hat{A}]\hat{B} \equiv (\hat{A} - \text{Tr}[\hat{A} \hat{B}])\hat{B} + \text{H.c.} \quad (2.12)$$

Averaging over the measurement results would give

$$d\rho = dt \mathcal{D}[\hat{c}]\rho, \quad (2.13)$$

which does not necessarily preserve $\langle \hat{X} \rangle_{\rho(t)} = \text{Tr}[\hat{X}\rho(t)]$ because \hat{c} need not commute with \hat{X} .

For continuous measurements we wish to consider a sequence of measurements as described above. However, for the purposes of deriving weak values, we wish to assume we have just a single such weak measurement at time t . (This unrealistic assumption will be removed in Sec. III.) Say the system was prepared at time 0 in state $|\psi(0)\rangle$ and evolved unitarily to state $|\psi\rangle$ at time t . Let us use $\rho(t)$ for $|\psi\rangle\langle\psi|$. The infinitely weak (i.e., infinitesimally strong) measurement is then performed [22]. Then at time T a final projective measurement is made, and we wish to keep only those values of X obtained for which the final result corresponds to the state $|\phi(T)\rangle$. Let $|\phi\rangle$ be that state evolved unitarily backwards in time from T to just after t . Dropping the time arguments, we have in the notation of Sec. I,

$$\phi\langle X_w \rangle_\psi = \text{E}[X_w | \phi, \psi] = \sum_x x \text{Prob}[X_w = x | \phi, \psi] \quad (2.14)$$

$$= \sum_x x \text{Prob}[X_w = x, \phi | \psi] / \text{Prob}[\phi | \psi]. \quad (2.15)$$

Here we have used a sum rather than an integral merely for convenience.

Now because the measurement is infinitely weak, it does not affect the denominator $D = \text{Prob}[\phi | \psi]$, which equals $|\langle \phi | \psi \rangle|^2$. The numerator meanwhile evaluates to

$$N = \sum_x x \text{Prob}[\phi | X_w = x, \psi] \text{Prob}[X_w = x | \psi] \quad (2.16)$$

$$= \sum_x x \langle \phi | \rho_x(t+dt) | \phi \rangle \text{Prob}[X_w = x | \psi] \quad (2.17)$$

$$= \text{E}[X_w \langle \phi | \rho_{X_w}(t+dt) | \phi \rangle]. \quad (2.18)$$

Now from the above formulas (2.7)–(2.10) it is easy to see that this evaluates to

$$\langle \phi | \{ \rho(t) \langle \hat{X} \rangle_{\rho(t)} + \mathcal{H}[\hat{c}]\rho(t) \} | \phi \rangle + O(dt). \quad (2.19)$$

Expanding the superoperator \mathcal{H} and ignoring infinitesimals gives

$$\langle \phi | \{ \hat{c}\rho(t) + \rho(t)\hat{c}^\dagger \} | \phi \rangle. \quad (2.20)$$

Substituting this into Eq. (2.15), and using $\rho(t) = |\psi\rangle\langle\psi|$, gives

$$\phi\langle X_w \rangle_\psi = \frac{\langle \phi | \hat{c} | \psi \rangle \langle \psi | \phi \rangle + \text{H.c.}}{|\langle \phi | \psi \rangle|^2} \quad (2.21)$$

$$= 2 \text{Re} \frac{\langle \phi | \hat{c} | \psi \rangle}{\langle \phi | \psi \rangle}. \quad (2.22)$$

In the case of quantum-nondemolition (QND) measurements where $\hat{c} = \hat{X}/2$, this reduces to the AAV formula (1.6). Putting in the unitary time evolution explicitly and setting $T = t + \tau$,

$$\phi\langle X_w(t) \rangle_\psi = 2 \text{Re} \frac{\langle \phi(T) | \hat{U}(\tau) \hat{c} \hat{U}(t) | \psi(0) \rangle}{\langle \phi(T) | \hat{U}(T) | \psi(0) \rangle}. \quad (2.23)$$

III. WEAK VALUES: THE GENERAL THEORY

We now generalize the above theory by removing many of the assumptions. First, we allow the initial system state $\rho(0)$ to be mixed. Second, we allow arbitrary Markovian evolution [20] of ρ between measurements,

$$\dot{\rho} = \mathcal{L}\rho \equiv -i[\hat{H}, \rho] + \sum_\mu \mathcal{D}[\hat{c}_\mu]. \quad (3.1)$$

Here \hat{H} is an Hermitian operator while $\{\hat{c}_\mu\}_\mu$ is a set of arbitrary operators (which strictly should be bounded [24]). It would be natural for one of them to be the operator \hat{c} introduced in Sec. II. Then at any time between preparation and final measurement, the conditional evolution of the system would be

$$d\rho_{X_w(s)} = ds \mathcal{L}\rho + dW(s) \mathcal{H}[\hat{c}]\rho. \quad (3.2)$$

Finally, we allow for the final measurement at time T to be described by a positive operator $\hat{E}(T)$, which is not necessarily a projector $|\phi(T)\rangle\langle\phi(T)|$. As long as $\hat{1} - \hat{E}(T) \geq 0$, $\hat{E}(T)$ is an effect for some final measurement.

Using E and ρ to denote the events of successful preparation and successful final measurement, the weak value at time t we wish to calculate now is

$${}_E\langle X_w \rangle_\rho = \text{E}[X_w | E, \rho] = \sum_x x \text{Prob}[X_w = x | E, \rho] \quad (3.3)$$

$$= \sum_x x \text{Prob}[X_w = x, E | \rho] / \text{Prob}[E | \rho]. \quad (3.4)$$

As before, the denominator is unaffected by the weak measurement and is given by

$$D = \text{Tr}[\hat{E}(T) e^{\mathcal{L}T} \rho(0)]. \quad (3.5)$$

The numerator can be manipulated similarly to yield

$$N = \text{E}[X_w(t) \text{Tr}[\hat{E}(T) e^{\mathcal{L}\tau} \rho_{X_w}(t+dt)]] \quad (3.6)$$

$$= \text{Tr}[\hat{E}(T) e^{\mathcal{L}\tau} \{ \hat{c} e^{\mathcal{L}t} \rho(0) + \text{H.c.} \}]. \quad (3.7)$$

Equation (2.23) thus generalizes to

$${}_E\langle X_w(t) \rangle_\rho = 2 \operatorname{Re} \frac{\operatorname{Tr}[\hat{E}(T)e^{\mathcal{L}\tau}\{\hat{c}e^{\mathcal{L}t}\rho(0)\}]}{\operatorname{Tr}[\hat{E}(T)e^{\mathcal{L}T}\rho(0)]}. \quad (3.8)$$

The above result looks more like a normalized correlation function, as is familiar in quantum optics, than the unexpected AAV formula (1.6). Indeed, it can be written in the Heisenberg picture [23] as

$${}_E\langle S_w \rangle_\rho = \frac{\langle \hat{E}(T)\hat{c}(t) + \hat{c}^\dagger(t)\hat{E}(T) \rangle}{\langle \hat{E}(T) \rangle}, \quad (3.9)$$

with initial conditions supplied by $\rho(0)$ and the bath state implied by the master equation (3.1). However, we can formulate Eq. (3.8) more like Eq. (1.6) as follows. First we denote, in analogy to $|\psi\rangle = U(t)|\psi(0)\rangle$,

$$\rho = e^{\mathcal{L}t}\rho(0). \quad (3.10)$$

This is the usual forward time evolution by the master equation (3.1). Next we define a *retrodictive* [25] positive operator \hat{E} according to

$$\operatorname{Tr}[\hat{E}\hat{A}] \equiv \operatorname{Tr}[\hat{E}(T)e^{\mathcal{L}\tau}\hat{A}], \quad (3.11)$$

where \hat{A} is an arbitrary operator.

It is easy to verify that this \hat{E} is $\hat{E}(T)$ evolved forward by time τ according to the *retrodictive* master equation

$$\frac{d\hat{E}}{dt} = +i[\hat{H}, \hat{E}] + \sum_\mu [\hat{c}_\mu^\dagger \hat{E} \hat{c}_\mu - \frac{1}{2}\{\hat{c}_\mu^\dagger \hat{c}_\mu, \hat{E}\}]. \quad (3.12)$$

Note that this differs slightly from the retrodictive master equation in Ref. [26]. This is because we do not require the norm of \hat{E} to be preserved, as it appears in both the numerator and denominator of Eq. (3.8). That Eq. (3.12) does generate a positive operator is obvious from the solution as a ‘‘Dyson-like’’ sum constructed in the manner of Eq. (7.24) of Ref. [14]. Using these definitions, we can write the postselected weak value as

$${}_E\langle X_w \rangle_\rho = 2 \operatorname{Re} \frac{\operatorname{Tr}[\hat{E}\hat{c}\rho]}{\operatorname{Tr}[\hat{E}\rho]}. \quad (3.13)$$

Thus retrodiction, quantum trajectories, and weak values are all united in this expression.

IV. WEAK VALUES IN THE RECENT CAVITY-QED EXPERIMENT

The preceding section showed that postselected weak values can be thought of as a correlation function (3.9). In quantum optics weak measurements of the type described in Sec. II are realized through homodyne detection [14,15], where \hat{c} is proportional to the lowering operator of the radiating system. The typical correlation functions thus found are of the form (in the Heisenberg picture)

$$\begin{aligned} \langle : \hat{X}(t+\tau)\hat{X}(t): \rangle &= \langle \hat{c}(t+\tau)\hat{c}(t) + \hat{c}^\dagger(t+\tau)\hat{c}(t) \\ &\quad + \hat{c}^\dagger(t)\hat{c}(t+\tau) + \hat{c}^\dagger(t)\hat{c}^\dagger(t+\tau) \rangle. \end{aligned} \quad (4.1)$$

Usually they are evaluated at steady state (ss), where

$$\rho(0) = \rho_{\text{ss}} \propto \mathcal{N}(\mathcal{L}), \quad (4.2)$$

where \mathcal{N} is the null space.

Note that this is not of the form of Eq. (3.9), so that this correlation function cannot be thought of as a weak value in the AAV sense.

A notable exception to this typicality is the recent cavity-QED experiment [17]. This involved a damped cavity, very weakly driven, through which a beam of resonant atoms passed. The master equation for the system can be approximated as

$$\begin{aligned} \dot{\rho} &= \varepsilon[\hat{a}^\dagger - \hat{a}, \rho] + 2\kappa\mathcal{D}[\hat{a}]\rho + \sum_j^N \{g_j[\hat{a}^\dagger\hat{\sigma}_j^\dagger - \hat{\sigma}_j^\dagger\hat{a}, \rho] \\ &\quad + 2\gamma_\perp\mathcal{D}[\hat{\sigma}_j]\rho\}. \end{aligned} \quad (4.3)$$

Here $\hat{\sigma}_j = |g\rangle_j\langle e|$ is the lowering operator for the j th atom and \hat{a} is the annihilation operator for the field. The output from the cavity is split into two beams by an 85-15 beam splitter. One beam is detected by homodyne detection (with net efficiency $\eta_h < 85\%$) and the other is detected using a photon counter (with net efficiency $\eta_c < 15\%$).

Choosing the slightly odd convention of measuring time in units of $1/(2\kappa\eta_h)$, the homodyne measurement gives a current proportional to

$$X_w(t) = \operatorname{Tr}[\rho(t)\hat{X}] + \xi(t), \quad (4.4)$$

where $\hat{X} = \hat{a} + \hat{a}^\dagger$. It conditions the system state ρ by adding the term

$$dW(t)\mathcal{H}[\hat{a}]\rho \quad (4.5)$$

to $d\rho$. Thus the homodyne measurement gives a continuous weak measurement as in Sec. II, with $\hat{c} = \hat{a}$. The photon counting also gives a continuous measurement that is weak in the sense that the average change in the conditioned system in time δt is of order δt . However, unlike homodyne detection, sometimes the change is great. This can be seen from the set of effects $\{\hat{E}_0, \hat{E}_1\}$ describing this measurement,

$$\hat{E}_0 = 1 - \delta t \kappa \eta_c \hat{a}^\dagger \hat{a}, \quad (4.6)$$

$$\hat{E}_1 = \delta t \kappa \eta_c \hat{a}^\dagger \hat{a}. \quad (4.7)$$

These give the probabilities for detecting zero or one photon over a short time δt . The latter result reveals a lot about the system, and its effect (and measurement operator) is not close to unity.

In the recent cavity-QED experiment, the correlation between the homodyne current $X_w(t)$ and the photon count increment at time $T = t + \tau$ was measured in order to demon-

strate an aspect of wave-particle duality. In the experiment τ ranged from negative to positive, and the measured correlation was proportional to (in the Heisenberg picture)

$$h(t-T) = \frac{\langle :(\hat{a}^\dagger \hat{a})(T) \hat{X}(t): \rangle_{ss}}{\langle \hat{a}^\dagger \hat{a} \rangle_{ss}}. \quad (4.8)$$

This is a ‘‘fractional’’ (3/2)-order correlation function [in contrast to the first-order function in Eq. (4.1)]. Here we are interested only in the case $\tau > 0$ in which case the correlation function can be written as [27]

$$h(-\tau) = \frac{\text{Tr}[\hat{E}_1(T) e^{\mathcal{L}\tau} (\hat{a} \rho_{ss} + \rho_{ss} \hat{a}^\dagger)]}{\text{Tr}[\hat{E}_1(T) \rho_{ss}]}. \quad (4.9)$$

Defining \hat{E}_1 to be $\hat{E}_1(T)$ retrodicted (as in Sec. III) from time T to time t , we can rewrite this correlation function as

$$E_1 \langle X_w \rangle_{\rho_{ss}} = 2 \text{Re} \frac{\text{Tr}[\hat{E}_1 \hat{a} \rho_{ss}]}{\text{Tr}[\hat{E}_1 \rho_{ss}]}. \quad (4.10)$$

It is now clear that the correlation function measured in Ref. [17] is a weak value, preselected by the system being in its stationary state, and postselected on the final measurement result E_1 (a photon detected at time T). Actually, since what was measured was a correlation *function* (i.e., a function of τ), this experiment showed the *dynamics* of a weak value over time. In this case, the spectrum of \hat{X} is the real line, so there is no chance of observing a weak value outside of this range. Nevertheless, the weak values measured in the experiment were, for a range of times τ , very far away from the stationary average value of \hat{X} . This led to the violation of various classical inequalities [28,17]. In hindsight, the strangeness of the weak values in this experiment is not surprising, since the condition cited by Aharonov and Vaidman [3] is fulfilled. That is, the postselection is done on a rare event (the detection of one photon rather than zero photons).

There are a number of features of the above system that let us transform the generalized weak value (4.10) into a form almost identical to the special form (2.22) derived in Sec. II. Because the driving is very weak ($\epsilon \ll \kappa, \gamma_\perp$), the ‘‘jump’’ terms in the master equation (4.3) have a very small effect. This means that it is possible to approximate the Liouvillian as

$$\mathcal{L} = \mathcal{H}[-i\hat{H}_{\text{eff}}], \quad (4.11)$$

where

$$\begin{aligned} -i\hat{H}_{\text{eff}} = & \epsilon(\hat{a}^\dagger - \hat{a}) - \kappa \hat{a}^\dagger \hat{a} + \sum_j^N [g(\hat{a}^\dagger \hat{\sigma}_j^\dagger - \hat{\sigma}_j^\dagger \hat{a}) \\ & - \gamma_\perp \hat{\sigma}_j^\dagger \hat{\sigma}_j]. \end{aligned} \quad (4.12)$$

Ignoring normalization, this generates nonunitary evolution according to the operator

$$N(t) = \exp(-i\hat{H}_{\text{eff}}t). \quad (4.13)$$

The stationary solution is, to a very good approximation, pure [29],

$$\rho_{ss} = |\psi_{ss}\rangle \langle \psi_{ss}|, \quad (4.14)$$

where $|\psi_{ss}\rangle$ is the eigenstate of $-i\hat{H}_{\text{eff}}$ with an eigenvalue having the largest real part.

These results mean that we can approximate Eq. (4.10) by

$$E_1 \langle X_w \rangle_{\psi_{ss}} = 2 \text{Re} \frac{\langle \psi_{ss} | \hat{a}^\dagger \hat{a} N(\tau) \hat{a} | \psi_{ss} \rangle}{\langle \psi_{ss} | \hat{a}^\dagger \hat{a} N(\tau) | \psi_{ss} \rangle}. \quad (4.15)$$

Again because of the weak excitation, $|\psi_{ss}\rangle$ is, to a good approximation, given by

$$|\psi_{ss}\rangle = |0\rangle + \alpha|1\rangle + \beta|1'\rangle. \quad (4.16)$$

Here α and β scale with ϵ . The state $|0\rangle$ is the state with no excitations, $|1\rangle$ that with one photon but all atoms in the ground state, and $|1'\rangle$ a state with one collective atomic excitation but no photons. The same sort of expansion holds for $\hat{a}|\psi_{ss}\rangle/\sqrt{\langle \hat{a}^\dagger \hat{a} \rangle_{ss}}$ (although the presence of a nonzero $|1\rangle$ terms here indicates that $|\psi_{ss}\rangle$ does have higher-order terms). Furthermore, to leading order, $N(t)$ preserves this expansion (apart from normalization). For these reasons, we can replace $\hat{a}^\dagger \hat{a}$, which is really $\hat{a}^\dagger \hat{a} \otimes \hat{1}_{\text{atoms}}$, by $|1\rangle\langle 1|$. That is, postselection on the detection of a photon is equivalent to postselection on projection into the state $|1\rangle$. The final result thus is

$$E_1 \langle X_w \rangle_{\psi_{ss}} = 2 \text{Re} \frac{\langle 1 | N(\tau) \hat{a} | \psi_{ss} \rangle}{\langle 1 | N(\tau) | \psi_{ss} \rangle}. \quad (4.17)$$

This is completely analogous to Eq. (1.6) when we recognize $N^\dagger(\tau)|1\rangle$ as the unnormalized retrodicted final state.

When $|\psi_{ss}\rangle$ and $N(\tau)$ are substituted into Eq. (4.17), and only the terms of leading order in the driving ϵ are kept, the result is

$$\frac{E_1 \langle X_w \rangle_{\psi_{ss}}}{\langle \psi_{ss} | \hat{X} | \psi_{ss} \rangle} = 1 + \zeta e^{-\eta\tau} \left[\cos \Omega \tau + \frac{\eta}{\Omega} \sin \Omega \tau \right], \quad (4.18)$$

where $\eta = (\kappa + \gamma_\perp)/2$ and where ζ and Ω are functions of κ, γ_\perp , and the coupling coefficients $\{g_j\}_j$. This is exactly the same as the analytical result in Ref. [17], where this result was derived by considering the correlation function (4.8) with $T < t$. That is, with the photodetection preceding the homodyne measurement. In that case, the correlation function is

$$h(-\tau) = h(|\tau|) = \frac{\text{Tr}[\hat{X} e^{\mathcal{L}|\tau|} \hat{a} \rho_{ss} \hat{a}^\dagger]}{\text{Tr}[\hat{a} \rho_{ss} \hat{a}^\dagger]}. \quad (4.19)$$

In Ref. [17] the symmetry of $h(\tau)$ was established by assuming Gaussian fluctuations. The analysis in this paper shows that the symmetry (to leading order) of $h(\tau)$ can be demonstrated explicitly by calculating both cases. The case of negative τ is found simply from the usual predictive quantum

mechanics with wave-function collapse [29], and the case of positive τ equally simply by retrodictive quantum mechanics with weak values, as here.

Finally, it is important to point out that the above form for the weak value as a function of time Eq. (4.18) matches well with the experiment [17]. This is so even though the experimental conditions are far from ideal, with fluctuations in the number of atoms, transverse motion of the atoms, imperfect beam alignment, and so on. This shows that a weak value is not as fragile as one might have thought, and does give quantitative predictions for the behavior of the system.

V. DISCUSSION

In this paper I have combined the quantum-trajectory theory and the concepts of weak values and retrodiction in order to show the relation between weak values and certain correlation functions as can be measured in quantum optics. This required a generalization of the weak value theory of Aharonov, Albert, and Vaidman. However, in a special case of considerable interest, the wave-particle cavity-QED experiment, I show that the correlation function reduces to a form as simple, and almost the same, as that originally derived by AAV. Moreover, this experiment measured not a weak value at a particular time, but the change in a weak

value over time. That is, it showed the *dynamics* of a weak value.

Weak values are of considerable intrinsic theoretical and philosophical interest. Moreover, the fact that they obey certain well-defined rules [3,5] imply that they are also heuristic tools for predicting the results of experiments. Of course the results could be obtained by other, more laborious methods. But the power of having several fonts of intuition should not be undervalued. The analysis of Sec. IV is a case in point. This work should help to identify other instances in which the theory of weak values can be used in quantum optics and related areas. It may even suggest new experiments that provide further illustrations and applications for this thought-provoking area of quantum physics.

ACKNOWLEDGMENTS

I would like to thank the Stony Brook group, in particular Luis Orozco and Joe Reiner, for their hospitality and for discussions without which I could never have done this work. Discussions with David Pegg about retrodiction, and with Klaus Mølmer about weak values, are also gratefully acknowledged. This work was supported by the Australian Research Council.

-
- [1] Y. Aharonov, D. Albert, and L. Vaidman, *Phys. Rev. Lett.* **60**, 1351 (1988).
- [2] Y. Aharonov, O. G. Bergman, and J. L. Lebowitz, *Phys. Rev.* **134**, B1410 (1964).
- [3] Y. Aharonov and L. Vaidman, *Phys. Rev. A* **41**, 11 (1990).
- [4] L. Hardy, *Phys. Rev. Lett.* **68**, 2981 (1992).
- [5] Y. Aharonov *et al.*, e-print quant-ph/0104062.
- [6] K. Mølmer, e-print quant-ph/0109042.
- [7] Although, by deliberately constructing a weak measurement, the unusual predictions of AAV [1] were experimentally verified not long after they were proposed by N. W. M. Ritchie, J. G. Story, and R. G. Hulet, *Phys. Rev. Lett.* **66**, 1107 (1991).
- [8] V. P. Belavkin, *Nondemolition Measurement and Nonlinear Filtering of Quantum Stochastic Processes*, edited by A. Blaquière, Lecture Notes in Control and Information Sciences Vol. 121 (Springer, Berlin, 1988), pp. 256–266.
- [9] V. P. Belavkin and P. Staszewski, *Phys. Rev. A* **45**, 1347 (1992).
- [10] A. Barchielli, *Quantum Opt.* **2**, 423 (1990).
- [11] A. Barchielli, *Int. J. Theor. Phys.* **32**, 2221 (1993).
- [12] J. Dalibard, Y. Castin, and K. Mølmer, *Phys. Rev. Lett.* **68**, 580 (1992).
- [13] C. W. Gardiner, A. S. Parkins, and P. Zoller, *Phys. Rev. A* **46**, 4363 (1992).
- [14] H. J. Carmichael, *An Open Systems Approach to Quantum Optics* (Springer-Verlag, Berlin, 1993).
- [15] H. M. Wiseman and G. J. Milburn, *Phys. Rev. A* **47**, 1652 (1993).
- [16] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer, Berlin, 1994), and references therein.
- [17] G. T. Foster, L. A. Orozco, H. M. Castro-Beltran, and H. J. Carmichael, *Phys. Rev. Lett.* **85**, 3149 (2000).
- [18] K. Kraus, *States, Effects, and Operations: Fundamental Notions of Quantum Theory* (Springer, Berlin, 1983).
- [19] V. B. Braginsky and F. Y. Khalili, *Quantum Measurement* (Cambridge University Press, Cambridge, 1992).
- [20] C. W. Gardiner and P. Zoller, *Quantum Noise* (Springer, Berlin, 2000).
- [21] H. M. Wiseman and L. Diósi, *J. Chem. Phys.* **268**, 91 (2001).
- [22] Of course in a real experiment, such as Ref. [17], the measurement is not infinitely weak, otherwise an infinitely large ensemble would be required to obtain a sensible average. The infinitesimal dt is actually a finite but small $\delta t = 1/B$, where B is the bandwidth of the apparatus (including the detector, amplifier, and often a deliberately introduced filter).
- [23] Note that $\hat{E}(T)$ in the Heisenberg picture is not necessarily the same as $\hat{E}(T)$ in the Schrödinger picture.
- [24] G. Lindblad, *Commun. Math. Phys.* **48**, 199 (1976).
- [25] S. M. Barnett, D. T. Pegg, and J. Jeffers, *J. Mod. Opt.* **47**, 1779 (2000).
- [26] S. M. Barnett, D. T. Pegg, J. Jeffers, and O. Jedrkiewicz, *Phys. Rev. Lett.* **86**, 2455 (2001).
- [27] Note that the τ I am using is the negative of the τ in Ref. [17], which is why I write their correlation function as $h(-\tau)$.
- [28] H. J. Carmichael, H. M. Castro-Beltran, G. T. Foster, and L. A. Orozco, *Phys. Rev. Lett.* **85**, 1855 (2000).
- [29] H. J. Carmichael, R. J. Brecha, and P. R. Rice, *Opt. Commun.* **82**, 73 (1991).