

Sympathetic cooling and collisional properties of a Rb-Cs mixture

M. Anderlini, E. Courtade, M. Cristiani, D. Cossart, D. Ciampini, C. Sias, O. Morsch, and E. Arimondo
INFN, Dipartimento di Fisica E. Fermi, Università di Pisa, Largo B. Pontecorvo 3, I-56127 Pisa, Italy
 (Received 28 February 2005; published 1 June 2005)

We report on measurements of the collisional properties of a mixture of ^{133}Cs and ^{87}Rb atoms in a magnetic trap at μK temperatures. By selectively evaporating the Rb atoms using a radio-frequency field, we achieved sympathetic cooling of Cs down to a few μK . The interspecies collisional cross section was determined through rethermalization measurements, leading to good agreement with a theoretical prediction of $595a_0$ for the triplet s -wave scattering length for Rb in the $|F=2, m_F=2\rangle$ and Cs in the $|F=4, m_F=4\rangle$ magnetic states. We briefly speculate on the prospects for reaching the Bose-Einstein condensation of Cs inside a magnetic trap through sympathetic cooling.

DOI: 10.1103/PhysRevA.71.061401

PACS number(s): 32.80.Pj, 34.20.Cf

Laser cooling of neutral atoms, combined with evaporative cooling in conservative (magnetic and optical) traps, has led to a number of important breakthroughs in atomic physics, most notably the observation of Bose-Einstein condensation (BEC) in a dilute gas of alkali-metal atoms in 1995 [1]. In recent years, research on ultracold atoms has expanded into the realms of atomic mixtures. Adding a second atomic species has, among other things, opened up the possibility to sympathetically cool one atomic species through collisional energy exchange with the other species [2]. On the one hand, this has proved an invaluable tool for reaching the quantum degeneracy regime with fermionic atoms for which Pauli blocking reduces the evaporative cooling efficiency at low temperatures. On the other hand, ultracold atomic mixtures are also interesting in their own right. In particular, the possibility of creating cold heteronuclear molecules could be an important ingredient in neutral atom quantum computing due to the expected large permanent dipole moment of such molecules. Furthermore, ultracold mixtures can lead to interesting quantum phases when loaded into an optical lattice [3]. A number of mixtures has been studied in magneto-optical traps (MOTs) [4,5], and recent experiments by Kerman *et al.* [6] have yielded information about the rovibrational structure of the $^{85}\text{Rb}^{133}\text{Cs}$ molecule. Furthermore, a few combinations of ultracold atoms have also been experimentally investigated in conservative traps, among them Li-Cs [7], K-Rb [8,9], and Na-Li [10].

In this paper we study the collisional properties of a mixture of ultracold ^{87}Rb and ^{133}Cs atoms in a magnetic trap. Both Rb and Cs have been used in laser cooling of atoms for more than 15 years now and are important as time and frequency standards. However, not much is known about their interatomic potentials and collisional properties. While for the lighter alkali metals the interatomic potentials can be calculated relatively easily, it has been difficult to do the same for the Rb-Cs potential. Jamieson *et al.* [11] calculated the collisional parameters using several similar-looking choices for the interatomic potentials, and found that even for small changes in the potential the s -wave scattering length varies by up to two orders of magnitude. Furthermore, the adverse collisional properties (large inelastic cross sections) of Cs in the magnetically trappable states make experimental studies of Rb-Cs mixtures challenging.

The experimental apparatus used for this work is similar to our Rb-BEC setup described in detail elsewhere [12]. We use a double-chamber vacuum system with a two-dimensional (2D) collection MOT in the upper chamber and a six-beam MOT in the lower chamber. In order to trap and cool both Rb and Cs atoms, the trapping and repumping light for the two species is superimposed and the same optics (mirrors, lenses, and wave plates) is used to create the beams for both MOTs. Once the atoms have been transferred into the lower MOT, after a brief compressed MOT and molasses phase, the trapping beams are switched off and the atoms are optically pumped into the $|F=2, m_F=2\rangle$ and $|F=4, m_F=4\rangle$ magnetically trappable states of Rb and Cs, respectively. Immediately after that, the time-averaged orbiting potential (TOP) magnetic trap is switched on. Since the two atomic species have different equilibrium positions in the magnetic trap (due to gravitational sag), the positions of the MOTs have to be adjusted accordingly in order to avoid subsequent oscillations in the TOP trap. This is achieved by tuning the radiation pressure in the MOT using the (wavelength-selective) quarter-wave plates inserted into the optical path of the trapping beams.

In order to demonstrate sympathetic cooling, after loading the atoms into the magnetic trap, we first performed circle-of-death evaporative cooling (which exploits the rotating zero-of-field locus on which spin-flips can occur) by continuously reducing the strength of the rotating bias field [13]. Because this cooling technique is not species selective, cooling one or the other species separately (i.e., loading only that species into the MOT) made little difference.

We then proceeded to apply a radio-frequency field resonant with a $\Delta m_F = \pm 1$ Zeeman transition in the Rb atoms. By ramping down the frequency of the RF field, forced evaporative cooling is induced as the radius of the surface on which atoms are transferred into untrapped states shrinks. Radio-frequency evaporation is species selective as it depends on the Zeeman sublevel spacing. Therefore, only the Rb atoms were evaporatively cooled in this way. Nevertheless, the measured temperature of the Cs atoms exactly followed the Rb temperature down to a few μK , clearly indicating that sympathetic cooling was taking place [see Fig. 1(a)]. In fact, repeating the experiment with Cs atoms only, a much smaller cooling rate, consistent with residual circle-of-




FIG. 1. (a) Sympathetic cooling of Cs atoms (filled circles) through RF evaporation of Rb (open circles). Removing the Rb atoms before the RF sweep, Cs atoms are only cooled slightly by residual circle-of-death evaporation at a fixed bias field (triangles). (b) A comparison of experiment and Monte Carlo simulations results for Cs: no interspecies collisions (dotted line), data set C (dash-dotted line), data set A (solid line), data set B (dashed line). Rb temperatures were only slightly sensitive of the cross section.

death evaporation, was observed. Initially, the atom numbers in the trap were $N_{\text{Rb}} \approx 2 \times 10^6$ and $N_{\text{Cs}} \approx 10^5$. Towards the end of the ramp, the number of Rb atoms had dropped by a factor of 10. As expected, sympathetic cooling was efficient as long as the number of Rb atoms was larger than the number of Cs atoms; once there were fewer Rb than Cs atoms, sympathetic cooling stopped.

In order to understand the dynamics of sympathetic cooling and to extract information on the interspecies scattering properties, we solved the kinetic equations for the atomic dynamics inside the magnetic trap in the presence of interatomic collisions, treating the collisions through a Monte Carlo approach [14] (in the following to be called “Monte Carlo simulations”). Those simulations were extended to samples of two species with different masses, atom numbers, and trap frequencies. Since the known values of the C_6 coefficient for the Rb-Cs molecular potential [15] give a threshold energy for p -wave scattering of about 50 μK [16], we took into account both the s -wave and the p -wave contributions to the interspecies elastic cross section. Using the effective range approximation [17,18], the s -wave term was written in terms of a scattering length a and an effective range r_e , while the p -wave term was expressed in terms of a p -wave volume A_1 [19]




FIG. 2. Rethermalization in a Rb-Cs mixture. The initial difference between the Rb temperature (open circles) and the Cs temperature (filled circles) was created by a fast RF evaporation of the Rb atoms. The solid lines are best fits obtained by a numerical integration of the differential equations describing the rethermalization process (see text).

$$\sigma_{\text{tot}} = \frac{4\pi a^2}{\left(1 - \frac{1}{2}ar_e k^2\right)^2 + a^2 k^2} + \frac{12\pi A_1^2 k^4}{1 + A_1^2 k^6}, \quad (1)$$

where k is the modulus of the wave vector of the relative atomic motion. The angular dependence of the $l=1$ partial wave does not affect the efficiency of the p -wave contribution in the interspecies thermalization process (in contrast to single-species cross-dimensional relaxation, where it reduces the p -wave contribution by a factor 3/5 [20]). In order to analyze the experimental results, we considered the theoretical sets of scattering parameters $\{a, r_e, A_1\}$ for the triplet interaction reported in [11], corresponding to data set A = $\{595.2a_0, 190.2a_0, -168.5 \times 10^4 a_0^3\}$, data set B = $\{177.2a_0, 126.4a_0, -4681 \times 10^4 a_0^3\}$, data set C = $\{-317.6a_0, 424.2a_0, -112.3 \times 10^4 a_0^3\}$, and data set D = $\{-45.37a_0, 3075a_0, -84.22 \times 10^4 a_0^3\}$ (where a_0 is the Bohr radius). Numerical simulations using those parameters showed that only the two sets A and B were consistent with the observed efficiency of the sympathetic cooling [Fig. 1(b)]. However, systematic effects mainly related to the uncertainties in the efficiency of the evaporation processes prevented an accurate extraction of the scattering parameters from this data.

In order to obtain a more accurate measurement of the scattering cross section, we performed rethermalization measurements. First, Rb and Cs were cooled down to temperatures between 5 and 60 μK by circle-of-death evaporative cooling. Thereafter, a radio-frequency ramp was applied in such a way that the Rb atoms were cooled, but with a fast enough sweep so that there was no energy exchange between Cs and Rb during the evaporation time. At the end of the RF ramp there was thus a temperature difference between the Rb and Cs atoms [21]. The mixture was then held in the magnetic trap for up to 20 s and the temperatures of both species were measured as a function of time. These measurements were repeated for various mean temperatures of the mixture.

Figure 2 shows the result of such a rethermalization mea-

surement. In order to suppress systematic effects as much as possible, for each experimental run we measured both the Rb and Cs temperatures *without* the other species by eliminating one or the other through a resonant flash before the rethermalization process started. Typically, we observed single-species heating rates of up to 350 nK s^{-1} .

Rethermalization techniques have been extensively used to measure collisional scattering lengths and p -wave cross sections [8,18,22–25]. For mixtures of two clouds prepared at different temperatures, the relaxation of the temperatures due to elastic collisions with a certain constant cross section proceeds exponentially with a rate that can be calculated analytically through the model of [7]. From the rates obtained by exponential fits of observed thermalizations one can generally extract the value of the cross section. We extended that model to cross sections that explicitly depend on the energy of the colliding particles [as in Eq. (1)]. In this case the relaxation rate of the temperatures in the mixture depends linearly on the effective cross section σ_e [18,26,27], which, in the case of Eq. (1), is

$$\sigma_e = \int_0^\infty dx x^5 e^{-x^2} \left(\frac{4\pi a^2}{\left(1 - \frac{1}{2} C a r_e x^2\right)^2 + C a^2 x^2} + \frac{12\pi C^2 A_1^2 x^4}{1 + C^3 A_1^2 x^6} \right), \quad (2)$$

where $C = 2\mu k_B(m_1 T_2 + m_2 T_1) / (\hbar^2 M)$, μ is the reduced mass, and $M = m_1 + m_2$.

We checked that exponential fits to this model and to Monte Carlo simulations gave the same thermalization rates to within a few percent [28]. However, due to the decay of the number of atoms during the thermalization and to the observed intrinsic heating independent of the interspecies interactions, we could not determine the effective cross section through a simple exponential fit to the data. Those cross sections were determined by running a numerical simulation using the model discussed above for different σ_e and initial temperatures of the two species, taking into account the experimental single-species heating rates and atom number decay. We then compared the results of these simulations with the experimental data, and from the combination of parameters giving the least χ^2 we finally determined σ_e . The results of this analysis are plotted in Fig. 3 as a function of the initial weighted average temperature of the mixed sample.

Having measured the effective scattering cross sections, we calculated the effective cross sections corresponding to the scattering parameters predicted by Jamieson *et al.* [11] for the temperature range relevant to our experiment. The agreement with our data is best for the scattering parameters of data set A (see Fig. 3).

We note here that due to imperfect optical pumping (and, possibly, other depolarizing processes during the evaporation cycle), both the Rb and Cs clouds had admixtures of atoms in other Zeeman sublevels. In the case of Rb, around 90% of the atoms were in the desired state $|F=2, m_F=2\rangle$, with around 10% in the $|F=2, m_F=1\rangle$ sublevel, while for Cs we measured (by performing a Stern-Gerlach-type experiment to separate the Zeeman levels in time of flight) relative populations of 70–80% in the $|F=4, m_F=4\rangle$ sublevel and 20–30%




FIG. 3. Effective elastic scattering cross sections for Rb-Cs collisions as a function of temperature. The theoretical predictions by Jamieson *et al.* [11] plotted in this graph are: data set B (dotted line), data set A (solid line), data set C (dashed line), and data set D (dash-dotted line). The error bars on the experimental points are derived from the χ^2 fit (see text).

in $|F=4, m_F=3\rangle$. This posed two problems in interpreting our data. Firstly, the scattering cross sections for atoms in the various Zeeman sublevels are not necessarily the same. Since we could not eliminate the populations in the other sublevels, we can only quote them here to indicate the possible error involved in our determination of the Rb $|F=2, m_F=2\rangle$ – Cs $|F=4, m_F=4\rangle$ cross section. Secondly, the presence of other Zeeman sublevels distorted the density profiles from which we calculated the Rb and Cs temperatures. This problem was solved by fitting a double-Gaussian curve with a fixed separation and extracting the temperature from the widths of these two Gaussians.

Concerning the *inelastic* collisional properties of the Rb-Cs mixture, we found that in the temperature range $5\text{--}40 \mu\text{K}$ and for typical densities of $(0.5\text{--}2) \times 10^{10} \text{ cm}^{-3}$, no additional losses in either Rb or Cs in the presence of the other species occurred. We can, therefore, put the upper limit of $\sim 10^{-12} \text{ cm}^3 \text{ s}^{-1}$ on the inelastic coefficient for Rb-Cs collisions in magnetic fields in the range $4\text{--}40 \text{ G}$. In the same range we also performed a slow sweep of the bias field, but observed no pronounced losses. This rules out the possibility of broad interspecies Feshbach resonances (as observed recently for a Na-Li mixture [10]).

Finally, the fact that Cs can be sympathetically cooled by Rb and that the scattering length for interspecies collisions is large leads us to speculate whether it might be possible to reach Bose-Einstein condensation of Cs inside a magnetic trap using a sympathetic cooling approach. Although Cs was recently condensed inside an optical trap [29], it would still be interesting to achieve condensation in a magnetic trap, thus avoiding the complicated setup of [29]. Since in our current experiment the number of Rb atoms we could initially trap was too small to extend the sympathetic cooling below a few μK , we conducted Monte Carlo simulations with larger numbers of atoms. Our simulations took into account the inelastic losses especially of Cs, which in the $|F=4, m_F=4\rangle$ Zeeman sublevel has a zero-energy resonance [30] associated with a large inelastic collision rate responsible for the failure of all attempts so far to reach Bose-

Einstein condensation in a magnetic trap. Running simulations with up to 2×10^7 Rb atoms in the $|F=2, m_F=2\rangle$ sublevel, 10^5 Cs atoms in $|F=4, m_F=4\rangle$ and an initial temperature of $10 \mu\text{K}$ for both species clearly showed that even with a large (but still realistic) number of Rb atoms and very low final trap frequencies (chosen so as to reduce the Cs density and hence inelastic losses), it is not possible to reach quantum degeneracy in such a scheme. However, using the $|F=1, m_F=-1\rangle$ and $|F=3, m_F=-3\rangle$ sublevels for Rb and Cs, respectively, we found that with the same numbers of atoms as in the first simulation, the same initial temperature and similarly large scattering lengths (i.e., $>300a_0$), the threshold for condensation of Cs was reached with roughly 5×10^4 Cs atoms left in the mixture. While we have no precise knowledge of the scattering length involved in the collisions between atoms in those sublevels it still seems likely that the effective scattering length will be sufficiently large to make the condensation of Cs possible. Nevertheless, further work needs to be done to experimentally verify the scattering properties of those sublevels.

In conclusion, we have demonstrated sympathetic cooling of Cs atoms down to very low temperatures in a Rb-Cs mixture, even in the presence of large inelastic losses in Cs. We have characterized the ultracold collisions between the two species by measuring rethermalization rates. Monte Carlo simulations yielded a precise reproduction of the overall collision processes. Our results are consistent with a large interspecies triplet *s*-wave scattering length. If the scattering length between atoms in the $|1, -1\rangle$ state of Rb and $|3, -3\rangle$ state of Cs is similarly large, it appears feasible to reach Bose-Einstein condensation of Cs in a magnetic trap through sympathetic cooling with Rb.

We thank T. Bergeman, D. Guéry-Odelin, M. Holland, and M. Jamieson for useful discussions, and J.H. Müller and N. Malossi for help in the early stages of the experiment. This research was supported by the INFM (PRA Photonmatter), by Progetto MIUR-COFIN 2004 and by the EU Network Cold Quantum Gases, Contract No. HPRN-CT-2000-00125.

-
- [1] K. B. Davis *et al.*, Phys. Rev. Lett. **75**, 3969 (1995); C. C. Bradley *et al.*, *ibid.* **75**, 1687 (1995); M. H. Anderson *et al.*, Science **269**, 198 (1995).
- [2] For a general review of ultracold atomic collisions, see J. Weiner *et al.*, Rev. Mod. Phys. **71**, 1 (1999); J. Weiner, *Cold and Ultracold Collisions in Quantum Microscopic and Mesoscopic Systems* (Cambridge University Press, Cambridge, 2003).
- [3] M. Lewenstein *et al.*, Phys. Rev. Lett. **92**, 050401 (2004).
- [4] G. D. Telles *et al.*, Phys. Rev. A **63**, 033406 (2001).
- [5] M. W. Mancini *et al.*, Phys. Rev. Lett. **92**, 133203 (2004).
- [6] A. J. Kerman *et al.*, Phys. Rev. Lett. **92**, 153001 (2004); **92**, 033004 (2004).
- [7] A. Mosk *et al.*, Appl. Phys. B **73**, 791 (2001).
- [8] G. Ferrari *et al.*, Phys. Rev. Lett. **89**, 053202 (2002).
- [9] J. Goldwin *et al.*, Phys. Rev. A **70**, 021601(R) (2004).
- [10] C. A. Stan *et al.*, Phys. Rev. Lett. **93**, 143001 (2004).
- [11] M. J. Jamieson *et al.*, J. Phys. B **36**, 1085 (2003).
- [12] J. H. Müller *et al.*, J. Phys. B **33**, 4095 (2000).
- [13] W. Petrich *et al.*, Phys. Rev. Lett. **74**, 3352 (1995).
- [14] H. Wu and C. J. Foot, J. Phys. B **29**, L321 (1996); H. Wu *et al.*, Phys. Rev. A **56**, 560 (1997).
- [15] M. Marinescu and H. R. Sadeghpour, Phys. Rev. A **59**, 390 (1999).
- [16] B. De Marco and D. S. Jin, Phys. Rev. A **58**, R4267 (1998).
- [17] C. J. Joachain, *Quantum Collision Theory* (North Holland, Amsterdam, 1983).
- [18] P. O. Schmidt *et al.*, Phys. Rev. Lett. **91**, 193201 (2003).
- [19] G. Gutiérrez *et al.*, Phys. Rev. B **29**, 5211 (1984).
- [20] B. De Marco *et al.*, Phys. Rev. Lett. **82**, 4208 (1999).
- [21] At the end of this fast evaporation the rubidium cloud is not in thermal equilibrium, having a temperature in the *x* and *y* directions different from the temperature in the *z* direction. Monte Carlo simulations showed that the relaxation of the average temperature $T=(2T_x+T_z)/3$ for a cloud out of equilibrium is in fact identical to the temperature relaxation of a cloud at equilibrium. In the following, the rubidium temperature is understood to be the average temperature.
- [22] M. Arndt *et al.*, Phys. Rev. Lett. **79**, 625 (1997).
- [23] R. Legere and K. Gibble, Phys. Rev. Lett. **81**, 5780 (1998).
- [24] G. Delannoy *et al.*, Phys. Rev. A **63**, 051602(R) (2001).
- [25] I. Bloch *et al.*, Phys. Rev. A **64**, 021402(R) (2001).
- [26] G. M. Kavoulakis *et al.*, Phys. Rev. A **61**, 053603 (2000).
- [27] P. J. J. Tol *et al.*, Phys. Rev. A **70**, 013404 (2004).
- [28] The validity of this model is restricted to mixtures in the *collisionless* regime, i.e., where the harmonic trap frequencies are much bigger than the thermalization rates, as was the case for our experimental conditions. In other regimes a more complex model, taking into account space-velocity correlations, is needed [D. Guéry-Odelin (private communication)].
- [29] T. Weber *et al.*, Science **299**, 5604 (2003).
- [30] J. Söding *et al.*, Phys. Rev. Lett. **80**, 1869 (1998).