

Bose-Einstein-condensation temperature of a gas of weakly dissociated diatomic molecules

L. M. Jensen

Department of Physics, Nanoscience Center, University of Jyväskylä, P. O. Box 35, FI-40014 Jyväskylä, Finland

H. Mäkelä

Department of Physics, University of Turku, FI-20014 Turku, Finland

C. J. Pethick

Nordita, Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark

(Received 16 October 2006; published 15 March 2007)

We consider the properties of a gas of bosonic diatomic molecules in the limit when few of the molecules are dissociated. Taking into account the effects of dissociation and scattering among molecules and atoms, we calculate the dispersion relation for a molecule, and the thermal depletion of the condensate. We calculate the dependence of the Bose-Einstein-condensation (BEC) temperature of a uniform gas on the atom-atom scattering length, and conclude that, for a broad Feshbach resonance, the condensation temperature increases as the molecular state becomes less strongly bound, thereby giving rise to a maximum in the transition temperature in the BEC-BCS crossover. We also argue on general grounds that, for a gas in a harmonic trap and for a narrow Feshbach resonance, the condensation temperature will decrease with increasing scattering length.

DOI: [10.1103/PhysRevA.75.033606](https://doi.org/10.1103/PhysRevA.75.033606)

PACS number(s): 03.75.Ss, 03.75.Hh, 05.30.Jp

I. INTRODUCTION

The crossover between BCS pairing of two species of fermions and Bose-Einstein-condensation (BEC) of diatomic molecules with changing atom-atom scattering length has been studied by many authors, beginning with the works of Eagles [1], Leggett [2], and Nozières and Schmitt-Rink [3]. One of the surprising results of the calculations of Ref. [3] and also found in Refs. [4,5] is that, in the crossover region, the transition temperature has a maximum as the strength of the atom-atom interaction is varied. However, it is unclear whether the maximum is real, or merely an unphysical artifact of either the assumptions made in the model or the numerical procedures. The early calculations only accounted for quadratic fluctuations around the mean-field solution and it was generally expected that the fluctuation-induced peak was an artifact of the Gaussian approximation. This view was furthermore supported by a self-consistent field-theoretical calculation in Ref. [6] where T_c was found to be a monotonic function of the dimensionless coupling constant. However, the most recent self-consistent calculations of T_c [7] do indicate the presence of a peak in T_c . Most calculations for broad resonances for a uniform gas give a peak in T_c for both one-channel and two-channel models for the interaction, and for a variety of approximations for the atom propagators, which are in some cases replaced by either bare or dressed propagators [8–10]. In calculations that use a combination of bare and dressed propagators, T_c initially decreases coming from the far BEC limit before turning around to give a local maximum near resonance [11]. In calculations for a harmonically trapped gas in the local density approximation, the peak vanishes for both one-channel and two-channel models for the interaction, and for a variety of approximations for the atom propagators [12]. The peak vanishes in calculations for sufficiently narrow resonances (for a recent review, see [13]). A summary of the present situation is that microscopic calculations point to there being

a maximum in T_c for broad resonances for the uniform gas, but not for harmonically trapped gases or sufficiently narrow resonances.

The purpose of this paper is to study the properties of a weakly dissociated gas of diatomic molecules composed of fermionic atoms with a view to determining whether or not the transition temperature T_c to a paired state should exhibit a maximum in the BEC-BCS crossover. We shall consider the molecule effective mass and the reduction of the density of molecules due to dissociation, as well as the interaction between molecules. As a consequence of critical fluctuations, molecule-molecule interactions lead to an increase in T_c away from the BEC limit and this effect dominates the decreases in T_c we find due to changes in the effective mass and the number of molecules. For a trapped gas, we shall argue that T_c should decrease away from the BEC limit due to the reduction of the central density in the trap as a consequence of the increasingly repulsive molecule-molecule interaction.

This paper is organized as follows. In Sec. II we consider the uniform system, and calculate the effective mass of a molecule and thermal depletion of the number of molecules. The transition temperature is discussed in Sec. III, first for a uniform gas and then for a trapped gas. Finally, Sec. IV contains concluding remarks.

II. THE UNIFORM SYSTEM

We investigate the properties of a system containing equal numbers N of two species of fermions, labeled by an index $\sigma = \pm$, that can form a diatomic molecular bound state in the limit when the system consists mainly of molecules, with a small admixture of atoms. We shall assume that the two species of fermions have the same mass m . In the spirit of Landau, we write the effective low-energy Hamiltonian for the system as

$$\begin{aligned}
 H = & \sum_{\mathbf{p}} \left(-\varepsilon_b + \frac{p^2}{4m} \right) \hat{b}_{\mathbf{p}}^\dagger \hat{b}_{\mathbf{p}} + \sum_{\mathbf{q}\sigma} \frac{q^2}{2m} \hat{a}_{\mathbf{q},\sigma}^\dagger \hat{a}_{\mathbf{q},\sigma} \\
 & + \frac{\pi\hbar^2 a_{mm}}{mV} \sum_{\mathbf{p},\mathbf{p}',\mathbf{q}} \hat{b}_{\mathbf{p}}^\dagger \hat{b}_{\mathbf{p}'}^\dagger \hat{b}_{\mathbf{p}'+\mathbf{q}} \hat{b}_{\mathbf{p}-\mathbf{q}} \\
 & + \frac{6\pi\hbar^2 a_{am}}{mV} \sum_{\mathbf{p},\mathbf{p}',\mathbf{q}\sigma} \hat{a}_{\mathbf{p},\sigma}^\dagger \hat{b}_{\mathbf{p}'}^\dagger \hat{b}_{\mathbf{p}'+\mathbf{q}} \hat{a}_{\mathbf{p}-\mathbf{q},\sigma} \\
 & + \frac{4\pi\hbar^2 a_{aa}}{mV} \sum_{\mathbf{p},\mathbf{p}'} \hat{a}_{\mathbf{p},+}^\dagger \hat{a}_{\mathbf{p}',-}^\dagger \hat{a}_{\mathbf{p}'+\mathbf{q},-} \hat{a}_{\mathbf{p}-\mathbf{q},+} \\
 & + \frac{g}{V^{1/2}} \sum_{\mathbf{p},\mathbf{q}} (\hat{b}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{q}+\mathbf{p}/2,+} \hat{a}_{-\mathbf{q}+\mathbf{p}/2,-} + \text{H.c.}), \quad (1)
 \end{aligned}$$

where the operators \hat{a}_{σ}^\dagger , \hat{a}_{σ} , \hat{b}^\dagger , and \hat{b} create and destroy atoms and molecules, respectively. Here V is the volume of the system, g is the matrix element for decay of a molecule into two atoms, a_{ij} is the scattering length between a particle i and a particle j , and subscript ‘‘a’’ stands for an atom and ‘‘m’’ for a molecule. The quantity a_{aa} is the background scattering length for atom-atom scattering due to nonresonant processes. In the case of strong coupling between the molecule and the atoms, e.g., for a broad Feshbach resonance, and in the limit of a deeply bound molecular state it has been shown that [14,15]

$$a_{ma} \approx 1.2a_{aa}, \quad a_{mm} \approx 0.6a_{aa}. \quad (2)$$

We begin by considering a uniform gas, and we first calculate the molecule spectrum. The leading contributions to the energy of a molecule are due to the molecule-molecule interactions, which give a shift $2\pi\hbar^2 n_m a_{mm}/m$ to the self-energy of a molecule in the mean field of the other molecules, and a contribution one-half of this to the average energy per molecule. Here n_m is the number density of molecules. Atom-molecule interactions contribute $6\pi\hbar^2 n_a a_{am}/m$ to the molecule self-energy from the scattering of atoms by molecules, and from dissociation of molecules into atoms an amount

$$\begin{aligned}
 \delta\varepsilon_{\mathbf{p}} &= g^2 \int \frac{d^3q}{(2\pi\hbar)^3} \frac{n_{\mathbf{p}/2+\mathbf{q}} + n_{\mathbf{p}/2-\mathbf{q}}}{\varepsilon_b + q^2/m} \\
 &= 2g^2 \int \frac{d^3q}{(2\pi\hbar)^3} \frac{n_{\mathbf{q}}}{\varepsilon_b + (\mathbf{q} + \mathbf{p}/2)^2/m}. \quad (3)
 \end{aligned}$$

The term independent of the atom distribution function $n_{\mathbf{q}}$ is not included, since this is already contained in the binding energy ε_b of two fermions in the absence of a medium. The contribution (3) is due to Pauli blocking of dissociation of molecules, which raises the energy of a molecule. As a consequence of Galilean invariance, in the absence of a medium, the effective mass of a molecule remains equal to the bare mass $M=2m$. The binding energy of a molecule with momentum zero is thus reduced by an amount

$$\delta\varepsilon_b = 2g^2 \int \frac{d^3q}{(2\pi\hbar)^3} \frac{n_{\mathbf{q}}}{\varepsilon_b + q^2/m} \approx 2g^2 \frac{n_a}{\varepsilon_b}, \quad (4)$$

where n_a is the density of atoms of both species. In the second expression we have assumed that the molecule is deeply bound in the sense that $\varepsilon_b \gg kT$. The molecule spectrum varies over momentum scales of order $(m\varepsilon_b)^{1/2}$, and therefore for the momenta of interest in the problem of Bose-Einstein condensation, which are of order $(mkT)^{1/2}$, it is sufficient to calculate only to order p^2 . The result is

$$\varepsilon_{\mathbf{p}} \approx -\varepsilon_b + \frac{2\pi\hbar^2 n_m a_{mm}}{m} + \frac{6\pi\hbar^2 n_a a_{am}}{m} + 2g^2 \frac{n_a}{\varepsilon_b} + \frac{p^2}{2M^*}, \quad (5)$$

where the effective mass of a molecule is increased by an amount given by

$$\frac{1}{M^*} = \frac{1}{2m} \left(1 - 2g^2 \frac{n_a}{\varepsilon_b^2} \right). \quad (6)$$

We now turn to the transition temperature of the gas.

III. TRANSITION TEMPERATURE

A. Uniform system

For a uniform gas of elementary bosons, the transition temperature is unaffected by the Hartree mean-field energy, since this gives a shift to the energy of a boson and to the chemical potential which are independent of the momentum of the particle. However, it has been demonstrated that, due to critical fluctuations, the transition temperature is increased, by an amount which is given for small $n_m^{1/3} a_{mm}$ by [16,17]

$$\frac{\delta T_c}{T_c} = b n_m^{1/3} a_{mm}, \quad (7)$$

where b is a positive coefficient, which has been estimated to be approximately 1.3 [18–20]. This effect will tend to be counteracted by the increase of the effective mass and the reduction of the density of molecules due to dissociation, both of which tend to decrease the transition temperature, since for an ideal Bose gas $T_c \sim \hbar^2 n_m^{2/3}/2M^*$. The number density of molecules is given by

$$n_m = \frac{N}{V} - \frac{n_a}{2}. \quad (8)$$

The two latter effects give changes in the transition temperature which are proportional to $n_a/n_m \propto e^{-\varepsilon_b/kT_c}$. Thus, for a deeply bound molecule, the effects of critical fluctuations dominate, and the transition temperature initially increases away from the BEC limit. Since in the BCS limit the transition temperature tends to zero, we conclude that the transition temperature must have a local maximum, provided that the transition temperature is a continuous function of the scattering length.

B. Trapped cloud

We now turn to the case of a cloud in a harmonic trap, and we shall assume that the trapping potentials for the two spe-

cies of fermions are the same, and equal to one-half of the trapping potential for a molecule. Previously, the transition temperature for a trapped gas of bosons has been calculated with allowance for the nonzero spacing of the single-particle levels in the oscillator potential and for the distortion of the density profile by the interaction, and the result is [21]

$$\frac{\delta T_c}{T_c} \approx -0.73 \frac{\omega_m}{\bar{\omega}} N^{-1/3} - 1.33 \frac{a_{\text{mm}}}{\bar{a}} N^{1/6}, \quad (9)$$

where ω_m is the algebraic mean of the trap frequencies for the three principal axes of the trap, $\bar{\omega}$ is their harmonic mean, and $\bar{a} = (\hbar/m\bar{\omega})^{1/2}$ is the mean harmonic oscillator length.

We now estimate the effect of critical fluctuations by making a local density approximation, that is, by inserting in Eq. (7) the central density $n_m(0)$ at T_c , which is given in the absence of interactions by

$$n_m(0) = \zeta(3/2) \left(\frac{MkT_c}{2\pi\hbar^2} \right)^{3/2}. \quad (10)$$

Since $kT_c = [N/\zeta(3)]^{1/3} \hbar \bar{\omega}$, it follows that the contribution to the change in T_c due to critical fluctuations is

$$\frac{\delta T_c}{T_c} \approx \frac{1.3}{(2\pi)^{1/2}} \frac{[\zeta(3/2)]^{1/3}}{[\zeta(3)]^{1/6}} N^{1/6} \frac{a_{\text{mm}}}{\bar{a}}. \quad (11)$$

Detailed calculations indicate that critical fluctuations are suppressed in traps by positive powers of $n_m^{1/3} a_{\text{mm}}$ [18,22], so the local density approximation gives an upper limit to the increase in T_c . Adding this to the earlier result (9) without fluctuations, we find for the total change in T_c

$$\frac{\delta T_c}{T_c} \approx -0.73 \frac{\omega_m}{\bar{\omega}} N^{-1/3} - 0.64 \frac{a_{\text{mm}}}{\bar{a}} N^{1/6}. \quad (12)$$

Thus, in contrast to the uniform case, the change in T_c for small a_{mm} is *negative*, since the effect of critical fluctuations is overwhelmed by the reduction of the central density.

The discussion we have given above applies to broad Feshbach resonances, for which the energy dependence of the atomic interactions may be neglected. For narrow Fesh-

bach resonances, critical fluctuations are suppressed because the magnitude of the contribution from processes in which the argument of the interaction contains a frequency which is to be summed over will be reduced because of the strong reduction of the interaction away from resonance, as discussed in Ref. [23].

IV. CONCLUSION

In this paper we have identified a number of effects that produce changes in the transition temperature away from the BEC limit. Two of them are due to thermal dissociation of molecules, and they lead to a decrease in T_c . However, for a uniform gas, these effects are overwhelmed by a rise in T_c due to the increasingly repulsive molecule-molecule interaction. We therefore conclude, quite generally, that T_c must display a global maximum between the BEC and BCS limits for a broad Feshbach resonance. For trapped gases, we have argued that T_c will initially decrease away from the BEC limit. Thus, on the basis of the arguments given here, there need not be a maximum in T_c but it cannot be excluded. Likewise, for a narrow Feshbach resonance, critical fluctuations are suppressed, and consequently T_c is expected to decrease away from the BEC limit.

The framework we have described in this paper could be a useful tool for analysis of detailed crossover calculations, since it points to a number of quantities that could be calculated in order to test our fundamental understanding.

ACKNOWLEDGMENTS

This work was carried out under the auspices of a Nordita Nordic Project on ultracold atoms, and H. M. wishes to thank Nordita for hospitality. L.M.J. was supported by Nordita and by the Academy of Finland and EUROHORCS (EURYI, Academy Projects Nos. 106299, 205470, and 207083). H.M. was supported by the Finnish Academy of Science and Letters (Ville, Yrjö and Kalle Väisälä Foundation) and the Academy of Finland (Grant No. 206108). We are grateful to G. Watanabe and H.M. Nilsen for useful conversations and S. Giorgini for helpful comments.

-
- [1] D. M. Eagles, *Phys. Rev.* **186**, 456 (1969).
 [2] A. J. Leggett, in *Modern Trends in the Theory of Condensed Matter*, edited by A. Pekalski and R. Przystawa (Springer-Verlag, Berlin, 1980), p. 13.
 [3] P. Nozières and S. Schmitt-Rink, *J. Low Temp. Phys.* **59**, 195 (1985).
 [4] M. Drexler and W. Zwerger, *Ann. Phys.* **1**, 15 (1992).
 [5] C. A. R. Sá de Melo, M. Randeria, and J. R. Engelbrecht, *Phys. Rev. Lett.* **71**, 3202 (1993).
 [6] R. Haussmann, *Phys. Rev. B* **49**, 12975 (1994).
 [7] R. Haussmann, W. Rantner, S. Cerrito, and W. Zwerger, *Phys. Rev. A* **75**, 023610 (2007).
 [8] J. N. Milstein, S. J. J. M. F. Kokkelmans, and M. J. Holland, *Phys. Rev. A* **66**, 043604 (2002).
 [9] Y. Ohashi and A. Griffin, *Phys. Rev. Lett.* **89**, 130402 (2002).
 [10] Y. Ohashi and A. Griffin, *Phys. Rev. A* **67**, 033603 (2003).
 [11] Q. Chen, J. Stajic, and K. Levin, *Phys. Rep.* **412**, 1 (2005).
 [12] A. Perali, P. Pieri, L. Pisani, and G. C. Strinati, *Phys. Rev. Lett.* **92**, 220404 (2004).
 [13] G. M. Falco and H. T. C. Stoof, eprint cond-mat/0603270, *Phys. Rev. A* (to be published).
 [14] G. Skorniyakov and K. A. Ter-Martirosian, *Zh. Eksp. Teor. Fiz.* **31**, 775 (1956) [*Sov. Phys. JETP* **4**, 648 (1957)].
 [15] D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, *Phys. Rev. Lett.* **93**, 090404 (2004).
 [16] G. Baym, J.-P. Blaizot, M. Holzmann, F. Laloë, and D. Vautherin, *Phys. Rev. Lett.* **83**, 1703 (1999).

- [17] The importance of critical fluctuations in tending to increase T_c has also been pointed out in Ref. [7].
- [18] P. Arnold and G. Moore, Phys. Rev. Lett. **87**, 120401 (2001).
- [19] V. A. Kashurnikov, N. V. Prokof'ev, and B. V. Svistunov, Phys. Rev. Lett. **87**, 120402 (2001).
- [20] In the calculations of the critical fluctuations, the energy dependence of the effective interaction is neglected and consequently the result is appropriate for a broad Feshbach resonance. We shall later return to the case of a narrow resonance.
- [21] S. Giorgini, L. P. Pitaevskii, and S. Stringari, Phys. Rev. A **54**, R4633 (1996).
- [22] O. Zobay, G. Metikas, and H. Kleinert, Phys. Rev. A **71**, 043614 (2005).
- [23] A. Schwenk and C. J. Pethick, Phys. Rev. Lett. **95**, 160401 (2005).