# Quantum origin of quantum jumps: Breaking of unitary symmetry induced by information transfer in the transition from quantum to classical

Wojciech Hubert Zurek

Theory Division, MS B213, LANL Los Alamos, New Mexico 87545, USA

(Received 26 March 2007; revised manuscript received 10 September 2007; published 16 November 2007)

Measurements transfer information about a system to the apparatus and then, further on, to observers and (often inadvertently) to the environment. I show that even imperfect copying essential in such situations restricts possible unperturbed outcomes to an orthogonal subset of all possible states of the system, thus breaking the unitary symmetry of its Hilbert space implied by the quantum superposition principle. Preferred outcome states emerge as a result. They provide a framework for "wave-packet collapse," designating terminal points of quantum jumps and defining the measured observable by specifying its eigenstates. In quantum Darwinism, they are the progenitors of multiple copies spread throughout the environment—the fittest quantum states that not only survive decoherence, but subvert the environment into carrying information about them—into becoming a witness.

DOI: 10.1103/PhysRevA.76.052110

PACS number(s): 03.65.Ta, 03.65.Yz, 03.67.-a

### I. INTRODUCTION: QUANTUM AXIOMS

The quantum principle of superposition applies to isolated systems, but is famously violated in the course of measurements: A quantum system can exist in any superposition, but a measurement forces it to choose from a limited set of outcomes represented by an orthogonal set of states [1]. In textbook discussions of quantum theory these states are preassigned as "the eigenstates of the measured observable." They are the only possible states that can ever be detected (and therefore prepared) by that measuring device. As Dirac puts it [2], "...a measurement always causes the system to jump into an eigenstate of the dynamical variable that is being measured...."

I show, using ideas that parallel the no-cloning theorem, that this restriction (usually imposed "by decree," by the collapse postulate) can be derived when a transfer of information essential for both measurement and decoherence is modeled as a unitary quantum process that leads to records with predictive significance. This sheds new light on quantum measurements, on the quantum-classical transition and on the selection of preferred pointer states by the environmentinduced decoherence [3-6]: It shows that a restriction to a limited set of orthogonal states (but, of course, not the nonunitary collapse *per se*) can be deduced in a setting that does not assume a priori existence of macroscopic apparatus usually invoked to define the measured observable [1,2,7]. This connection between the transfer of information and the selection of preferred states is also of crucial importance for quantum Darwinism [5], where the capacity of the environment to acquire multiple records of selected states of the system is essential. Resilient states that can withstand monitoring by the environment without getting disturbed become endowed with objective reality: They are simultaneously accessible to many observers through the imprints they leave in the environment.

Quantum theory is founded on several simple postulates [2,5-10]. The first two encapsulate the quantum principle of superposition and decree unitarity of evolutions. Thus the axioms (i) *The quantum state of a system is represented by* 

a vector in its Hilbert space and (ii) quantum evolutions are unitary (e.g., generated by the Schrödinger equation) describe the mathematical formalism of the theory. To make contact with the real world one needs to relate abstract unitarily evolving quantum states to experiments. The next postulate, (iii) *immediate repetition of a measurement yields the same outcome*, is the first of the "measurement postulates." It is uncontroversial—states of classical systems satisfy it. It establishes a predictive utility of quantum states that have already entered through postulate (i). One can rephrase the predictability postulate by noting that when a state is known (for example, because of an earlier measurement), then one can choose a measurement that will confirm it but leave it intact.

These first three postulates indicate no bias—they treat every state in the Hilbert space of the system on equal footing. In particular, postulates (i) and (ii) emphasize linearity: Any superposition of quantum states is a legal quantum state, and any evolution of such a superposition leads to a superposition of evolved "ingredients." By contrast, the last two postulates, (iv) measurement outcomes are restricted to an orthonormal set { $|s_k\rangle$ } of eigenstates of the measured observable (i.e., measurement does not reveal the state of the system because it limits possible outcomes to the preassigned outcome states) and (v) the probability of finding a given outcome is  $p_k = |\langle s_k || \psi \rangle|^2$ , where  $|\psi\rangle$  is the preexisting state of the system, are at the heart of the long-standing disputes on the interpretation of quantum theory [1–10].

As a consequence of (iv) (the "collapse postulate") it is impossible to find out an unknown state of a quantum system: In contrast to the egalitarian postulates (i)–(iii), explicit symmetry breaking implied by (iv) defies the superposition principle by allowing only certain states as outcomes. The aim of this paper is to point out that already the (symmetric and uncontroversial) postulates (i)–(iii) necessarily imply selection of *some* preferred set of *orthogonal* states—that they impose the broken symmetry that is at the heart of the collapse postulate (iv)—although they stop short of specifying what this set of outcome states is, and obviously cannot result in anything explicitly nonunitary (e.g., the actual "collapse").

## **II. PREDICTABILITY AND SYMMETRY BREAKING**

To see how the symmetry-breaking restriction to a set of orthonormal outcome states arises from symmetric and uncontroversial assumptions (i)–(iii) consider a system S, to be measured by an apparatus A. For simplicity, we start with the smallest possible S with a two-dimensional Hilbert space  $\mathcal{H}_S$  (although our derivation will be extended to arbitrarily large systems). In a two-dimensional  $\mathcal{H}_S$  any state of the system can be written as a superposition of a pair of linearly independent states,  $|\psi_S\rangle = \alpha |v\rangle + \beta |w\rangle$ . This freedom to choose any two linearly independent  $|v\rangle$  and  $|w\rangle$  that need *not* be orthogonal is guaranteed by the principle of superposition implicit in postulate (i). A measurement of S by an apparatus A that starts in a "ready-to-measure" state  $|A_0\rangle$ ,

$$|\psi_{\mathcal{S}}\rangle|A_{0}\rangle = (\alpha|v\rangle + \beta|w\rangle)|A_{0}\rangle \Rightarrow \alpha|v\rangle|A_{v}\rangle + \beta|w\rangle|A_{w}\rangle = |\Phi_{\mathcal{S}\mathcal{A}}\rangle,$$
(1)

yields the desired information transfer from S to A: The state of A now contains a record of S. Above we assumed postulate (iii) (there are states that are left intact,  $|v\rangle|A_0\rangle \rightarrow |v\rangle|A_v\rangle$ ,  $|w\rangle|A_0\rangle \rightarrow |w\rangle|A_w\rangle$ ) and linearity that follows from postulate (ii). We have also recognized that the state of a composite quantum system (here SA) is a vector in the tensor product of constituent Hilbert spaces [i.e., has a form of Eq. (1)], an observation often regarded as an additional "complexity postulate (o)."

The norm of the state of the composite system must be preserved: Postulate (i) demands it (Hilbert space is a *normed* linear space), and postulate (ii) ensures that this demand is met. Thus, simple algebra and the recognition that  $\langle A_0 || A_0 \rangle = \langle A_w || A_w \rangle = 1$  yields

$$\langle \psi_{\mathcal{S}} || \psi_{\mathcal{S}} \rangle - \langle \Phi_{\mathcal{S}\mathcal{A}} || \Phi_{\mathcal{S}\mathcal{A}} \rangle = 2 \operatorname{Re} \alpha^* \beta \langle v || w \rangle (1 - \langle A_v || A_w \rangle) = 0.$$
(2a)

Equation (2a) involves the real part only, but must hold for arbitrary relative phase between  $\alpha$  and  $\beta$ . This leads to

$$\langle v \| w \rangle (1 - \langle A_v \| A_w \rangle) = 0.$$
 (2b)

This equality is the basis for our further discussion.

Depending on the overlap  $\langle v || w \rangle$  there are two possibilities. Let us first suppose that  $\langle v || w \rangle \neq 0$  (but is otherwise arbitrary). In this case one is forced to conclude that the state of  $\mathcal{A}$  cannot be affected by the process above. That is, the transfer of information from  $\mathcal{S}$  to  $\mathcal{A}$  must have failed completely, since  $\langle A_v || A_w \rangle = 1$  must hold: The apparatus can bear no imprint that distinguishes between the components of the superposition  $|\psi_{\mathcal{S}}\rangle$ —the prospective outcome states of the system.

The second possibility is that  $\langle v | | w \rangle = 0$ . This allows for an arbitrary  $\langle A_v | | A_w \rangle$ , including a perfect record,  $\langle A_v | | A_w \rangle = 0$ . Thus, outcome states  $|v\rangle$  and  $|w\rangle$  must be orthogonal if—in accordance with postulate (iii)—they are to survive intact a successful information transfer in general or a quantum measurement in particular, so that the immediate remeasurement can yield the same result. The same derivation can be carried out for S with a Hilbert space of dimension  $\mathcal{N}$  starting with a system state vector  $|\psi_S\rangle = \sum_{k=1}^{\mathcal{N}} \alpha_k |s_k\rangle$ . As before, a priori

the set of basis states  $\{|s_k\rangle\}$  need not be orthogonal, but only linearly independent.

The simple derivation above leads to a very decisive conclusion: Orthogonality of outcome states of the system is absolutely essential for them to exert distinct influences—to imprint even a minute difference—on the state of any other system while retaining their identity: The overlap  $\langle v || w \rangle$ must be 0 *exactly* for  $\langle A_v || A_w \rangle$  to differ from unity. Thus, also sloppy and accidental information transfers (e.g., to the environment during decoherence [3–6]) will define preferred sets of mutually orthogonal states provided that the crucial predictability demand of postulate (iii) is met. This "had to be so:" When an imperfect measurement can be repeated many times without affecting the original, collective records will come arbitrarily close to orthogonality.

Selection of an orthonormal basis induced by information transfer—the need for a spontaneous symmetry breaking that arises from the otherwise symmetric axioms (i)—(iii)—is a general and intriguing result. Our derivation parallels the proof of the no-cloning theorem [11–13]: We have employed the assumption of linearity and started with Eq. (1), as does the proof of no-cloning in Refs. [11,12]. Moreover, linearity and preservation of the norm follow from the unitarity postulate (ii) used in the alternative proof of no-cloning [see Ref. [13]; the only difference is that when copies are "clones"  $\langle v || w \rangle = \langle A_v || A_w \rangle$  in Eq. (2b)].

Similar reasoning appears in a proof of the security of cryptographic protocols [14]. There, however, the focus is on the ability to detect eavesdropper through perturbations her measurements inflict on the transmitted nonorthogonal states, which is rather different from the questions considered here. Nevertheless, connections between quantum prohibition on cloning, information gain and disturbance trade-off (exemplified by Ref. [14]), and our proof of orthogonality of outcome states are hard to miss. However, the implications we are led to are quite different.

The scope of our result is closer to the study of measurements due to Wigner [7]. He argued that record states of a *classical* apparatus must be orthogonal. He inferred this orthogonality from the "Copenhagen assumption"—that an apparatus must be classical [1]—and translated classical distinguishability of pointer positions into their orthogonality into  $|\langle A_v || A_w \rangle| = 0$ . With the additional assumption of real eigenvalues, Wigner concluded that observables must correspond to self-adjoint operators. We did not use such a strong "Copenhagen-motivated" assumption— $|\langle A_v || A_w \rangle| \neq 1$  suffices to establish our result. Therefore, when we assume that eigenvalues should be real, the Hermitian nature of observables immediately follows from axioms (i)–(iii), without any need to appeal to the classicality of measuring devices.

The reader may be concerned about idealizations we have made to represent information transfer by Eq. (1). We now briefly explore these assumptions, see how they can be relaxed, and consider what this implies about terminal states for "quantum jumps."

Let us start from the most obvious: The apparatus (or the environment monitoring the system in course of decoherence) is usually *not* in a pure state at the beginning, so one should represent it with a density matrix,  $\rho_0^{\mathcal{A}} = \sum_k p_k |a_k\rangle \langle a_k|$ .

To deal with this we note that such a mixed state can be always "purified" by allowing  $\mathcal{A}$  to have a "ghost partner"  $\mathcal{A}'$ , so that the combined initial state of the two is given by  $|\mathbf{A}_0\rangle = \sum_k \alpha_k |a_{0,k}\rangle |a'_{0,k}\rangle$  in the obvious notation, and with  $|\alpha_k|^2 = p_k$ . Orthogonality of the outcome states can be now established starting with the equation

$$\langle v \| w \rangle (1 - \langle \mathbf{A}_v \| \mathbf{A}_w \rangle) = 0.$$
 (2c)

As before, this leads to the conclusion that  $|v\rangle$  and  $|w\rangle$  [or any two outcome states that satisfy postulates (i)–(iii)] must be orthogonal if the information transfer is to succeed. Therefore, relaxing this assumption does not change our conclusions. This is also the case when the environment  $\mathcal{E}$  is influenced directly by  $\mathcal{S}$ .

In particular, we did not invoke postulate (v)—Born's rule—which is necessary to attribute physical significance to reduced density matrices employed in the study of decoherence [3–6]. This is obvious when  $\mathcal{A}$  is pure. Still, one might worry that "purification" relating a mixed state of  $\mathcal{A}$  to a pure state of enlarged  $\mathcal{A}\mathcal{A}'$  uses Born's rule "in the reverse direction." This is "only mathematics" ( $p_k$  need not be regarded as probabilities), and it is not essential: It is easy to see that *any* prescription relating pure states of  $\mathcal{A}\mathcal{A}'$  to mixed states of  $\mathcal{A}$  will do, as long as it is based on a one-to-one correspondence between eigenvalues of the density operator of  $\mathcal{A}$  and absolute values of amplitudes in the entangled state of  $\mathcal{A}\mathcal{A}'$ , and as long as it is used in both directions. So even  $|\alpha_k|^2 = p_k$  (probability interpretation not-withstanding) is not needed.

This independence of unitary symmetry breaking responsible for quantum jumps from Born's rule is especially important when one aims to establish the extent to which "controversial" postulates (iv) and (v) follow from the generally accepted axioms (o)–(iii). In this context it is reassuring that the proof can be carried out without "purification." To this end, we note that unitary evolution preserves scalar product of operators defined as Tr  $\rho\tilde{\rho}$ . Therefore, in the obvious notation, Tr  $|v\rangle\langle v|\rho_{A_0}|w\rangle\langle w|\rho_{A_0}=\text{Tr} |v\rangle\langle v|\rho_{A_u}|w\rangle\langle w|\rho_{A_u}$ . Thus,

$$|\langle v||w\rangle|^2 (\operatorname{Tr} \rho_{\mathcal{A}_0}^2 - \operatorname{Tr} \rho_{\mathcal{A}_v} \rho_{\mathcal{A}_w}) = 0.$$
 (2d)

So either  $\langle v || w \rangle = 0$  (we recover orthogonality of outcomes) or Tr  $\rho_{\mathcal{A}_0}^2 = \text{Tr} \rho_{\mathcal{A}_v} \rho_{\mathcal{A}_w}$ . This last equality holds if and only if  $\rho_{\mathcal{A}_v} = \rho_{\mathcal{A}_w}$ . This is because  $\rho_{\mathcal{A}_0} = \sum_k p_k |a_{k0}\rangle \langle a_{k0}|$ , as well as  $\rho_{\mathcal{A}_v} = \sum_k p_k |a_{kv}\rangle \langle a_{kv}|$ , and  $\rho_{\mathcal{A}_w} = \sum_k p_k |a_{kw}\rangle \langle a_{kw}|$  have the same eigenvalues  $\{p_k\}$ —they are related to one another by a unitary evolution. Therefore, only their eigenvectors can differ. But then  $\text{Tr}(\rho_{\mathcal{A}_0}^2 - \rho_{\mathcal{A}_v}\rho_{\mathcal{A}_w}) = \frac{1}{2}\text{Tr}(\rho_{\mathcal{A}_v} - \rho_{\mathcal{A}_w})^2$ , which is nonnegative and vanishes iff  $\rho_{\mathcal{A}_v} = \rho_{\mathcal{A}_w}$ —only when there is no record left in  $\mathcal{A}$ .

This establishes that only orthogonal states can leave imprints on other systems without getting disrupted also in the case when the state of that apparatus or the environment is mixed and represented by a Hermitian operator. Born's rule is not invoked: We make no claim that  $p_k$ 's are probabilities or even relate them to amplitudes. We only recognize that—in addition to pure states given by state vectors—there are mixed states given by Hermitian operators. There is no appeal to the physical significance of reduced density matrices. The whole proof parallels our original pure state case step by step. And, as before, it leads to the special set of orthogonal stable states, reminiscent of pointer states. It is good to derive them without any appeal to probabilities: One can then use them—without any danger of circularity—as "events" while deriving Born's rule.

On the other hand, we cannot really drop any of the first three postulates. This is obvious for the superposition principle of postulate (i) and for the unitarity postulate (ii)—they define quantum theory [although (ii) could be weakened, e.g., by allowing antilinear and antiunitary evolutions]. One can, however, relax the demands of postulate (iii). It is one of the standard axioms, and, in principle, its demands can be always met. Moreover, the ability to reconfirm outcomes of measurements encapsulates predictability that is essential to introduce the very concept of a "state" to describe S. So it is appropriate to rely on (iii) in a discussion of quantum foundations. However, its central demand—that measurements should not perturb the measured system—is only rarely met in laboratory experiments. It is therefore interesting to consider more general measurement schemes—e.g.,

$$(\alpha | v \rangle + \beta | w \rangle) | A_0 \rangle \Rightarrow \alpha | \tilde{v} \rangle | A_v \rangle + \beta | \tilde{w} \rangle | A_w \rangle = | \Phi_{SA} \rangle.$$
(3)

When  $|\tilde{v}\rangle$  and  $|\tilde{w}\rangle$  are related with their progenitors by a transformation that preserves scalar product (e.g., by a dynamical evolution in a closed system) proof of orthogonality goes through unimpeded. Both unitary and antiunitary transformations are in this class. We can also imagine situations when this is not the case, and  $\langle v \| w \rangle \neq \langle \tilde{v} \| \tilde{w} \rangle$ . An extreme example of this arises when the state of the measured system retains no memory of what it was beforehand (e.g.,  $|v\rangle \Rightarrow |0\rangle$ ,  $|w\rangle \Rightarrow |0\rangle$ ). Then the apparatus can (and, indeed, by unitarity, has to) "inherit" the information that was contained in the state of the system. Clearly, the need for orthogonality of outcomes disappears. Of course, such measurements do not fulfill axiom (iii)-they are not repeatable. An example of this situation is encountered in quantum optics. Photons are usually absorbed by detectors, and coherent states (which are not orthogonal) are the outcome states.

It is also interesting to consider sequences of information transfers:

$$|v\rangle|A_{0}\rangle|B_{0}\rangle\cdots|E_{0}\rangle \Rightarrow |\tilde{v}\rangle|A_{v}\rangle|B_{0}\rangle\cdots|E_{0}\rangle$$
$$\Rightarrow \cdots \Rightarrow |\tilde{v}\rangle|\tilde{A}_{v}\rangle|\tilde{B}_{v}\rangle\cdots|E_{v}\rangle, \qquad (4a)$$

$$|w\rangle|A_{0}\rangle|B_{0}\rangle\cdots|E_{0}\rangle \Rightarrow |\tilde{w}\rangle|A_{w}\rangle|B_{0}\rangle\cdots|E_{0}\rangle$$
$$\Rightarrow \cdots \Rightarrow |\tilde{w}\rangle|\tilde{A}_{w}\rangle|\tilde{B}_{w}\rangle\cdots|E_{w}\rangle. \tag{4b}$$

Such "von Neumann chains" [10] appear in quantum measurements, environment-induced decoherence, amplification, and in quantum Darwinism. As information is passed along this chain, links can be perturbed (as indicated by tilde). Unitarity implies that, at each stage, products of overlaps must be the same,

$$\langle v || w \rangle = \langle \widetilde{v} || \widetilde{w} \rangle \langle \widetilde{A}_{v} || \widetilde{A}_{w} \rangle \langle \widetilde{B}_{v} || \widetilde{B}_{w} \rangle \cdots \langle E_{v} || E_{w} \rangle.$$
(5)

When a logarithm of both side is taken,

$$\ln|\langle v||w\rangle|^{2} = \ln|\langle \tilde{v}||\tilde{w}\rangle|^{2} + \ln|\langle \tilde{A}_{v}||\tilde{A}_{w}\rangle|^{2} + \ln|\langle \tilde{B}_{v}||\tilde{B}_{w}\rangle|^{2} + \cdots + \ln|\langle E_{v}||E_{w}\rangle|^{2}.$$
(6)

Therefore, when  $\langle v \| w \rangle \neq 0$ , as the information about the outcome is passed along the two von Neumann chains, the quality of the records must suffer: The sum of logarithms above must equal  $\ln |\langle v \| w \rangle|^2$ , and the overlap of the two states is a measure of their distinguishability. For orthogonal states there is no need for such deterioration of the quality of records, as  $\ln |\langle v \| w \rangle|^2 = -\infty$ , so arbitrarily many orthogonal records can be made.

### **III. DISCUSSION**

Selection of a preferred basis implied by postulates (i)– (iii) brings to mind einselection—the choice of preferred pointer states in decoherence [15]. Moreover, amplification is precisely such a multiplication of records. Quantum Darwinism [5,16,17] is based on the observation that decoherence typically leads to amplification. The main difference between the above discussion and einselection is the point of departure: Our results follow from very basic assumptions and lead to very general conclusions. Moreover, we only establish that *some* orthonormal basis is needed to define measurement outcomes, while einselection shows what specific pointer basis will emerge given, e.g., the coupling with the environment.

Note that such pointer states einselected for their resilience (in accordance with the original definition [15] or with its generalization in terms of the predictability sieve [5,19]) are *not* guaranteed to be the eigenstates of the reduced density matrix  $\rho_S = \text{Tr}_{\mathcal{E}} |\Psi_{S\mathcal{E}}\rangle \langle \Psi_{S\mathcal{E}}|$  of the system, where  $|\Psi_{S\mathcal{E}}\rangle = \alpha |v\rangle |\varepsilon_v\rangle + \beta |w\rangle |\varepsilon_w\rangle$  by analogy with Eqs. (1) and (3). This is easy to see: States that are left unperturbed by the information transfer will coincide with the Schmidt states of  $|\Psi_{S\mathcal{E}}\rangle$  (which are on the diagonal of  $\rho_S$ ) only when their copies in the environment are perfect—only when scalar products  $\langle \varepsilon_v || \varepsilon_w \rangle$  of the corresponding "record states" vanish.

We take this to mean that only in that case can one safely attribute the usual probability interpretation to events associated with pointer states [22,23]. This can be illustrated in a setting where the system S is first entangled with the apparatus and then A is decohered by its environment. Orthogonality that is implied by the resilience of pointer states in spite of the immersion of A in  $\mathcal{E}$  is then imposed on the pointer states of the apparatus (so that they can retain measurement records). Decoherence will suppress off-diagonal terms representing quantum coherences between records stored in pointer states [15]. This allows for the description of, e.g., AS correlations in terms of classical probabilities, as is indicated by the vanishing of *quantum discord* [18]—of the difference between two classically equivalent definitions of mutual information that arises in quantum setting.

Resilience of pointer states is especially important in the "environment as a witness" paradigm [5,16,17] based on recognition of the indirect manner used by the real world observers (such as the reader of this text) to acquire informa-

tion. Such an observer does not perform direct measurements-the reader is not *directly* interacting with this text. Rather, he or she acquires information by intercepting a small fraction of the photon environment that was emitted by (or scattered from) the text appearing on the computer screen (or printed on a sheet of paper). Thus, von Neumann's model of direct measurement [10] involving a quantum system and isolated quantum apparatus is a gross oversimplification. Even inclusion of the environment to account for decoherence (clearly a step in the right direction) does not capture the essential role of  $\mathcal{E}$  for the transition from quantum to classical: Environment is not just a "garbage disposal" for the unwanted quantum coherence, but a communication channel and a witness to the state of the system. Observers acquire their information about the universe by intercepting fragments of  $\mathcal{E}$ .

Quantum Darwinism—proliferation of selected information about the system throughout the environment—is a frequent byproduct of decoherence. Decoherence is caused by monitoring of the system by its environment. Quantum Darwinism leads to a deposition of redundant records—multiple "copies" of the selected states of the system. And only states that survive this monitoring intact can lead to such *redundant* imprinting in the environment. The existence of multiple records accounts for the objective nature of pointer observables and explains (through arguments put forward elsewhere [5,16,17,22]) why superpositions of pointer states are inaccessible. Exploring these connections promises to be fruitful and is especially relevant in the context of interpretations of quantum mechanics that do not invoke explicit "collapse" [20–22].

In the presence of einselection the reduced density matrix is *locally* equivalent to a probability distribution of events defined by pointer states: Local observers cannot tell when such a mixture represents a state that is already known to someone else (and in that sense definite) or when it is a part of a larger entangled whole (and, hence, unknown in principle to anyone [22,23]). This "quantum principle of local equivalence" can be compared to the famous equivalence of the gravitational pull and *bona fide* acceleration: When it is locally impossible to distinguish two globally distinct situations (e.g., speeding up in a rocket and being at rest near a mass), such local symmetry implies equivalence of all local physical consequences.

There is a different setting in which symmetric assumptions lead to asymmetric outcomes: In symmetry breaking phase transitions, a symmetric initial state and a symmetrypreserving equation lead to a configuration that breaks that symmetry. In thermodynamic transitions this choice can be attributed to thermal fluctuations. In quantum phase transitions spontaneous symmetry breaking is sometimes blamed on "quantum fluctuations," but this is a meaningless statement. Classical (e.g., thermal) fluctuations can break the symmetry of the problem, so in an ensemble of thermodynamic systems average over all possible fluctuations can be symmetric, but each member can be a subject to an "asymmetric" fluctuation. By contrast, "quantum noise" is a symmetric superposition of all possibilities, so choice cannot be explained any more than "collapse"-selection of a specific outcome—can be explained by postulates (i)-(iii).

In quantum phase transitions—e.g., in quantum Ising model (Ref. [24]), where the decrease of external bias leads from a unique polarized state  $|\rightarrow\rightarrow\rightarrow\rightarrow\rightarrow\rightarrow\rightarrow\ldots\rangle$  aligned with a bias field to two "ferromagnetic" alternatives selected by the Ising coupling,  $|\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\ldots\rangle$  and  $|\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\ldots\rangle$ —the true ground state is degenerate. Any superposition of these two alternatives will do for an infinite system, although in a finite case symmetric  $|\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\uparrow\ldots\rangle+|\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\downarrow\ldots\rangle$  is energetically favored.

Our analysis of symmetry breaking—the essence of the collapse postulate (iv)—fits a similar framework. Postulates (i)–(iii), when used to express information transfer, imply symmetry breaking: They show that many copies can be made, but only of a certain (orthogonal) subset of all possible states in the Hilbert space. These alternatives provide a "menu"—a set of choices—but [as a consequence of linearity implied by postulate (ii)] quantum theory stops short of selecting any one of them.

Similar conclusions hold in the less ordered but qualitatively similar case of *branching states* [17], e.g.,  $|\varphi\rangle = \alpha |v\rangle |\varepsilon_v^{(1)}\rangle |\varepsilon_v^{(2)}\rangle \dots |\varepsilon_v^{(n)}\rangle + \beta |w\rangle |\varepsilon_w^{(1)}\rangle |\varepsilon_w^{(2)}\rangle \dots |\varepsilon_w^{(n)}\rangle$ . Such states arise naturally [e.g., via Eq. (1)] in the course of unitary evolutions that lead to decoherence [5,16,17,22]. Differences between  $|\phi\rangle$  and  $|\varphi\rangle$  are merely quantitative. For instance, in  $|\varphi\rangle$ , the "copies" deposited in the subenvironments will be generally imperfect, so it will take more measurements to find out the branch. Moreover, information will often spread beyond the primary environment-for instance, information about text printed on a page may reside both in the photon environment (in an easy to extract form) and in the scattered air molecules (where it is scrambled by scattering). This may affect the ease with which one can extract information about the "system of interest" from these two environments, but it does not change our basic conclusion: The essence of the controversial collapse postulate (iv)-symmetry breaking that makes it impossible to detect an unknown preexisting quantum state-arises from the uncontroversial and generally accepted quantum postulates (principle of superposition, unitarity of evolutions, and predictability) in a manner that settles much of the long-standing confusion about the role and origin of "quantum jumps" and inevitability of "collapse" in quantum measurements.

### ACKNOWLEDGMENT

I thank Richard Jozsa for discussion that stimulated this paper and for comments on its early version.

- [1] N. Bohr, Nature (London) **121**, 580 (1928).
- [2] P. A. M. Dirac, *The Principles of Quantum Mechanics* (Oxford University Press, New York, 1958).
- [3] W. H. Zurek, Phys. Today 44(10), 36 (1991); see also an update, e-print arXiv:quant-ph/0306072.
- [4] E. Joos, H. D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu, *Decoherence and the Appearancs of a Classical World in Quantum Theory* (Springer, Berlin, 2003).
- [5] W. H. Zurek, Rev. Mod. Phys. 75, 715 (2003).
- [6] M. Schlosshauer, Rev. Mod. Phys. 76, 1267 (2004); Decoherence and the Quantum-to-Classical Transition (Springer, Berlin, 2007).
- [7] E. P. Wigner, in *Quantum Theory and Measurement*, edited by J. A. Wheeler and W. H. Zurek (Princeton University Press, Princeton, 1983), pp. 260–314.
- [8] E. Farhi, J. Goldstone, and S. Guttmann, Ann. Phys. (N.Y.) 24, 118 (1989).
- [9] Quantum Theory and Measurement, edited by J. A. Wheeler and W. H. Zurek (Princeton University Press, Princeton, 1983).
- [10] J. von Neumann, *Mathematical Foundations of Quantum Theory* (Princeton University Press, Princeton, 1955).
- [11] W. K. Wootters and W. H. Zurek, Nature (London) 299, 802 (1982).

- [12] D. Dieks, Phys. Lett. **92A**, 271 (1982).
- [13] H. P. Yuen, Phys. Lett. **113A**, 405 (1986).
- [14] C. H. Bennett, G. Brassard, and N. D. Mermin, Phys. Rev. Lett. 68, 557 (1992).
- [15] W. H. Zurek, Phys. Rev. D 24, 1516 (1981); 26, 1862 (1982).
- [16] H. Ollivier, D. Poulin, and W. H. Zurek, Phys. Rev. Lett. 93, 220401 (2004); Phys. Rev. A 72, 042113 (2005).
- [17] R. Blume-Kohout and W. H. Zurek, Phys. Rev. A 73, 062310 (2006).
- [18] H. Ollivier and W. H. Zurek, Phys. Rev. Lett. 88, 017901 (2001).
- [19] J.-P. Paz and W. H. Zurek, in *Coherent Atomic Matter Waves*, *Les Houches Lectures*, edited by R. Kaiser, C. Westbrook, and F. David (Springer, Berlin, 2001).
- [20] H. Everett, III, Rev. Mod. Phys. 29, 454 (1957).
- M. Gell-Mann and J. B. Hartle, Phys. Rev. D 47, 3345 (1993);
   R. B. Griffiths, *Consistent Quantum Theory* (Cambridge University Press, Cambridge, England, 2002).
- [22] W. H. Zurek, e-print arXiv:0707.2832.
- [23] W. H. Zurek, Phys. Rev. Lett. 90, 120404 (2003); Phys. Rev. A 71, 052105 (2005).
- [24] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, Cambridge, England, 1999).