

Preservation of positivity by dynamical coarse graining

Gernot Schaller* and Tobias Brandes

Institut für Theoretische Physik, Technische Universität Berlin, Hardenbergstrasse 36, 10623 Berlin, Germany

(Received 22 April 2008; revised manuscript received 17 June 2008; published 13 August 2008)

We compare different quantum master equations for the time evolution of the reduced density matrix. The widely applied secular approximation (rotating wave approximation) applied in combination with the Born-Markov approximation generates a Lindblad-type master equation ensuring for completely positive and stable evolution and is typically well applicable for optical baths. For phonon baths however, the secular approximation is expected to be invalid. The usual Markovian master equation does not generally preserve positivity of the density matrix. As a solution we propose a coarse-graining approach with a dynamically adapted coarse-graining time scale. For some simple examples we demonstrate that this preserves the accuracy of the integro-differential Born equation. For large times we analytically show that the secular approximation master equation is recovered. The method can in principle be extended to systems with a dynamically changing system Hamiltonian, which is of special interest for adiabatic quantum computation. We give some numerical examples for the spin-boson model of cases where a spin system thermalizes rapidly, and other examples where thermalization is not reached.

DOI: [10.1103/PhysRevA.78.022106](https://doi.org/10.1103/PhysRevA.78.022106)

PACS number(s): 03.65.Yz, 03.67.Ac, 87.19.1o

I. INTRODUCTION

The discovery that quantum computers would have much stronger capabilities for solving certain kinds of problems (such as number factoring [1] or database search [2]) than their classical counterparts has initiated a lot of research in quantum-information theory [3].

Unfortunately, the fragile quantum coherence necessary for the superior performance of quantum computers is very sensitive to the inevitable interaction with the environment such that theoretical understanding of this process—called decoherence—is absolutely necessary [4].

For simple models (such as, e.g., a single spin or harmonic oscillator coupled to a thermalized bath of harmonic oscillators) and for sufficiently complex couplings to the reservoir after some time the system will equilibrate in a thermal state with the bath temperature [4]. This behavior would be consistent with our classical expectations.

A recent idea is to protect the quantum information by encoding the solution to a given problem in the (unknown) ground state of a problem Hamiltonian H_p . Given sufficient experimental control of the system Hamiltonian and a reservoir at sufficiently low temperature $k_B T \ll \Delta E_{\min}$ (where ΔE_{\min} denotes the energy gap above the ground-state energy of H_p), the ground state should be robust against decoherence in the sense that decoherence would always drive the system toward its ground state. In principle, one could then prepare the quantum system in any accessible state and wait sufficiently long until the equilibration has taken place. Unfortunately, this mere-cooling approach is not expected to be very efficient, since the relaxation rate may be very small [5] or (in extreme cases) the system might get stuck in a local minimum [6].

A possible solution was proposed with the concept of adiabatic quantum computation [7]: The system is initially

subject to a simple Hamiltonian H_I and is prepared in its (easily accessible) ground state. Then, the Hamiltonian is slowly deformed into the problem Hamiltonian H_p . The adiabatic theorem states that if this transformation proceeds slowly enough, the quantum state will closely follow the instantaneous ground state [8]. Finally, for a nearly adiabatic evolution the system state approximates the system ground state to a high degree. Consequently, the maximum transformation rate (where the final excitations are acceptably small) corresponds to the computational complexity of the adiabatic algorithm. For closed systems, it is related to the spectral properties of the time-dependent system Hamiltonian [9,10]. For a reservoir at sufficiently low temperatures, this scheme is thought to be robust against decoherence [6] and might even be aided by it [11].

Unfortunately, the standard framework of deriving master equations relies on some prerequisites that are not always fulfilled in realistic systems. For example, the Markovian approximation widely used is usually only formulated for time-independent system Hamiltonians. In addition, it sometimes leads to master equations that do not preserve positivity of the system density matrix [12,13]. Together with trace preservation, positivity grants stability of the density matrix eigenvalues and is thus necessary for probability interpretation (cf. [14]) of the density matrix. Consequently, observables obtained from nonpositive density matrices may become unphysical [15]. This problem can be cured by the secular approximation. Combined with the Markovian approximation, it leads to Lindblad-type [16] master equations that generically preserve positivity of the density matrix. Unfortunately, the secular approximation is rather valid for quantum-optical but not for phonon baths [4]. Note that there exist non-Markovian master equations that are not of Lindblad type but nevertheless preserve positivity by construction. These models however are either phenomenologic [17,18] in the sense that their parameters are not derived from a microscopic Hamiltonian or they only grant positivity on a restricted set of initial states [19]. Especially in view of an experimental optimization of decoherence effects it is,

*schaller@itp.physik.tu-berlin.de

however, necessary to relate the parameters in the master equation to those in the microscopic Hamiltonian [20]. For example, for a realistic implementation of an adiabatic (or gate-model) quantum computer one would expect the qubits to be coupled to phonon degrees of freedom as well, such that a general treatment is advised. The present paper shall present a further step in that direction.

It has been noted [21,23,24] that coarse graining may ensure for positive evolution of the reduced density matrix. However, the coarse-graining time scale so far must be much larger than the inverse of the bath density of states cutoff. In this paper we argue that by adaptively changing the coarse-graining time scale one does not have to obey this constraint. Beyond this, we show that for infinitely large coarse-graining times we reproduce the widely used secular approximation. We show analytically that for any fixed coarse-graining time scale $\tau \geq 0$, the resulting master equations are of Lindblad form, i.e., they preserve positivity and thereby also stability of the density matrix.

This paper is organized as follows: In Sec. II we introduce our notation and in Sec. III we compare the standard procedure of deriving quantum master equations from microscopic models with the proposed adaptive coarse-graining scheme. We make our method explicit by the example of the spin-boson model in Sec. IV.

II. GENERAL PREREQUISITES

We will consider Hamiltonians which can be divided into three parts

$$H(t) = H_S(t) + H_{SB} + H_B, \quad (1)$$

where $H_S(t)$ describes the system part, H_B the part acting on the bath (with $[H_S(t), H_B] = 0$) and

$$H_{SB} = \lambda \sum_A A_A \otimes B_A \quad (2)$$

denotes the interaction Hamiltonian with the small dimensionless coupling parameter $\lambda \ll 1$ and system operators A_A as well as bath operators B_A (differing coupling constants can be absorbed in the operator definitions). Hermiticity is only required for the complete sum ($H_{SB} = H_{SB}^\dagger$), but by splitting operators into Hermitian and anti-Hermitian parts one can always redefine them such that

$$A_A = A_A^\dagger, \quad B_A = B_A^\dagger, \quad (3)$$

which will be assumed further on.

The density matrix of the complete system is thought to evolve according to ($\hbar=1$ throughout) the von Neumann equation of motion

$$\dot{\rho}(t) = -i[H_S(t) + H_{SB} + H_B, \rho(t)]. \quad (4)$$

Denoting the time evolution operators of system and reservoir by $U(t)$ and $V(t)$, respectively, we can switch to the interaction picture

$$\rho(t) = U^\dagger(t)V^\dagger(t)\rho(t)V(t)U(t),$$

$$\begin{aligned} \mathbf{H}_{SB}(t) &= U^\dagger(t)V^\dagger(t)H_{SB}V(t)U(t) \\ &= \lambda \sum_A [U^\dagger(t)A_A U(t)] \otimes [V^\dagger(t)B_A V(t)] \\ &\equiv \lambda \sum_A A_A(t) \otimes B_A(t). \end{aligned} \quad (5)$$

We will denote all operators in the interaction picture by bold symbols throughout. In the interaction picture, the equation of motion for the density operator (4) transforms into

$$\dot{\rho}(t) = -i[\mathbf{H}_{SB}(t), \rho(t)], \quad (6)$$

where one can exploit the smallness of the coupling λ to apply perturbation theory.

Equation (6) can be formally integrated to yield $\rho(t) = \rho(0) - i \int_0^t [\mathbf{H}_{SB}(t'), \rho(t')] dt'$ and reinserting this result in Eq. (6) one obtains the following exact equation:

$$\dot{\rho}(t) = -i[\mathbf{H}_{SB}(t), \rho(0)] - \int_0^t [\mathbf{H}_{SB}(t), [\mathbf{H}_{SB}(t'), \rho(t')]] dt' \quad (7)$$

for the density operator.

III. QUANTUM MASTER EQUATIONS

We will first state the results of the Born-Markov approximation without secular approximation (in Sec. III A) and with the secular approximation (Sec. III B) in our notation. Afterwards, we will consider the coarse-graining approach in Sec. III C.

With the usual assumptions (compare Appendix A) involving initial factorization of the density matrix and neglecting any change in the reservoir part of the density matrix in Eq. (7) we obtain the Born equation

$$\begin{aligned} \dot{\rho}_S &= -i \text{Tr}_B \{ [\mathbf{H}_{SB}(t), \rho_S(0) \rho_B^0] \} \\ &\quad - \int_0^t \text{Tr}_B \{ [\mathbf{H}_{SB}(t), [\mathbf{H}_{SB}(t'), \rho_S(t') \rho_B^0]] \} dt' + O\{\lambda^3\}, \end{aligned} \quad (8)$$

where $\text{Tr}_B\{\cdot\}$ denotes the trace over the reservoir degrees of freedom. Evaluating the traces leads to the definition of the reservoir correlation functions

$$C_{AB}(\tau) \equiv \text{Tr}_B \{ e^{+iH_B \tau} B_A e^{-iH_B \tau} B_B \rho_B^0 \} = C_{BA}^*(-\tau), \quad (9)$$

and we obtain with $\langle B_A \rangle = 0$ (which can always be achieved by a suitable transformation [22])

$$\begin{aligned} \dot{\rho}_S &= \lambda^2 \sum_{AB} \int_0^t \{ [A_B(t') \rho_S(t'), A_A(t)] C_{AB}(t-t') + \text{H.c.} \} dt' \\ &\quad + O\{\lambda^3\}, \end{aligned} \quad (10)$$

where H.c. denotes the Hermitian conjugate. The integro-differential character of above equation complicates its solution, since analytical solutions are only possible in very

simple cases [25,26] (see also Sec. IV B), and numerical solutions are hampered by the fact that the complete history of $\rho_S(t')$ must be stored in order to evolve $\rho_S(t)$.

A. Markovian approximation scheme

In the usual [4] Markovian approximation (see Appendix B) one obtains for constant system Hamiltonians [$H_S(t)=H_S$] with the half-sided Fourier transforms

$$\Gamma_{AB}(\omega) \equiv \int_0^\infty C_{AB}(\tau) e^{i\omega\tau} d\tau \quad (11)$$

of the reservoir correlation functions (9) the time-local Born-Markov (BM) master equation (here given in the Schrödinger picture)

$$\begin{aligned} \dot{\rho}_S = & -i[H_S, \rho_S(t)] + \lambda^2 \sum_{abcd} \sum_{AB} \{ \Gamma_{AB}(E_b - E_a) \langle a|A_B|b \rangle \\ & \times \langle c|A_A|d \rangle^* [(|a\rangle\langle b|)\rho_S(t), (|c\rangle\langle d|)^\dagger] + \text{H.c.} \}, \end{aligned} \quad (12)$$

where $H_S|a\rangle = E_a|a\rangle$ denote the orthonormal energy eigenbasis. By construction, Eq. (12) does preserve trace and Hermiticity of ρ_S . Note however, that positivity of its solution $\rho_S(t)$ is not generally preserved [12,27], see Sec. IV D for some counterexamples.

B. Secular approximation

In order to restore preservation of positivity in the BM approximation in general, it is necessary to perform the secular approximation (see Appendix C). Typically, this approximation is known to be well satisfied for quantum-optical systems, where it is also known as rotating wave approximation [28]. In this case, one can combine

$$\begin{aligned} \gamma_{AB}(\omega) &= \Gamma_{AB}(\omega) + \Gamma_{BA}^*(\omega) = \int_{-\infty}^{+\infty} C_{AB}(\tau) e^{i\omega\tau} d\tau, \\ \sigma_{AB}(\omega) &= \Gamma_{AB}(\omega) - \Gamma_{BA}^*(\omega) = \int_{-\infty}^{+\infty} C_{AB}(\tau) \text{sgn}(\tau) e^{i\omega\tau} d\tau, \end{aligned} \quad (13)$$

which yields the Born-Markov-Secular (BMS) approximation (in the Schrödinger picture)

$$\begin{aligned} \dot{\rho}_S = & -i[H_S, \rho_S(t)] - i \left[\sum_{ab} \tilde{\sigma}_{ab} |a\rangle\langle b|, \rho_S(t) \right] \\ & + \sum_{abcd} \tilde{\gamma}_{ab,cd} \left((|a\rangle\langle b|)\rho_S(t)(|c\rangle\langle d|)^\dagger \right. \\ & \left. - \frac{1}{2} \{ (|c\rangle\langle d|)^\dagger (|a\rangle\langle b|), \rho_S(t) \} \right), \end{aligned}$$

$$\tilde{\sigma}_{ab} = \frac{\lambda^2}{2i} \sum_c \sum_{AB} \sigma_{AB}(E_a - E_c) \delta_{E_b, E_a} \langle c|A_A|a\rangle^* \langle c|A_B|b\rangle,$$

$$\tilde{\gamma}_{ab,cd} = \lambda^2 \sum_{AB} \gamma_{AB}(E_b - E_a) \delta_{E_a - E_c, E_b - E_a} \langle a|A_B|b\rangle \langle c|A_A|d\rangle^*. \quad (14)$$

Equation (14) has many favorable properties (compare, e.g., Chap. 3.3 in [4]): By construction, it preserves trace and Hermiticity of the system density matrix ρ_S . Since it is of Lindblad [16] form [the matrix $\gamma_{AB}(\omega)$ is positive semidefinite], it preserves positivity of the density matrix.

For a thermalized reservoir characterized by the inverse temperature β , the Fourier transforms of the bath correlation functions can be used to show that the system thermal equilibrium state with the same temperature

$$\rho_S^{\text{th}} = \frac{e^{-\beta H_S}}{\text{Tr}_S\{e^{-\beta H_S}\}} \quad (15)$$

is a stationary state. If the spectrum of the system Hamiltonian H_S is nondegenerate (implying that $\delta_{E_a, E_b} = \delta_{ab}$), the equations for the diagonal elements of ρ_S in the eigenbasis of H_S completely decouple from the equations for the off diagonals, and one obtains the same transition rates between the populations as with the Fermi golden rule.

C. Coarse-graining approach

Equation (6) is formally solved by $\rho(t_2) = \mathbf{W}(t_2, t_1) \rho(t_1) \mathbf{W}^\dagger(t_2, t_1)$ with the interaction picture time evolution operator

$$\mathbf{W}(t_2, t_1) = \mathcal{T} \exp \left(-i \int_{t_1}^{t_2} \mathbf{H}_{\text{SB}}(t') dt' \right), \quad (16)$$

where the time dependence of $\mathbf{H}_{\text{SB}}(t)$ necessitates the time ordering [28] $\mathcal{T} \mathbf{H}_{\text{SB}}(t_1) \mathbf{H}_{\text{SB}}(t_2) \equiv \mathbf{H}_{\text{SB}}(t_1) \mathbf{H}_{\text{SB}}(t_2) \Theta(t_1 - t_2) + \mathbf{H}_{\text{SB}}(t_2) \mathbf{H}_{\text{SB}}(t_1) \Theta(t_2 - t_1)$, with $\Theta(x)$ denoting the Heavyside step function. Now, by expanding $\mathbf{W}(t+\tau, t)$ up to second order in λ we obtain a second-order approximation to the full density matrix

$$\begin{aligned} \rho(t+\tau) = & \rho(t) - i \left[\int_t^{t+\tau} \mathbf{H}_{\text{SB}}(t_1) dt_1, \rho(t) \right] \\ & - i \left[\frac{1}{2i} \int_t^{t+\tau} \int_t^{t+\tau} [\mathbf{H}_{\text{SB}}(t_2), \mathbf{H}_{\text{SB}}(t_1)] \right. \\ & \left. \times \Theta(t_2 - t_1) dt_1 dt_2, \rho(t) \right] \\ & + \int_t^{t+\tau} \int_t^{t+\tau} \left[\mathbf{H}_{\text{SB}}(t_1) \rho(t) \mathbf{H}_{\text{SB}}(t_2) \right. \\ & \left. - \frac{1}{2} \{ \mathbf{H}_{\text{SB}}(t_2) \mathbf{H}_{\text{SB}}(t_1), \rho(t) \} \right] dt_1 dt_2 + \mathcal{O}\{\lambda^3\}. \end{aligned} \quad (17)$$

Since such a truncated finite-order approximation to the time evolution operator $\mathbf{W}(t+\tau)$ is still unitary, the above map (17) preserves Hermiticity, trace, and positivity of the full density matrix ρ . An equivalent expression can be obtained

from iterative solution of Eq. (7) by keeping only terms up to $O\{\lambda^2\}$. We will proceed with this second-order approximation and derive from this fully unitary map a nonunitary map for the system part of the density matrix that preserves positivity.

If we neglect the back-action of the system on the bath and assume factorization (Born approximation) as described in Appendix A we can perform the partial trace over the reservoir degrees of freedom. By inserting the definition (2) in Eq. (17) and employing the definition of the reservoir correlation functions (9) we obtain (again working in a frame where $\langle B_A \rangle = 0$ [22])

$$\begin{aligned} \rho_S(t + \tau) &= \rho_S(t) - i \frac{\lambda^2}{2i} \sum_{AB} \int_t^{t+\tau} \int_t^{t+\tau} C_{AB}(t_2 - t_1) \text{sgn}(t_2 - t_1) \\ &\quad \times [A_A(t_2) A_B(t_1), \rho_S(t)] dt_1 dt_2 \\ &\quad + \lambda^2 \sum_{AB} \int_t^{t+\tau} \int_t^{t+\tau} C_{AB}(t_2 - t_1) \left(A_B(t_1) \rho_S(t) A_A(t_2) \right. \\ &\quad \left. - \frac{1}{2} \{A_A(t_2) A_B(t_1), \rho_S(t)\} \right) dt_1 dt_2 + O\{\lambda^3\} \\ &\equiv \rho_S(t) + \tau L^\tau(t) \rho_S(t) + O\{\lambda^3\}, \end{aligned} \quad (18)$$

which defines the action of the Liouvillian on the reduced density matrix in the interaction picture. If one rearranges the matrix elements of the density matrix as a N^2 -dimensional vector, the Liouvillian superoperator can be understood as an $N^2 \times N^2$ (generally non-Hermitian) matrix acting on ρ_S . In the interaction picture the Liouvillian is small, i.e., since $\langle B_A \rangle = 0$ we even have $L^\tau = O\{\lambda^2\}$.

Unfortunately, the above map $\rho_S(t) \rightarrow \rho_S(t + \tau)$ has some shortcomings: It does not generally preserve positivity of the reduced density matrix ρ_S and in addition, if one applies the above equation recursively n times with small time steps $\Delta\tau$ such that $n\Delta\tau = \tau$ and compares the result with a single iteration of Eq. (18), the difference between the two solutions is much larger than $O\{\lambda^3\}$, i.e., the solution depends on the choice of the step size.

By defining time averaging of an operator $\hat{O}(t)$ over a time interval $[t, t + \tau]$ as

$$\langle\langle \hat{O} \rangle\rangle_{[t, t+\tau]} \equiv \frac{1}{\tau} \int_t^{t+\tau} \hat{O}(t') dt', \quad (19)$$

we can write Eq. (18) as

$$\langle\langle \dot{\rho}_S \rangle\rangle_{[t, t+\tau]} = \frac{\rho_S(t + \tau) - \rho_S(t)}{\tau} = L^\tau(t) \rho_S(t) + O\{\lambda^3\}. \quad (20)$$

1. Explicit Liouvillian

It is convenient to insert the even and odd Fourier transforms from Eq. (13) of the reservoir correlation functions

$$\begin{aligned} C_{AB}(\tau) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \gamma_{AB}(\omega) e^{-i\omega\tau} d\omega, \\ C_{AB}(\tau) \text{sgn}(\tau) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \sigma_{AB}(\omega) e^{-i\omega\tau} d\omega, \end{aligned} \quad (21)$$

and to expand the system operators in the interaction picture in the orthonormal energy eigenbasis $H_S|a\rangle = E_a|a\rangle$ of the system Hamiltonian (here assumed to be constant)

$$\begin{aligned} A_A(t_2) &= \sum_{ab} \langle a|A_A|b\rangle e^{i(E_a - E_b)t_2} |a\rangle\langle b|, \\ A_B(t_1) &= \sum_{cd} \langle c|A_B|d\rangle e^{i(E_c - E_d)t_1} |c\rangle\langle d|. \end{aligned} \quad (22)$$

Then, we can use the relation

$$\int_t^{t+\tau} e^{i\alpha t'} dt' = \tau e^{i\alpha t} e^{i\alpha\tau/2} \text{sinc}\left(\frac{\alpha\tau}{2}\right)$$

with $\text{sinc}(x) \equiv \frac{\sin(x)}{x}$ together with Eqs. (21) and (22) in order to make the Liouvillian in Eq. (20) explicit. With denoting the energy differences of the system Hamiltonian by $\Omega_{ab} \equiv E_a - E_b$ we arrive at

$$\begin{aligned} L^\tau(t) \rho_S(t) &= -i \left[\sum_{ab} e^{i\Omega_{ab}t} \sigma_{ab}(\tau) |a\rangle\langle b|, \rho_S(t) \right] + \sum_{abcd} e^{i(\Omega_{cd} - \Omega_{ab})t} \gamma_{cd,ba}(\tau) \left((|c\rangle\langle d|) \rho_S(t) (|b\rangle\langle a|)^\dagger - \frac{1}{2} \{ (|b\rangle\langle a|)^\dagger (|c\rangle\langle d|), \rho_S(t) \} \right), \\ \sigma_{ab}(\tau) &= \frac{\lambda^2 \tau}{4\pi i} e^{i\Omega_{ab}\tau/2} \sum_c \sum_{AB} \langle c|A_A|a\rangle^* \langle c|A_B|b\rangle \int_{-\infty}^{+\infty} \sigma_{AB}(\omega) \text{sinc}\left((w + \Omega_{ca})\frac{\tau}{2}\right) \text{sinc}\left((w + \Omega_{cb})\frac{\tau}{2}\right) d\omega, \\ \gamma_{cd,ba}(\tau) &= \frac{\lambda^2 \tau}{2\pi} e^{i(\Omega_{cd} - \Omega_{ba})\tau/2} \sum_{AB} \langle b|A_A|a\rangle^* \langle c|A_B|d\rangle \int_{-\infty}^{+\infty} \gamma_{AB}(\omega) \text{sinc}\left((w + \Omega_{ba})\frac{\tau}{2}\right) \text{sinc}\left((w + \Omega_{cd})\frac{\tau}{2}\right) d\omega. \end{aligned} \quad (23)$$

The coarse-grained derivative on the left-hand side of Eq. (20) generates a time evolution that can be compared with its usual, first-order differential equation counterpart (we denote the corresponding density operators by an overbar and the continuous index τ referring to the Liouvillian chosen)

$$\frac{d}{dt}\bar{\rho}_S^\tau(t) = L^\tau(t)\bar{\rho}_S^\tau(t). \quad (24)$$

If we initialize this differential equation with a known density matrix ρ_S^0 and evaluate its formal solution at time $t = \tau$ (i.e., after exactly the coarse-graining time scale), we find to first order in L^τ ,

$$\bar{\rho}_S^\tau(\tau) \approx \left(1 + \int_0^\tau L^\tau(t')dt'\right)\rho_S^0. \quad (25)$$

This means that when considering the same initial condition, the difference between the two time evolutions resulting from Eqs. (24) and (20) is given by

$$\begin{aligned} \bar{\rho}_S^\tau(\tau) - \rho_S(\tau) &= \eta(\tau) + O\{(L^\tau)^2\}, \\ \eta(\tau) &= \left(\int_0^\tau L^\tau(t')dt' - \tau L^\tau(0)\right)\rho_S^0, \end{aligned} \quad (26)$$

i.e., essentially by the difference between the coarse-graining-averaged Liouvillian and its initial value.

It follows from Eq. (23) that especially for short coarse-graining times $\tau \ll \frac{1}{|\Delta E_{\max}|}$ (with ΔE_{\max} denoting the maximum energy difference of H_S) the difference will be negligible $\eta(\tau) \approx 0$, such that the two solutions $\bar{\rho}_S^\tau(\tau)$ and $\rho_S(\tau)$ are equivalent. Since we have so far not made any assumption on separating time scales, this also implies that non-Markovian effects (that are expected for short times where the Markovian approximation does not hold) are within reach of the adaptive coarse-graining approach, where the coarse-graining time is chosen to match the physical time.

In the limit of very large coarse-graining times we also obtain $\lim_{\tau \rightarrow \infty} \eta(\tau) \approx 0$, since then the $\text{sinc}(\cdots)$ functions in Eq. (23) begin to act like δ functions, see below. Also for intermediate coarse-graining times it is easy to see from the structure of the Liouvillian (23) that the difference $\eta(\tau)$ is bounded throughout.

From now on, we will omit the overbar and write $\rho_S^\tau(t)$ for the solutions of Eq. (24).

2. Infinite- τ limit

In the limit $\tau \rightarrow \infty$ one should note that for discrete a, b and under an integral over $d\omega$ with another integrable function one has in a distributive sense (see Appendix F)

$$\begin{aligned} f(\omega, a, b) &\equiv \lim_{\tau \rightarrow \infty} \tau \text{sinc}\left((w+a)\frac{\tau}{2}\right) \text{sinc}\left((w+b)\frac{\tau}{2}\right) \\ &\asymp 2\pi \delta_{ab} \delta(w+a), \end{aligned} \quad (27)$$

where δ_{ab} is a Kronecker symbol and $\delta(w+a)$ denotes the Dirac δ distribution. Therefore, we obtain for the Liouvillian matrix elements (23) in this limit

$$\sigma_{ab}^\infty = \frac{\lambda^2}{2i} \delta_{E_a, E_b} \sum_c \sum_{AB} \langle c|A_A|a\rangle^* \langle c|A_B|b\rangle \sigma_{AB}(E_a - E_c),$$

$$\gamma_{cd,ba}^\infty = \lambda^2 \delta_{E_b - E_a, E_c - E_d} \sum_{AB} \langle b|A_A|a\rangle^* \langle c|A_B|d\rangle \gamma_{AB}(E_a - E_b), \quad (28)$$

and we recover the secular approximation (14). Naturally, this also implies that in the large-time limit, the solution $\rho_S^{\tau=t}(t)$ captures all the favorable properties of Eq. (14).

3. Positivity

The most striking advantage of the coarse-graining procedure is that not only for very large, but for any fixed coarse-graining time $\tau \geq 0$, the resulting first-order differential equations (24) are all of Lindblad form [16] and thus intrinsically preserve the positivity of the density matrix. This is easily seen by switching back to the Schrödinger picture, where the time-dependent phases in Eq. (23) cancel

$$\begin{aligned} \dot{\rho}_S^\tau(t) \equiv L^\tau \rho_S^\tau(t) &= -i[H_S, \rho_S^\tau(t)] - i \left[\sum_{ab} \sigma_{ab}(\tau) |a\rangle\langle b|, \rho_S^\tau(t) \right] \\ &+ \sum_{abcd} \gamma_{cd,ba}(\tau) \left((|c\rangle\langle d|) \rho_S^\tau(t) (|b\rangle\langle a|)^\dagger \right. \\ &\left. - \frac{1}{2} \{ (|b\rangle\langle a|)^\dagger (|c\rangle\langle d|), \rho_S^\tau(t) \} \right), \end{aligned} \quad (29)$$

which implies that the Schrödinger picture Liouvillian L^τ only depends on the coarse-graining time scale τ .

First of all, the first two commutator terms correspond to commutators with a Hermitian operator, i.e., the second term accounts for the unitary action of decoherence (sometimes called Lamb shift [4]). Note that Hermiticity of the corresponding effective Hamiltonian follows from the definition of the odd Fourier transform versus the half-sided Fourier transform (13).

In order to show that Eq. (29) is a Lindblad form, it remains to be shown that the matrix $\gamma_{cd,ba}(\tau)$ is positive semidefinite. In order to see this we introduce the double indices $i=(cd)$ and $j=(ba)$ running from 1 to N^2 for an N -dimensional system Hilbert space. Then, we can use the shorthand notation $w_i = E_c - E_d$, $A_B^i = \langle c|A_B|d\rangle$, $w_j = E_b - E_a$, $A_A^j = \langle b|A_A|a\rangle$ and consider for arbitrary complex-valued numbers x_j in Eq. (23),

$$\sum_{ij} x_i^* \gamma_{ij} x_j = \frac{\tau}{2\pi} \int_{-\infty}^{+\infty} \sum_{AB} z_A^*(\omega) \gamma_{AB}(\omega) z_B(\omega) d\omega, \quad (30)$$

where we have

$$z_A(\omega) \equiv \sum_j A_A^j \text{sinc}[(w + w_j)\tau/2] e^{i\omega_j \tau/2} x_j. \quad (31)$$

Now, since the Fourier-transform matrix $\gamma_{AB}(\omega)$ is positive semidefinite (compare also Appendix C) one has $z_A^*(\omega) \gamma_{AB}(\omega) z_B(\omega) \geq 0$. The integral over a strictly positive function can only yield positive results, and since also $\tau \geq 0$ it follows that $\gamma_{ij}(\tau) = \gamma_{cd,ba}(\tau)$ is a positive semidefinite matrix.

This also implies that solutions of the form $\rho_S^t(t)$ are always positive density matrices, since they correspond to an interpolation along Lindblad density matrices.

D. Time-dependent generalization

Within the coarse-graining approach, it was in Eq. (22) where it was used for the first time that the system Hamiltonian H_S was time independent and had a discrete spectrum. Here we will show that coarse graining generally leads to a time-inhomogeneous (i.e., one with time-dependent operators) Lindblad form of master equations and thus always preserves positivity of the density matrix. With introducing the notation

$$\tilde{A}_A(t, \omega) \equiv A_A(t) e^{i\omega t} \quad (32)$$

we can use Eqs. (21) and (19) in Eq. (18) to obtain

$$\begin{aligned} \langle\langle \dot{\rho}_S \rangle\rangle_{[t, t+\tau]} = & -i \left[\frac{\lambda^2 \tau}{4\pi i} \int_{-\infty}^{+\infty} \sum_{AB} \sigma_{AB}(\omega) \right. \\ & \times \langle\langle \tilde{A}_A(\omega) \rangle\rangle_{[t, t+\tau]}^\dagger \langle\langle \tilde{A}_B(\omega) \rangle\rangle_{[t, t+\tau]} d\omega, \rho_S(t) \left. \right] \\ & + \frac{\lambda^2 \tau}{2\pi} \int_{-\infty}^{+\infty} \sum_{AB} \gamma_{AB}(\omega) \\ & \times \left[\langle\langle \tilde{A}_B(\omega) \rangle\rangle_{[t, t+\tau]} \rho_S(t) \langle\langle \tilde{A}_A(\omega) \rangle\rangle_{[t, t+\tau]}^\dagger \right. \\ & \left. - \frac{1}{2} \{ \langle\langle \tilde{A}_A(\omega) \rangle\rangle_{[t, t+\tau]}^\dagger \langle\langle \tilde{A}_B(\omega) \rangle\rangle_{[t, t+\tau]}, \rho_S(t) \} \right]. \quad (33) \end{aligned}$$

With the replacements $\langle\langle \dot{\rho}_S \rangle\rangle_{[t, t+\tau]} \rightarrow \dot{\rho}_S^\tau$ and $\rho_S(t) \rightarrow \rho_S^\tau(t)$ this becomes a time-inhomogeneous Lindblad master equation, since the positivity of the $\gamma_{AB}(\omega)$ matrix at every ω is guaranteed for reservoirs in thermal equilibrium. Intuitively, the time dependence of the Lindblad operators does not destroy positivity, since at any fixed time t one may approximate the time-dependent operators by a constant-operator Lindblad form, see Appendix E for a more explicit discussion. Since the case of slowly varying system Hamiltonians is of special interest in the context of adiabatic quantum computation [6], we outline in Appendix D how one could in principle calculate the time-averaged operators

$$\begin{aligned} \langle\langle \tilde{A}_A(\omega) \rangle\rangle_{[t, t+\tau]} &= \frac{1}{\tau} \int_t^{t+\tau} A_A(t') e^{+i\omega t'} dt' \\ &= \frac{1}{\tau} \int_t^{t+\tau} U^\dagger(t') A_A U(t') e^{+i\omega t'} dt', \quad (34) \end{aligned}$$

in the adiabatic limit.

IV. SPIN-BOSON MODEL

We will make our method explicit at the example of the spin-boson model in the following. We will also give some

numerical solutions to the master equations used: The eigenvalues of the density matrices have been calculated with the LAPACK package [29]. The half-sided Fourier transforms (11) and odd Fourier transforms (13) were calculated numerically from the full Fourier transform by consecutive application of backward and forward integral transforms. This was implemented by an integration algorithm based on the discrete Fourier transform optimized for oscillating integrands [30]. For the integration of partial differential equations, a fourth-order Runge-Kutta method with an adaptive step size [30] was used. Trace and Hermiticity of the density matrix were always preserved within machine accuracy. Likewise, whenever a Lindblad-type master equation was integrated, non-negativity of the smallest eigenvalue of the density matrix was preserved within numerical accuracy defined by the accuracy of the Fourier transform.

A. Microscopic derivation

We consider a system Hamiltonian with discrete energy eigenvalues that can be described by a quadratic form of Pauli matrices for a system of n spins,

$$\begin{aligned} H_S = & \gamma \mathbf{1} + \sum_{i=1}^n (\gamma_i^x \sigma_i^x + \gamma_i^y \sigma_i^y + \gamma_i^z \sigma_i^z) \\ & + \sum_{i=1}^n \sum_{j=i+1}^n \sum_{\alpha, \beta=x, y, z} \gamma_{ij}^{\alpha\beta} \sigma_i^\alpha \sigma_j^\beta, \quad (35) \end{aligned}$$

where $\vec{\sigma}_i$ denotes the Pauli matrices acting on spin i . In the worst case, this Hamiltonian is defined by $1-3/2n+9/2n^2$ real parameters. Note that when considering explicit examples, we will not give the dimension of these parameters which implies that all times have their inverse dimensions. This system Hamiltonian is nontrivial in the sense that even in the case of time-independent parameters considered here, the time evolution of the operators in the interaction picture cannot be solved analytically without using exponential resources. The system is coupled to a bosonic bath

$$H_B = \sum_k \omega_k \left(b_k^\dagger b_k + \frac{1}{2} \right) \quad (36)$$

with the usual bosonic commutation relations. The coupling between system and bath is realized by the quite general interaction Hamiltonian

$$H_{SB} = \lambda \sum_{i=1}^n \sum_k (\vec{n}_{ik} \cdot \vec{\sigma}_i \otimes b_k + \vec{n}_{ik}^* \cdot \vec{\sigma}_i \otimes b_k^\dagger), \quad (37)$$

with $\lambda \ll 1$ where the frequency dependence is contained in the complex-valued coupling coefficients \vec{n}_{ik} .

1. Bath correlation functions

In order to obtain a rather simple form of the master equation, we will make the assumption of a collective coupling, where the frequency dependence of the coupling strengths factorizes with the different spin positions and spin directions, i.e.,

$$\vec{n}_{ik} = \vec{n}_i h_k \quad (38)$$

for some function h_k . This implies that the distance between the spins is smaller than the correlation length of the reservoir oscillators. This approximation is not crucial for the further procedure but simplifies the resulting system of equations considerably, since the coupling to the reservoir can be described by just two effective spin operators. In this case, the interaction Hamiltonian (37) can be written as

$$H_{\text{SB}} = \lambda \Sigma_R \otimes (B + B^\dagger) + \lambda \Sigma_I \otimes i(B - B^\dagger) \quad (39)$$

with the composed operators

$$\begin{aligned} \Sigma_R &= \sum_{i=1}^n \text{Re}(\vec{n}_i) \cdot \vec{\sigma}_i, & \Sigma_I &= \sum_{i=1}^n \text{Im}(\vec{n}_i) \cdot \vec{\sigma}_i, \\ B &= \sum_k h_k b_k, \end{aligned} \quad (40)$$

where we can identify from Eq. (2) the Hermitian operators

$$\begin{aligned} A_1 &= \Sigma_R, & A_2 &= \Sigma_I, \\ B_1 &= B + B^\dagger, & B_2 &= i(B - B^\dagger). \end{aligned} \quad (41)$$

From the bath Hamiltonian (36) we obtain $e^{iH_B \tau} b_k e^{-iH_B \tau} = e^{-i\omega_k \tau} b_k$ and the Hermitian conjugate, respectively. We will consider the limit of a large bath with a continuous spacing of oscillator frequencies. In this limit, the sums over k can be approximated by an integral

$$\sum_k |h_k|^2 f(\omega_k) \rightarrow \int_0^\infty g(\omega) f(\omega) d\omega, \quad (42)$$

where $g(\omega)$ is defined by the distribution of bath oscillators (spectral density) as well as the frequency-dependent coupling strengths h_k . With the bosonic expectation value for a thermalized reservoir $\langle N_k \rangle = \text{Tr}_B \{ b_k^\dagger b_k \rho_B^0 \} = [\exp\{\beta \omega_k\} - 1]^{-1}$ at temperature $\beta = (k_B T)^{-1}$ this can be used to determine the bath correlation functions as

$$\begin{aligned} C_{11}(\tau) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{2\pi g(|\omega|)}{|1 - e^{-\beta\omega}|} e^{-i\omega\tau} d\omega, \\ C_{12}(\tau) &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} \frac{2\pi(-i)g(|\omega|)\text{sgn}(\omega)}{|1 - e^{-\beta\omega}|} e^{-i\omega\tau} d\omega, \\ C_{21}(\tau) &= -C_{12}(\tau), & C_{22}(\tau) &= C_{11}(\tau), \end{aligned} \quad (43)$$

where we can directly read off the Fourier transform matrix

$$\gamma(\omega) = \frac{2\pi g(|\omega|)}{|1 - e^{-\beta\omega}|} \begin{pmatrix} 1 & -i \text{sgn}(\omega) \\ +i \text{sgn}(\omega) & 1 \end{pmatrix}. \quad (44)$$

Note that it is positive semidefinite (compare Appendix C) at every ω ensuring positive evolution of the Lindblad form master equation (14).

In some examples given, we will phenomenologically parametrize [31,32] the density of states as

$$g(\omega) = \left(\frac{\omega}{\omega_{\text{ph}}} \right)^s e^{-\omega/\omega_{\text{ct}}}, \quad (45)$$

where the exponent s determines the behavior near small frequencies, ω_{ph} is some physical frequency of the bath and ω_{ct} is a cutoff frequency necessary for normalization.

2. Exact solution for pure dephasing

The limit of pure dephasing of a single qubit is defined by considering $n=1$ with

$$\gamma = \frac{1}{2}, \quad \gamma^z = -\frac{1}{2}, \quad (46)$$

and all other coefficients to vanish in Eq. (35) as well as the simple coupling $\vec{n} = (0, 0, 1)$ in Eq. (37)—in this case also the collective coupling assumption (38) becomes exact. Then the time evolution operator in the interaction picture (16) can be calculated exactly [4,24,33] to all orders and one obtains that in the eigenbasis of the system Hamiltonian the diagonal elements of the density matrix remain unchanged and the off-diagonal elements simply decay as [cf. Eq. (82) in [24] in the limit of a continuous bath spectrum]

$$\rho_{01}(t) = e^{-\Gamma(t)} \rho_{01}(0),$$

$$\Gamma(t) = 8\lambda^2 \int_0^\infty g(\omega) \frac{\sin^2(\omega t/2)}{\omega^2} \coth\left(\frac{\beta\omega}{2}\right) d\omega, \quad (47)$$

i.e., in the pure dephasing limit one does not obtain thermalization. Likewise, since $e^{+iH_S t} A e^{-iH_S t} = A$ it is evident that BM and BMS approximations are equivalent for this case (compare also Appendix B and Appendix C).

B. Non-Markovian solutions

For simple system and interaction Hamiltonians

$$H_S = \frac{1}{2}[\mathbf{1} - \sigma^z], \quad H_{\text{SB}} = \lambda \sigma^a \otimes B_1, \quad (48)$$

where $a \in \{x, y, z\}$ and $B_1 = B_1^\dagger$ is a bath operator, the non-Markovian Born equation (10) is analytically solvable [15,25,26] in the special case of an exponentially decaying correlation function

$$C_{11}(\tau) \equiv \text{Tr}_B \{ B_1(\tau) B_1 \rho_B^0 \} = \frac{1}{2\tau_b} e^{-|\tau|/\tau_b}. \quad (49)$$

By considering large τ_b one can thus model reservoirs with a long-term memory, and the BM limit (where the BM approximation becomes exact) is obtained by considering $\lim_{\tau_b \rightarrow 0} C_{11}(\tau) = \delta(\tau)$. We obtain for the even and odd Fourier transforms in Eq. (13),

$$\gamma_{11}(\omega) = \frac{1}{1 + (\omega\tau_b)^2}, \quad \sigma_{11}(\omega) = \frac{i\omega\tau_b}{1 + (\omega\tau_b)^2}, \quad (50)$$

where it becomes visible (for later comparison with the spin-boson model) from Eq. (43) that this case can in principle be reproduced by a bosonic reservoir in the large temperature

limit with a Drude-like slowly decaying (but temperature-dependent) spectral coupling density

$$g(\omega) = \frac{1}{2\pi} \frac{\beta\omega}{1 + (\omega\tau_b)^2}. \quad (51)$$

Note however that by assuming a sum of many exponentials and allowing for phases, the method can in principle be generalized also to the low-temperature limit [25,34]. Inserting the operator definitions in the Born equation (10) we obtain

$$\eta_a(t) \equiv \int_0^t e^{+i/2(t-t')\sigma^z} \sigma^a \rho_S(t') e^{-i/2(t-t')\sigma^z} \frac{e^{-(t-t')/\tau_b}}{2\tau_b} dt',$$

$$\dot{\rho}_S = \frac{i}{2}[\sigma^z, \rho_S(t)] + \lambda^2[\eta_a(t) - \eta_a^\dagger(t), \sigma^a], \quad (52)$$

where it becomes evident that by taking the time derivative of the operator η^a one simply obtains a coupled set of first-order differential equations

$$\rho_S = \frac{i}{2}[\sigma^z, \rho_S(t)] + \lambda^2[\bar{\eta}_a(t), \sigma^a],$$

$$\dot{\bar{\eta}}_a = \frac{i}{2}[\sigma^z, \bar{\eta}_a(t)] - \frac{1}{\tau_b} \bar{\eta}_a(t) + \frac{1}{2\tau_b}[\sigma^a, \rho_S(t)]. \quad (53)$$

for the operators $\bar{\eta}_a(t) = \eta_a(t) - \eta_a^\dagger(t)$ and $\rho_S(t)$. Evidently, trace and Hermiticity of ρ_S and anti-Hermiticity of $\bar{\eta}_a$ are always preserved. Due to the initial condition $\bar{\eta}_a(0) = \mathbf{0}$, the trace of $\bar{\eta}_a$ will always vanish. Therefore, it suffices to parametrize ρ_S and $\bar{\eta}$ by just six real variables,

$$\rho_S \equiv \begin{pmatrix} \rho_{00} & \rho_x + i\rho_y \\ \rho_x - i\rho_y & 1 - \rho_{00} \end{pmatrix},$$

$$\bar{\eta} \equiv \begin{pmatrix} i\eta_{00} & \eta_x + i\eta_y \\ -\eta_x + i\eta_y & -i\eta_{00} \end{pmatrix}. \quad (54)$$

In the limit $\tau_b \rightarrow 0$ we simply obtain

$$\dot{\rho}_S = \frac{i}{2}[\sigma^z, \rho_S(t)] + \lambda^2[\sigma^a \rho_S(t) \sigma^a - \rho_S(t)] \quad (55)$$

for the system density matrix. In the BM approximation (with finite τ_b) we obtain, along the lines of Appendix B for $a=z$ (pure dephasing), $\eta_z^{\text{BM}}(t) = \frac{1}{2}\sigma^z \rho_S(t)$ and thereby

$$\dot{\rho}_S^{\text{BM}} = \frac{i}{2}[\sigma^z, \rho_S(t)] + \lambda^2[\sigma^z \rho_S(t) \sigma^z - \rho_S(t)] \quad (56)$$

which coincides with Eq. (55), i.e., the dependence on τ_b vanishes. In contrast, for the more interesting dissipative case ($a=x$) we obtain $\eta_x^{\text{BM}}(t) = \frac{1}{2} \frac{1}{1+\tau_b^2} \sigma^x \rho_S(t) - \frac{1}{2} \frac{\tau_b}{1+\tau_b^2} \sigma^y \rho_S(t)$, which yields

$$\dot{\rho}_S^{\text{BM}} = \frac{i}{2}[\sigma^z, \rho_S(t)] + \frac{\lambda^2}{2} \frac{1}{1+\tau_b^2} [[\sigma^x, \rho_S(t)], \sigma^x]$$

$$- \frac{\lambda^2}{2} \frac{\tau_b}{1+\tau_b^2} [[\sigma^y, \rho_S(t)], \sigma^x]. \quad (57)$$

1. Pure dephasing

In the pure dephasing case one has $a=z$. Inserting the matrix elements in Eq. (53) one finds that ρ_S^{00} and ρ_S^{11} are time independent (as is also known from the full solution) and that the time evolution of the off-diagonal matrix element is governed by

$$\frac{d}{dt} \begin{pmatrix} \rho_S^{01}(t) \\ \bar{\eta}^{01}(t) \end{pmatrix} = \begin{pmatrix} i & -2\lambda^2 \\ +1/\tau_b & i - 1/\tau_b \end{pmatrix} \begin{pmatrix} \rho_S^{01}(t) \\ \bar{\eta}^{01}(t) \end{pmatrix}.$$

With the initial condition $\bar{\eta}^{01}(0) = 0$ one finds

$$\rho_S^{01}(t) = \rho_S^{01}(0) e^{it} e^{-t/(2\tau_b)} \left[\cosh\left(\sqrt{1-8\lambda^2\tau_b} \frac{t}{2\tau_b}\right) + \frac{1}{\sqrt{1-8\lambda^2\tau_b}} \sinh\left(\sqrt{1-8\lambda^2\tau_b} \frac{t}{2\tau_b}\right) \right], \quad (58)$$

which reproduces the decay of the off-diagonal elements (the factor e^{it} is a consequence of the Schrödinger picture). In the high-temperature limit and for the corresponding density of states (51) leading to exponentially decaying correlation functions (49), the decay rate of the exact solution (47) becomes

$$\Gamma(t) = \frac{8\lambda^2}{\pi} \int_0^\infty \frac{\sin^2(\omega t/2)}{\omega^2 [1 + (\omega\tau_b)^2]} d\omega$$

$$= 4\lambda^2 \tau_b \left[\cosh^2\left(\frac{t}{2\tau_b}\right) - 1 - \sinh\left(\frac{t}{2\tau_b}\right) \cosh\left(\frac{t}{2\tau_b}\right) + \frac{t}{2\tau_b} \right], \quad (59)$$

which reduces in the limit $\tau_b \rightarrow 0$ to $\Gamma(t) \approx 2\lambda^2 t$. Similarly, we find that in this limit, Eq. (58) reduces to $\rho_S^{01}(t) = \rho_S^{01}(0) e^{it} e^{-2\lambda^2 t}$. This can be understood as the limit $\tau_b \rightarrow 0$ also corresponds to an infinitely fast relaxation of the reservoir, where also the Born approximation becomes exact. Likewise, it is straightforward to see that $|\rho_S^{01}(t)| - e^{-T(t)} \rho_S^{01}(0) = \mathcal{O}\{\lambda^4\}$. We will therefore not further discuss the pure dephasing case with exponentially decaying correlation functions and compare with the exact solution (47) instead later on.

2. Dissipative coupling

Another important case is reproduced by choosing the dissipation coupling $a=x$ in Eq. (52). Inserting the ansatz (54) into Eq. (53) one finds two 3×3 systems

$$\begin{pmatrix} \dot{\rho}_{00} \\ \dot{\eta}_x \\ \dot{\eta}_y \end{pmatrix} = \begin{pmatrix} 0 & 2\lambda^2 & 0 \\ -\tau_b^{-1} & -\tau_b^{-1} & -1 \\ 0 & 1 & -\tau_b^{-1} \end{pmatrix} \begin{pmatrix} \rho_{00} \\ \eta_x \\ \eta_y \end{pmatrix} + \frac{1}{2\tau_b} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix},$$

$$\begin{pmatrix} \dot{\rho}_x \\ \dot{\rho}_y \\ \dot{\eta}_{00} \end{pmatrix} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 2\lambda^2 \\ 0 & -\tau_b^{-1} & -\tau_b^{-1} \end{pmatrix} \begin{pmatrix} \rho_x \\ \rho_y \\ \eta_{00} \end{pmatrix}, \quad (60)$$

which have an analytic solution that is too lengthy to be reproduced here. At first glance, these systems seem to be




FIG. 1. (Color online) Eigenvalue evolution of the system density matrix (only one eigenvalue shown since trace conservation implies a symmetric evolution) initialized in the pure state $\langle i | \rho_S^0 | j \rangle = 1/2$ for single-qubit dephasing. As predicted by Eq. (63), the solutions of the coarse-graining master equations (thin dashed lines) intersect the exact solution (solid line) at $t = \tau$ (vertical dashed lines). The secular approximation (thick dashed line) corresponds to $\tau \rightarrow \infty$ and does not correctly cover the short time behavior of the exact solution. Parameters were chosen as follows: $\beta = 1$, $\omega_{\text{ph}} = 1$, $\omega_{\text{ct}} = 5$, $s = 1$, $\lambda = 0.1$.

completely independent but note that the condition of initial validity of the density matrix relates their initial conditions. The steady-state solution for the density matrix corresponds to the thermalized Gibbs state (15) for high temperatures ($\beta \rightarrow 0$).

C. Single-qubit coarse graining

1. Pure dephasing

In order to determine the master equations (29) for the pure dephasing limit discussed in Sec. IV A 2, we use

$$A_1 = \sigma^z, \quad B_1 = \sum_k (h_k b_k + h_k^* b_k^\dagger) \quad (61)$$

to obtain that the Lamb shift Hamiltonian in Eq. (23) is proportional to the identity matrix and thus has no effect. From the dissipative part however we obtain a nonvanishing contribution from Eq. (23), such that the master equation (24) in the interaction picture is given by

$$\begin{aligned} \dot{\rho}_S^\tau &= \bar{\Gamma}(\tau) [|0\rangle\langle 0| \rho_S^\tau(t) |0\rangle\langle 0| + |1\rangle\langle 1| \rho_S^\tau(t) |1\rangle\langle 1| \\ &\quad - |0\rangle\langle 0| \rho_S^\tau(t) |1\rangle\langle 1| - |1\rangle\langle 1| \rho_S^\tau(t) |0\rangle\langle 0|] - \bar{\Gamma}(\tau) \rho_S^\tau(t), \\ \bar{\Gamma}(\tau) &= \frac{\lambda^2 \tau}{2\pi} \int_{-\infty}^{+\infty} \gamma_{11}(\omega) \text{sinc}^2\left(\frac{\omega\tau}{2}\right) d\omega. \end{aligned} \quad (62)$$

With assuming exponentially decaying correlation functions (49) we can use Eq. (50) to recover the BM approximation (56), but the above equation for pure dephasing of course holds also for more general correlation functions. It is straightforward to show that the off-diagonal elements of the

density matrix will decay as $\langle 0 | \rho_S^\tau(t) | 1 \rangle = e^{-\bar{\Gamma}(\tau)t} \langle 0 | \rho_S^\tau(0) | 1 \rangle$. A closer inspection of the decay rate yields

$$\begin{aligned} \bar{\Gamma}(\tau)t &= 8\lambda^2 \frac{t}{\tau} \int_{-\infty}^{+\infty} \frac{\sin^2(\omega\tau/2)}{\omega^2} \frac{g(|\omega|)}{|1 - e^{-\beta\omega}|} d\omega \\ &= 8\lambda^2 \frac{t}{\tau} \int_0^{+\infty} \frac{\sin^2(\omega\tau/2)}{\omega^2} g(\omega) \coth(\beta\omega/2) d\omega = \frac{t}{\tau} \Gamma(\tau), \end{aligned} \quad (63)$$

i.e., we reproduce the result of [24] that for pure dephasing of a single qubit, the solution $\rho_S^t(t)$ yields the exact solution $\rho_S(t)$, see also Fig. 1. Naturally, this is also equivalent with the Born equation up to $O\{\lambda^2\}$. Note also that for large coarse-graining times, Fig. 1 shows that the coarse-graining master equations approach the secular approximation master equation—which can also be seen as a numerical confirmation of Eq. (27).

2. Dissipative coupling

With considering $A_1 = \sigma^x$, an exponentially decaying correlation function (49), and $H_S = \frac{1}{2}[\mathbf{1} - \sigma^z]$ one obtains a more complicated master equation for $\rho_S^\tau(t)$ in the Schrödinger picture

$$\begin{aligned} \dot{\rho}_S^\tau(t) &= \left[\frac{i}{2} \sigma^z - i \sigma_{00}(\tau) |0\rangle\langle 0| - i \sigma_{11}(\tau) |1\rangle\langle 1|, \rho_S^\tau \right] \\ &\quad + \gamma_{01,01}(\tau) \left[|0\rangle\langle 1| \rho_S^\tau |1\rangle\langle 0| - \frac{1}{2} \{ |1\rangle\langle 1|, \rho_S^\tau \} \right] \\ &\quad + \gamma_{10,10}(\tau) \left[|1\rangle\langle 0| \rho_S^\tau |0\rangle\langle 1| - \frac{1}{2} \{ |0\rangle\langle 0|, \rho_S^\tau \} \right] \\ &\quad + \gamma_{01,10}(\tau) |0\rangle\langle 1| \rho_S^\tau |0\rangle\langle 1| + \gamma_{10,01}(\tau) |1\rangle\langle 0| \rho_S^\tau |1\rangle\langle 0|. \end{aligned} \quad (64)$$

From the nonvanishing matrix elements in Eq. (24) one can deduce that (unlike in the pure dephasing case) the Lamb-shift term will contribute, since (although diagonal) it is not proportional to the identity matrix anymore. Likewise, we also observe here a decoupled evolution of diagonal

$$\begin{aligned} \rho_{S_{00}}^\tau(t) &= \frac{\gamma_{01,01}}{\gamma_{01,01} + \gamma_{10,10}} (1 - e^{-(\gamma_{01,01} + \gamma_{10,10})t}) \\ &\quad + e^{-(\gamma_{01,01} + \gamma_{10,10})t} \rho_{S_{00}}^\tau(0) \end{aligned} \quad (65)$$

and off-diagonal matrix elements

$$\dot{\rho}_{S_{01}}^\tau = [i - \Gamma(\tau)] \rho_{S_{01}}^\tau + \gamma_{01,10}(\tau) (\rho_{S_{01}}^\tau)^*, \quad (66)$$

with $\Gamma(\tau) = \frac{1}{2}(\gamma_{01,01} + \gamma_{10,10}) - i(\sigma_{11} - \sigma_{00})$ (we have omitted the τ dependence). Considering even and odd Fourier transforms of the form (50) corresponding to exponentially decaying correlation functions, we can now compare the solution (60) of the Born equation with the solutions (64) of the coarse-graining approach.

When one initializes the density matrix as $\rho_S(0) = |0\rangle\langle 0|$ one finds that for infinite coarse-graining times the diagonal terms will equilibrate to value 1/2 (which corresponds to




FIG. 2. (Color online) Evolution of the ρ_{11} -matrix element of the density matrix for dissipative coupling. The solid line represents the solution of the Born equation (60) and the thin dashed lines correspond to solutions of Eq. (64) for different coarse-graining times τ . Up to second order in λ , a perfect numerical agreement is found for the matrix elements of $\rho^i(t)$ (symbols) and the Born solution (solid line). Parameters were chosen as follows: $\lambda=0.1$, $\tau_b=1$.

thermalization at infinite temperatures), see Fig. 2. In this case, the off-diagonal terms will evidently vanish throughout. Note that for the small coupling chosen ($\lambda=0.1$), the solution of the Born equation $\rho_S(t)$ is approximated by the adaptive coarse-graining solutions $\rho_S^i(t)$ with extraordinary accuracy.

In contrast, when initializing the density matrix as $\rho_S(0) = \frac{1}{2}[|0\rangle\langle 0| + |0\rangle\langle 1| + |1\rangle\langle 0| + |1\rangle\langle 1|]$ one observes that the




FIG. 3. (Color online) Evolution of the $|\rho_{01}|$ matrix element of the density matrix for pure dissipation. The solid line represents the solution of the Born equation (60) and the thin dashed lines correspond to solutions of Eq. (64) for different coarse-graining times τ . Up to second order in λ , a perfect agreement is found for the matrix elements of $\rho^i(t)$ (symbols) and the Born solution (solid line). The difference between Born solution and BM as well as BMS approximations is too small to be visible. Parameters were chosen as follows: $\lambda=0.1$, $\tau_b=1$.




FIG. 4. (Color online) Evolution of the eigenvalues of the density matrix for a single qubit initialized in the pure state $\langle i|\rho_S^0|j\rangle=1/2$. For long times the BMS solution (thick dashed line) and the adaptive coarse-graining solution (thick solid line) predict thermalization in the thermal equilibrium state (15), whereas a different equilibrium state is assumed for short coarse-graining times (thin dashed lines). Parameters were chosen as follows: $\beta=1$, $\omega_{ph}=1$, $\omega_{ct}=5$, $s=1$, $\lambda=0.1$.

diagonal entries will remain unchanged and the off-diagonals will decay. The actual behavior of the decay of the off-diagonal elements is depicted in Fig. 3. Again we observe a strikingly good agreement between the Born solution $\rho_S(t)$ and the adaptive coarse-graining solutions $\rho_S^i(t)$.

It is also instructive to compare for the adaptive coarse-graining approach the limit of infinite coarse-graining times (BMS approximation)

$$\begin{aligned} \dot{\rho}_S = & \left(\frac{i}{2} + \frac{i}{2} \frac{\lambda^2 \tau_b}{1 + \tau_b^2} \right) [\sigma^z, \rho_S] + \frac{\lambda^2}{1 + \tau_b^2} [|0\rangle\langle 1| \rho_S |1\rangle\langle 0| \\ & + |1\rangle\langle 0| \rho_S |0\rangle\langle 1| - \rho_S] \end{aligned} \quad (67)$$

with the BM approximation master Eq. (57), where one can see that only the equations for the diagonals match. Likewise, in the limit $\tau_b \rightarrow 0$ one obtains

$$\begin{aligned} \dot{\rho}_S^\tau = & \frac{i}{2} [\sigma^z, \rho_S^\tau] + \lambda^2 [|0\rangle\langle 1| \rho_S^\tau |1\rangle\langle 0| + |1\rangle\langle 0| \rho_S^\tau |0\rangle\langle 1| - \rho_S^\tau] \\ & + \lambda^2 e^{-i\tau} \text{sinc } \tau |0\rangle\langle 1| \rho_S^\tau |0\rangle\langle 1| + \lambda^2 e^{i\tau} \text{sinc } \tau |1\rangle\langle 0| \rho_S^\tau |1\rangle\langle 0|. \end{aligned} \quad (68)$$

Here, one finds that the BMS approximation ($\tau \rightarrow \infty$) yields a different master equation than the BM limit (55) and also the BM approximation (57). However, also within the coarse-graining approach the limit $\tau_b \rightarrow 0$ (with finite τ) leads to the same steady state as the BMS approximation $\tau \rightarrow \infty$ and the BM limit (55)—only the relaxation rates may differ. Note however that the BM approximation (57) may even lead to instable behavior of the off-diagonal matrix elements for large λ (compare also Sec. IV D), whereas the coarse-graining approach always generates Lindblad forms.

3. General coupling

For a more complex coupling of the spin to the reservoir

$$H_S = \frac{1}{2}[\mathbf{1} - \sigma^z], \quad \vec{n}_1 = \frac{1}{\sqrt{2}}(1 + i, 1 + i, 1 + i), \quad (69)$$

and also more realistic spectral coupling densities of type (45) we find thermalization as predicted by the BMS approximation for long times. For short times however, the solution $\rho_S^t(t)$ may strongly differ from the BMS solution, see Fig. 4. For example, if coarse-graining times are chosen too small, a nonthermalized steady state is obtained. For large coarse-graining times, the thermalized steady state is reached, but then in the short time regime, large differences between fixed graining and the adaptive graining solution are found. Again, we numerically confirm Eq. (27), since for large coarse-graining times the secular approximation is reproduced.

D. Staying reasonable

Frequently, the BM approximation (12) is used although it does not generally guarantee for positive evolution. This is certainly tolerable if the solution obtained approximates the exact solution well (and thus violates positivity only slightly). In addition, if the interaction Hamiltonian used in such models implicitly corresponds to a secular approximation [6,35], one will obtain a Lindblad form master equation and positivity as well as stability of the density matrix will be granted throughout. In general however, this will not be the case. Here we will analytically consider a single qubit $H_S = \frac{1}{2}[\mathbf{1} - \sigma^z]$ with the simple coupling $H_{SB} = \sigma^x \otimes B$. Denoting with $\Gamma(\omega)$ the half-sided Fourier transform (11) of the reservoir correlation function, one can write Eq. (12) in the form $\dot{\rho}_S = L\rho_S(t)$ and calculate the eigenvalues of the Liouvillian.

If one is only interested in the subspace of diagonal density matrices one finds the two corresponding eigenvalues $\sigma_0 = 0$ and $\sigma_1 = -2\lambda^2 A$ with $A = \text{Re}[\Gamma(+1) + \Gamma(-1)]$. For physically motivated bath correlation functions C_{11} in Eq. (43) one obtains that $\sigma_1 < 0$, such that the evolution in this subspace may not lead to unstable behavior—although positivity may be violated.

In contrast, in the off-diagonal subspace one obtains the two eigenvalues $\lambda_{2/3} = -2\lambda^2 A \pm \sqrt{-1 + 2B\lambda^2 + A^2\lambda^4}$ with A defined as above and $B = \text{Im}[\Gamma(-1) - \Gamma(+1)]$. Given $B > 0$ [which can be achieved with correlation functions of the form (43)], one of these will pick a positive real part as soon as $\lambda^2 > (2B)^{-1}$, which corresponds to an unstable evolution, see also Fig. 5. Numerically, we observe the same for $n=2$ mutually uncoupled qubits (with similar system Hamiltonians and couplings). In this case, the expectation value of operators that are not diagonal in the system Hamiltonian will not yield any meaningful results and the Born-Markov approximation is questionable.

E. Thermalization

In the pure dephasing limit, one observes a rapid (i.e., exponential) decay of the off-diagonal elements of the den-




FIG. 5. (Color online) Largest eigenvalue of the density matrix for one (solid) and two (dashed) qubits initialized with $\langle i | \rho_S^0 | j \rangle = 1/2^n$ calculated from Eq. (12). For large λ , not only positivity is violated (negative eigenvalues not shown—trace conservation), but the solution even becomes unstable. The parameters in Eq. (43) were chosen as $\beta=1$, $\omega_{ph}=1$, $\omega_{ct}=5$, $s=2$, such that the transition from stable to unstable behavior occurs at $\lambda_{crit} \approx 0.231294$ for $n=1$.

sity matrix. Also, if there are no degeneracies in the spectrum, the rotating wave approximation (14) predicts a decoupled evolution of the diagonal elements of the density matrix in the eigenbasis of H_S , i.e., from the corresponding evolution equation $\dot{\rho}_{ii}(t) = \sum_j A_{ij} \rho_{jj}(t)$ one might expect an exponential decay into the eigenvector of the matrix A which has eigenvalue 0 (steady state). The corresponding subspace can still be degenerate (many stationary states) or there may exist exponentially many other eigenvectors with very small eigenvalues, such that thermalization does not necessarily happen very fast. Here we will consider some specific realizations of the spin-boson model (35) and solve Eq. (14) numerically.

For example, if the system Hamiltonian has degeneracies and the coupling to the reservoir does not lift these, the system may relax into states that are not even diagonal in the system Hamiltonian basis. In these cases, the initial density matrix may determine which steady state is actually reached, see Fig. 6. As a first example we consider

$$\gamma = 1, \quad \gamma_1^z = \gamma_2^z = -\frac{1}{2},$$

$$\vec{n}_1 = \vec{n}_2 = \frac{1}{\sqrt{2}}(1 + i, 1 + i, 1 + i), \quad (70)$$

and all other coefficients of Eq. (35) vanishing, such that there exists a twofold degeneracy in the spectrum of H_S . In this section, we will consider spectral densities of type (45).

For the low reservoir temperatures assumed in Fig. 6, thermalization corresponds to relaxation into the ground state and one can see that it may depend on the initial state of the density matrix and possibly lifted degeneracies whether thermalization takes place. In reality, the degeneracy might always be lifted by some imperfect Hamiltonian implementa-




FIG. 6. (Color online) Top: Eigenvalue evolution for a 2-qubit system according to the secular approximation master equation (14) with a twofold degeneracy for different initial states (solid and dashed lines) and with a near twofold degeneracy (dotted lines). Vertical dashed lines indicate the times at which the bottom row snapshots have been taken. Bottom rows: Absolute values of the matrix elements of the corresponding 4×4 density matrix in the ordered energy eigenbasis (such that the top left-hand corner corresponds to the ground state). Color coding ranges from blue (value 0) to red (value 1). In the upper row, the system is initialized in a nonthermal density matrix $\langle i | \rho_{ij}^0 | j \rangle = 1/4$ and relaxes into a thermalized state (15) with $\beta_{\text{sys}} = \beta$, i.e., for the low reservoir temperature chosen ($\beta = 10$) essentially into the ground state. In contrast, in the middle row the initial state was a thermalized one with initial system temperature $\beta_{\text{sys}}^0 = 1$, and the system does not relax into a thermal equilibrium state. In the lowest row, the degeneracy was broken by choosing $\gamma_1^z = -1/2 + 0.01$ and $\gamma_2^z = -1/2 - 0.01$. Other parameters were chosen as $\beta = 10$, $\omega_{\text{ph}} = 1$, $\omega_{\text{ct}} = 5$, $s = 1$, $\lambda = 0.1$.

tions. In addition, the coupling to the reservoir may be more complex than in Eq. (70) thus making the thermalized system state (15) the only stationary state of the rotating wave master equation (14), compare also [4]. However, it is to be expected that thermalization in this case will take rather long times, especially for large and complicated systems (see the next section).

F. Solving problems by cooling

It is known that Hamiltonians of the form (35) can be used to encode solutions to computationally hard problems

in their ground state. This is, for example, exploited in adiabatic quantum computation [7]. For a system made of five spins, we will compare an enlarged version of the previous example (70),

$$\gamma = \frac{5}{2}, \quad \gamma_{i=1..5}^z = -\frac{1}{2} \quad (71)$$

with the ground-state encoding of *Exact Cover 3*—a specific NP-complete problem.

The Exact Cover 3 problem can be introduced as follows [7]: Given a set of m constraints where each constraint contains the positions of three bits $C_j = (p_j^1, p_j^2, p_j^3)$ (with evidently $1 \leq p_j^i \leq n$), one is looking for the n -bit bit string b_1, \dots, b_n (with $b_i \in \{0, 1\}$) that fulfills for each constraint $b_{p_j^1} + b_{p_j^2} + b_{p_j^3} = 1$ (where the plus sign denotes the integer sum). One way to encode the solution to this problem into the ground state of a Hamiltonian of the type (35) is given by [36,37]

$$\gamma = m, \quad \tilde{\gamma}_i^z = -\frac{n_i}{2}, \quad \tilde{\gamma}_{ij}^{zz} = +\frac{n_{ij}}{2} \quad (72)$$

and all other coefficients vanishing—the differing prefactor of $\tilde{\gamma}_{ij}^{zz}$ in comparison to [37] results from the absence of double counting in ij in Eq. (35). In the above equation, m denotes the total number of clauses, n_i the number of clauses involving bit i , and n_{ij} the number of clauses involving both bits i and j . Specifically, we will consider the four clauses $C_1 = (2, 3, 4)$, $C_2 = (1, 2, 5)$, $C_3 = (1, 4, 5)$, and $C_4 = (3, 4, 5)$. This implies the nonvanishing coefficients $\gamma = 4$, $\tilde{\gamma}_1^z = \tilde{\gamma}_2^z = \tilde{\gamma}_3^z = -1$, $\tilde{\gamma}_4^z = \tilde{\gamma}_5^z = -3/2$, $\tilde{\gamma}_{12}^{zz} = \tilde{\gamma}_{14}^{zz} = \tilde{\gamma}_{23}^{zz} = \tilde{\gamma}_{24}^{zz} = \tilde{\gamma}_{25}^{zz} = \tilde{\gamma}_{35}^{zz} = 1/2$, and $\tilde{\gamma}_{15}^{zz} = \tilde{\gamma}_{34}^{zz} = \tilde{\gamma}_{45}^{zz} = 1$.

As can be easily checked, this problem has the unique solution $|10100\rangle$ (with energy $E_0 = 0$) and the first excited states (with a sixfold degeneracy) are given by $|00101\rangle, |10010\rangle, |00001\rangle, |00010\rangle, |01001\rangle, |01010\rangle$ with energies $E_1 = \dots = E_6 = 1$. The first excited states have Hamming distances (i.e., number of bit flips necessary for transformation) to the solution of 2, 2, 3, 3, 4, 4, respectively. This already indicates the hardness of such problems. Simple coupling Hamiltonians such as Eq. (37) that are only linear in the Pauli matrices will to first order only yield transitions between states with Hamming distance 1, since $\langle a | \sigma_i^{x/y} | b \rangle = 0$ if $|a\rangle$ and $|b\rangle$ have Hamming distance larger than 1. Of course, this does not completely prohibit transitions between states with a larger Hamming distance, but such tunneling processes will have to pass through energetically less favorable states and are therefore strongly suppressed. Accordingly, one may expect that the process of thermalization is strongly hampered. This is also observed in Fig. 7, where we have assumed a reservoir temperature much smaller than the fundamental energy gap. Whereas for the simple qubit system (71) the system rapidly relaxes into the ground state, this is very different for the example (72).

V. CONCLUSION

We have compared different procedures of deriving master equations from microscopic models and have shown that

by using a coarse-graining approach one always obtains master equations in Lindblad form. This ensures for positivity and stability of the density matrix. In contrast, the usual Markovian approximation scheme may sometimes lead to a non-positive and even unstable behavior, where in the latter case there is no hope of approximating the exact solution. The coarse-grained master equations depend on a parameter—the coarse-graining time scale τ . For short coarse-graining times that are adaptively matched with the physical time, the solution $\rho_S^\tau(\tau)$ must approximate the result of the Born approximation by construction. For large coarse-graining times and time-independent system Hamiltonians, we reproduce the secular approximation. For all intermediate coarse-graining times, a positive evolution of the system density matrix is ensured by the Lindblad form of the resulting differential equations governing the time evolution. For the special case of pure dephasing of a single qubit we reproduce the analytical result by [24] that $\rho_S^t(t) = \rho_S(t)$ yields the exact solution (which is of course equivalent in the weak-coupling limit to the Born approximation). For the simple example of exponentially decaying correlation functions we find by numerical simulation a surprisingly perfect agreement between solutions of the integro-differential Born equation and solutions of the adaptive coarse-graining approach. Given an interest in the system density matrix at time t , we therefore propose to match the coarse-graining time with the physical time $\tau = t$. In this case one must calculate the Liouvillian matrix elements (23) only once (for the desired time t) and then evolve the density matrix using Eq. (24) until that time t . In terms of computational complexity, this is more efficient than an evolving integro-differential equation. A similar advantage is given when the resulting master equations are so simple such that an analytical solution in terms of the damping matrix elements is possible. If in contrast the system is very complex and one is interested in the density matrix at all times, this advantage is destroyed. From a computational perspective, it would therefore be interesting to find the general map

$$\dot{\rho}_S(t) = \mathcal{L}(t)\rho_S(t) = \lim_{\Delta t \rightarrow 0} \frac{\rho_S(t + \Delta t) - \rho_S(t)}{\Delta t} \quad (73)$$

which fulfills $\rho_S(t) = \rho_S^t(t)$. Formally, such a map can be found by inserting the solution $\rho_S^\tau(t) = e^{L^\tau t} \rho_S(0)$ into the derivative in Eq. (73),

$$\mathcal{L}(t) = \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} (e^{L^{t+\Delta t}(t+\Delta t)} e^{-L^t t} - 1) = \left(\frac{d}{dt} e^{L^t t} \right) e^{-L^t t}. \quad (74)$$

We have given various examples of simple qubit systems coupled to a bosonic reservoir which have demonstrated that thermalization of spin systems coupled to a bosonic bath may depend on a plethora of factors such as the initial state, complexity of the system Hamiltonian, and complexity of the coupling. Future research will consider the importance of the coarse-graining approach for different scenarios.

ACKNOWLEDGMENTS

One of the authors (G.S.) is indebted to G. Kiesslich and M. Vogl for fruitful discussions.




FIG. 7. (Color online) Top: Eigenvalue evolution for a 5-qubit system according to the secular approximation master equation (14) for different system Hamiltonians given by Eq. (71) (solid lines) and Eq. (72) with an Exact Cover 3-problem (dashed lines), respectively. In both cases, the density matrix has been initialized as $\langle i | \rho_S^0 | j \rangle = 1/32$. Whereas for the uncoupled spin system the system rapidly relaxes into the ground state, the final state for the exact cover problem is very different. Bottom: Absolute values of the matrix elements of the 7×7 top left-hand submatrix of the full 32×32 density matrix in the ordered energy eigenbasis. On the left-hand side (corresponding to the solid line on top), the system relaxes to the ground state (red spot), whereas for the exact cover problem (right-hand side, dashed line on top), the system state is not thermalized. Other parameters were chosen as $\beta = 10$, $\omega_{ph} = 1$, $\omega_{cl} = 5$, $s = 1$, $\lambda = 0.1$.

APPENDIX A: NEGLECT OF BACK-ACTION

Throughout the paper, we make the following simplifying assumptions:

Initial factorization of the density matrix. By assuming

$$\rho(0) = \rho_S(0) \otimes \rho_B^0 = \rho_S(0) \otimes \rho_B^0 \quad (A1)$$

we implicitly demand that one can at $t=0$ prepare the system in a product state, which requires sufficient experimental control.

Born approximation. For a bath much larger than the system and weak coupling it is reasonable to assume that the back-action of the system onto the bath is small, such that the bath part of the density matrix is hardly changed from its initial value, i.e.,

$$\rho(t') = \rho_S(t') \otimes \rho_B^0 + O\{\lambda\}. \quad (A2)$$

Clearly, inserting this approximation in the third term in Eq. (7) is consistent up to second order in λ .

Reservoir in stationary state. We will assume that the reservoir is in a stationary state of the bath Hamiltonian, which

implies $[\rho_B^0, H_B]=0$. One possibility of such a stationary state is to assume the thermal equilibrium state ($\beta^{-1}=k_B T$ at temperature T)

$$\rho_B^0 = \rho_B = \frac{e^{-\beta H_B}}{\text{Tr}_B\{e^{-\beta H_B}\}}. \quad (\text{A3})$$

APPENDIX B: MARKOV APPROXIMATION

Starting from the Born equation (8), the Markovian approximation in [4] is performed by replacing (in the interaction picture) $\rho_S(t') \rightarrow \rho_S(t)$ under the integral, substituting $\tau = t - t'$ and extending the integration to infinity. This is usually motivated by fast decaying reservoir correlation functions (9). By doing so, we obtain the BM master equation

$$\begin{aligned} \dot{\rho}_S = & -i \text{Tr}_B\{[H_{SB}(t), \rho_S(0)\rho_B^0]\} \\ & - \int_0^\infty \text{Tr}_B\{[H_{SB}(t), [H_{SB}(t-\tau), \rho_S(t)\rho_B^0]]\}d\tau + O\{\lambda^3\}, \end{aligned} \quad (\text{B1})$$

where one can now insert the decomposition (2) of the interaction Hamiltonian to obtain

$$\begin{aligned} \dot{\rho}_S = & -i\lambda \sum_A \langle B_A \rangle [A_A(t), \rho_S(0)] \\ & + \lambda^2 \left(\sum_{AB} \int_0^\infty [A_B(t-\tau)\rho_S(t), A_A(t)] C_{AB}(\tau) d\tau + \text{H.c.} \right) \\ & + O\{\lambda^3\}, \end{aligned} \quad (\text{B2})$$

with the reservoir correlation functions (9). We will use $\langle B_A \rangle = 0$ throughout, since this case can always be generated with a suitable transformation [22]. In order to evaluate the time dependence of the operators in Eq. (B2), it is useful to expand them into eigenoperators of the system Hamiltonian (assuming H_S to be time independent),

$$\begin{aligned} A_A &= \sum_\omega A_A(\omega) = \sum_\omega A_A^\dagger(\omega) = A_A^\dagger, \\ A_A(\omega) &= \sum_{ab} \delta_{(E_b - E_a), \omega} |a\rangle \langle a| A_A |b\rangle \langle b|, \end{aligned} \quad (\text{B3})$$

where the variable ω ranges over all energy differences of H_S , $H_S|a\rangle = E_a|a\rangle$, and δ is a Kronecker symbol. These operators $A_A(\omega)$ have the advantageous properties

$$\begin{aligned} [H_S, A_A(\omega)] &= -\omega A_A(\omega), \\ [H_S, A_A^\dagger(\omega)] &= +\omega A_A^\dagger(\omega), \\ [H_S, A_A^\dagger(\omega)A_B(\omega)] &= 0, \end{aligned} \quad (\text{B4})$$

which implies in the interaction picture

$$A_A(t) = \sum_\omega e^{-i\omega t} A_A(\omega) = \sum_\omega e^{+i\omega t} A_A^\dagger(\omega). \quad (\text{B5})$$

With inserting the half-sided Fourier transform of the reservoir correlation functions (11) we obtain for the master equation in Eq. (B2) with $\langle B_A \rangle = 0$,

$$\begin{aligned} \dot{\rho}_S \approx & \lambda^2 \left(\sum_{AB} \sum_{\omega, \omega'} e^{i(\omega' - \omega)t} \Gamma_{AB}(\omega) [A_B(\omega)\rho_S(t), A_A^\dagger(\omega')] \right. \\ & \left. + \text{H.c.} \right). \end{aligned} \quad (\text{B6})$$

Finally, we observe that by reinserting the definition of the eigenoperators (B3) in Eq. (B6) and switching back to the Schrödinger picture the time-dependent phase factors vanish and one obtains the Born-Markov master equation (12).

APPENDIX C: SECULAR APPROXIMATION

For large times, the terms with an oscillating prefactor in Eq. (B6) will average out and by inserting

$$e^{i(\omega' - \omega)t} \rho_S(t) \approx \delta_{\omega, \omega'} \rho_S(t) \quad (\text{C1})$$

in Eq. (B6) and switching to the Schrödinger picture (compare also, e.g., Chap. 3.3 in [4]) we obtain

$$\begin{aligned} \dot{\rho}_S = & -i[H_S, \rho_S(t)] + \lambda^2 \sum_\omega \sum_{AB} \{\Gamma_{AB}(\omega) [A_B(\omega)\rho_S, A_A^\dagger(\omega)] \\ & + \text{H.c.}\}. \end{aligned} \quad (\text{C2})$$

The advantage of the secular approximation is that we can now combine the half-sided Fourier transforms to full (even and odd) Fourier transforms (13) of the reservoir correlation functions. Inserting these definitions into Eq. (C2), one obtains a Lindblad [16] form

$$\begin{aligned} \dot{\rho}_S = & -i[H_S, \rho_S(t)] - i \left[\frac{\lambda^2}{2i} \sum_\omega \sum_{AB} \sigma_{AB}(\omega) A_A^\dagger(\omega) A_B(\omega), \rho_S(t) \right] \\ & + \lambda^2 \sum_\omega \sum_{AB} \gamma_{AB}(\omega) \left(A_B(\omega) \rho_S A_A^\dagger(\omega) \right. \\ & \left. - \frac{1}{2} \{A_A^\dagger(\omega) A_B(\omega), \rho_S(t)\} \right), \end{aligned} \quad (\text{C3})$$

where the positivity of the $\gamma_{AB}(\omega)$ matrix is guaranteed by Bochner's theorem [4,38], which states that the Fourier transform of a function of positive type (as are the reservoir correlation functions) gives rise to a positive definite matrix. In the above equation, the second commutator corresponds to the unitary action of decoherence (Lamb shift). Finally, we can insert the operator definitions (B3) in (C3) to obtain Eq. (14). Naturally, if $[H_S, A_A] = 0$ (pure dephasing), BM and BMS approximations are equivalent.

APPENDIX D: ADIABATIC APPROXIMATION

With inserting the ansatz

$$U(t) = \sum_{ab} u_{ab}(t) \exp\left(-i \int_0^t E_a(t') dt'\right) |a(t)\rangle \langle b(0)|, \quad (\text{D1})$$

where the $|a(t)\rangle$ span an instantaneous basis (chosen to be complete and orthonormal) of the system Hilbert space defined via $H_S(t)|a(t)\rangle = E_a(t)|a(t)\rangle$ in the evolution equation for

the time evolution operator $\hat{U}(t) = -iH_S(t)U(t)$, one obtains an equation for the expansion coefficients

$$\dot{u}_{ab} + u_{ab}\langle a|\dot{a}\rangle = -\sum_{c \neq a} u_{cb} e^{-i\int_0^t g_{ca}(t') dt'} \langle a|\dot{c}\rangle \quad (\text{D2})$$

with the energy gap

$$g_{ca}(t') = E_c(t') - E_a(t'). \quad (\text{D3})$$

With introducing the Berry phase [39]

$$\gamma_a(t) = i \int_0^t \langle a(t')|\dot{a}(t')\rangle dt' \quad (\text{D4})$$

this can also be written as

$$\frac{d}{dt}(u_{ab} e^{-i\gamma_a(t)}) = -\sum_{c \neq a} u_{cb} e^{-i\gamma_a(t) - i\int_0^t g_{ca}(t') dt'} \langle a|\dot{c}\rangle,$$

which gives the general time evolution of the expansion coefficients u_{ab} for any (also nonadiabatic) system Hamiltonian $H_S(t)$ if one uses the initial condition $u_{ab}(0) = \delta_{ab}$. The full adiabatic approximation essentially consists in setting the right-hand side of above equation to zero: For slowly varying system Hamiltonians, the change of the eigenvectors will be negligible such that $\langle a|\dot{c}\rangle \approx 0$. Note however, that in the vicinity of avoided crossings, the condition of adiabaticity relates the maximum speed of the time evolution with the spectral properties of the system Hamiltonian

$$\langle a|\dot{c}\rangle = \frac{\langle a|H_S|\dot{c}\rangle}{E_c - E_a}, \quad (\text{D5})$$

see also, e.g., [9,10]. If $\langle a|\dot{c}\rangle \approx 0$, one obtains with the adiabatic approximation $u_{ab}^{\text{ad}}(t) = \delta_{ab} e^{i\gamma_a(t)}$ which implies for the time evolution operator in the adiabatic limit

$$U(t) \approx \sum_a e^{i\gamma_a(t) - i\int_0^t E_a(\tau) d\tau} |a(t)\rangle \langle a(0)|. \quad (\text{D6})$$

Therefore, we obtain for the time-averaged operator in Eq. (34),

$$\begin{aligned} \langle \langle \tilde{A}_{\mathcal{A}}(\omega) \rangle \rangle_{[t, t+\tau]} &\approx \sum_{ab} |a(0)\rangle \langle b(0)| \frac{1}{\tau} \int_t^{t+\tau} \langle a(t')|A_{\mathcal{A}}|b(t')\rangle \\ &\times e^{-i[\gamma_{ab}(t') - \omega t' - \int_0^{t'} g_{ab}(\tau) d\tau]} dt' \end{aligned} \quad (\text{D7})$$

with $\gamma_{ab}(t') = \gamma_a(t') - \gamma_b(t')$ and $g_{ab}(\tau) = E_a(\tau) - E_b(\tau)$.

APPENDIX E: POSITIVITY-PRESERVING MASTER EQUATIONS

Here we will show that master equations of the form

$$\begin{aligned} \dot{\rho} = &-i[\mathcal{H}(t), \rho(t)] + \sum_{\alpha\beta=1}^K \Gamma_{\alpha\beta}(t) \left(\mathcal{L}_{\alpha}(t) \rho(t) \mathcal{L}_{\beta}^{\dagger}(t) \right. \\ &\left. - \frac{1}{2} \{ \mathcal{L}_{\beta}^{\dagger}(t) \mathcal{L}_{\alpha}(t), \rho(t) \} \right) \end{aligned} \quad (\text{E1})$$

generally preserve the positive semidefiniteness of an initial condition $\rho(0)$ if the matrix $\Gamma_{\alpha\beta}(t)$ is positive semidefinite and the operator $\mathcal{H}(t) = \mathcal{H}^{\dagger}(t)$ is Hermitian at all times (smoothness of all time dependencies provided). With discretizing the time derivative and by introducing new operators

$$W_1(t) = \mathbf{1} = W_1^{\dagger}(t),$$

$$W_2(t) = i\mathcal{H}(t) + \frac{1}{2} \sum_{\alpha\beta} \Gamma_{\alpha\beta}(t) \mathcal{L}_{\beta}^{\dagger}(t) \mathcal{L}_{\alpha}(t),$$

$$W_3(t) = \mathcal{L}_1(t), \quad \dots, \quad W_{K+2}(t) = \mathcal{L}_K(t) \quad (\text{E2})$$

we obtain

$$\rho(t + \Delta t) = \sum_{\alpha\beta=1}^{K+2} w_{\alpha\beta}(t) W_{\alpha}(t) \rho(t) W_{\beta}^{\dagger}(t), \quad (\text{E3})$$

where the $w_{\alpha\beta}(t)$ matrix is given by

$$w(t) = \left(\begin{array}{cc|ccc} 1 & -\Delta t & 0 & \dots & 0 \\ -\Delta t & 0 & 0 & \dots & 0 \\ \hline 0 & 0 & & & \\ \vdots & \vdots & & \Delta t \Gamma(t) & \\ 0 & 0 & & & \end{array} \right). \quad (\text{E4})$$

This matrix has a simple block structure and it is therefore straightforward to relate the eigenvalues of w to those of Γ . In particular, one obtains the eigenvalues $w_1 = 1/2(1 - \sqrt{1 + 4\Delta t^2})$, $w_2 = 1/2(1 + \sqrt{1 + 4\Delta t^2})$, and $w_{i \geq 3}(t) = \Delta t \gamma_{i-2}(t)$, where $\gamma_{i-2}(t)$ are the non-negative eigenvalues of $\Gamma(t)$. With diagonalizing the matrix in Eq. (E3) via $w_{\alpha\beta}(t) = \sum_{\gamma} \mu_{\alpha\gamma}(t) u_{\beta\gamma}^*(t) w_{\gamma}(t)$ with a suitable (time-dependent) unitary transformation $U(t)$ we obtain from Eq. (E3)

$$\rho(t + \Delta t) = \sum_{\gamma} w_{\gamma}(t) \tilde{W}_{\gamma}(t) \rho(t) \tilde{W}_{\gamma}^{\dagger}(t),$$

$$\tilde{W}_{\gamma}(t) = \sum_{\alpha} u_{\alpha\gamma}(t) W_{\alpha}(t). \quad (\text{E5})$$

Assuming that at time t we have a valid density matrix with $0 \leq \rho_{\delta}(t) \leq 1$ we obtain by inserting the spectral decomposition $\rho(t) = \sum_{\delta} \rho_{\delta}(t) |\Phi_{\delta}(t)\rangle\langle\Phi_{\delta}(t)|$,

$$\begin{aligned} \langle\Phi|\rho(t + \Delta t)|\Phi\rangle &= \sum_{\gamma\delta} w_{\gamma}(t) \rho_{\delta}(t) |\langle\Phi|\tilde{W}_{\gamma}|\Phi_{\delta}(t)\rangle|^2 \\ &\geq \frac{1}{2} (1 - \sqrt{1 + 4\Delta t^2}) \sum_{\delta} \rho_{\delta}(t) |\langle\Phi|\tilde{W}_1|\Phi_{\delta}(t)\rangle|^2 \\ &\geq -\Delta t^2 \sum_{\delta} \rho_{\delta}(t) |\langle\Phi|\tilde{W}_1|\Phi_{\delta}(t)\rangle|^2 \stackrel{\Delta t \rightarrow 0}{\geq} 0, \quad (\text{E6}) \end{aligned}$$

such that in the limit $\Delta t \rightarrow 0$ [which defines the original differential Eq. (E1)], the smallest eigenvalue of the density matrix at time $t + \Delta t$ approaches zero faster than the discretization width Δt . Therefore, in this limit the matrix $w(t)$ becomes positive semidefinite and the differential equation (E3) becomes a positivity-preserving map.

APPENDIX F: SINC DISTRIBUTION

For discrete a, b , and continuous ω we would like to analyze

$$\begin{aligned} f(\omega, a, b) &\equiv \lim_{\tau \rightarrow \infty} \tau \operatorname{sinc}\left((\omega + a)\frac{\tau}{2}\right) \operatorname{sinc}\left((\omega + b)\frac{\tau}{2}\right) \\ &= \lim_{\tau \rightarrow \infty} \frac{4 \sin\left((\omega + a)\frac{\tau}{2}\right) \sin\left((\omega + b)\frac{\tau}{2}\right)}{(\omega + a)(\omega + b)\tau}. \quad (\text{F1}) \end{aligned}$$

Under an integral with a smooth function $g(\omega)$ one can consider the case $a \neq b$ by a partial fraction expansion where

without loss of generality we find for the first term (due to the symmetry the second term can be treated in a completely analogous way)

$$\begin{aligned} I_{a \neq b} &= \int_{-\infty}^{+\infty} g(\omega) \frac{\sin\left((\omega + a)\frac{\tau}{2}\right) \sin\left((\omega + b)\frac{\tau}{2}\right)}{\tau(\omega + a)} d\omega \\ &= \cos\left((b - a)\frac{\tau}{2}\right) \int_{-\infty}^{+\infty} g(\omega) \frac{\sin^2\left((\omega + a)\frac{\tau}{2}\right)}{\tau(\omega + a)} d\omega \\ &\quad + \frac{1}{2} \sin\left((b - a)\frac{\tau}{2}\right) \int_{-\infty}^{+\infty} g(\omega) \frac{\sin[(\omega + a)\tau]}{\tau(\omega + a)} d\omega, \end{aligned}$$

where we have inserted $(\omega + b) = (\omega + a) + (b - a)$ to use trigonometric relations for $\sin(\alpha + \beta)$. With a suitable transformation, this becomes

$$\begin{aligned} I_{a \neq b} &= \frac{\cos\left((b - a)\frac{\tau}{2}\right)}{\tau} \int_{-\infty}^{+\infty} g\left(\frac{x}{\tau} - a\right) \frac{\sin^2(x/2)}{x} dx \\ &\quad + \frac{\sin\left((b - a)\frac{\tau}{2}\right)}{2\tau} \int_{-\infty}^{+\infty} g\left(\frac{x}{\tau} - a\right) \frac{\sin(x)}{x} dx, \quad (\text{F2}) \end{aligned}$$

where for large τ , the dominant integral contribution evaluates the (smooth) function $g(\omega)$ near $g(-a)$ such that this value can be taken out of the integral and we obtain $\lim_{\tau \rightarrow \infty} I_{a \neq b} = 0$. Using the representation

$$\lim_{\tau \rightarrow \infty} \tau \operatorname{sinc}^2(\Omega\tau) = \pi\delta(\Omega) \quad (\text{F3})$$

one can consider the general case via

$$f(\omega, a, b) \asymp \delta_{ab} \lim_{\tau \rightarrow \infty} \tau \operatorname{sinc}^2\left((\omega + a)\frac{\tau}{2}\right) = \delta_{ab} \pi\delta\left(\frac{\omega + a}{2}\right), \quad (\text{F4})$$

which yields Eq. (27).

-
- [1] P. W. Shor, *SIAM J. Comput.* **26**, 1484 (1997).
[2] L. K. Grover, *Phys. Rev. Lett.* **79**, 325 (1997).
[3] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
[4] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
[5] V. Khidekel, *Phys. Rev. E* **52**, 2510 (1995).
[6] A. M. Childs, E. Farhi, and J. Preskill, *Phys. Rev. A* **65**, 012322 (2001).
[7] E. Farhi *et al.*, *Science* **292**, 472 (2001).
[8] M. S. Sarandy, L.-A. Wu, and D. A. Lidar, *Quantum Inf. Process.* **3**, 331 (2004).
[9] S. Jansen, M. B. Ruskai, and R. Seiler, *J. Math. Phys.* **48**, 102111 (2007).
[10] G. Schaller, S. Mostame, and R. Schützhold, *Phys. Rev. A* **73**, 062307 (2006).
[11] M. H. S. Amin, P. J. Love, and C. J. S. Truncik, *Phys. Rev. Lett.* **100**, 060503 (2008).
[12] Y. Zhao and G. H. Chen, *Phys. Rev. E* **65**, 056120 (2002).
[13] S. Stenholm and M. Jakob, *J. Mod. Opt.* **51**, 841 (2004).
[14] P. Pechukas, *Phys. Rev. Lett.* **73**, 1060 (1994); R. Alicki, *ibid.* **75**, 3020 (1995); P. Pechukas, *ibid.* **75**, 3021 (1995).
[15] T. Yu, L. Diosi, N. Gisin, and W. T. Strunz, *Phys. Lett. A* **265**, 331 (2000).
[16] G. Lindblad, *Commun. Math. Phys.* **48**, 119 (1976).
[17] W. J. Munro and C. W. Gardiner, *Phys. Rev. A* **53**, 2633 (1996).
[18] J. Wilkie, *Phys. Rev. E* **62**, 8808 (2000).
[19] A. Tameshtit and J. E. Sipe, *Phys. Rev. Lett.* **77**, 2600 (1996).

- [20] R. J. C. Spreeuw and T. W. Hijmans, *Phys. Rev. A* **76**, 022306 (2007).
- [21] R. Karrlein and H. Grabert, *Phys. Rev. E* **55**, 153 (1997).
- [22] The general case $\langle B_A \rangle \neq 0$ can be reduced to the case $\langle B'_A \rangle = 0$ by using the transformations $H_S \rightarrow H'_S = H_S + \lambda g_A \sum_A A_A$ and $B_A \rightarrow B'_A = B_A - g_A \mathbf{1}$ in the Schrödinger picture, where the real constant must be adapted such that $g_A = \langle B_A \rangle$.
- [23] B. Vacchini, *Phys. Rev. Lett.* **84**, 1374 (2000).
- [24] D. A. Lidar, Z. Bihary, and K. B. Whaley, *Chem. Phys.* **268**, 35 (2001).
- [25] U. Kleinekathöfer, *J. Chem. Phys.* **121**, 2505 (2004).
- [26] M. Schröder, M. Schreiber, and U. Kleinekathöfer, *J. Chem. Phys.* **126**, 114102 (2007).
- [27] R. S. Whitney, *J. Phys. A: Math. Theor.* **41**, 175304 (2008).
- [28] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 2002).
- [29] E. Anderson *et al.*, *LAPACK Users' Guide*, SIAM, 1999, <http://www.netlib.org/lapack>.
- [30] W. H. Press *et al.*, *Numerical Recipes in C* (Cambridge University Press, Cambridge, 1994).
- [31] T. Brandes and T. Vorrath, *Phys. Rev. B* **66**, 075341 (2002); T. Brandes, *Phys. Rep.* **408**, 315 (2005).
- [32] F. Nesi, E. Paladino, M. Thorwart, and M. Grifoni, *Phys. Rev. B* **76**, 155323 (2007); F. Nesi *et al.*, *Europhys. Lett.* **80**, 40005 (2007).
- [33] W. G. Unruh, *Phys. Rev. A* **51**, 992 (1995).
- [34] G. Burkard, e-print arXiv:0803.0564v1.
- [35] R. Schützhold and M. Tiersch, *J. Opt. B: Quantum Semiclassical Opt.* **7**, S120 (2005).
- [36] M. C. Banuls, R. Orus, J. I. Latorre, A. Perez, and P. Ruiz-Femenia, *Phys. Rev. A* **73**, 022344 (2006).
- [37] R. Schützhold and G. Schaller, *Phys. Rev. A* **74**, 060304(R) (2006).
- [38] R. Alicki and K. Lendi, *Quantum Dynamical Semi-Groups and Applications*, Lecture Notes in Physics No. 286 (Springer-Verlag, Berlin, 1987).
- [39] C.-P. Sun, *J. Phys. A* **21**, 1595 (1988).