

Noise-induced interference fringes in trapped ultracold bosonic gases

S. Anderloni,^{1,2} F. Benatti,^{1,2} R. Floreanini,² and G. G. Guerreschi¹

¹*Dipartimento di Fisica Teorica, Università di Trieste, 34014 Trieste, Italy*

²*Istituto Nazionale di Fisica Nucleare, Sezione di Trieste, 34014 Trieste, Italy*

(Received 12 August 2008; published 21 November 2008)

We study the dynamics of ultracold atoms trapped in optical double-well potentials in presence of noise generated by an external environment. When prepared in a Fock number state, the system shows phase coherence in the averaged density profile obtained using standard absorption image techniques, that disappears in absence of noise. This effect indicates that also in many-body systems an external environment may enhance quantum coherence, instead of destroying it.

DOI: [10.1103/PhysRevA.78.052118](https://doi.org/10.1103/PhysRevA.78.052118)

PACS number(s): 03.65.Yz, 03.75.Gg

I. INTRODUCTION

Cold atoms trapped in a periodic optical potential have become a preferred test ground for the study of quantum effects in many-body physics, e.g., in quantum phase transition, and matter interference phenomena, and also for applications in quantum information (for instance, see Refs. [1,2], and references therein).

In particular, the study of phase coherence between fractions of a same condensate confined in a double well potential has been the focus of much theoretical and experimental analysis as a powerful technique to obtain information about the state of a trapped condensate. In general, phase coherence cannot be directly observed within the trap; nevertheless, it can be put into evidence by releasing the confining potential and letting the condensate freely expand, so that the atoms from the two wells may interfere with each other: as a consequence, fringe patterns are formed in the spatial density. The latter can be obtained by illuminating the expanding clouds with a probe light and then by collecting the corresponding absorption image. In particular, information about the initial relative phase can be extracted from the interference pattern [3–13].

One expects to observe interference fringes in the density profile only if there is phase coherence between the wells. Instead, they appear even starting from a state with random relative phase, such as a Fock number state.¹ In this case, however, the interference patterns have different offset positions in successive shots; as a consequence, by averaging over multiple realizations of the experiment the interference pattern disappears.

In the following, we shall study how this picture changes in the presence of an external environment, weakly affecting the dynamics of the condensate in the double-well trap. Usually, the environment is a source of noise and dissipation that are expected to lead to decoherence. Instead, we shall show that, the environment may create phase coherence between two initially uncorrelated fractions of the condensate, an ef-

fect that may be traced to the presence of an environment induced current between the two wells [14].

We will consider a system consisting of N atoms confined in a double-well trap with a very high barrier; in absence of noise, the Fock number states with k atoms in one well and $N-k$ in the other are equilibrium states. The presence of an environment changes the situation and makes the system an open quantum system; its dynamics is no longer unitary and is characterized by the presence of noise and dissipation, so that the Fock number states are no longer stationary. We will assume that the interaction between the atoms and the environment be weak, with very fast decaying environment correlations. In this case, the reduced dynamics of the trapped gas admits a Markovian (i.e., memoryless) time evolution described by a family of maps forming a quantum dynamical semigroup [15–17]. An explicit example of such an environment has been proposed in Ref. [18], where cold bosons in an optical lattice are coupled to the Bogoliubov excitations of another condensate, which mediates transitions between the lowest and the first excited lattice Bloch bands. Alternatively, one may consider the coupling of the atoms to an external classical stochastic field, with white noise correlations.

Using open quantum system techniques, we shall analyze below the averaged density resulting from the free expansion of the gas after trap release. We shall see that, even starting from a Fock number state, with no definite relative phase between the states in the two wells, interference fringes may appear in the averaged density profile; they disappear in absence of the environment, thus showing the purely noisy origin of the effect. We shall also see that this phenomenon may be related to the presence of a dissipative current between the two wells [14].

II. COLD ATOMS IN A DOUBLE-WELL POTENTIAL

The usual description of cold atoms trapped in a double well potential uses a two-mode Bose-Hubbard model [20,21] with Hamiltonian

$$H_{\text{BH}} = \varepsilon_1 a_1^\dagger a_1 + \varepsilon_2 a_2^\dagger a_2 + U[(a_1^\dagger a_1)^2 + (a_2^\dagger a_2)^2] - T(a_1^\dagger a_2 + a_1 a_2^\dagger), \quad (1)$$

where $a_{1,2}$, $a_{1,2}^\dagger$ annihilate and create atom states in the first,

¹This phenomenon has been the focus of various investigations, aiming to explain single shot interference formation via the detection process (see Refs. [6–13], and references therein). An alternate explanation is, however, possible: as further discussed in Sec. III, it makes use of a projection onto fixed-phase, coherentlike states.

second well, respectively, and satisfy the Bose commutation relations $[a_i, a_j^\dagger] = \delta_{ij}$. Of the contributions to H_{BH} , the last one corresponds to a hopping term depending on the tunneling amplitude T , the first two are due to the trapping potential and are proportional to the depth of the wells: in the following we will consider the case of a symmetric trap, i.e., $\varepsilon_1 = \varepsilon_2$. Finally, the third term, quadratic in the number operators $a_i^\dagger a_i$, takes into account repulsive Coulomb interactions inside each well.

Let $\{|\phi_i\rangle\}_{i=1}^\infty$ be a complete set of orthonormal single-particle atom states; the creation operator $\hat{\psi}^\dagger(x)$ of an atom at position x can then be decomposed as

$$\hat{\psi}^\dagger(x) = \sum_{i=1}^{\infty} \phi_i^*(x) a_i^\dagger, \quad (2)$$

where a_i^\dagger creates an atom in the state $|\phi_i\rangle = a_i^\dagger |\text{vac}\rangle$ with wave function $\phi_i(x) = \langle x | \phi_i \rangle$. The Bose-Hubbard Hamiltonian (1) results from a tight binding approximation, where only the first two of the basis vector are relevant; in this case $\phi_{1,2}(x)$ are orthogonal functions, $\phi_1(x)$ localized within the first well, $\phi_2(x)$ within the second one.

The total number N of atoms is conserved by Eq. (1). Therefore, the Hilbert space of the system is $(N+1)$ -dimensional and can be spanned by Fock number states of the form

$$|k, N-k\rangle := \frac{(a_1^\dagger)^k (a_2^\dagger)^{N-k}}{\sqrt{k!} \sqrt{(N-k)!}} |\text{vac}\rangle, \quad (3)$$

describing the situation in which the first well is filled with k atoms, while the other one contains $N-k$ particles. They are obtained by the action of the creation operators on the vacuum state. These states turn out to be eigenstates of the Bose-Hubbard Hamiltonian when the tunneling term can be neglected.

Alternatively, one can consider the so-called phase states

$$|\varphi, \xi; N\rangle = \frac{1}{\sqrt{N!}} (\sqrt{\xi} e^{i(\varphi/2)} a_1^\dagger + \sqrt{1-\xi} e^{-i(\varphi/2)} a_2^\dagger)^N |\text{vac}\rangle, \quad (4)$$

depending on two real parameters, an amplitude $\xi \in [0, 1]$ and a phase $\varphi \in [0, 2\pi]$. These states describe a physical situation where all N atoms are in a same superposition of a two-mode single particle state with relative phase φ ; from a many-body point of view, such states correspond to a macroscopic wave function obeying the Gross-Pitaevski equation [3]. Moreover, the parameter ξ is related to the mean value of the relative occupation number $\langle \varphi, \xi; N | (a_1^\dagger a_1 - a_2^\dagger a_2) | \varphi, \xi; N \rangle = N(2\xi - 1)$.

These states form an overcomplete set

$$\int_N = \frac{N+1}{2\pi} \int_0^1 d\xi \int_0^{2\pi} d\varphi |\varphi, \xi; N\rangle \langle \varphi, \xi; N|. \quad (5)$$

Although not orthogonal, for sufficiently large N (a very common situation in actual experiments), the phase states become nearly so,

$$\langle \varphi, \xi; N | \varphi', \xi'; N \rangle \approx \frac{1}{N} \delta(\xi - \xi') \delta(\varphi - \varphi'). \quad (6)$$

One can then decompose any N atom state in terms of these states; in particular, for Fock number states, one can write

$$|k, N-k\rangle = \frac{N+1}{2\pi} \int_0^1 d\xi \int_0^{2\pi} d\varphi \langle \varphi, \xi; N | k, N-k \rangle |\varphi, \xi; N\rangle, \quad (7)$$

where the overlapping functions are explicitly given by

$$\langle \varphi, \xi; N | k, N-k \rangle = \binom{N}{k}^{1/2} \xi^{k/2} (1-\xi)^{(N-k)/2} e^{-i\varphi(k-N/2)}. \quad (8)$$

Further, the action of $\hat{\psi}(x)$ on the phase states can be easily derived using Eq. (2),

$$\hat{\psi}(x) |\varphi, \xi; N\rangle = \sqrt{N} [\sqrt{\xi} \phi_1(x) e^{i(\varphi/2)} + \sqrt{1-\xi} \phi_2(x) e^{-i(\varphi/2)}] \times |\varphi, \xi; N-1\rangle. \quad (9)$$

III. DENSITY PROFILE AFTER FREE EXPANSION

As explained in the Introduction, our aim is to study the spatial density profile obtained by releasing the trap and by subsequently letting the cold atoms freely expand for a time τ . If τ is large enough, the two fractions of cold atoms interfere with each other thus providing a spatial fringe pattern that one can probe by light absorption.

Suppose the trap is released at $\tau=0$ when the cold atoms are in the phase state (4). In order to describe the free expansion, disregarding atom-atom interactions, we make use of the following model: we let the single-particle states evolve under the action of an unitary operator U_τ ; notice that, by the unitarity of U_τ the basis states $|\phi_i\rangle$ evolve in another set of basis states

$$|\phi_i\rangle_\tau = U_\tau |\phi_i\rangle := a_i^\dagger(\tau) |\text{vac}\rangle, \quad a_i^\dagger(\tau) \equiv U_\tau a_i^\dagger U_\tau^\dagger. \quad (10)$$

Thanks to the absence of interactions during the expansion, the evolution of the phase states follows immediately from the evolution of the single particle states

$$|\varphi, \xi; N\rangle_\tau = \frac{1}{\sqrt{N!}} [\sqrt{\xi} e^{i(\varphi/2)} a_1^\dagger(\tau) + \sqrt{1-\xi} e^{-i(\varphi/2)} a_2^\dagger(\tau)]^N |\text{vac}\rangle, \quad (11)$$

while both the relations (5) and (6) still hold for any τ [22]. The annihilation operator (which does not evolve in time) could be again decomposed over the new basis

$$\hat{\psi}(x) = \sum_{i=0}^{\infty} \phi_i(x, \tau) a_i(\tau), \quad \phi_i(x, \tau) = \langle x | \phi_i \rangle_\tau, \quad (12)$$

and therefore the action on the freely evolved phase state reads

$$\hat{\psi}(x) |\varphi, \xi; N\rangle_\tau = \sqrt{N} [\sqrt{\xi} \phi_1(x, \tau) e^{i(\varphi/2)} + \sqrt{1-\xi} \phi_2(x, \tau) e^{-i(\varphi/2)}] \times |\varphi, \xi; N-1\rangle_\tau. \quad (13)$$

If the two fractions of the condensate expand independently, we can explicitly evaluate the relative phase gained during the ballistic expansion. Writing $\phi_i(x, \tau) = |\phi_i(x, \tau)| e^{i\theta_i(x, \tau)}$, the phase $\theta_i(x, \tau)$ can be obtained, for large τ , in terms of the classical behavior of the velocity $\vec{v} = \frac{\hbar}{m} \vec{\nabla} \theta_i = \frac{\vec{r}}{\tau}$ [3]. We shall focus upon the direction x , hence upon the phase modulation

$$\theta_{1,2}(x, \tau) = \frac{1}{2} \frac{m(x - x_{1,2})^2}{\hbar \tau},$$

where x_i is the position of the i th well; it follows that the overall relative phase after a time of flight τ can be well approximated by

$$\theta_1(x, \tau) - \theta_2(x, \tau) = \frac{md}{\hbar \tau} x, \quad (14)$$

where the trap is considered to be located along the x axis, centered at $x=0$ with the two wells at position $x_1 = -d/2$, $x_2 = d/2$, respectively.

Then, starting from a phase state, the one-particle spatial density explicitly reads

$$\begin{aligned} n_{\varphi, \xi}(x, \tau) &= {}_{\tau} \langle \varphi, \xi; N | \hat{\psi}^\dagger(x) \hat{\psi}(x) | \varphi, \xi; N \rangle_{\tau} \\ &= N \left[\xi |\phi_1(x, \tau)|^2 + (1 - \xi) |\phi_2(x, \tau)|^2 \right. \\ &\quad \left. + 2\sqrt{\xi(1 - \xi)} |\phi_1(x, \tau)| |\phi_2(x, \tau)| \cos\left(\frac{md}{\hbar \tau} x + \varphi\right) \right]. \end{aligned} \quad (15)$$

The last term in this equation exhibits the presence of fringes that are equally separated by a distance $\ell \propto \frac{\hbar \tau}{md}$ with an offset position fixed by the initial phase φ .

The situation is completely different if one starts instead from a Fock number state; in this case the mean value of the density operator after the free expansion can be calculated by decomposing the state using the analog² of Eq. (7) at time τ ,

$$|k, N - k\rangle_{\tau} = \frac{N + 1}{2\pi} \int_0^1 d\xi \int_0^{2\pi} d\varphi \langle \varphi, \xi; N | k, N - k \rangle | \varphi, \xi; N \rangle_{\tau}. \quad (16)$$

Experimentally, the light-absorption image relative to a Fock number state shows the same fringe pattern (15) of the superfluid state, whereas one would expect no interference fringes at all. However, if one repeats the experiment with the same initial conditions, the fringe patterns are obtained with randomly distributed offset positions. Therefore, by averaging over them, the fringes cancel out and the averaged density profile does not exhibit any interference pattern [3–13].

In order to interpret this effect, notice that the absorption image amounts to a measurement of the initial relative phase φ , for it gives a one-shot density profile of the form (15).

Furthermore, starting from a phase-state $|\varphi, \xi; N\rangle$, every single repetition of the experiment will always give the same interference pattern $n_{\varphi, \xi}(x, \tau)$ [11]. On the contrary, when the initial state at $\tau=0$ is a Fock number state $|k, N - k\rangle$, each single shot will correspond to a random selection of one of the phase states (11) in the expansion (16), with probability densities that are independent of the relative phase and proportional to

$$|\langle \varphi, \xi; N | k, N - k \rangle|^2 = \binom{N}{k} \xi^k (1 - \xi)^{N - k}. \quad (17)$$

In other words, every single-shot measurement of the density profile in the state $|k, N - k\rangle_{\tau}$ resulting from the Fock number state $|k, N - k\rangle$, randomly projects out the density profile of one of the phase states $|\varphi, \xi; N\rangle_{\tau}$ with weight (17). As a consequence, the averaged density profile reads

$$\begin{aligned} n_k(x, \tau) &:= \frac{N + 1}{2\pi} \int_0^1 d\xi \int_0^{2\pi} d\varphi |\langle \varphi, \xi; N | k, N - k \rangle|^2 n_{\varphi, \xi}(x, \tau) \\ &= \frac{N}{N + 2} [(k + 1) |\phi_1(x, \tau)|^2 + (N - k + 1) |\phi_2(x, \tau)|^2] \end{aligned} \quad (18)$$

and shows no interference pattern (see Ref. [22] for further details).

The density profile at time τ after trap release depends on the state of the N -atom system at time $\tau=0$. Since we are interested in studying the effects of a dissipative open dynamics during the interval of time t spent by the system within the confining double-well potential, we will describe the system state at time t by a generic density matrix ρ_t :

$$\rho_t = \sum_{k, q=0}^N R_{kq}^t |k, N - k\rangle \langle q, N - q|. \quad (19)$$

Notice that the information about the dynamics during the time t spent by the atoms within the double-well trap is contained in the coefficients R_{kq}^t .

In order to construct the averaged density profile in such a case, one considers the ensemble of profiles $n_{\varphi, \xi}(x, \tau) = {}_{\tau} \langle \varphi, \xi; N | \hat{\psi}^\dagger(x) \hat{\psi}(x) | \varphi, \xi; N \rangle_{\tau}$ that are obtained in many repetitions of the experiment, with the same initial conditions. Each image shows interference fringes of the form (15) and appears in the statistical ensemble of all collected images with weights proportional to

$$\begin{aligned} \langle \varphi, \xi; N | \rho_t | \varphi, \xi; N \rangle &= \sum_{k, q=0}^N R_{kq}^t \sqrt{\binom{N}{k} \binom{N}{q}} \\ &\quad \times \xi^{(k+q)/2} (1 - \xi)^{N - (k+q)/2} e^{-i\varphi(k-q)}. \end{aligned} \quad (20)$$

The resulting one-particle spatial density profile is then given by the corresponding statistical average

²Note that the scalar products $\langle \varphi, \xi; N | k, N - k \rangle$ in Eq. (16) do not depend on time.

$$\begin{aligned}
n_{\rho_t}(x, \tau) &= \frac{N+1}{2\pi} \int_0^1 d\xi \int_0^{2\pi} d\varphi \langle \varphi, \xi | \rho_t | \varphi, \xi \rangle n_{\varphi, \xi}(x, \tau) \\
&= \frac{N}{N+2} \left\{ \sum_{k=0}^N R_{kk}^t [(k+1) |\phi_1(x, \tau)|^2 + (N-k+1) \right. \\
&\quad \times |\phi_2(x, \tau)|^2] + \sum_{k=0}^{N-1} R_{kk+1}^t \sqrt{(k+1)(N-k)} |\phi_1(x, \tau)| \\
&\quad \left. \times |\phi_2(x, \tau)| e^{-i(md/\hbar)\tau x} + \text{c.c.} \right\}. \quad (21)
\end{aligned}$$

The presence of visible interference fringes is strictly related to the next-to-diagonal entries of the density matrix R_{kk+1}^t (and their Hermitian conjugates).³

Notice that the previous average can be computed as the mean value of the density operator $\hat{\psi}^\dagger(x)\hat{\psi}(x)$ with respect to a density matrix $\tilde{\rho}_t$ which results from the freely evolved state $U_{\tau}\rho_t U_{\tau}^\dagger$ by the action of a generalized measurement, a so-called positive operator valued measure (POVM) [23,24], consisting of the family of projectors $P_{\varphi, \xi; N} := |\varphi, \xi; N\rangle\langle\varphi, \xi; N|$. Explicitly, one has

$$\tilde{\rho}_t = \frac{N+1}{2\pi} \int_0^1 d\xi \int_0^{2\pi} d\varphi P_{\varphi, \xi; N} U_{\tau}\rho_t U_{\tau}^\dagger P_{\varphi, \xi; N}. \quad (22)$$

The map from ρ_t to $\tilde{\rho}_t$ corresponds to the canonical description of the quantum measurement process associated with the POVM $\{P_{\varphi, \xi; N}\}$. One notices that it does not correspond to a measurement of the number operator in the state $U_{\tau}\rho_t U_{\tau}^\dagger$ which gives an average number of the form

$$\begin{aligned}
&\text{Tr}[\hat{\psi}^\dagger(x)\hat{\psi}(x)U_{\tau}\rho_t U_{\tau}^\dagger] \\
&= N \int dx_2 \cdots dx_N \langle x, x_2, \dots, x_N | U_{\tau}\rho_t U_{\tau}^\dagger | x, x_2, \dots, x_N \rangle \\
&= \sum_{k=0}^N R_{kk}^t [k |\phi_1(x, \tau)|^2 + (N-k) |\phi_2(x, \tau)|^2] \\
&\quad + \left\{ \sum_{k=0}^{N-1} R_{kk+1}^t \sqrt{(k+1)(N-k)} |\phi_1(x, \tau)| \right. \\
&\quad \left. \times |\phi_2(x, \tau)| e^{-i(md/\hbar)\tau x} + \text{c.c.} \right\}, \quad (23)
\end{aligned}$$

which is often used to fit experimental data; this average corresponds to a POVM consisting of projections onto eigenstates of the operator $\hat{\psi}^\dagger(x)\hat{\psi}(x)$, not onto phase states.

For N large, the difference between Eqs. (23) and (21) are very small for all practical purposes. Furthermore, the difference does not show up in the oscillating terms. In the fol-

lowing we will make use of the the description of the absorption images in terms of a POVM consisting of the projections onto the phase states, since this appears to describe more adequately the absorption image formation [22].

IV. NOISY DYNAMICS WITHIN THE TRAP

In this section we study the dynamics of the N atoms within a trap immersed in an external environment; in particular, we shall focus upon its effects on the ground state $|N/2, N/2\rangle$ of Eq. (1) with a very high potential barrier. The density matrix of the total system, atoms plus environment, ρ_{SE} evolves unitarily under the action of a total Hamiltonian which can be written in the form

$$H_{SE} = H_{\text{BH}} \otimes 1_E + 1 \otimes H_E + H_I, \quad (24)$$

where H_{BH} is as in Eq. (1) and describes the motion of the system in absence of the environment, H_E is the Hamiltonian of the environment, while H_I describes the interaction between them and can be taken to have the general form

$$H_I = \sum_i V_i \otimes B_i \quad (25)$$

where V_i and B_i are suitable system and environment hermitian operators, respectively.

The evolution of the system density matrix $\rho = \text{Tr}_E(\rho_{SE})$ can be obtained by tracing out the environment degrees of freedom from the standard unitary evolution of the total system initial state ρ_{SE} . In general this operation leads to a master equation affected by memory terms and nonlinearities. The latter can be avoided by choosing an initially uncorrelated state of the system and the bath, $\rho_{SE} = \rho \otimes \rho_E$; this is a common situation in experimental contexts when the environment is supposed to be in a reference equilibrium state ρ_E .

On the other hand, memory effects can be considered a short time phenomenon; if we are not interested in this transient regime, they can be neglected by studying the effective evolution of the system on a slower timescale. This amounts to a Markov approximation [15–17] which is only plausible when a neat separation exists between system and environment time scales. Necessary conditions for this approximation are, for instance, satisfied within the so-called singular coupling limit [25–27], where the characteristic decay times of the environment correlations

$$G_{ij}(t) = \langle B_i(t)B_j \rangle \equiv \text{Tr}[B_i(t)B_j\rho_E], \quad B_i(t) = e^{iH_E t} B_i e^{-iH_E t} \quad (26)$$

are very small when compared to the typical time scale of the system. This condition is satisfied by a thermal bath in equilibrium at very high temperature or by an external stochastic white-noise classical field; in this latter case, the matrix $[G_{ij}(t)]$ turn out to be real symmetric. In Ref. [18], a suitable bath fulfilling the above conditions has been proposed, where the system is coupled with the Bogoliubov excitations of another condensate, which drive a transition of the trapped

³The evolution of the atoms within the trap affects only the coefficient R_{kk}^t and depends on the time t spent inside the trap; the time of flight τ refers instead to the time interval between the releasing of the trap and the snapshot of the density and it is a parameter of the experimental setup.

atoms to the first excited lattice Bloch band, with a characteristic transition rate κ .⁴

In the singular-coupling limit, the dissipative time-evolution of the open system density matrix is generated by a master equation of the form

$$\frac{\partial \rho(t)}{\partial t} = \mathbb{L}[\rho(t)] \equiv -i[H_{BH} + H^{(2)}, \rho(t)] + \mathbb{D}[\rho(t)], \quad (27)$$

where the noise contributes with an effective correction $H^{(2)}$ to the free Hamiltonian and with a linear operator $\mathbb{D}[\cdot]$ which cannot be recast in Hamiltonian form. Their explicit expressions take the standard Kossakowski-Lindblad form [15–17]

$$\mathbb{D}[\rho] = \sum_{ij=1}^4 c_{ij} \left[V_j^\dagger \rho V_i - \frac{1}{2} \{V_i V_j^\dagger, \rho\} \right], \quad (28)$$

$$H^{(2)} = \sum_{ij=1}^4 s_{ij} V_i V_j^\dagger. \quad (29)$$

The operators V_i are the operators involved in the interaction Hamiltonian (25), while the entries of the two Hermitian matrices $[c_{ij}]$ and $[s_{ij}]$ embody all the information about the effects induced by the environment and are obtained from the correlation functions of the environment

$$\int_0^\infty dt G_{ij}(t) = \frac{1}{2} c_{ij} + i s_{ij}, \quad (30)$$

$$c_{ij} = \int_{-\infty}^{+\infty} dt G_{ij}(t) \equiv \mathcal{F}G_{ij}(\omega)|_{\omega=0},$$

$$s_{ij} = \frac{1}{2\pi} \mathcal{P} \int_{-\infty}^{+\infty} d\omega \frac{\mathcal{F}G_{ij}(\omega)}{\omega}, \quad (31)$$

where \mathcal{F} denotes Fourier transforming and \mathcal{P} the principal value.

In particular $[c_{ij}]$ is known as Kossakowski matrix and turns out to be positive definite, a key requirement to ensure the positivity of the eigenvalues of the evolved density matrix ρ_t at all times and therefore the consistency of the dynamics. For instance, in the case of the environment proposed in Ref. [18], the matrix $[c_{ij}]$ takes the explicit expression⁵

⁴It is interesting to notice that another possible mechanism inducing dissipation is provided by inelastic collisions among atoms on the same lattice site, that can be tuned on by a nonvanishing imaginary part of the interaction parameter U [19]. These effects would nevertheless lead to atom loss, i.e., to nonconservation of N , requiring an extension of the formalism to be properly treated.

⁵The dissipative effects induced by inelastic atom collision discussed in Ref. [19] can similarly be described in terms of a master equation of the form (27); the magnitude of the elements of the corresponding Kossakowski matrix is determined by $\text{Im}(U)$.

$$[c_{ij}] = \begin{pmatrix} 2\kappa & -2\kappa & 0 & i\kappa \\ -2\kappa & 2\kappa & 0 & -i\kappa \\ 0 & 0 & 0 & 0 \\ -i\kappa & i\kappa & 0 & \kappa/2 \end{pmatrix}. \quad (32)$$

The operator $\mathbb{L}[\cdot]$ in Eq. (27) generates a family of maps $\gamma_t = e^{t\mathbb{L}}$, which forms a quantum dynamical semigroup [15–17], fulfilling the forward in time composition law $\gamma_t \circ \gamma_s = \gamma_{t+s}$, $t, s \geq 0$, and describing the irreversible character of the dissipative evolution.

Quite in general, the presence of an environment will affect both the tunneling amplitude and the minima of the confining potential; the environment operators B_i will then be coupled to the following bilinear atom operators appearing in the system Hamiltonian (1) $a_1^\dagger a_1$, $a_2^\dagger a_2$, $a_1^\dagger a_2$, and $a_2^\dagger a_1$, or more precisely to the four Hermitian operators

$$V_i = \{a_1^\dagger a_1, a_2^\dagger a_2, (a_1^\dagger a_2 + a_2^\dagger a_1), i(a_1^\dagger a_2 - a_2^\dagger a_1)\}. \quad (33)$$

As a final remark, notice that the action of the environment ceases as a consequence of the trap release.

V. EFFECTS OF NOISE IN THE DENSITY PROFILE

As mentioned before, we shall consider an initial atom state of the form

$$\rho_0 = |N/2, N/2\rangle\langle N/2, N/2|. \quad (34)$$

In absence of the environment and for a high potential barrier, this state is the ground state of Eq. (1); as a consequence, the components R_{kq}^t in Eq. (19) are time independent and $R_{kq} = 0$ if k and q differ from $N/2$. Therefore in both Eqs. (21) and (23) no interference fringes appear.

In order to study the effects induced by the presence of noise, it is sufficient to consider the small time dynamics resulting from Eq. (27), whose solution up to first order in time then reads

$$\rho_t = \rho_0 + it[H_{BH} + H^{(2)}, \rho_0] + t\mathbb{D}[\rho_0] + \mathcal{O}(t^2). \quad (35)$$

Since $[H_{BH}, \rho_0] = 0$, only the dissipative contributions in Eq. (35) need to be studied. The explicit expressions of $\mathbb{D}[\rho_0]$ and $i[H^{(2)}, \rho_0]$ are collected in the Appendix. Using them, to first order in t , one finally gets

$$\begin{aligned} n_{\rho_t}(x, \tau) = & \frac{N}{2} [|\phi_1(x, \tau)|^2 + |\phi_2(x, \tau)|^2] + t \frac{N^2}{4} \left\{ 4 \text{Im}(c_{34}) \right. \\ & \times [|\phi_2(x, \tau)|^2 - |\phi_1(x, \tau)|^2] + |\phi_1(x, \tau)| |\phi_2(x, \tau)| \\ & \times \left[2 \text{Re}(s_{24} - s_{14}) + \text{Im}(c_{14} - c_{24}) \right] \cos\left(\frac{md}{\hbar\tau} x\right) \\ & \left. + [2 \text{Re}(s_{23} - s_{13}) + \text{Im}(c_{13} - c_{23})] \sin\left(\frac{md}{\hbar\tau} x\right) \right\}. \end{aligned} \quad (36)$$

The last two lines of the last equation contain oscillating terms which give rise to interference fringes spaced by a distance $l \propto \hbar\tau/md$; notice that these fringes disappear in absence of noise (i.e., when $c_{ij} = 0 = s_{ij}$). While the interference

pattern is the same as in Eq. (15) and is essentially due to the ballistic expansion, the amplitude of these oscillations is proportional to the entries of the matrices $[c_{ij}]$ and $[s_{ij}]$, which depend on the strength of the environment correlation functions.

Since for large enough τ the atomic clouds coming from the two wells overlap, i.e., $|\phi_1(x, \tau)| \approx |\phi_2(x, \tau)|$, the magnitude of the dissipative terms goes as $tN|c_{ij}|$ with respect to the standard, noise independent contribution. Despite the large number N of atoms, because the system environment coupling is weak and thus the constants c_{ij} and s_{ij} are small [25–27], the first-order expansion in Eq. (35) is meaningful even for sufficiently long system environment interaction times t .

For instance, if the Kossakowski matrix $[c_{ij}]$ is as in Eq. (32), the magnitude of its entries is proportional to the transition rate κ ; for typical values of this parameter and of N , t can be large enough to allow a direct experimental observation of the interference fringes. Finally, notice that if instead the system is coupled to a stochastic classical field, the matrix $[c_{ij}]$ is real symmetric while all the entries of $[s_{ij}]$ vanishes [25]; therefore, the oscillating terms in Eq. (36) disappear; this means that classical correlations do not lead to any environment induced interference phenomena.

VI. RELATIONS WITH DISSIPATIVE CURRENT

To better understand the physical origin of the environment induced interference pattern, let us consider the mean value of the following current operator [14]:

$$\hat{J} = i(a_1^\dagger a_2 - a_2^\dagger a_1), \quad (37)$$

which describes the motion of the barycenter of the atom gas in the double well. Its mean value is clearly zero in the initial state ρ_0 , while, in the evolved state up to first order in t , it becomes

$$\langle \hat{J} \rangle_{\rho_t} = N \left(\frac{N}{2} + 1 \right) [\text{Im}(c_{23} - c_{13}) + 2 \text{Re}(s_{13} - s_{23})]. \quad (38)$$

By comparing the last expression with Eq. (36) one notices that, if the current operator has nonzero mean value, then interference fringes can be observed; however the structure of the interference pattern is in general richer, as also the tunneling operator $a_1^\dagger a_2 + a_2^\dagger a_1$ contributes to it. Indeed, one finds

$$\begin{aligned} \mathbb{D}[\rho_0] = & \frac{N}{2} \left(\frac{N}{2} + 1 \right) \left\{ [c_{33} + c_{44} + i(c_{34} - c_{34}^*)] \left| \frac{N}{2} + 1 \right\rangle \left\langle \frac{N}{2} + 1 \right| \right. \\ & + [c_{33} + c_{44} - i(c_{34} - c_{34}^*)] \left| \frac{N}{2} - 1 \right\rangle \left\langle \frac{N}{2} - 1 \right| - 2(c_{33} + c_{44}) \left| \frac{N}{2} \right\rangle \left\langle \frac{N}{2} \right| \Big\} \\ & + \left\{ \sqrt{\frac{N}{2} \left(\frac{N}{2} + 1 \right)} \left[\frac{N}{4} (c_{13}^* - ic_{14}^* + c_{23}^* - ic_{24}^*) - \frac{1}{2} (c_{13}^* - ic_{14}^* - c_{23}^* + ic_{24}^*) \right. \right. \\ & \left. \left. - \frac{N}{4} (c_{13} - ic_{14} + c_{23} - ic_{24}) \right] \right\} \left| \frac{N}{2} \right\rangle \left\langle \frac{N}{2} + 1 \right| \end{aligned}$$

$$\langle a_1^\dagger a_2 + a_2^\dagger a_1 \rangle_{\rho_t} = N \left(\frac{N}{2} + 1 \right) [\text{Im}(c_{14} - c_{24}) + 2 \text{Re}(s_{24} - s_{14})]. \quad (39)$$

This is precisely the situation that occurs for the Kossakowski matrix (32); it leads to a vanishing mean value of the current operator, while instead the mean value of the tunneling operator is proportional to κ .

VII. OUTLOOK

We have seen that, even in the ground state ρ_0 of the Hamiltonian (1), the presence of a noisy environment may give rise to interference effects in the averaged density profile. The periodic structure of these fringes comes from the ballistic evolution of the gas and does not depend on the noise; their oscillations amplitude are, however, directly proportional to the characteristic parameters of the noise and vanish in absence of it.

This phenomenon can be related to a dissipative current between the wells, so that the study of the interference fringes in the averaged density profile gives information about this current.

The system we analyzed can in principle be engineered in the laboratory, for instance, using a second condensed gas as an environment as proposed in Ref. [18]: the resulting interferences fringes can be detected by subtracting the density pattern in absence of noise. This effect can also be useful to further understand the role of noise in double well potentials and in particular for the possibility of driving the system out of the ground state ρ_0 without lowering the barrier.

Similar conclusions can be easily extended to the case of an optical lattice. In this situation the capability of suitably engineered environment to create correlations between different sites of the lattice could be practically very useful in order to create specific classes of correlated states.

ACKNOWLEDGMENTS

This work is supported by the MIUR project ‘‘Quantum Noise in Mesoscopic Systems.’’ S.A. acknowledges Eurotech SPA for financial support.

APPENDIX

By setting $|k\rangle \equiv |k, N-k\rangle$ for the sake of simplicity, the explicit expressions of $\mathbb{D}[\rho_0]$ and $i[H^{(2)}, \rho_0]$ read

$$\begin{aligned}
& + \sqrt{\frac{N}{2}\left(\frac{N}{2}+1\right)} \left[\frac{N}{4}(c_{13}^* + ic_{14}^* + c_{23}^* + ic_{24}^*) + \frac{1}{2}(c_{13}^* + ic_{14}^* - c_{23}^* - ic_{24}^*) - \frac{N}{4}(c_{13} + ic_{14} + c_{23} + ic_{24}) \right] \left| \frac{N}{2} \right\rangle \left\langle \frac{N}{2} - 1 \right| \\
& + \frac{1}{2} \sqrt{\left(\frac{N}{2}+2\right)\left(\frac{N}{2}+1\right)\left(\frac{N}{2}-1\right)} \frac{N}{2} (c_{44} - c_{33}) \left[\left| \frac{N}{2} \right\rangle \left\langle \frac{N}{2} + 2 \right| + \left| \frac{N}{2} \right\rangle \left\langle \frac{N}{2} - 2 \right| \right] \\
& + \frac{N}{2} \left(\frac{N}{2}+1\right) [c_{33} - c_{44} + i(c_{34} + c_{34}^*)] \left| \frac{N}{2} + 1 \right\rangle \left\langle \frac{N}{2} - 1 \right| + \text{H.c.} \Big\}, \tag{A1}
\end{aligned}$$

$$\begin{aligned}
i[H^{(2)}, \rho_0] = & i \sqrt{\frac{N}{2}\left(\frac{N}{2}+1\right)} \left\{ \left[\left(\frac{N}{2}+1\right) s_{13} + \frac{N}{2} s_{13}^* + i\left(\frac{N}{2}+1\right) s_{14} + i\frac{N}{2} s_{14}^* \right. \right. \\
& \left. \left. + \left(\frac{N}{2}-1\right) s_{23} + \frac{N}{2} s_{23}^* + i\left(\frac{N}{2}-1\right) s_{24} + \frac{N}{2} s_{24}^* \right] \left| \frac{N}{2} + 1 \right\rangle \left\langle \frac{N}{2} \right| \right. \\
& \left. + \left[\left(\frac{N}{2}-1\right) s_{13} + \frac{N}{2} s_{13}^* - i\left(\frac{N}{2}-1\right) s_{14} - i\frac{N}{2} s_{14}^* + \left(\frac{N}{2}+1\right) s_{23} + \frac{N}{2} s_{23}^* - i\left(\frac{N}{2}+1\right) s_{24} - \frac{N}{2} s_{24}^* \right] \left| \frac{N}{2} - 1 \right\rangle \left\langle \frac{N}{2} \right| \right. \\
& \left. + 2i \sqrt{\left(\frac{N}{2}+1\right)\left(\frac{N}{2}+2\right)} (s_{34} - s_{34}^*) \left| \frac{N}{2} + 2 \right\rangle \left\langle \frac{N}{2} \right| - 2i \sqrt{\left(\frac{N}{2}-1\right)\frac{N}{2}} (s_{34} - s_{34}^*) \left| \frac{N}{2} - 2 \right\rangle \left\langle \frac{N}{2} \right| \right\} + \text{H.c.} \tag{A2}
\end{aligned}$$

-
- [1] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen, and U. Sen, *Adv. Phys.* **56**, 243 (2007).
- [2] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [3] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Oxford University Press, Oxford, 2003).
- [4] C. J. Pethick and H. Smith, *Bose-Einstein Condensation in Dilute Gases* (Cambridge University Press, Cambridge, 2004).
- [5] S. Haroche and J.-M. Raimond, *Exploring the Quantum: Atoms, Cavities and Photons* (Oxford University Press, Oxford, 2006).
- [6] J. Javanainen and Sun Mi Yoo, *Phys. Rev. Lett.* **76**, 161 (1996).
- [7] M. Naraschewski, H. Wallis, A. Schenzle, J. I. Cirac, and P. Zoller, *Phys. Rev. A* **54**, 2185 (1996).
- [8] J. I. Cirac, C. W. Gardiner, M. Naraschewski, and P. Zoller, *Phys. Rev. A* **54**, R3714 (1996).
- [9] T. Wong, M. J. Collett, and D. F. Walls, *Phys. Rev. A* **54**, R3718 (1996).
- [10] Y. Castin and J. Dalibard, *Phys. Rev. A* **55**, 4330 (1997).
- [11] A. J. Leggett, *Rev. Mod. Phys.* **73**, 307 (2001).
- [12] W. J. Mullin, R. Krotkov, and F. Laloë, *Am. J. Phys.* **74**, 880 (2006).
- [13] A. Imambekov, V. Gritsev, and E. Demler, e-print arXiv:cond-mat/0703766.
- [14] S. Anderloni, F. Benatti, and R. Floreanini, *Open Syst. Inf. Dyn.* **15**, 143 (2008).
- [15] V. Gorini, A. Frigerio, M. Verri, A. Kossakowski, and E. C. G. Sudarshan, *Rep. Math. Phys.* **13**, 149 (1978).
- [16] R. Alicki and K. Lendi, *Quantum Dynamical Semigroups and Applications*, Vol. 717 of Lecture Notes in Physics (Springer-Verlag, Berlin, 2007).
- [17] F. Benatti and R. Floreanini, *Int. J. Mod. Phys. B* **19**, 3063 (2005).
- [18] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. P. Büchler, and P. Zoller, e-print arXiv:0803.1482.
- [19] N. Syassen, D. M. Bauer, M. Lettner, T. Volz, D. Dietze, J. J. Garcia-Ripoll, J. I. Cirac, G. Rempe, and S. Dürr, *Science* **320**, 1329 (2008).
- [20] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls, *Phys. Rev. A* **55**, 4318 (1997).
- [21] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller, *Phys. Rev. Lett.* **81**, 3108 (1998).
- [22] S. Anderloni, F. Benatti, R. Floreanini, and A. Trombettoni, e-print arXiv:0811.0925
- [23] A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer, Dordrecht, 1993).
- [24] R. Alicki and M. Fannes, *Quantum Dynamical Systems* (Oxford University Press, Oxford, 2001).
- [25] V. Gorini and A. Kossakowski, *J. Math. Phys.* **17**, 1298 (1976).
- [26] A. Frigerio and V. Gorini, *J. Math. Phys.* **17**, 2123 (1976).
- [27] A. Frigerio, C. Novellone, and M. Verri, *Rep. Math. Phys.* **12**, 279 (1977).