

Strengthened Bell inequalities for entanglement verification

Pavel Lougovski¹ and S. J. van Enk^{1,2}

¹*Department of Physics and Oregon Center for Optics, University of Oregon, Eugene, Oregon 97403, USA*

²*Institute for Quantum Information, California Institute of Technology, Pasadena, California 91125, USA*

(Received 3 August 2009; published 30 September 2009)

Bell inequalities were meant to test quantum mechanics vs local hidden variable models, but can also be used to verify entanglement. For entanglement verification purposes one assumes the validity of quantum mechanics as well as quantum descriptions of one's measurements. With the help of these assumptions it is possible to derive a *strengthened* Bell inequality whose violation implies entanglement. We generalize known examples of such inequalities by relating the expectation value of the Bell operator to a particular quantitative measure of entanglement, namely, the negativity. Moreover, we obtain statistics illustrating the fact that violating a given (strengthened or not) Bell inequality is a much more rare feat for a quantum state of two qubits than it is to be entangled.

DOI: [10.1103/PhysRevA.80.034302](https://doi.org/10.1103/PhysRevA.80.034302)

PACS number(s): 03.67.Mn, 03.65.Ud

Bell introduced his famous inequality as a way of testing predictions of quantum mechanics versus those of all possible local hidden variable (LHV) theories [1]. But the inequality (or its experimentally useful generalization, the Clauser-Horne-Shimony-Holt (CHSH) inequality [2]) can also be used to verify the presence of entanglement under the specific assumption that quantum mechanics is correct, if the inequality is violated. In fact, since the Bell-CHSH inequalities are derived from classical probability theory, without depending at all on quantum mechanics, such tests are safe in the sense of avoiding pitfalls resulting from assuming too much about the quantum state generated [3].

Let us consider a scenario where (a multitude of copies of) a quantum state ρ is shared by two parties A and B. Moreover, each party has a choice of two local measurements with dichotomic outcomes ± 1 , say, A_1, A_2 for party A and B_1, B_2 for party B. Then one can define the Bell-CHSH [2] operator,

$$\mathcal{B} = A_1 \otimes (B_1 + B_2) + A_2 \otimes (B_1 - B_2). \quad (1)$$

In order to calculate the expectation value of \mathcal{B} for a quantum state ρ the standard quantum-mechanical rule should be applied, i.e.,

$$\langle \mathcal{B} \rangle_{QM} = \text{Tr}(\mathcal{B}\rho). \quad (2)$$

On the other hand, if we assume that a LHV model is correct, then expectation values should be calculated differently [and the tensor product in Eq. (1) should be replaced by an ordinary product], namely, by

$$\langle \mathcal{B} \rangle_{LHV} = \int_{\Omega} d\omega p(\omega) \mathcal{B}(\omega), \quad (3)$$

where ω is a set of “hidden” variables and $p(\omega)$ is the probability density for those variables. The CHSH inequality reads

$$|\langle \mathcal{B} \rangle_{LHV}| \leq 2. \quad (4)$$

This inequality holds true for all LHV models but can be violated by some quantum-mechanical states for certain choices of operators $A_{1,2}, B_{1,2}$. That is, there are states for

which $|\langle \mathcal{B} \rangle_{QM}| > 2$. Such quantum states must be entangled, as all separable states satisfy the inequality. Hence, the violation of the CHSH inequality provides a sufficient criterion for entanglement.

The largest possible violation of the CHSH inequality allowed by quantum mechanics is given by Cirel'son's bound [4]:

$$|\langle \mathcal{B} \rangle_{QM}| \leq 2\sqrt{2}. \quad (5)$$

On the other hand, it was recently shown by Uffink and Seevinck [5] that, in the special case that the measurements $A_{1,2}$ and $B_{1,2}$ correspond to spin measurements in locally orthogonal spatial directions [6] (on spin-1/2 systems, which we will refer to as qubits), a significantly stronger inequality can be found for separable states. From that inequality one can derive a “strengthened Bell inequality”

$$|\langle \mathcal{B} \rangle_{QM,sep,\perp}| \leq \sqrt{2} \quad (6)$$

(derived explicitly first, as far as we know, by Roy in Ref. [7]; the inequality is also implicit in, e.g., Ref. [8]). We will refer to this inequality as the Roy-Uffink-Seevinck (RUS) bound. The subscript \perp reminds us that it holds only for spin measurements in orthogonal directions and the subscript *sep* reminds us the inequality refers to separable states. The RUS bound demonstrates the known fact that one certainly does not have to violate the CHSH inequality (4) in order to conclude one has entanglement. In other words, all data from an experiment measuring the Bell correlations in which the RUS bound is violated, but the CHSH inequality is not, can be reproduced by a LHV model, although the underlying quantum state *must* be entangled, if orthogonal spin directions were measured.

We now generalize the RUS and Cirel'son's bounds. We consider, after Uffink and Seevinck [5], orthogonal spin measurements $\{A_1, A_2\}$ and $\{B_1, B_2\}$ on a spin-1/2 system (qubit) for each party. For this choice of local measurements the Bell-CHSH operator \mathcal{B} in Eq. (1) has only two nonzero eigenvalues $\pm 2\sqrt{2}$, with the corresponding eigenvectors being two orthogonal maximally entangled two-qubit states. Let us

now maximize the expectation value of \mathcal{B} over all possible pairs of orthogonal spin measurement directions,

$$\langle \mathcal{B} \rangle_{\max} = \max\{\text{Tr}(U\mathcal{B}U^{-1}\rho), U = U_A \otimes U_B\}. \quad (7)$$

Calculating the trace in the basis where \mathcal{B} is diagonal we immediately obtain

$$\langle \mathcal{B} \rangle_{\max} = \max[2\sqrt{2}(\langle \psi|U^{-1}\rho U|\psi\rangle - \langle \phi|U^{-1}\rho U|\phi\rangle)], \quad (8)$$

where $|\psi\rangle$ and $|\phi\rangle$ are maximally entangled states, the eigenstates of \mathcal{B} with eigenvalues $\pm 2\sqrt{2}$. Now we recall that for an arbitrary two-qubit quantum state ρ we can define a fidelity F [9],

$$F(\rho) = \max\langle \Psi|U_A \otimes U_B \rho (U_A \otimes U_B)^{-1}|\Psi\rangle, \quad (9)$$

where $|\Psi\rangle$ is a maximally entangled state and U_A, U_B are local unitaries. Combining these two definitions shows that the absolute value of $\langle \mathcal{B} \rangle_{\max}$ is bounded from above by

$$|\langle \mathcal{B} \rangle_{\max}| \leq 2\sqrt{2}F(\rho). \quad (10)$$

Furthermore, for any mixed two-qubit state ρ we have the inequality (proven in Ref. [10])

$$F(\rho) \leq [1 + N(\rho)]/2, \quad (11)$$

where $N(\rho)$ is the negativity [11,12], defined as

$$N(\rho) = 2 \sum_k \max(0, -\lambda_k), \quad (12)$$

in terms of the eigenvalues λ_k of the partial transpose of ρ . We thus arrive at the announced generalization of the RUS inequality for two-qubit states:

$$|\langle \mathcal{B} \rangle_{QM,\perp}| \leq \sqrt{2}[1 + N(\rho)]. \quad (13)$$

It is straightforward to see that the last inequality, Eq. (13), contains both the RUS and Cirel'son's bounds. Namely, when $N(\rho)=0$, ρ is a separable state (this is true for our two-qubit states, although not in general) and we recover the RUS bound. On the other hand, for maximally entangled states $N(\rho)=1$ and Cirel'son's bound is recovered. Moreover, Eq. (13) provides a lower bound on the degree of entanglement in terms of negativity, if one violates the RUS bound. The bound complements the relations found in [13] between Bell inequalities and the concurrence and purity of mixed states.

The next question we address is this: if one measures spin in two fixed orthogonal spatial directions, then what is the probability that a randomly picked two-qubit state violates the original CHSH inequality? Or what is the probability it violates the RUS bound? How do these two probabilities compare to the probability of the state being entangled?

In order to answer these questions, we have to specify how to "randomly" pick a two-qubit state. In fact, this boils down to choosing a measure on the space of all two-qubit density matrices. This choice is rather arbitrary. The only (rather weak) property we wish the random distribution of states to have is that a non-negligible fraction is entangled

and that a non-negligible fraction is separable. We pick then, for no other reason in particular, a parametrization based on Ref. [12] (see also [14]). A random sample of 3×10^7 states showed a fraction $(36.437 \pm 0.010)\%$ to be entangled *a priori* [as compared to $(36.8 \pm 0.2)\%$ obtained in [12] for the same measure], thus fulfilling our one weak condition.

The following statistics apply to a *fixed* set of four correlation measurements involving spin measurements in two *fixed* orthogonal spatial directions on each of the two qubits. We note that in this case we can construct four Bell-CHSH operators from the four measured correlations:

$$\mathcal{B}_1 = A_1 \otimes (B_1 + B_2) + A_2 \otimes (B_1 - B_2),$$

$$\mathcal{B}_2 = A_1 \otimes (B_1 + B_2) - A_2 \otimes (B_1 - B_2),$$

$$\mathcal{B}_3 = A_1 \otimes (B_1 - B_2) + A_2 \otimes (B_1 + B_2),$$

$$\mathcal{B}_4 = A_1 \otimes (-B_1 + B_2) + A_2 \otimes (B_1 + B_2). \quad (14)$$

From our numerical investigations we obtained the following statistics for random mixed two-qubit states:

(i) Only a tiny fraction $(3.31 \pm 0.03) \times 10^{-4}$ violates at least one of the four CHSH inequalities that can be constructed from the four Bell-CHSH operators [Eq. (14)].

(ii) Only a small fraction $(1.244 \pm 0.003)\%$ violates at least one of the four RUS inequalities that can be constructed from the same operators [Eq. (14)].

We also checked the case of a tomographically complete measurement, where one measures each spin in three fixed spatial directions in each location: from these data 36 Bell-CHSH operators can be constructed. For this sort of measurements we find the following:

(i) The probability to violate at least one of the 36 CHSH inequalities is still small, $(0.249 \pm 0.0008)\%$.

(ii) The probability to violate at least one RUS bound out of 36 is $(5.690 \pm 0.004)\%$.

Finally, what if we construct a set of pure states (randomly distributed according to the Haar measure)? First of all, all pure states, except for a set of measure zero, are entangled. Furthermore, we find the following:

(i) A fraction $(9.908 \pm 0.005)\%$ violates at least one of the four CHSH inequalities that can be constructed from operators (14).

(ii) A fraction $(46.627 \pm 0.010)\%$ violates at least one of the four RUS inequalities that can be constructed from the same operators.

These statistics (although the exact numbers are measure dependent, of course) illustrate the fact that violating a CHSH inequality or violating the RUS bound for a fixed correlation measurement is far from necessary for entanglement (it is always sufficient, though); being entangled is much less rare than violating a given CHSH or RUS inequality.

This research was supported by the Disruptive Technologies Office (DTO) of the DNI.

- [1] J. S. Bell, *Speakable and Unsayable in Quantum Mechanics* (Cambridge University Press, Cambridge, 1987).
- [2] J. Clauser, M. Horne, A. Shimony, and R. Holt, Phys. Rev. Lett. **23**, 880 (1969); J. F. Clauser and M. Horne, Phys. Rev. D **10**, 526 (1974); J. F. Clauser and A. Shimony, Rep. Prog. Phys. **41**, 1881 (1978).
- [3] S. J. van Enk, N. Lütkenhaus, and H. J. Kimble, Phys. Rev. A **75**, 052318 (2007).
- [4] B. S. Cirel'son, Lett. Math. Phys. **4**, 93 (1980).
- [5] J. Uffink and M. Seevinck, Phys. Lett. A **372**, 1205 (2008).
- [6] No relation is assumed between the directions in location A and those in B .
- [7] S. M. Roy, Phys. Rev. Lett. **94**, 010402 (2005).
- [8] G. A. Durkin and C. Simon, Phys. Rev. Lett. **95**, 180402 (2005).
- [9] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **54**, 3824 (1996).
- [10] F. Verstraete and H. Verschelde, Phys. Rev. A **66**, 022307 (2002).
- [11] G. Vidal and R. F. Werner, Phys. Rev. A **65**, 032314 (2002).
- [12] K. Zyczkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, Phys. Rev. A **58**, 883 (1998).
- [13] F. Verstraete and M. M. Wolf, Phys. Rev. Lett. **89**, 170401 (2002).
- [14] T. Tilma, M. Byrd, and E. C. G. Sudarshan, J. Phys. A **35**, 10445 (2002).