

Probing phase separation in Bose-Fermi mixtures by the critical superfluid velocity

Jacob Linder and Asle Sudbø

Department of Physics, Norwegian University of Science and Technology, N-7491 Trondheim, Norway

(Received 4 August 2009; published 25 January 2010)

We investigate the effect exerted by spin-polarized fermions on the interaction between superfluid bosons for a Bose-Fermi mixture residing on an optical lattice, with particular emphasis on the possibility of an induced phase separation. Using a set of microscopic parameters relevant to a ^{40}K - ^{87}Rb mixture, we show how the phase-separation criterion may be directly probed by means of the critical superfluid velocity of the bosonic condensate. We report quantitative results for the magnitude of the superfluid velocity and its dependence on the trap depth, the boson-fermion interaction, and the fermionic filling fraction. All of these parameters can be controlled experimentally in a well-defined manner. We propose an experimental setup for probing the critical superfluid velocity.

DOI: [10.1103/PhysRevA.81.013622](https://doi.org/10.1103/PhysRevA.81.013622)

PACS number(s): 03.75.Lm, 05.30.Jp, 05.30.Fk

I. INTRODUCTION

The scenario of trapped ultracold atoms residing on tunable optical lattices offers a fertile arena for exploration of fundamental physics. One of the most intriguing features of such systems is the possibility to exert experimental control over the environment where the atoms are located. This is accomplished by means of tuning the frequency of the lasers generating the optical lattice, a feature that may be used to induce phase transitions in the system. Trapped ultracold atoms host phases including supersolidity, Mott insulation, and superfluidity, and have been studied extensively (see Refs. [1–3] and references therein).

The atoms on the optical lattice may be either bosons, fermions, or a mixture of both. In particular, it is experimentally possible to generate Bose-Fermi mixtures where the spins of the fermions are frozen due to the influence of the confining magnetic trap. Depending on the relative strength of the intersite hopping and interaction parameters, respectively, the system enters into a specific quantum phase. A key observation in this context is that the interaction between the fermions and the bosons may strongly influence the preferred ground state of the system.

A convenient way of treating Bose-Fermi mixtures theoretically is to integrate out the fermionic sector, thus obtaining an effective interacting boson theory. This is possible when the fermion spin is frozen, leading to a vanishing fermionic onsite interaction. It turns out that the resulting effective boson-boson interaction U_b is very sensitive to the presence of a fermionic density [4]. In fact, the boson-fermion interaction may render the Bose system thermodynamically unstable and lead to phase separation provided the effective boson-boson interaction becomes attractive. Such a phase separation is certainly interesting in its own right, but also represents a serious obstacle for observing novel quantum phases arising out of an interacting mixture of bosons and fermions, since it narrows the parameter range in which the bosons and fermions coexist.

From an experimental point of view, the phase-separation criterion may be probed by means of monitoring the critical superfluid velocity in the Bose-Fermi mixture. This has previously been accomplished experimentally in Bose-Einstein condensates by means of stirring the trapped gas with a

blue-detuned laser [5,6]. The superfluid quantum state then becomes energetically unstable at a critical magnitude of the velocity. Previously, several aspects of the critical superfluid velocity have been investigated in the context of single- and multicomponent Bose-Einstein condensates [7–15] as well as in Fermi superfluids [16–18].

However, an analysis of the critical velocity for the bosonic superfluid phase in a Bose-Fermi mixture is still lacking. Of particular interest is the question of how the fermion-boson interaction influences the critical velocity in a Bose-Fermi mixture. Very recently, it was shown in Ref. [19] how the fermion-boson interaction can be tuned over a wide range using a Feshbach resonance, allowing for both an attractive or repulsive character. This finding opens up new possibilities in terms of probing the various quantum phases that may arise in such Bose-Fermi mixtures [20].

In this article, we calculate quantitatively the critical superfluid velocity v_c in a Bose-Fermi mixture using a set of realistic experimental parameters pertaining to a ^{40}K - ^{87}Rb mixture. We focus especially on how v_c depends on the trap depth, the boson-fermion interaction, and the fermionic filling fraction, which all are parameters that can be tuned experimentally in a controllable fashion. Our results yield numbers that are similar in magnitude to the critical velocity obtained experimentally in a Bose-Einstein condensate [5], namely of order $\mathcal{O}(\text{mm/s})$. We also propose an experimental setup for probing the critical superfluid velocity in a Bose-Fermi mixture.

This work is organized as follows. In Sec. II, we introduce the theoretical framework and previously obtained results that we will rely on in our study of the critical superfluid velocity. In Sec. III, we present our main results, which is a study of how v_c is influenced by the trap depth, the boson-fermion interaction, and the fermionic chemical potential. We discuss our results in Sec. IV, suggesting also a possible experimental setup that may probe the predicted effects, and finally conclude in Sec. V. Note that to obtain quantitative results for v_c , we will not use units such that $\hbar = c = 1$, but instead use their actual values.

II. THEORY

To begin with, we briefly account for the route employed to obtain our main results. A general Hamiltonian describing

interacting fermions and bosons reads

$$H = H_f + H_b + H_{bf}, \quad (1)$$

where we have defined

$$H_f = \int d\mathbf{r} \psi_f^\dagger(\mathbf{r}) [-\hbar^2 \nabla^2 / (2m_f) + V_f(\mathbf{r})] \psi_f(\mathbf{r}), \quad (2)$$

and $f \rightarrow b$ for H_b . The interaction term is

$$H_{bf} = \int d\mathbf{r} [g_{bf} \psi_b^\dagger(\mathbf{r}) \psi_b(\mathbf{r}) \psi_f^\dagger(\mathbf{r}) \psi_f(\mathbf{r}) + g_b \psi_b^\dagger(\mathbf{r}) \psi_b^\dagger(\mathbf{r}) \psi_b(\mathbf{r}) \psi_b(\mathbf{r})]. \quad (3)$$

Above, m_α and V_α denote the mass and optical lattice potential for $\alpha = \{f, b\}$, whereas g_b and g_{bf} are the boson-boson and boson-fermion interactions, respectively. It is implicitly assumed above that we are dealing with a fully spin-polarized fermion system. By expanding the field operators ψ_α in Bloch wave functions $\{u_k, v_k\}$ for a periodic potential [21],

$$\begin{aligned} \psi_f^\dagger(\mathbf{r}) &= \frac{1}{\sqrt{N}} \sum_k v_k(\mathbf{r}) c_k^\dagger, \\ \psi_b^\dagger(\mathbf{r}) &= \frac{1}{\sqrt{N}} \sum_k u_k(\mathbf{r}) b_k^\dagger, \end{aligned} \quad (4)$$

we arrive at an effective lattice Hamiltonian

$$H = \sum_k \varepsilon_{k,b} b_k^\dagger b_k + \sum_k \varepsilon_{k,f} c_k^\dagger c_k + \frac{U_b}{2N} \sum_{\{k_j\}} b_{k_1}^\dagger b_{k_2}^\dagger b_{k_3} b_{k_4} + \frac{U_{bf}}{N} \sum_{\{k_j\}} b_{k_1}^\dagger b_{k_2} c_{k_3}^\dagger c_{k_4}. \quad (5)$$

This procedure is justified when the optical potential is strong enough, typically $V_\alpha > E_\alpha^{\text{rec}}$, where

$$E_\alpha^{\text{rec}} = 2\hbar^2 \pi^2 / (\lambda^2 m_\alpha) \quad (6)$$

is the atom recoil energy and λ is the wavelength of the laser light. For later purposes, we define the optical trap depth $s_\alpha = V_\alpha / E_\alpha^{\text{rec}}$. To evaluate the critical superfluid velocity explicitly from the microscopic parameters of an experimental setup, we make use of the following expression for the hopping and interaction parameters [20]:

$$\begin{aligned} t_\alpha &= \frac{2(E_\alpha^{\text{rec}} V_\alpha^3)^{1/4}}{\sqrt{\pi} e \sqrt{4V_\alpha / E_\alpha^{\text{rec}}}}, \\ U_b &= \frac{\sqrt{32\pi} (E_b^{\text{rec}} V_b^3)^{1/4} a_b}{\lambda}. \end{aligned} \quad (7)$$

For the fermion-boson interaction, one has

$$U_{bf} = \frac{8\sqrt{\pi} (E_f^{\text{rec}} V_b^3 V_f^3)^{1/4} (1 + m_f / m_b) a_{bf}}{\lambda (\sqrt{V_b} + \sqrt{V_f E_b^{\text{rec}} / E_f^{\text{rec}}})^{3/2}}. \quad (8)$$

It is also useful to introduce the scattering lengths $\{a_b, a_{bf}\}$, which are related to the interaction parameters in Eq. (3) as follows:

$$a_b = \frac{g_b m_b}{4\pi \hbar^2}, \quad a_{bf} = \frac{g_{bf} m_f m_b}{2\pi (m_f + m_b) \hbar^2}. \quad (9)$$

The onsite potentials in Eqs. (7) and (8) are obtained by relating them directly to the Wannier functions $\{\mathcal{U}(\mathbf{r}), \mathcal{V}(\mathbf{r})\}$ used to

approximate the wave functions in the lowest Bloch band. For instance, one has [21]

$$U_{bf} = g_{bf} \int d\mathbf{r} |\mathcal{U}(\mathbf{r})|^2 |\mathcal{V}(\mathbf{r})|^2, \quad (10)$$

where we have defined

$$\begin{aligned} \mathcal{U}(\mathbf{r} - \mathbf{R}) &= \frac{1}{N} \sum_k u_k(\mathbf{r}) e^{-i\mathbf{R}\cdot\mathbf{k}}, \\ \mathcal{V}(\mathbf{r} - \mathbf{R}) &= \frac{1}{N} \sum_k v_k(\mathbf{r}) e^{-i\mathbf{R}\cdot\mathbf{k}}. \end{aligned} \quad (11)$$

The energy dispersions are dictated by the geometry of the optical lattice and are proportional to the nearest-neighbor matrix elements t_α , whereas the summation over $\{\mathbf{k}_j\}$ should be taken such that momentum is conserved in the scattering process.

The Hamiltonian Eq. (5) is now quadratic in the fermion sector, which allows us to integrate out the fermions in the partition function by using a functional integral formulation. After doing so, one identifies an effective boson-boson interaction U_b of the form [4]

$$U_b = U_b + U_{bf}^2 \chi(T, \mathbf{q}), \quad \chi(T, \mathbf{q}) = \frac{1}{N} \sum_k \frac{F(\varepsilon_{k,f}) - F(\varepsilon_{k+q,f})}{\varepsilon_{k,f} - \varepsilon_{k+q,f} + i\delta}, \quad \delta \rightarrow 0. \quad (12)$$

Here, $\chi(T, \mathbf{q})$ is the Lindhard function describing the fermionic polarization-bubble response, and $F(\varepsilon) = [1 + e^{\beta(\varepsilon - \mu)}]^{-1}$, $\beta = (k_B T)^{-1}$ is the Fermi distribution function.

To proceed analytically, we restrict ourselves to the weak-coupling regime and employ a Bogoliubov mean-field theory for superfluidity [22] to arrive at the bosonic quasiparticle excitation spectrum

$$\mathcal{E}_{\mathbf{q},b} = \sqrt{\varepsilon_{\mathbf{q},b} \{ \varepsilon_{\mathbf{q},b} + 2n_b [U_b + U_{bf}^2 \chi(T, \mathbf{q})] \}}. \quad (13)$$

The phase-separation criterion (the point at which the bosonic excitation energies cease to be real) thus reads

$$U_b < -U_{bf}^2 \lim_{q \rightarrow 0} \chi(T, \mathbf{q}). \quad (14)$$

Phase separation is triggered by the effective boson-boson interaction becoming attractive, which leads to a negative compressibility and an unstable homogeneous superfluid state [23]. Note that the critical value of U_{bf} where phase-separation sets in is independent of the sign of the interaction U_{bf} . As an example of how the phase separation may be manifested, it was shown in Ref. [24] how the bosonic density in a Bose-Fermi mixture confined in a three-dimensional harmonic trap would be strongly enhanced in the center of the trap surrounded by a fermionic density shell in the phase-separated regime.

Denoting the Fermi level by ε_f , one finds that

$$\chi(T, \mathbf{q} \rightarrow 0) = \int d\varepsilon N(\varepsilon) \partial_\varepsilon F(\varepsilon) = -N(\varepsilon_f). \quad (15)$$

The effective boson-boson interaction then takes the form $U_{\text{eff}} = U_b - U_{bf}^2 N(\varepsilon_f)$, and remains at a constant positive or negative value when varying the temperature in the regime $T \ll T_f$, where T_f is the Fermi temperature. In a two-dimensional (2D) lattice structure, the energy bands feature

saddle points at distinct wave vectors, thus giving rise to well-known van Hove singularities. In the vicinity of a van Hove singularity, the density of states (DOS) is not a smooth function of the energy and Eq. (15) no longer holds. When the fermionic chemical potential is tuned to match the van Hove singularity, the Lindhard function diverges logarithmically as follows [25]:

$$\chi(T \rightarrow 0, 0) = -\chi_0 \ln\left(\frac{C t_f}{k_B T}\right), \quad (16)$$

in the zero-temperature limit $T \rightarrow 0$. Here, χ_0 is a prefactor of dimension inverse energy, whereas C is a numerical prefactor. For a square lattice, one finds $\chi_0 = 1/(2\pi^2 t_f)$ and $C \simeq 18.08$, whereas, for example, a triangular lattice one would find $\chi_0 = 3/(4\pi^2 t_f)$ and $C = 9.04$ [25]. It should be noted that we have considered the static limit $i\varepsilon_n \rightarrow 0$ for the Lindhard function, where ε_n is a bosonic Matsubara frequency. This approximation is valid for a scenario where the fermion response time is much faster than the bosonic equivalent, which means that one can disregard retardation effects [25].

III. RESULTS

From now on, we will consider a simple square lattice for concreteness, which is the easiest setup to realize experimentally. In this case, the van Hove singularity is located at $\varepsilon = 0$ and the DOS has a bandwidth of $W = 8t_f$. We find that the energy dispersion in the long wavelength limit reads

$$\mathcal{E}_b(T, \mathbf{q} \rightarrow 0) = \sqrt{4n_b t_b a^2 [U_b + U_{bf}^2 \chi(T, 0)]} |\mathbf{q}|. \quad (17)$$

The critical superfluid velocity v_c is obtained in the standard way

$$v_c = \min\left(\frac{\mathcal{E}_b}{\hbar |\mathbf{q}|}\right), \quad (18)$$

leading to

$$v_c = \sqrt{4n_b t_b a^2 [U_b + U_{bf}^2 \chi(T, 0)]} / \hbar. \quad (19)$$

Some properties of Bose-Fermi mixtures with a fermionic chemical potential tuned to the van Hove singularity were discussed in Refs. [4,21]. Here, we will consider a situation of a nonzero chemical potential, thus moving away from half-filling. To model a realistic experiment, we will employ the following parameters for a ^{40}K - ^{87}Rb Bose-Fermi mixture [26,27]: $T_{\text{BEC}} = 100$ nK, $m_f = 6.64 \times 10^{-26}$ kg, $m_b = 1.44 \times 10^{-25}$ kg, $a_b = 98a_0$. Here, $a_0 \simeq 52.9 \times 10^{-3}$ nm is the Bohr radius. To ensure equal lattice depths $s_\alpha \equiv s$ for the fermions and bosons, measured relative to their respective recoil energies E_α^{rec} , we fix $\lambda = 755$ nm [19]. In general, the effective potentials seen by the fermions and bosons can be tuned by detuning the lattice wavelength relative to the wavelengths $\lambda_{f(b)}$ of the fermions (bosons) according to [28]

$$\frac{V_b}{V_f} = \frac{\Gamma_f \lambda_f^4 \Delta \lambda_b}{\Gamma_b \lambda_b^4 \Delta \lambda_f}, \quad (20)$$

where Γ_α is the natural linewidth, $\alpha = \{f, b\}$. We set the zero-temperature condensate fraction to $n_B(T = 0) = 0.5$, and employ a mean-field approximation for its temperature




FIG. 1. (Color online) Fermionic density of states on a square lattice, featuring a van Hove singularity at $\varepsilon = 0$. We consider two filling fractions characterized by $\mu/t_f = 0.5$ and $\mu/t_f = 3.5$, respectively. Here, $N_0 = 1/(2\pi^2 t_f)$.

dependence $n_B(T)$. We will fix the temperature at $T/T_{\text{BEC}} = 0.6$, which should be feasible to reach experimentally and still within the regime of validity for a mean-field approximation [29]. The remaining parameters that must be specified are the chemical potential and the boson-fermion scattering length. As shown in Fig. 1, we will consider two fermion fillings characterized by $\mu/t_f = 0.5$ (close to the van Hove singularity) and $\mu/t_f = 3.5$ (close to the band edge), respectively. The influence of inhomogeneous density distributions and finite-size effects, leading to a spatially dependent chemical potential, will be discussed in detail in Sec. IV. The approximation of constant chemical potential is here expected to be useful to obtain experimentally relevant results, as we shall argue later. The boson-fermion scattering length a_{bf} is tunable, as shown recently in Ref. [19]. By using a Feshbach resonance, scattering lengths in a range $\pm 800a_0$ were reached. We shall therefore consider both positive and negative scattering lengths, reaching up to several hundreds of a_0 . Finally, it should be noted that we here restrict our attention to values of the trap depth $s < 8$ in light of a recent experiment [30] where a 2D Bose gas was realized with a Mott insulating phase above $s \simeq 8$.

To evaluate the critical superfluid velocity, we employ a numerical solution of the expression:

$$v_c = \left[4n_b t_b a^2 \left(U_b - \frac{\beta}{8\pi^2 t_f} U_{bf}^2 \int_{-4t_f}^{4t_f} \int_0^{\pi/2} \frac{d\varepsilon d\gamma}{\sqrt{[\cos^2 \gamma + (\varepsilon \sin \gamma / 4t_f)^2] \cosh[\beta(\varepsilon - \mu)]}} \right) \right]^{1/2}. \quad (21)$$

As seen from Eq. (21), it becomes necessary to account properly for the finite temperature T to describe the physical properties of Bose-Fermi mixtures, unlike the purely bosonic case.

In what follows, we will investigate how v_c depends on the fermion-boson interaction parameter a_{bf} and the trap depth s ,




FIG. 2. (Color online) Plot of the critical superfluid velocity v_c for the bosons and its dependence on the trap depth for several values of the boson-fermion scattering length $a = a_{bf}$. Here, we have set (a) $\mu/t_f = 0.5$ and (b) $\mu/t_f = 3.5$. All other parameter values are given as follows: $T_{\text{BEC}} = 100$ nK, $m_f = 6.64 \times 10^{-26}$ kg, $m_b = 1.44 \times 10^{-25}$ kg, $a_b = 98a_0$, $\lambda = 755$ nm, and $T/T_{\text{BEC}} = 0.6$.

using the set of experimentally realistic parameters described above. Consider first its dependence on the trap depth s , as shown in Fig. 2. One of the main features is that v_c exhibits a robustness toward the trap depth for relatively low values of a_{bf} . For high values of s , one would expect a transition into a Mott insulating state for commensurate fillings. When the interaction a_{bf} becomes strong compared to the intrinsic bosonic repulsion a_b , any increase in trap depth s is much more efficient in suppressing the critical velocity. We have distinguished between two fermionic fillings corresponding to $\mu/t_f = 0.5$ and $\mu/t_f = 3.5$ in Fig. 2 to compare the cases with a chemical potential close to the van Hove singularity and close to the band edge, respectively. As seen, the difference is minor except at large values of the interaction a_{bf} , where




FIG. 3. (Color online) Plot of the critical superfluid velocity v_c for the bosons and its dependence on the fermion-boson interaction for several values of the trap depth s . Here, we have set (a) $\mu/t_f = 0.5$ and (b) $\mu/t_f = 3.5$. All other parameter values are given as follows: $T_{\text{BEC}} = 100$ nK, $m_f = 6.64 \times 10^{-26}$ kg, $m_b = 1.44 \times 10^{-25}$ kg, $a_b = 98a_0$, $\lambda = 755$ nm, and $T/T_{\text{BEC}} = 0.6$.

v_c is substantially reduced with $\sim 50\%$ for a given trap depth, in addition to a much smaller critical trap depth s where the superfluid velocity vanishes.

Next, we consider how the critical superfluid velocity v_c is influenced by the fermion-boson interaction a_{bf} . The result is shown in Fig. 3. Since the critical velocity in Eq. (21) depends on U_{bf}^2 , the sign of the interaction is irrelevant for the magnitude of v_c . Upon increasing the magnitude of the interaction a_{bf} , the superfluid velocity is strongly reduced and eventually vanishes, indicating a phase-separated regime. This may be understood physically by noting that the contribution from the Lindhard function is negative in Eq. (21), meaning that the fermionic contribution to the induced boson-boson




FIG. 4. (Color online) Plot of the critical superfluid velocity as a function of the intrinsic boson-boson scattering length a_b and the boson-fermion interaction a_{bf} . We have set $\mu/t_f = 0.5$ and considered a trap depth $s = 5$. All other parameter values are given as follows: $T_{\text{BEC}} = 100$ nK, $m_f = 6.64 \times 10^{-26}$ kg, $m_b = 1.44 \times 10^{-25}$ kg, $\lambda = 755$ nm, and $T/T_{\text{BEC}} = 0.6$.

interaction is attractive. As the Bose-condensed phase is unstable toward attractive interactions, the critical velocity vanishes when the fermionic contribution eventually overtakes the intrinsic bosonic repulsion. In Ref. [19], it was very recently experimentally demonstrated how the fermion-boson interaction a_{bf} could be tuned in a well-defined manner over a wide range $\pm 800a_0$ by exploiting a Feshbach resonance. It should therefore be experimentally viable to monitor the critical velocity v_c as a function of the interaction a_{bf} by using such techniques.

Finally, the mutual dependence on the intrinsic boson-boson interaction and the fermion-boson interaction is plotted in Fig. 4, setting $\mu/t_f = 0.5$ and $s = 5$. While the critical velocity is suppressed with increasing a_{bf} , it is enhanced by increasing a_b . The reason for this is that the effective boson-boson interaction becomes more repulsive, in favor of the phase-coexistent state. It should nevertheless be emphasized that above a critical magnitude for the effective interaction U_b , a phase transition from superfluid to Mott insulator takes place. As shown in Ref. [22], the present mean-field Bogoliubov approach does not capture this transition as it treats the interaction only in a weak-coupling regime. Therefore, the results reported here are obviously only valid inside the superfluid regime.

IV. DISCUSSION

The experimental detection of a critical superfluid velocity requires measurements at temperatures well below T_{BEC} , thus in the nano-Kelvin regime. One possible route to probing the critical velocity was described in Ref. [5]. There, dissipation in a Bose-Einstein condensed gas was monitored by means of moving a laser beam through the condensate at different velocities (see Fig. 5). The laser effectively plays the role of a massive macroscopic object, which creates a moving boundary




FIG. 5. (Color online) Proposed experimental setup for probing the critical superfluid velocity in a Bose-Fermi mixture. A laser beam serves as a macroscopic object flowing through the condensate, thereby creating a moving boundary condition for the quasiparticle excitations.

condition. The main finding in Ref. [5] was that strong heating was observed only above a critical velocity, and the laser was enabled to move back and forth through the condensate at a constant velocity by applying a triangular waveform to an acousto-optic deflector.

In the treatment of the critical superfluid velocity, it is implicitly assumed that the bosons are in the superfluid phase for the relevant parameter regime. To verify this, a full numerical self-consistent solution is required. Our main purpose here is to report on the magnitude of the critical velocity and its dependence on tunable parameters, using a set of realistic parameters employed in real experiments [19,26,29] in which the bosons indeed were in the condensed state, which should justify our assumption. We have restricted our attention to trap depths satisfying $s < 8$ [30], as should be reasonable for the superfluid regime. Finally, we note that the results we have obtained quantitatively agree with previous measurements for the critical velocity in Bose-Einstein condensates. In particular, $v_c \sim 1.6$ mm/s was reported in Ref. [5].

In our treatment of the critical velocity, we have employed a mean-field treatment, which thus neglects phase-fluctuations of the superfluid order parameter near the phase separation point. In this sense, the magnitude of the critical velocity estimated by our mean-field theory should be somewhat larger than the real critical velocity. Nevertheless, to obtain analytical results and understand the role of the different interactions with respect to the critical velocity, we have assumed in our manuscript that the mean-field theory gives a reasonable quantitative measure for v_c . It should then be noted that critical fluctuations near the phase-separation point, assuming thus a second-order phase transition, could render the actual v_c to be lower than the value estimated here. We also note that in the presence of an inhomogeneous environment such as a confinement trap of the condensate setup by utilizing magnetic fields, the local chemical potential for the fermions and bosons may be expected to vary locally depending on the position. From an experimental point of view, measurements on the condensate are often performed after the confinement trap has been switched off such that the condensate will expand and the chemical potential should over time relax to a slower spatial variation. Granted, the chemical potential will most likely still feature some local variations and one might, for instance, expect a critical velocity that depends on the radial distance

from the trap center. However, as our results show, the specific value of the chemical potential does not seem to have a large impact on the results, whereas, for example, the boson-fermion interaction strongly modifies the critical velocity. It therefore seems reasonable to expect that local variations in the chemical potential should not drastically change our results. Moreover, one may in any case envision a situation where the critical superfluid velocity is measured in a region where the chemical potential is largely homogeneous. In fact, a recent study [31] used large-scale Monte Carlo simulations to investigate the influence of density inhomogeneities and finite size effects of a trapped Bose-Einstein condensate. It was concluded in Ref. [31] that predictions based on a uniform density could have robust finite-size realizations in actual experimental, inhomogeneous trapped systems. In other words, although a trapping potential effectively models a spatially dependent chemical potential, leading to a density that typically is high in the center of the trap, the approximation of a uniform chemical potential turns out to be a reasonable approximation as verified directly in Ref. [31].

V. SUMMARY

In conclusion, we have studied how the fermion-boson interaction for a Bose-Fermi mixture residing on an optical

lattice modifies the effective interaction between the superfluid bosons. In particular, we have investigated how the phase-separation criteria is manifested through the critical superfluid velocity. Employing a set of microscopic parameters relevant to a ^{40}K - ^{87}Rb mixture [19,26,29], we report quantitative results for the magnitude of the superfluid velocity and its dependence on the trap depth, the boson-fermion interaction, and the fermionic filling fraction. All of these parameters can be tuned experimentally by means of the laser intensity and by exploiting Feshbach resonances. We find that the overall tendency of the boson-fermion interaction is to suppress v_c , and our quantitative results are of similar magnitude as previous measurements of the critical velocity in Bose-Einstein condensates, where $v_c \simeq 1.6$ mm/s was estimated [5]. We have proposed an experimental setup for probing the critical superfluid velocity, which may serve as a direct tool to monitor a phase-separation scenario in a Bose-Fermi mixture.

ACKNOWLEDGMENTS

J. Linder thanks I. B. Sperstad for helpful comments. This work was supported by the Research Council of Norway under Grant Nos. 158518/431 and 158547/431 (NANOMAT), and Grant No. 167498/V30 (STORFORSK).

-
- [1] O. Morsch and M. Oberthaler, *Rev. Mod. Phys.* **78**, 179 (2006).
 - [2] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
 - [3] S. Giorgini, L. P. Pitaevskii, and S. Stringari, *Rev. Mod. Phys.* **80**, 1215 (2008).
 - [4] H. P. Büchler and G. Blatter, *Phys. Rev. Lett.* **91**, 130404 (2003).
 - [5] C. Raman, M. Köhl, R. Onofrio, D. S. Durfee, C. E. Kulewicz, Z. Hadzibabic, and W. Ketterle, *Phys. Rev. Lett.* **83**, 2502 (1999).
 - [6] R. Onofrio, C. Raman, J. M. Vogels, J. R. Abo-Shaer, A. P. Chikkatur, and W. Ketterle, *Phys. Rev. Lett.* **85**, 2228 (2000).
 - [7] C. Menotti, M. Kraeämer, A. Smerzi, L. Pitaevskii, and S. Stringari, *Phys. Rev. A* **70**, 023609 (2004).
 - [8] D. Boers, C. Weiss, and M. Holthaus, *Europhys. Lett.* **67**, 887 (2004).
 - [9] V. M. Kaurov, A. B. Kuklov, and A. E. Meyerovich, *Phys. Rev. Lett.* **95**, 090403 (2005).
 - [10] D. V. Fil and S. I. Shevchenko, *Phys. Rev. A* **72**, 013616 (2005).
 - [11] J. Ruostekoski and Z. Dutton, *Phys. Rev. A* **76**, 063607 (2007).
 - [12] X. P. Liu, *Phys. Rev. A* **76**, 053615 (2007).
 - [13] L. Yu. Kravchenko and D. V. Fil, *J. Low Temp. Phys.* **150**, 612 (2008).
 - [14] J. Linder and A. Sudbø, *Phys. Rev. A* **79**, 063610 (2009).
 - [15] U. Shrestha, J. Javanainen, and J. Ruostekoski, *Phys. Rev. Lett.* **103**, 190401 (2009).
 - [16] T. K. Ghosh and K. Machida, *Phys. Rev. A* **73**, 013613 (2006).
 - [17] R. B. Diener, R. Sensarma, and M. Randeria, *Phys. Rev. A* **77**, 023626 (2008).
 - [18] T. N. De Silva, *Phys. Rev. A* **79**, 013612 (2009).
 - [19] T. Best, S. Will, U. Schneider, L. Hackermüller, D. van Oosten, I. Bloch, and D.-S. Lühmann, *Phys. Rev. Lett.* **102**, 030408 (2009).
 - [20] K. Sengupta, N. Dupuis, and P. Majumdar, *Phys. Rev. A* **75**, 063625 (2007).
 - [21] H. P. Büchler and G. Blatter, *Phys. Rev. A* **69**, 063603 (2004).
 - [22] D. van Oosten, P. van der Straten, and H. T. C. Stoof, *Phys. Rev. A* **63**, 053601 (2001).
 - [23] A. A. Abrikosov, L. P. Gorkov, and I. E. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics* (Dover, New York, 1963).
 - [24] K. Mølmer, *Phys. Rev. Lett.* **80**, 1804 (1998).
 - [25] P. P. Orth, D. L. Bergman, and K. Le Hur, *Phys. Rev. A* **80**, 023624 (2009).
 - [26] G. Roati, F. Riboli, G. Modugno, and M. Inguscio, *Phys. Rev. Lett.* **89**, 150403 (2002).
 - [27] E. G. M. van Kempen, S. J. J. M. F. Kokkelmans, D. J. Heinzen, and B. J. Verhaar, *Phys. Rev. Lett.* **88**, 093201 (2002).
 - [28] F. Illuminati and A. Albus, *Phys. Rev. Lett.* **93**, 090406 (2004).
 - [29] M. Modugno, F. Ferlaino, F. Riboli, G. Roati, G. Modugno, and M. Inguscio, *Phys. Rev. A* **68**, 043626 (2003).
 - [30] I. B. Spielman, W. D. Phillips, and J. V. Porto, *Phys. Rev. Lett.* **100**, 120402 (2008).
 - [31] S. Kragset, E. Babaev, and A. Sudbø, *Phys. Rev. Lett.* **97**, 170403 (2006); *Phys. Rev. A* **77**, 043605 (2008).