

Temporal compression of quantum-information-carrying photons using a photon-echo quantum memory approach

S. A. Moiseev

Institute for Quantum Information Science, and Department of Physics & Astronomy, University of Calgary, Canada and Kazan Physical-Technical Institute of the Russian Academy of Sciences, Russia

W. Tittel

Institute for Quantum Information Science, and Department of Physics & Astronomy, University of Calgary, Canada
(Received 9 April 2010; published 12 July 2010)

We study quantum compression and decompression of light pulses that carry quantum information using a photon-echo quantum memory technique with controllable inhomogeneous broadening of an isolated atomic absorption line. We investigate media with differently broadened absorption profiles, transverse and longitudinal, finding that the recall efficiency can be as large as unity and that the quantum information encoded into the photonic qubits can remain unperturbed. Our results provide insight into reversible light-atom interaction and are interesting in view of future quantum communication networks, where pulse compression and decompression may play an important role in increasing the qubit rate or in mapping quantum information from photonic carriers with large optical bandwidth into atomic memories with smaller bandwidth.

DOI: [10.1103/PhysRevA.82.012309](https://doi.org/10.1103/PhysRevA.82.012309)

PACS number(s): 03.67.Hk, 32.80.Qk, 42.50.Md

I. INTRODUCTION

As any communication system, quantum communication relies on preparing carriers of (quantum) information, transmitting those carriers in a reliable way, and processing the information. Obviously, in order to maximize the communication rate, the duration of the carriers, naturally photons, should be as short as possible. However, while quantum communication channels such as optical fibers or free-space channels allow transmission of broadband, subnanosecond photons, it is often impossible to generate or process such carriers. Of particular concern are key elements for quantum repeaters [1] such as certain entangled photon-pair sources [2–4], or quantum memories [5], whose bandwidth is often limited by material constraints.

In this article, we study temporal or bandwidth conversion as a quantum-information-preserving transformation for photonic information carriers. On the one hand, increasing the optical bandwidth in an efficient way, that is, decreasing the duration of an information carrier, will allow increasing the transmission rate over a single quantum channel when time-multiplexing several small bandwidth photon sources or photon pair sources. On the other hand, decreasing the bandwidth will allow mapping of large-bandwidth photonic carriers into quantum memories with limited spectral width. In view of this transformation, we investigate a photon-echo-type quantum memory approach based on controlled reversible inhomogeneous broadening (CRIB) of a large ensemble of atomic absorbers [6–10]. Relaxing the requirement of symmetric inversion of atomic detunings $\Delta \rightarrow -\Delta$ during absorption of the light and recall, respectively, in a more general version with nonsymmetric inversion $\Delta \rightarrow -\eta\Delta$ with compression factor $\eta \neq 1$, we find accelerated (decelerated) rephasing of atomic coherences and thus temporal compression (decompression) of the re-emitted light field. We analytically analyze the proposed scheme in optically thick atomic media with transverse and longitudinal inhomogeneous broadenings for *recall efficiency*, *fidelity*, and *gain* of the transmission rate

over a single channel through multiplexing. In transverse inhomogeneously broadened media, the atomic resonance frequencies vary normal (transverse) to the spatial coordinate z measured along the propagation direction of the light, and the absorption profile is independent of z . In longitudinal inhomogeneously broadened media, the resonance frequencies depend linearly on z . Assuming large optical depth for storage and retrieval, we find, for the case of transverse broadening, that the recall efficiency is limited by the compression factor η , while it reaches unity in the case of a longitudinal broadened medium. We also find, for transverse broadening, that the fidelity of a recalled photonic time-bin qubit with the original qubit is one, regardless the compression factor, but that it is limited in the case of longitudinal broadening. We point out that optical pulse compression has previously been considered using traditional photon echos and chirped excitation pulses [11–13]. However, similar to data storage [14,15], this approach is not suitable for temporal compression of quantum data. Quantum compression using CRIB was first discussed in [16], and the first observations as well as numerical studies for the case of longitudinal broadening have recently been reported [17,18].

This article is organized in the following way. We first discuss the figures of merit chosen to assess the performance of quantum compression. We then introduce photonic *time-bin qubits* and present two ways of describing this basic unit of quantum information. This part is followed by a description of the standard CRIB-based quantum memory protocol, which employs a hidden symmetry in the equations describing the atom-light interaction during storage and recall. In order to assess the change in the recalled photonic quantum state for CRIB-based quantum compression, which does not involve time-reversed quantum dynamics, we have to solve the equations of motion. This is done in the next section, where we also derive the efficiency, fidelity, and gain of quantum compression for the examples of transverse and longitudinal broadening. This is the main part of this article. It is followed by a brief feasibility study of our protocol in

rare-earth-ion-doped crystals. The article terminates with a conclusion.

II. FIGURES OF MERIT

Temporal compression of quantum data is of importance to quantum communication, similar to temporal compression of classical data and classical (tele) communication. However, the criteria imposed on a “good” compression procedure are much more severe in the quantum case: Reduced efficiency impacts irreversibly on the quantum information rate through photon loss, in opposition to amplitude loss in the classical case, which can be compensated by means of optical amplifiers (note that amplification is unsuitable for quantum communication, as stated in the no-cloning theorem [19,20]). Furthermore, unpredictable modification of the input photonic quantum state $|\psi_{\text{in}}\rangle_p$ during compression results in an increased quantum bit error rate (QBER) or requires compensation via not-yet-practical quantum error correction [21,22]. In opposition, classical information, due to its digital nature, is much more tolerant to noise. In this article we therefore use the efficiency ϵ , the fidelity F (which specifies the unpredictable change of an input quantum state), and the gain G (which derives from the efficiency and the compression parameter) as figures of merit to analyze the performance of quantum compression.

In the following, we assume pure photonic qubit input states. The compression operation yields a (possibly mixed) compressed photonic output state that is obtained from the total density matrix by tracing over the degrees of freedom related to the atomic system and possibly noncompressed photonic modes. Ignoring vacuum components, this results in a generally not normalized density matrix $\rho_{\text{out},p}$. This mathematical procedure reflects the possibility of experimentally restricting detection to photons in compressed modes (for other examples involving nonnormalized density matrices, see, e.g., [23]). Accordingly, we define the efficiency ϵ of the compression as

$$\epsilon = \text{tr}(\rho_{\text{out},p}). \quad (1)$$

The index p denotes “photon” and is added henceforth to avoid confusion of photonic with atomic states. Furthermore, we use the following definition of the fidelity F :

$$F = {}_p \langle \psi_{\text{in}} | \rho'_{\text{out},p} | \psi_{\text{in}} \rangle_p, \quad (2)$$

where $\rho'_{\text{out},p}$ is the renormalized and unitarily transformed density matrix

$$\rho'_{\text{out},p} = \frac{1}{\epsilon} U \rho_{\text{out},p} U^\dagger. \quad (3)$$

Note that we allowed for deterministic unitary operations U composed of rotations around σ_x , σ_y , or σ_z in the qubit Hilbert space to maximize the fidelity. This is similar to quantum teleportation [24], where, depending on the result of the Bell state measurement, bit flip, phase flip, or bit and phase flip operations have to be applied to the teleported state to recover the initial state.

Finally, in order to characterize the usefulness of quantum compression in view of enhanced qubit transmission rate, we

define the gain as

$$G = \epsilon \eta. \quad (4)$$

This reflects that the detection rate of qubits increases linearly with both the compression efficiency ϵ as well as the factor η by which a qubit can be compressed, that is, the number of qubits per unit time.

III. PHOTONIC TIME-BIN QUBITS

We are concerned with encoding of quantum information into qubits, that is, quantum states that are described by

$$|\psi\rangle = \alpha|0\rangle + e^{i\phi}\beta|1\rangle. \quad (5)$$

The coefficients α , β , and ϕ are real, $\alpha^2 + \beta^2 = 1$, and the kets $|0\rangle$ and $|1\rangle$ form an orthogonal basis in a two-dimensional Hilbert space ($\langle i | j \rangle = \delta_{ij}$, $i, j = [0, 1]$).

In this article, we are specifically interested in so-called *time-bin* qubits, where the basis states $|0\rangle$ and $|1\rangle$ describe photon wave packets localized at a particular position z at early and late times $t = z/c$ and $t' = z/c + \tau_o$, respectively. Differently stated, at a given time t , the photon is in a superposition of being at positions z and $z' = z - z_o$, with $z_o = c\tau_o$ (see Fig. 1). We use these two pictures interchangeably in our analysis. Time-bin qubits have been shown to be very well suited for quantum communication over telecommunication fibers [25,26]. In the following, we derive a description of time-bin qubits using a physical representation of the abstract qubit Hilbert space. We limit ourselves to one polarization mode.

A general photonic wave packet is described in *momentum space* as [27,28]

$$|\psi(t)\rangle_p = \int_{-\infty}^{\infty} dk \tilde{g}(k, t) \hat{a}_k^\dagger |\emptyset\rangle, \quad (6)$$

where \hat{a}_k^\dagger (\hat{a}_k) denotes the creation (annihilation) operator for a photon with wave vector k , $[\hat{a}_k, \hat{a}_{k'}^\dagger] = \delta(k - k')$, and $|\emptyset\rangle$ is the vacuum state of light. For a more suitable description of our qubit states, we define operators creating forward- and backward-propagating photons at a particular position:

$$\hat{a}_\sigma^\dagger(z) = \sqrt{1/2\pi} \int dk \hat{a}_{k+\sigma k_o}^\dagger \exp\{-i(\kappa + \sigma k_o)z\}, \quad (7)$$




FIG. 1. (Color online) Schematic representation of a photonic time-bin qubit $|\psi\rangle = \alpha|0\rangle + \beta e^{i\phi}|1\rangle$, where α and β are the amplitudes of photon wave packets propagating along the $+z$ axis with relative phase ϕ ; $\delta z = c\delta t$, and $z_o = c\tau_o$. δt and τ_o denote the temporal duration of the wave packets and the relative time delay, respectively.

where, $k_o = \omega_{eg}/c$, $\sigma = +$ denotes forward (f)-propagating light modes, $\sigma = -$ backward (b)-propagating modes, and $[\hat{a}_{\sigma'}(z'), \hat{a}_{\sigma}^{\dagger}(z)] \cong \delta_{\sigma',\sigma} \delta(z' - z)$.

The associated annihilation operators are defined by $\hat{a}_{\sigma}(z) = [\hat{a}_{\sigma}^{\dagger}(z)]^{\dagger}$, satisfying the usual bosonic commutation relations. With these new operators, the basis states of a time-bin qubit can be expressed via the center $z = \sigma ct$ and $z - z_o = \sigma c(t - \tau_o)$ of the wave packets, respectively, their extension δz and carrier frequency ω_{σ} :

$$\begin{aligned} |0\rangle_{\sigma,p} &= \int_{-\infty}^{\infty} dz' \exp\{-i\omega_{\sigma}(t - \sigma z'/c)\} \\ &\quad \times g(ct - \sigma z', \delta z) \hat{a}_{\sigma}^{\dagger}(z') |\emptyset\rangle, \\ &\equiv |t, \delta z, \omega_{\sigma}\rangle_{\sigma,p}, \end{aligned} \quad (8)$$

$$\begin{aligned} |1\rangle_{\sigma,p} &= \int_{-\infty}^{\infty} dz' \exp\{-i\omega_{\sigma}(t - \tau_o - \sigma z'/c)\} \\ &\quad \times g(ct - c\tau_o - \sigma z', \delta z) \hat{a}_{\sigma}^{\dagger}(z') |\emptyset\rangle, \\ &\equiv |t - \tau_o, \delta z, \omega_{\sigma}\rangle_{\sigma,p}, \end{aligned} \quad (9)$$

where $g(ct - z', \delta z)$ is the normalized envelope of the photon wave packet with length δz , which is related to its spectral function $\bar{g}(k, \delta k)$ by Fourier transformation: $g(z, \delta z) = \sqrt{1/2\pi} \int_{-\infty}^{\infty} dk \bar{g}(k, \delta k) \exp\{-ikz\}$. Note that the identification of the states $|t, \delta z, \omega_{\sigma}\rangle_{\sigma,p}$ and $|t - \tau_o, \delta z, \omega_{\sigma}\rangle_{\sigma,p}$ with $|0\rangle_{\sigma}$ and $|1\rangle_{\sigma}$, respectively, requires $\delta z \ll z_o = c\tau_o$ so that the states are orthogonal and normalized. Using this notation, the general photonic wave packet can then be expressed as

$$\begin{aligned} |\psi(t)\rangle_p &= |\psi(t)\rangle_{f,p} + |\psi(t)\rangle_{b,p} \\ &= \sqrt{1/2\pi} \sum_{\sigma} \int dz' A_{\sigma}(t, z') \\ &\quad \times \exp\{-i\omega_{\sigma}(t - \sigma z'/c)\} \hat{a}_{\sigma}^{\dagger}(z') |\emptyset\rangle. \end{aligned} \quad (10)$$

In the case of the initial, forward-propagating time-bin qubit, the amplitude A_+ takes on the form

$$\begin{aligned} A_+(t \rightarrow -\infty, z) &= \sqrt{2\pi} \{\alpha g(ct - z, \delta z) + \exp\{i\omega_+ \tau_o + i\phi\} \\ &\quad \times \beta g(ct - z - c\tau_o, \delta z)\} \end{aligned} \quad (11)$$

and $A_-(t \rightarrow -\infty, z) = 0$, with $\int_{-\infty}^{\infty} dz |A_{\sigma=\pm}(t \rightarrow -\infty, z)|^2 = 1$, identifying $|A_{\sigma=\pm}(t, z)|^2 dz$ as the probability of detecting a photon at time t within the space coordinates z and $z + dz$. Hence,

$$|\psi_{\text{in}}(t)\rangle_p = \alpha |t, \delta z, \omega_+\rangle_{f,p} + e^{i\phi} \beta |t - \tau_o, \delta z, \omega_+\rangle_{f,p}. \quad (12)$$

A schematical representation of a time-bin qubit assuming Gaussian shapes of the basis wave packets is given in Fig. 1.

IV. QUANTUM-STATE STORAGE BASED ON SYMMETRIC CRIB

An extensively investigated approach to quantum-state storage is based on CRIB of an isolated absorption line [6–10]. It has recently led to storage efficiencies up to 69% [29], and the possibility for random access quantum memory has been demonstrated [17]. In the ideal, standard CRIB scheme, the optical input data is launched along the forward (+)z

direction into a resonant, optically thick atomic medium ($\alpha L \rightarrow \infty$, where α is the resonant absorption coefficient and L is the length of the atomic medium). The spectral width $\delta\omega_{\text{in}}$ of the input light should be narrow in comparison with the inhomogeneous broadening Δ_{inh} of the resonant atomic transition $1 \leftrightarrow 2$, that is, $\delta\omega_{\text{in}} < \Delta_{\text{inh}}$, and larger than the homogeneous line width γ_{eg} of the optical transition, that is, $\delta\omega_{\text{in}} > \gamma_{eg}$.

All atoms are initially prepared in the pure state $|g\rangle = \prod_{j=1}^N \otimes |g\rangle_j$ [where $|g\rangle_j$ is a long-lived (ground) state of the j th atom]. Thus the initial light-atom state, denoted using the capital letter Ψ , (for $t \rightarrow -\infty$) is

$$|\Psi(t \rightarrow -\infty)\rangle_f = |\psi_{\text{in}}(t)\rangle_{f,p} \otimes |g\rangle. \quad (13)$$

After absorption of the light, the atomic coherences acquire phase factors $\exp\{-i(\Delta_j + \omega_{eg})t\}$, where Δ_j is the detuning of atom j as compared to the central absorption frequency ω_{eg} , leading to fast dephasing of the excited collective atomic coherence. The absorption process can be time reversed by inverting the atomic detunings at time t_1 (i.e., $\Delta_j \rightarrow -\Delta_j$) and applying a position-dependent phase shift of $2kz$. This leads to rephasing of the collective atomic coherence, hence irradiation of an echo signal at time t_{echo} in backward direction as a perfect time-reversed copy of the input data [6]. This reversibility is based on a hidden symmetry of the equations describing the evolution of the slowly varying light and atomic parameters during storage and retrieval [9].

Analyzing the evolution of the complete wave function (including the fast, time varying part $\exp\{-ick_{eg}t\}$, where ck_{eg} is the central frequency of the atomic transition), it was shown [30] that the photon operators (in momentum space) of the input (*forward*) and output (*backward*) light fields are related by $\exp\{-ic(k + k_{eg})t\} \hat{a}_{k+k_{eg}}^{\dagger} \rightarrow -\exp\{ic(k - k_{eg})(t - t_{\text{echo}})\} \hat{a}_{k-k_{eg}}^{\dagger}$. As we show in more detail in Appendix A, this results in input photonic time-bin qubit states [Eq. (12)] to emerge as

$$\begin{aligned} |\psi_{\text{out}}(t - t_{\text{echo}})\rangle_{b,p} &= -[\alpha |t, \delta z, \omega_-\rangle_{b,p} + e^{i(\phi + 2\omega_{eg}\tau_o)} \\ &\quad \times \beta |t + \tau_o, \delta z, \omega_-\rangle_{b,p}], \end{aligned} \quad (14)$$

where we have ignored a global phase shift and where $\omega_- = 2\omega_{eg} - \omega_+$ denotes the new carrier frequency. Hence, the recalled qubit state is associated with an exchange of the leading and trailing wave packets, and the now leading wave packet (with probability amplitude proportional to β) has acquired an additional phase that depends on the time delay τ_o between the wave packets and the atomic transition frequency ω_{eg} (see also [31]). Please note that not only does the order of the wave packets change, but each wave packet's temporal envelope is also time-reversed. For simplicity of notation, we henceforth restrict our investigation to time-symmetric, Gaussian-shaped basis wave packets described by $g(ct - z, \delta z) = \sqrt{2/[\pi(\delta z)^2]} \exp\{-(c\tau/\delta z)^2\}$, where $\tau = t - z/c$.

Returning to the abstract qubit notation, we find that the initial state $|\psi_{\text{in}}\rangle = \alpha |0\rangle + e^{i\phi} \beta |1\rangle$ is transformed in the quantum memory into $|\psi_{\text{out}}\rangle = \alpha |1\rangle + e^{i(\phi + 2\omega_{eg}\tau_o)} \beta |0\rangle$, that

is, that the two states are related by a deterministic unitary transformation T ,

$$|\psi_{\text{out}}\rangle = T|\psi_{\text{in}}\rangle = e^{i\omega_{eg}\tau_o}\sigma_x R_z(\theta)|\psi_{\text{in}}\rangle, \quad (15)$$

where $R_z(\theta) = e^{i\frac{\theta}{2}\sigma_z}$ denotes a rotation of $\theta = 2\omega_{eg}\tau_o$ around σ_z , and σ_x is the bit-flip operator.

Hence, for the ideal standard CRIB protocol, we find the efficiency to be one. Furthermore, from Eq. (3), we find $U = T^{-1}$; hence, $\rho'_{\text{out},p} = \rho_{\text{in},p}$ and thus $F = 1$. Please note that two subsequent storage sequences lead to compensation of the bit flip as well as the additional phase factor of $2\omega_{eg}\tau_o$.

Finally, being obvious in the case of “no compression” and unity efficiency, we note that the gain G is one.

V. QUANTUM COMPRESSION BASED ON GENERALIZED CRIB

In the CRIB protocol described previously, the efficiency, fidelity, and gain can be derived using arguments stemming from symmetries in the equations of motion. Relaxing the perfect reversibility of atomic detunings by introducing a more general relation between the initial ($t < t_1$) and output ($t > t_1$) spectral detunings,

$$\Delta_j(t > t_1) = -\eta\Delta_j(t < t_1), \quad (16)$$

with η being the compression parameter, this symmetry-based approach is no longer possible. Here we pursue the following approach: We start with a general photonic qubit state [Eq. (5)] described in the abstract two-dimensional Hilbert space. We then express the qubit state using a physical description in position space [Eqs. (8), (9), and (12), respectively]. Using the equations of motion, which we introduce in the following section, we then calculate the state of the combined atom-photon system after quantum compression. From Eq. (14), we find the single-photon density matrix. Redefining new basis states $|0\rangle$ and $|1\rangle$, determined by the compression parameter, we then express the output state in qubit state notation, which finally allows us to calculate our figures of merit [Eqs. (1), (2), and (4)] and the unitary operation U [see Eq. (3)].

A. Basic equations

First, we write the field Hamiltonian containing forward- and backward-propagating light modes as

$$\hat{H}_{ph} = \hbar \int dk \omega_k \hat{a}_k^\dagger \hat{a}_k = \hbar \sum_{\sigma=\pm} \int d\kappa |\kappa + \sigma k\rangle \hat{a}_{\kappa+\sigma k}^\dagger \hat{a}_{\kappa+\sigma k} \quad (17)$$

$$= \hat{H}_{ph}^o + \sum_{\sigma=\pm} \delta \hat{H}_{ph,\sigma}. \quad (18)$$

Using Eq. (7) and $\hat{a}_{\kappa+\sigma k}^\dagger = \sqrt{1/2\pi} \int dz \hat{a}_\sigma^\dagger(z) \exp\{i(\kappa + \sigma k)z\}$, we find

$$\begin{aligned} \hat{H}_{ph}^o &= \hbar k_o \sum_{\sigma=\pm} \int d\kappa \hat{a}_{\kappa+\sigma k}^\dagger \hat{a}_{\kappa+\sigma k} \\ &= \hbar k_o \sum_{\sigma=\pm} \int dz \hat{a}_\sigma^\dagger(z) \hat{a}_\sigma(z) \end{aligned} \quad (19)$$

and

$$\begin{aligned} \delta \hat{H}_{ph,\sigma} &= \sigma \hbar \int d\kappa \kappa \hat{a}_{\kappa+\sigma k}^\dagger \hat{a}_{\kappa+\sigma k} \\ &= -i\sigma \hbar \int dz \hat{a}_\sigma^\dagger(z) \frac{\partial}{\partial z} \hat{a}_\sigma(z). \end{aligned} \quad (20)$$

This allows us to write the total Hamiltonian, describing the atom-light interaction [27,28,30], as

$$\hat{H}(t) = \hat{H}_0 + \hat{H}_1(t), \quad (21)$$

where

$$\hat{H}_0 = \hbar \omega_{eg} \left[\sum_{\sigma=\pm} \int_{-\infty}^{\infty} dz \hat{a}_\sigma^\dagger(z) \hat{a}_\sigma(z) + \sum_{j=1}^N P_{ee}^j \right], \quad (22)$$

and

$$\begin{aligned} \hat{H}_1(t) &= -\hbar g \sqrt{2\pi} \sum_{j=1}^N \{ [\hat{a}_f(z_j) + \hat{a}_b(z_j)] \hat{P}_{eg}^j + \text{H.c.} \} \\ &\quad + \sum_{j=1}^N [\hbar \Delta_j(t) + \delta E_{eg}^j(t)] \hat{P}_{ee}^j \\ &\quad - i\hbar c \sum_{\sigma=\pm} \sigma \int_{-\infty}^{\infty} dz \hat{a}_\sigma^\dagger(z) \frac{\partial}{\partial z} \hat{a}_\sigma(z). \end{aligned} \quad (23)$$

\hat{H}_0 describes the total number of excitations in the system and commutes with the total Hamiltonian \hat{H} . The first term in $\hat{H}_1(t)$ describes the atom-field interaction with atomic operator $\hat{P}_{mn}^j = |m\rangle_{jj}\langle n|$. N denotes the number of atoms, $g(\omega) \cong g = id_{ge}(\frac{\omega_{eg}}{2\hbar\epsilon_0 S})^{1/2}$, d_{12} is the dipole moment of the atomic transition $|g\rangle \leftrightarrow |e\rangle$, S is the cross section of the light beams, ϵ_0 is the electric permittivity, and \hbar is Planck's constant divided by 2π . The second term in Eq. (23) describes inhomogeneous and homogeneous atomic line broadening within a unitary approach to quantum evolution. Indeed, it has been shown [32,33] that the decay rate of the atomic coherence γ_{eg} can be calculated by a statistical average over local stochastic phase fluctuations $\delta\phi_{eg}^j(t,t') = \int_{t'}^t dt'' \delta E_{eg}^j(t)/\hbar$ of the atomic transition,

$$\langle \exp\{-i\delta\phi_{eg}^j(t,t')\} \rangle = \exp\{-\gamma_{eg}(t-t')\}, \quad (24)$$

where $\langle \delta\phi_{eg}^j(t,t') \rangle = 0$. These fluctuations are due to interaction between the absorbers and its environment. Finally, the third term of $\hat{H}_1(t)$ arises from the fact that the here treated, localized, propagating photonic wave packet features a broad spectrum (see earlier discussion and [34]).

Using the total Hamiltonian (21), the initial photon-atom state (13) will evolve into

$$|\Psi(t)\rangle = |\Psi(t)\rangle_p + |\Psi(t)\rangle_a, \quad (25)$$

$$|\Psi(t)\rangle_p = |\psi(t)\rangle_p \otimes |g\rangle \quad (26)$$

[with $|\psi(t)\rangle_p$ as defined in Eq. (6)], and

$$|\Psi(t)\rangle_a = \sum_{j=1}^N b_j(t) \hat{P}_{eg}^j |g\rangle \otimes |\emptyset\rangle, \quad (27)$$

with $b_j(t)$ being the probability amplitude for atom j to be in the excited state. Initially, $b_j(t \rightarrow -\infty) = 0$, that is, all atoms

are in the ground state, and $A_-(t, \rightarrow -\infty, z) = 0$; that is, all backward modes are empty.

In the Schrödinger equation with the Hamiltonian introduced in [Eq. (21)], the quantum evolution during absorption of the forward-propagating light field ($\sigma = +$ for $t < t_1$) and for re-emission of the backward-propagating light field ($\sigma = -$ for $t > t_1$) is given by

$$\left(\frac{1}{c}\frac{\partial}{\partial t} + \sigma\frac{\partial}{\partial z}\right)A_\sigma(t, z) = i(\pi n_o S g^*/c)\exp\{i\omega_\sigma(t - \sigma z/c)\}b(t, z), \quad (28)$$

$$\begin{aligned} \frac{\partial}{\partial t}b_j(t) &= -i\{\omega_{eg} + \Delta_j(t) + \delta\Delta_{eg}^j(t)\}b_j(t) \\ &+ igA_\sigma(t, z_j)\exp\{-i\omega_\sigma(t - \sigma z_j/c)\}, \end{aligned} \quad (29)$$

where $b(t, z) = (n_o S)^{-1} \sum_{j=1}^N b_j(t)\delta(z - z_j)$ is a collective atomic variable describing the averaged, position-dependent atomic coherence, and n_o is the atomic density. For an explicit derivation of these formulas, see [35,36]. Using these equations, we now evaluate quantum compression of photonic time-bin qubits in atomic systems with *transverse* and *longitudinal* inhomogeneous broadening.

B. Transverse broadened media

In *transverse* broadened media, the atomic absorption profile is independent of the spatial coordinate z measured along the propagation direction of the light. In rare-earth-ion-doped inorganic crystals (RE crystals) with low site symmetry and sufficiently narrow absorption line, *controlled transverse broadening* can be introduced through the Stark effect [37] by applying an electric field gradient *transverse* to the propagation direction of light. For RE ions implemented in a glassy host, for example, an optical fiber, controlled transverse broadening is already obtained for uniform electric fields, due to the random orientation and magnitude of the electric dipole moments of the RE ions [38]. In the following, we assume a broadened medium with Lorentzian lineshape $G(\Delta/\Delta_{\text{inh}}) = \Delta_{\text{inh}}/[(\Delta_{\text{inh}}^2 + \Delta^2)\pi]$ with inhomogeneous spectral width $\Delta_{\text{inh}} \gg c\delta k$. The center of the first wave packet $|0\rangle_{f,p}$ enters the atomic medium at $t_0 = 0$ (see Fig. 2). During the interaction with the medium, the light field is partially or completely absorbed, depending on the optical depth. At time t_1 , after sufficient dephasing, we change the detuning Δ_j to $-\eta\Delta_j$, and we also apply a position-dependent phase shift of $(2k_{eg} - \delta k)z$, which allows phase matching for the retrieval of the light field in the backward direction [8,9,35,39]. δk describes a small deviation from the perfect phase matching. This leads to re-emission of the light field, with the center of the (now trailing) wave packet exiting the medium at time $t = t_{\text{echo}}$. The atom-light state for $t \gg t_{\text{echo}}$ is then given by

$$|\Psi(t \gg t_{\text{echo}})\rangle = |\psi_f(t)\rangle_p \otimes |g\rangle + |\Psi(t)\rangle_a + |\psi_b(t)\rangle_p \otimes |g\rangle. \quad (30)$$

As visualized in Figs. 2(c), 2(d), and 2(e), the first term describes the damped, nonabsorbed photonic component $|\psi_f(t)\rangle_p = \exp\{-\alpha_o L/2\}|\psi_{\text{in}}(t)\rangle_p$ that continued to travel in the forward direction behind the atomic medium, the second




FIG. 2. (Color online) Schematics showing different instances in the quantum compression protocol in the case of transverse broadening. The figure depicts forward- and backward-propagating photonic wave packets and atomic excitation.

term the remaining atomic excitation, and the third term denotes the now backward-moving, retrieved photon field, which is the subject of our investigation. α_o is the on-resonant absorption coefficient, L the length of the medium, and $\alpha_o L$ the optical depth. As shown in Appendix A, the amplitude of the retrieved photon field, assuming for simplicity a Gaussian spectral shape, is described in abstract qubit notation as:

$$|\psi_b(t)\rangle_p^{(t)} = R^{(t)}\{e^{-(1+1/\eta)\gamma_{eg}\tau_o}\alpha|t, \delta z', \omega_-\rangle_b + e^{i\phi'}\beta|t + \tau'_o, \delta z', \omega_-\rangle_b\}, \quad (31)$$

where the superscript (t) denotes transverse broadening, where we ignored a global phase shift, and where $\phi' = \phi + (1 + 1/\eta)\omega_{eg}\tau_o$. Furthermore,

$$|t^{(m)}, \delta z', \omega_-\rangle_b = \sqrt{\eta} \int_{-\infty}^{\infty} dz' \exp\{-i\omega_\sigma[t^{(m)} - t_{\text{echo}} + z'/c]\} \times g[ct^{(m)} - ct_{\text{echo}} + z', \delta z']\hat{a}_b^\dagger(z')|\emptyset\rangle, \quad (32)$$

where $t^{(m)}$ denotes the early or late wave packet ($t^{(m)} \in [t, t + \tau'_o]$), $t_{\text{echo}} = (1 + 1/\eta)t_1 - \delta t_R$, $\delta t_R = \frac{1}{2}\gamma_{eg}[1 + \eta](\delta z'/c)^2$, $\tau'_o = \tau_o/\eta$, and $\delta z' = \delta z/\eta$. The factor

$$R^{(t)} = \Gamma_{\gamma_{eg}}(t_1)M^{(t)}(\delta k, \alpha_o L)\epsilon_o^{1/2}(\eta) \quad (33)$$

combines different sources that affect the efficiency of the quantum compression through phase relaxation γ_{eg} , phase mismatch δk , and limited optical depth $\alpha_o L$. Its upper limit, given by ϵ_o (not to be confused with the electric permittivity ϵ_o), is determined by the compression parameter η . Specifically, we find

$$\Gamma_{\gamma_{eg}}(t_1) = \exp\left\{-(1 + 1/\eta)\gamma_{eg}\left(t_1 - \tau_o - \frac{1}{2}\eta\delta t_R\right)\right\}, \quad (34)$$

$$M^{(t)}(\delta k, \alpha_o L) = \frac{1 - \exp\left\{-\frac{1}{2}(1 + 1/\eta)\alpha_o L + i\delta k L\right\}}{1 - 2i(\delta k/\alpha_o)\eta/(\eta + 1)}, \quad (35)$$

$$\epsilon_o^{1/2}(\eta) = \frac{2\sqrt{\eta}}{(\eta + 1)}. \quad (36)$$

Let us now relate Eq. (31) with our figures of merit. First, it is important to note that Eq. (31) describes again a




FIG. 3. (Color online) Recall efficiency for transverse broadening as a function of the compression parameter η and the optical depth $\alpha_o L$ [see Eq. (39)].

photonic time-bin qubit with modified separation and width of basis states. Indeed, comparing the new with the old basis states [Eq. (12)], we find that the separation between the two basis wave packets has changed from z_o to $z'_o = z_o/\eta$, and their extension from δz to $\delta z' = \delta z/\eta$, while the wave-packet amplitude has changed $\propto \eta^{1/2}$, as visualized in Figs. 2(d) and 2(e).

Second, it is important to realize that the quantum information encoded into a qubit is independent of the physical realization of its abstract basis states $|0\rangle$ and $|1\rangle$. This allows us to relabel the basis states after compression, that is, $|t, \delta z', \omega_- \rangle \rightarrow |1\rangle$, and $|t + \tau'_o, \delta z', \omega_- \rangle \rightarrow |0\rangle$, resulting in

$$|\psi\rangle_p = R^{(t)}[e^{-(1+1/\eta)\gamma_{eg}\tau_o}|1\rangle + e^{i\phi'}\beta|0\rangle]. \quad (37)$$

Now, taking the trace of $\rho_{\text{out},p} = |\psi\rangle_{pp}\langle\psi|$, we find the efficiency according to Eq. (1):

$$\epsilon^{(t)} = [\Gamma_{\gamma_{eg}}(t_1)]^2 |M^{(t)}(\delta k, \alpha_o L)|^2 \epsilon_o(\eta) [\alpha^2 e^{-2(1+1/\eta)\gamma_{eg}\tau_o} + \beta^2]. \quad (38)$$

We now discuss our figures of merit for some particular cases. Ignoring phase relaxation (i.e., $\gamma_{eg} = 0$) and assuming perfect phase matching (i.e., $\delta k = 0$) we find the efficiency to be given by

$$\epsilon^{(t)} = \frac{4\eta}{(\eta+1)^2} \left[1 - \exp\left\{-\frac{1}{2}(1+1/\eta)\alpha_o L\right\} \right]^2. \quad (39)$$

It is depicted in Fig. 3 as a function of the optical depth $\alpha_o L$ and the compression parameter η . First, we note that for standard CRIB (without compression, i.e., $\eta = 1$), we find the previously published result $\epsilon^{(t)} = (1 - \exp\{-\alpha_o L\})^2$ [35,36]. Second, we point out that for a given compression factor, the maximum efficiency is always obtained for infinite optical depth, and hence the upper limit ϵ_{max} is only determined by the compression factor:

$$\epsilon_{\text{max}}^{(t)}(\eta) = \frac{4\eta}{(1+\eta)^2}. \quad (40)$$

To give an example, a quantum compression with $3 - 2\sqrt{2} \leq \eta \leq 3 + 2\sqrt{2}$ results in a recall efficiency limited to 50%.

To assess the fidelity, we compare Eq. (37) with the initial qubit state in Eq. (5). We find that the renormalized,




FIG. 4. (Color online) Gain for transverse broadening as a function of the compression parameter η and the optical depth $\alpha_o L$.

compressed qubit state $|\psi\rangle_p$ can be unitarily transformed into the initial state by a σ_z rotation of angle $\phi' - \phi = (1 + 1/\eta)\omega_{eg}\tau_o$ and a bit-flip operation σ_x . Furthermore, one should take into account an additional σ_x rotation to compensate for the amplitude reduction factor $\exp\{-(1 + 1/\eta)\gamma_{eg}\tau_o\}$ arising in the case of significant phase relaxation during time $(1 + 1/\eta)\tau_o$. This immediately implies that the fidelity as defined in Eq. (2) is always one, regardless the compression factor, phase relaxation, or phase mismatch. Note that the fact that atomic phase relaxation does not affect the fidelity in time qubit storage has been inferred from stimulated photon echo experiments with intense light pulses [40].

Figure 4 depicts the gain as a function of optical depth and compression parameter for transverse broadened media. As an example, assuming $\alpha_o L = 2$, we find an increased communication rate, that is, $G^{(t)} > 1$, for $\eta \gtrsim 1.7$. For large (infinite) optical depth [i.e., maximum efficiency as described by Eq. (40)], the gain is upper bounded by

$$G^{(t)}(\alpha L \rightarrow \infty) = 4\eta^2/(1+\eta)^2|_{\eta \gg 1} = 4, \quad (41)$$

a modest, yet significant improvement over quantum communication schemes without compression.

To finish this section, let us briefly inspect Eqs. (33)–(36) in view of symmetry between compression and decompression, that is, under exchange of η to $1/\eta$. First, we recall that the fidelity is one, regardless the compression factor: $F(\eta) = F(1/\eta) = 1$. Second, we find that the upper limit of the efficiency is symmetric with respect to compression or decompression: $\epsilon_{\text{max}}^{(t)}(\eta) = \epsilon_{\text{max}}^{(t)}(1/\eta) < 1$. However, limited optical depth, phase mismatch, and atomic phase relaxation break the symmetry in the efficiency. It should be possible to demonstrate this surprising behavior using only symmetry arguments and the equations of motion describing the recall efficiency, without having to solve the equations. This is likely to lead to a more profound understanding of the physical principles of the here studied compression or decompression protocol.

C. Longitudinally broadened media

Longitudinal broadening is an inhomogeneous broadening that cannot be found naturally. In this case, for each position z in the medium, the atomic absorption profile is given by a narrow line (here for simplicity assumed to be homogeneously




FIG. 5. (Color online) Schematic showing different instances in the quantum compression protocol in the case of longitudinal broadening. The figure depicts forward- and backward-propagating photonic wave packets and atomic excitation. Note the difference of the re-emission time t_{echo} and the different localization of atomic excitation compared to the transverse case (see Fig. 2). The change of color of the irradiated wave packets denotes additional frequency shifts [see Eq. (49)].

broadenend), whose center frequency varies monotonously with z . Longitudinal broadening can be realized through the Stark effect in non-centro-symmetric RE-ion-doped inorganic crystals by applying a electric field gradient *longitudinal* to the propagation direction of light [7]. In the following, we assume absorption lines whose detuning with respect to the light carrier frequency varies linearly with position z within the crystal, which extends from $z = -L/2$ to $z = +L/2$:

$$\Delta = -\chi z. \quad (42)$$

The initial inhomogeneous absorption line, obtained after integration over all atomic positions z , is assumed to be broad compared to the spectrum of the photonic wave packet: $\Delta_{\text{inh}} = \chi L \gg c\delta k$. A schematic of the compression procedure depicting relevant instances of the protocol is given in Fig. 5.

A similar approach as in the case of transverse broadening leads to the following wave function for the backward-emitted light field (for more details, see Appendix B):

$$|\psi_b(t)\rangle_p^{(l)} = R^{(l)} \left\{ e^{-(1+1/\eta)\gamma_{\text{eg}}\tau_o} \alpha |t, t - t_{\delta k}^{(l)}, \delta z', \omega_{-}\rangle_b + e^{i\phi'} \beta |t, t - t_{\delta k}^{(l)} + \tau_o', \delta z', \omega_{-}\rangle_b \right\}, \quad (43)$$

where the superscript “(l)” denotes longitudinal broadening and where we ignored again a global phase shift. Furthermore,

$$|t, t^{(m)}, \delta z', \omega_{-}\rangle_b = \sqrt{\eta} \int_{-\infty}^{\infty} dz' \exp\{i\Phi(t + z'/c, \eta)\} \times \exp\{-i\omega_{-}[t^{(m)} - t_{\text{echo}} + z'/c]\} \times g[ct^{(m)} - ct_{\text{echo}} + z', \delta z'] \hat{a}_b^{\dagger}(z') |\emptyset\rangle, \quad (44)$$

and $t^{(m)} \in [t - t_{\delta k}^{(l)} + \tau_o', t - t_{\delta k}^{(l)}]$, with

$$t_{\delta k}^{(l)} = \frac{1}{\eta \Delta_{\text{inh}}} [\delta k L + 2(\zeta/\chi)(1 - 1/\eta)]. \quad (45)$$

The parameters $\delta z' = \delta z/\eta$, $\tau_o' = \tau_o/\eta$, and ϕ' are the same as in the transverse case, and $\zeta = \pi n_o S g^2/c$. Furthermore,

$$R^{(l)} = \Gamma_{\gamma_{\text{eg}}} [t_1 + t_{\delta k}^{(l)}] M^{(l)}(\zeta/\chi, \eta), \quad (46)$$

with

$$M^{(l)}(\zeta/\chi, \eta) = \sqrt{\left(1 - \exp\left\{-\frac{2\pi\zeta}{\eta\chi}\right\}\right) \left(1 - \exp\left\{-\frac{2\pi\zeta}{\chi}\right\}\right)}. \quad (47)$$

Finally, the time-dependent phase $\Phi(\tau = t + z/c, \eta)$ is given by

$$\Phi(\tau, \eta) = \frac{\zeta}{\chi} \ln \left\{ \frac{\left(\frac{1}{2}\eta\Delta_{\text{inh}}\left|\tau - t_1 + \tau_m' - \frac{\delta k L}{\eta\Delta_{\text{inh}}}\right|\right)}{\left[\frac{1}{2}\eta\Delta_{\text{inh}}(\tau - t_1 + \tau_m')\right]^{1/\eta}} \right\}, \quad (48)$$

where $\tau_m' = \tau_m/\eta^2$ and $\tau_m = 2\frac{\zeta}{\chi\Delta_{\text{inh}}}$. For time-bin qubits with sufficiently narrow basis wave packets, the phase change (which is nonlinear in τ and η) translates into a different frequency change of each wave packet ($\delta\omega_0, \delta\omega_1$) as compared to the carrier frequency. From $\delta\omega(t) = d\Phi(t)/dt$ we find

$$\delta\omega_0 = \frac{\zeta}{\chi} \left[\frac{1}{t_1 - \tau_o + \tau_m + \delta k/\chi} - \frac{\eta}{t_1 - \tau_o + \tau_m} \right], \quad (49)$$

$$\delta\omega_1 = \frac{\zeta}{\chi} \left[\frac{1}{t_1 + \tau_m + \delta k/\chi} - \frac{\eta}{t_1 + \tau_m} \right].$$

Note that a given dephasing time \tilde{t} determines close-to-homogeneously broadened slices of length δl in the atomic medium with $\delta l \approx (\chi\tilde{t})^{-1}$, where coherence, hence radiation, remains. We believe that coupling between light and collective atomic coherence (as included in the analyzed equations) within these slices plays an important role for those frequency shifts, which are proportional to the number of atoms $n = N\delta l$ in each slice, and the square of the photon-atom coupling constant g (i.e., $\delta\omega \propto g^2 N\delta l$). The nonlinear phase modulation also leads to an additional, possibly substantial, phase difference between the two basic wave packets after recall: $\delta\varphi_{01} = \int_{t_{\text{echo}}}^{t_{\text{echo}} + \tau_o/\eta} dt \delta\omega(t) = \Phi(t_{\text{echo}} + \tau_o/\eta) - \Phi(t_{\text{echo}})$. The phase difference is depicted in Figs. 6 and 7. The effects of the nonlinear phase shift are enhanced with increasing effective optical depth $\kappa_{\text{eff}} = 2\pi\zeta/\chi$ and time delay τ_o between the wave packets and diminishes as the dephasing time t_1 increases and the compression parameter η approaches unity.




FIG. 6. (Color online) Phase shift $\delta\varphi_{01}$ for quantum compression as a function of the time delay τ_o and the compression parameter η , with dephasing time $t_1 = 20\delta t$, $\Delta_{\text{inh}}\delta t = 10$, and $2\pi\zeta/\chi = 6\pi$. δt denotes the temporal duration of the initial-basis wave packets, and $2\pi\zeta/\chi$ is the effective optical depth.




FIG. 7. (Color online) Phase shift $\delta\varphi_{01}$ for quantum decompression as a function of the time delay τ_o and the compression parameter η , with dephasing time $t_1 = 20\delta t$, $\Delta_{\text{inh}}\delta t = 10$, and $2\pi\zeta/\chi = 6\pi$.

Phase and frequency shifts in the here-analyzed case of backward emission are a result of the presence of phase mismatch [$\delta k^{(l)} \neq 0$], or compression or decompression ($\eta \neq 1$), or both, but they do not occur in the phase-matched, reversible case. However, in the case of forward emission, the phase and frequency shifts also arise for noncompressed recall ($\eta = 1$), provided the rephasing and dephasing times \tilde{t} are bounded [41,42].

Let us now discuss our figures of merit in the case of longitudinal broadening and compare it with transverse broadening. As mentioned earlier, the new basis states [Eqs. (44)] do not transfer into each other through time translation alone, in contrast to the case of transverse broadening. For instance, assuming $\delta t = 100$ ns, $\tau_o = 200$ ns, $2\pi\zeta/\chi = 6$, $\eta = 3$, and a sufficiently large dephasing time $t_1/\delta t = 20$, we find a frequency difference $|\delta\omega_0 - \delta\omega_1| \approx 2\pi \times 17.32$ kHz. As this value is small compared to the spectral width of the wave packet $c\delta k \approx 2\pi \times 1.59$ MHz, we can safely ignore this effect as a limitation to the fidelity. It is thus possible to map the new basis states onto the qubit basis states $|0\rangle$ and $|1\rangle$. Furthermore, as before, we can compensate for the additional relative phase arising during compression using an appropriate σ_z rotation. Hence, the fidelity in longitudinal compression can be close to one.

The recall efficiency $\epsilon^{(l)}$ is given by

$$\epsilon^{(l)} = [R^{(l)}]^2 = (\Gamma_{\gamma_{eg}}[t_1 + t_{\delta k}^{(l)}])^2 |M^{(l)}(\zeta/\chi, \eta)|^2 \times \{\alpha^2 e^{-2(1+1/\eta)\gamma_{eg}\tau_o} + \beta^2\}. \quad (50)$$

As compared to Eq. (38), we find the same function $(\Gamma_{\gamma_{eg}}[t_1 + t_{\delta k}^{(l)}])^2$, which characterized atomic decay during the storage process [Eq. (34)], only with slightly modified argument [now including an additional delay $t_{\delta k}^{(l)}$]. Note that phase mismatch only leads to a decrease of the efficiency when paired with significant atomic phase relaxation γ_{eg} , in contrast to the transverse case [Eq. (35)].

Figure 8 depicts the efficiency $\epsilon^{(l)}$ in the case of negligible phase relaxation (i.e., $\gamma_{eg} = 0$) as a function of effective optical depth $2\pi\zeta/\chi$ and compression parameter η . Furthermore, a comparison of efficiencies of the longitudinal and transverse compression schemes for various initial optical depths is shown in Fig. 9. For small initial optical depth ($\kappa_{\text{eff}} = 2\pi\zeta/\chi \lesssim 0.5\pi$ or $\kappa_{\text{eff}} = \alpha_o L \lesssim 0.5\pi$, respectively) and large




FIG. 8. (Color online) Recall efficiency for longitudinal broadening as a function of the compression parameter η and the effective optical depth $2\pi\zeta/\chi$ [see Eq. (50)].

compression parameter ($\kappa_{\text{eff}}/\eta < 1$), both schemes feature the same efficiency:

$$\epsilon^{(l)}(\eta, \kappa_{\text{eff}}) = \epsilon^{(t)}(\eta, \kappa_{\text{eff}}) \cong \kappa_{\text{eff}}^2/\eta. \quad (51)$$

This behavior is expected as reabsorption in transverse as well as longitudinally broadened media is negligible in the case of small optical depth. However, as the optical depth increases, reabsorption becomes more and more important in the case of transverse broadening, reflecting the departure from reversible light-atom interaction, while it is no issue in the case of longitudinal broadening, due to the correlation between atomic detuning Δ and position z [see Eq. (42)]. In particular, for longitudinal broadening, we find

$$\epsilon^{(l)}(\eta \rightarrow 0) = (1 - \exp\{-\kappa_{\text{eff}}\}) = 1|_{\kappa_{\text{eff}} \gg 1}, \quad (52)$$

while decompression with transverse broadening yields a monotonously decreasing efficiency,

$$\epsilon^{(t)}(\eta \rightarrow 0) = 4\eta|_{\kappa_{\text{eff}} \gg 1}, \quad (53)$$

and eventually becomes zero.




FIG. 9. (Color online) Recall efficiencies for compression schemes employing transverse (red dashed lines) and longitudinal broadening (blue solid lines). The initial optical depths $\kappa_{\text{eff}} = \alpha_o L = 2\pi\zeta/\chi$ are 0.1π , π , and 4π (bottom to top sets of curves, respectively). For small optical depth and sufficiently large compression factor η , the efficiencies $\epsilon^{(t)}(\eta)$ and $\epsilon^{(l)}(\eta)$ are equal, as described by Eq. (51). As the optical depth increases, schemes based on longitudinal broadening perform better.




FIG. 10. (Color online) Gain for longitudinal broadening as function of compression parameter η and effective optical depth $\kappa_{\text{eff}} = 2\pi\xi/\chi$.

Furthermore, the compression efficiency $\epsilon^{(l)}$ is only limited by the optical depth κ_{eff} during absorption, as well as by atomic relaxation. In particular, there is no compression-factor-dependent upper limit, in opposition to $\epsilon^{(l)}$ [see Eq. (40)]. Hence, provided sufficient optical depth $2\pi\xi/\chi > 1$ and $2\pi\xi/(\eta\chi) > 1$ (which may be difficult to achieve for large compression parameters), one can always achieve close-to-unit efficiency. This gives longitudinal broadening a clear advantage over transverse broadening for quantum compression.

Obviously, the improved performance is reflected in the gain. As shown in Fig. 10, we find that the gain can always be increased when increasing the effective optical depth or the compression parameter, as opposed to the case of transverse broadening where the gain was limited to four [see Eq. (41)]. Indeed, using Eqs. (4) and (50), and ignoring atomic relaxation (i.e., $\gamma_{eg} = 0$), we find

$$G^{(l)} = \eta(1 - \exp\{-\kappa_{\text{eff}}/\eta\})(1 - \exp\{-\kappa_{\text{eff}}\}). \quad (54)$$

Hence, assuming sufficiently large effective optical depth, we find

$$G^{(l)}(\eta/\kappa_{\text{eff}} \rightarrow 0) = \eta|_{\kappa_{\text{eff}} \gg 1}. \quad (55)$$

VI. FEASIBILITY STUDY

Before we conclude this article, let us briefly discuss possible implementations of our proposal. Quantum compression as considered here requires an atomic ensemble with (at least) three suitable energy levels. Two levels must be coupled resonantly by the photonic wave packet to be compressed. The third level, generally another ground state, is employed for temporal mapping of the excited optical coherence onto ground-state coherence using two counter-propagating π pulses. This results in the implementation of the $2kz$ phase shift, that is, in emission of light in the backward direction [6]. The width of the optical absorption line, which may be inhomogeneously broadened, should be sufficiently small. It should be possible to broaden the line in a controlled way via external fields, and the optical depth after broadening should be large. The maximum bandwidth of the optical wave packet to be absorbed or re-emitted

is given by the width of the externally broadened atomic transition, while the minimum bandwidth of the optical wave packet is limited by the atomic linewidth before broadening (more precisely, the minimum optical bandwidth is a small multiple of the atomic linewidth as some controlled broadening is required for controlled dephasing or rephasing). These conditions are similar to those required for quantum-state storage based on standard CRIB [10], but without the need for a long storage time, which considerably relaxes material requirements.

Any material that fulfills these requirements would be a suitable candidate for quantum compression. We note that some aspects of the here-presented theoretical study have recently been observed using atomic vapor, longitudinal broadening, and bright pulses of light [17]. In the following, we discuss RE-ion-doped solid-state material in view of quantum compression.

RE-doped solids have been studied extensively for data storage experiments based on stimulated photon echoes [43] and are currently being investigated for photon-echo quantum memory [5,10]. When implemented into crystals and cooled down to temperatures below 4 K, RE ions typically feature homogeneous linewidths (for optical transitions) of a few kHz, and values as small as 50 Hz have been reported [44]. Linewidths in glasses are larger, but can still be around or below 1 MHz [45–47]. The transitions in crystals and glasses are inhomogeneously broadened, with values ranging from 40 MHz [48] in crystals to hundreds of GHz in crystals or glasses [44,49]. The preparation of the initial, narrow absorption line therefore requires an initial preparation step via optical pumping [50–53]. In low-symmetry hosts, the RE ion's quantum states acquire permanent electric dipole moments, which makes it possible to shift their energy levels through the application of a dc electric field—the dc Stark shift (for a review on optical Stark spectroscopy of solids, see [37]). A different dipole moment for two energy levels consequently results in a shift of the associated transition frequency, making it possible, through the application of a field gradient, to broaden an isolated absorption line. Typical frequency shifts in RE-ion-doped materials are 10–100 kHz for an electric field of 1 V cm^{-1} .

Praseodymium-doped yttrium orthosilicate ($\text{Pr}^{3+}\text{Y}_2\text{SiO}_5$) is a very promising candidate for the demonstration of quantum compression in RE solids. This crystal has been employed for storage of light pulses using longitudinal broadening [41], and recall efficiencies up to 69% have recently been reported [29]. Taking into account a spectral width of the ${}^3H_4 \leftrightarrow {}^1D_2$ (605.9 nm wavelength) absorption line of 30 kHz after optical pumping [41], and a Stark coefficient of $112.1 \text{ kHz}/(\text{V cm}^{-1})$ [54], we find that $\text{Pr}^{3+}\text{Y}_2\text{SiO}_5$ allows absorption and re-emission of photonic wave packets with bandwidths between $\sim 100 \text{ kHz}$ and a few MHz, that is, temporal durations between $\sim 100 \text{ ns}$ and a few μs . This results in a maximum compression (scaling) factor η of ~ 100 (or $\eta \sim 1/100$). Taking into account the demonstrated high-efficiency storage [29], one can expect a gain $G^{(l)} > 10$. However, note that the maximum bandwidth for this transition is $\sim 10 \text{ MHz}$, due to limitations imposed by ground-state hyperfine splitting on the preparation of the initial absorption line via optical pumping [51].

For RE materials with larger ground-state splitting, allowing storage or recall of shorter wave packets, it is interesting to consider RE-ion-doped crystalline and amorphous waveguides. Due to the possibility of implementing electrodes with spacing as small as $\sim 10 \mu\text{m}$, large electric fields, that is, large Stark shifts, can be obtained through application of modest voltages. So far, waveguides in erbium- or thulium-doped lithiumniobate crystals and erbium-doped silicate fibers have been investigated in view of quantum-state storage [38,40,46,55], and Stark broadening up to ~ 100 MHz seems feasible, allowing for storage or recall of pulses as short as ~ 10 ns.

VII. CONCLUSION

In summary, we have studied quantum compression and decompression of photonic time-bin qubit states employing a generalized version of CRIB-based photon-echo quantum memory. Assuming high optical depth for storage and retrieval, we find, for the case of transverse broadening, that the recall efficiency is limited by the compression factor η , while it reaches unity in the case of a longitudinal broadened medium. We also find, for transverse broadening, that the fidelity of recalled photonic time-bin qubits with the original qubit is one, regardless the compression factor, but that it is limited in the case of longitudinal broadening. Taking into account realistic experimental data, we foresee that quantum compression will be useful for quantum communication and computation applications. In particular, it makes it possible to enhance the data rate in quantum communication schemes through temporal multiplexing and allows mapping of broadband photons into small-bandwidth quantum memory.

Our analysis reveals new aspects of coherent photon-atom interaction, specifically photon-echo type interactions, and highlights the advantage of schemes employing longitudinal broadening over transverse broadening. The theory can be generalized in a straightforward way to the encoding of quantum information into multiple ($n > 2$) (discrete) temporal modes of photons, including compression of intense light fields (provided the number of photons is smaller than number of resonant atoms). However, quantum compression or decompression of light carrying quantum information with continuous spectrum (continuous quantum variables) requires further theoretical investigations that take into account added quantum noise in the case of limited recall efficiency. Finally, we note that pulse compression has been observed in the context of frequency conversion based on Raman adiabatic transfer of optical states [56], a protocol based on electromagnetically induced transparency that may provide an interesting alternative to our approach.

ACKNOWLEDGMENTS

This work was supported by NSERC, iCORE (now part of Alberta Innovates - Technology Futures), GDC, as well as the Russian Foundation of Basic Research under Grant Nos. 08-07-00449 and 10-02-01348-a and government contract of RosNauka 02.740.11.01.03.

APPENDIX A: TRANSVERSE BROADENING

First, we find a solution of Eqs. (28) and (29) given an input photon amplitude $A_+(t \leq t_1, z)$ and excited atomic coherence $b_j(t \leq t_1)$. Assuming all atoms to be initially in the ground state [$b_j(\tau = t_o \rightarrow -\infty) \rightarrow 0$] and using a temporal Laplace transformation for the amplitude $\bar{A}_{ip,+}(z) = \int_{t_o}^{\infty} d\tau A_+(t, z) \exp\{-p(t - t_o)\}$, we find the formal solution of Eq. (29),

$$b_j(t < t_1) = ig \frac{1}{2\pi} \exp\{-i\omega_+(t - t_o)\} \int_{t_o}^t dt' \int_{-\infty}^{\infty} dv \bar{A}_{v,+}(z_j) \times \exp\{-ivt' - i(\omega_{eg} + \Delta_j - \omega_+)(t - t') - i\delta\phi_{eg}^j(t, t')\}, \quad (\text{A1})$$

where we have also used a backward Laplace transformation $A_+(t, z) = (2\pi)^{-1} \int_{-\infty}^{\infty} dv \bar{A}_{v,+}(z) \exp\{-iv(t - t_o)\}$ with $p = -iv$. Putting Eq. (A1) in Eq. (28) and taking into account phase relaxation as described in Eq. (24), we find the following expression for the sum of arbitrary atomic functions $F_j(t, z_j)$:

$$\sum_{j=1}^N F_j(t, z_j) \exp\left\{-i \int_{t_o}^{t_1} dt' \delta\Delta_{eg}^j(t')\right\} \delta(z - z_j) |_{N \gg 1} = (n_o S) \exp\{-\gamma_{eg}(t_1 - t_o)\} \int_{-\infty}^{\infty} d\Delta G(\Delta/\Delta_{inh}) F(\Delta, t, z). \quad (\text{A2})$$

The Fourier components of the input field are given by

$$\bar{A}_{v,+}(z) = \exp\{-\frac{1}{2}\alpha_+(v)z\} \bar{A}_{v,+}(0), \quad (\text{A3})$$

where

$$\alpha_+(v) = \alpha_o(\gamma_{eg} + \Delta_{inh}) \times \int_{-\infty}^{\infty} d\Delta \frac{G(\Delta/\Delta_{inh})}{[\gamma_{eg} - i(v + \omega_+ - \omega_{eg} - \Delta)]} \quad (\text{A4})$$

is the frequency-dependent absorption coefficient for an arbitrary, inhomogeneously broadened absorption line $G(\Delta/\Delta_{inh})$ of the transverse type and the on-resonant absorption coefficient $\alpha_o = 2\pi n_o S g^2 / [c(\gamma_{eg} + \Delta_{inh})]$. The amplitude $A_+(t > t_1, z = L)$ in Eq. (55) describes the light field at $z \geq L$ after interaction with the atomic medium. The Fourier component at the beginning of the medium $\bar{A}_{v,+}(0)$ is determined by the initial state of the input light at $z = 0$.

We now consider inhomogeneous broadening with Lorentzian lineshape $G(\Delta/\Delta_{inh}) = \frac{\Delta_{inh}}{\pi(\Delta^2 + \Delta_{inh}^2)}$, assuming the spectral width of the input field narrow compared to the atomic linewidth ($\delta\omega_+ \ll \Delta_{inh}$). This results in $\alpha_+(v) = \frac{\alpha_o(\gamma_{eg} + \Delta_{inh})}{[\gamma_{eg} + \Delta_{inh} + i(\omega_{eg} - \omega_+ - v)]}$ in Eq. (A4).

To allow for light re-emission in the backward direction, we change the phases of the atomic amplitudes through the application of two short π laser pulses that temporally map atomic coherence on the optically excited transition onto a ground-state transition [6,35], as discussed in Sec. IV. In general, this phase-matching operation leads to some phase mismatch between the atomic coherence and the irradiated echo signal that we quantify through an additional atomic phase shift δkz_j : $b_j[t_1 + (\epsilon \rightarrow 0)] = b_j(t_1) \exp\{i\delta kz_j\}$ in Eqs. (28) and (29) [with index $\sigma = -$ denoting echo signal emission

and $\Delta_j(t > t_1) = -\eta\Delta_j$. Taking into account the initial atomic amplitudes $b_j(t) = b_j(t_1) \exp\{i\delta kz_j + i\eta\Delta_j(t - t_1) - i\delta\phi_{eg}^j(t, t_1)\}$ and Eq. (A2), we find the irradiated light field to be described by

$$\frac{\partial}{\partial z} \bar{A}_{ip,-}(z) = (\pi n_o S g^2 / c) \left[\int_{t_1}^{\infty} d\tilde{\tau} \exp\{-p\tilde{\tau}\} \bar{P}_{\text{echo}}(\tilde{\tau}, z) + \frac{\bar{A}_{ip,-}(z)}{p + (\gamma_{eg} + \eta\Delta_{\text{inh}}) + i(\omega_{eg} - \omega_-)} \right], \quad (\text{A5})$$

where $\bar{P}_{\text{echo}}(\tilde{\tau}, z)$ is the rephased atomic coherence that serves as a source for the irradiated field:

$$\begin{aligned} \bar{P}_{\text{echo}}(\tilde{\tau}, z) &= \exp\{-(1 + \eta)\gamma_{eg}(\tilde{\tau} - t_1)\} \exp\{i[\delta kz + \omega_- - \omega_{eg} \\ &+ \eta(\omega_+ - \omega_{eg})\tilde{\tau} + i(\omega_{eg} - \omega_+)(1 + \eta)t_1]\} \\ &\times \int_{-\infty}^{\infty} d\nu \frac{2\Delta_{\text{inh}} \exp\{i\nu[\eta(\tilde{\tau} - t_1) - t_1]\}}{[(\omega_+ + \nu - \omega_{eg} + i\gamma_{eg})^2 + \Delta_{\text{inh}}^2]} \\ &\times \exp\left\{-\frac{\alpha_o z}{2[1 + i(\omega_{eg} - \omega_+ - \nu)/(\gamma_{eg} + \Delta_{\text{inh}})]}\right\} \bar{A}_{v,+}(0). \end{aligned} \quad (\text{A6})$$

We used again the temporal Laplace transformation for light and atom amplitudes within the temporal interval $[t_1, \infty]$ and

$$\frac{2}{\alpha_+(v) + \alpha_-(\Omega) - 2i\delta k} \cong \frac{2}{\alpha_o(\gamma_{eg} + \Delta_{\text{inh}})} \frac{[\gamma_{eg} + \chi\Delta_{\text{inh}} + i(\omega_{eg} - \omega_- + \Omega)][\gamma_{eg} + \Delta_{\text{inh}} + i(\omega_{eg} - \omega_+ - \nu)]}{[2\gamma_{eg} + (1 + \chi)\Delta_{\text{inh}} + i(2\omega_{eg} - \omega_+ - \omega_- - \nu + \Omega) - 2i\delta k \frac{(\gamma_{eg} + \chi\Delta_{\text{inh}})}{\alpha_o}]}. \quad (\text{A8})$$

The main part of the integration in Eq. (A7) over Ω in the complex plane is determined by the pole singularity at $\Omega = \Omega_1(\nu)$, while the other singularities describe fast-decaying signals associated with the large inhomogeneous broadening. After the integration, we obtain the following echo field amplitude at the medium output ($z = 0$):

$$A_-(\tilde{\tau}, z = 0) = \frac{\eta \exp\{i\varphi - (1 + \eta)\gamma_{eg}(\tilde{\tau} - t_1)\}}{\pi[(\eta + 1) + (1 - \eta)\gamma_{eg}/\Delta_{\text{inh}}]} \int_{-\infty}^{\infty} d\nu \frac{\exp\{i\nu[\eta(\tilde{\tau} - t_1) - t_1]\} \{1 - \exp[-\frac{1}{2}\alpha(v, \eta)L + i\delta kL]\}}{\left\{1 + i \frac{(\eta-1)(\nu+\omega_+-\omega_{eg})/\Delta_{\text{inh}}}{(\eta+1)+(1-\eta)\gamma_{eg}/\Delta_{\text{inh}}} - 2i \frac{(\delta k/\alpha_o)(\eta+\gamma_{eg}/\Delta_{\text{inh}})}{(\eta+1)+(1-\eta)\gamma_{eg}/\Delta_{\text{inh}}}\right\}} \bar{A}_{+,v}(0), \quad (\text{A9})$$

where $\alpha(v, \eta) = [\alpha_+(v) + \alpha_-(v)]$ describes the combined influence of absorption and dispersion effects on the echo field.

Equation (A9) makes it possible to calculate the compression efficiency. For instance, setting $\eta = 1$, we can derive the echo field in the presence of phase mismatch [35,39], which limits the recall efficiency. Setting $\delta k = 0$, we find the echo to be a perfect, temporally reversed copy of the input field, provided the optical depth is sufficiently large for all spectral components of the field ($\alpha(v, \eta)L \gg 1$). This confirms previously obtained results [9,35,36]. For sufficiently large inhomogeneous broadening, $\delta\omega_f/\Delta_{\text{inh}} \gg 1$, we find $\alpha(v, \eta) \cong (1 + 1/\eta)\alpha_o$ and Eq. (A9) can be simplified to

$$A_-(\tilde{\tau}, z = 0) = \epsilon_o(\eta)^{1/2} M^{(t)}(\delta k, \alpha_o L) \exp\{i\varphi - (1 + \eta) \times \gamma_{eg}(\tilde{\tau} - t_1)\} \sqrt{\eta} A_+[-\eta(\tilde{\tau} - t_1) + t_1], \quad (\text{A10})$$

new variables $\tilde{\tau} = t + z/c$ and $z = z$, and we have taken into account that initially $A_-(t_1, 0 < z < L) = 0$. The solution of Eq. (A2) has the form of a double integral

$$\begin{aligned} A_-(\tilde{\tau}, z) &= \frac{i}{2\pi^2} \int_{-\infty}^{\infty} d\nu \frac{\Delta_{\text{inh}} \bar{A}_{v,+}(0) \exp\{i(\varphi - \nu t_1)\}}{[(\omega_+ - \omega_o + \nu + i\gamma_{eg})^2 + \Delta_{\text{inh}}^2]} \\ &\times \int_{-\infty}^{\infty} d\Omega \frac{\exp\{i\Omega(\tilde{\tau} - \tilde{\tau}_o)\}}{[\alpha_+(v) + \alpha_-(\Omega) - 2i\delta k]} \\ &\times \frac{\alpha_o(\gamma_{eg} + \Delta_{\text{inh}})}{[\Omega - \Omega_1(\nu)]} \left\{ \exp\left[-\frac{1}{2}\alpha_+(v)z + i\delta kz\right] \right. \\ &\left. - \exp\left[-\frac{1}{2}\alpha_+(v)L + \frac{1}{2}\alpha_-(\Omega)(z - L) + i\delta kL\right] \right\}, \end{aligned} \quad (\text{A7})$$

where $\omega_- = \omega_{eg} + \eta(\omega_{eg} - \omega_+)$ is the carrier frequency of the echo field, $\varphi = (1 + \eta)(\omega_{eg} - \omega_+)t_1$, $\Omega_1(\nu) = (\omega_- - \omega_{eg}) + \eta(\omega_+ - \omega_{eg} + \nu) + i(1 + \eta)\gamma_{eg}$, and $\alpha_-(v) = \frac{\alpha_o(\gamma_{eg} + \Delta_{\text{inh}})}{[\gamma_{eg} + \eta\Delta_{\text{inh}} + i(\omega_{eg} - \omega_+ + \nu)]}$ is the absorption coefficient of the atomic system during echo emission. It is characterized by the compressed inhomogeneous broadening $\eta\Delta_{\text{inh}}$. Assuming again the inhomogeneous broadening to be large compared with the initial light spectrum, $\Delta_{\text{inh}} \gg \delta\omega_f$, we can simplify the dependence on the phase mismatch factor:

where $M^{(t)}(\delta k, \alpha_o L)$ and $\epsilon_o^{1/2}(\eta)$ are given in Eqs. (35) and (36). While we are only interested in the quantum state of the retrieved and compressed photonic qubit [as described by Eq. (31)], which we obtain through Eq. (A10) assuming initial photonic time-bin qubits [Eq. (11)] with Gaussian shape of the basis wave packets, we note that one can also derive the amplitude of the atomic state [Eq. (27)] after the echo signal emission by inserting Eq. (A7) into Eq. (29).

APPENDIX B: LONGITUDINAL BROADENING

As in the previously discussed case of transverse broadening, we are only interested in the retrieved photonic time-bin qubit state. Averaging Eq. (29) over the phase fluctuations, and after a variable transformation to a moving reference frame $\tau = t - z/c, z = z$, the light-atom equations for $t < t_1$ become

$$\frac{\partial}{\partial z} A_1(\tau, z) = i(\pi n_o S g^* / c) b_o(\tau, z), \quad (\text{B1})$$

$$\frac{\partial}{\partial \tau} b_o(\tau, z) = (i\chi z - \gamma_{eg})b_o(\tau, z) + igA_1(\tau, z), \quad (\text{B2})$$

where we used the substitution $b_j(\tau) = b_o(\tau, z_j) \exp\{-i\omega_{21}\tau\}$, with $b_o(\tau, z)$ describing averaged local atomic coherence, and $A_1(\tau, z) = A_+(\tau, z) \exp\{-i(\omega_+ - \omega_{eg})\tau\}$. The field amplitude $A_+(\tau, z)$ is given by Eq. (11) with additional factor $\exp\{i\phi_{lg}\}$, where $\phi_{lg} = \frac{1}{2}\omega_+L/c$.

The general solution of Eqs. (B1) and (B2) is obtained using a temporal Laplace transformation [42]. We are interested only in the atomic coherence excited by the input light, which gives rise to the echo field irradiated after further evolution, while the initial light field disappears in the medium [$A_1(\tau \gg \delta t, z) \rightarrow 0$]:

$$b_o(\tau_1, z) = -\frac{i}{2\pi} \int_{-\infty}^{\infty} d\omega e^{-i\omega\tau_1} \times \frac{(\omega - \chi L/2 + i\gamma_{eg})^{i\zeta/\chi}}{(\omega + \chi z + i\gamma_{eg})^{1+i\zeta/\chi}} \tilde{A}_1(\omega, -L/2). \quad (\text{B3})$$

$\tilde{A}_1(\omega, -L/2) = \int_{-\infty}^{\infty} d\tau e^{i\omega\tau} A_1(\tau; -L/2)$ is the field spectrum at the input of the atomic medium ($z = -L/2$), and $\zeta = \pi n_o S g^2/c$. Similar to the transverse scheme, we find the atomic coherence on the resonant optical transition $1 \rightarrow 2$ at the moment of re-emission of light using the phase mismatch factor δkz : $b_o[\tau_1 + (\epsilon \rightarrow 0)] = b_o(\tau_1) \exp\{i\delta kz_j\}$.

We now analyze the backward echo emission. After changing the atomic detunings $\Delta(\tau > \tau_1) = \chi'z$, we find the following system of equations for the atomic coherence and irradiated echo field $A_2(\tilde{\tau} > \tau_1, z)$ (where $\tilde{\tau} = t + z/c$):

$$\frac{\partial}{\partial \tilde{\tau}} A_2(\tilde{\tau}, z) = -i(\pi n_o S g^*/c)b_o(\tilde{\tau}, z), \quad (\text{B4})$$

$$\frac{\partial}{\partial \tilde{\tau}} b_o(\tilde{\tau}, z) = -(i\chi z + \gamma_{eg})b_o(\tilde{\tau}, z) + igA_2(\tilde{\tau}, z) \quad (\text{B5})$$

where $A_2(\tau, z) = A_-(\tau, z) \exp\{-i(\omega_- - \omega_{eg})\tilde{\tau}\}$ and ω_- is the new carrier frequency. Using a Laplace transformation in the solutions of Eqs. (B4) and (B5), we find $A_2(\tilde{\tau}, z < L/2)$ as a function of the stored atomic coherence $b_o(\tilde{\tau}_1, z)$:

$$A_2(\tau, z = -L/2) = \frac{\zeta}{(2\pi)^2} \int_{-\infty}^{\infty} d\omega e^{-i\omega(\tau-t_1)} \int_{-L/2}^{L/2} dz' \times \frac{(-\chi' L/2 - \omega - i\gamma_{eg})^{-i\frac{\zeta}{\chi'}}}{(\chi' z' - \omega - i\gamma_{eg})^{1-i\frac{\zeta}{\chi'}}} \exp\{i\delta kz'\} \times \int_{-\infty}^{\infty} d\omega' e^{-i\omega't_1} \frac{(\omega' - \chi L/2 + i\gamma_{eg})^{i\zeta/\chi}}{(\omega' + \chi z' + i\gamma_{eg})^{1+i\zeta/\chi}} \times \tilde{A}_1(\omega', -L/2), \quad (\text{B6})$$

where we used $\tau_1 = t_1$. Taking into account backward emission of the echo field, as well as large inhomogeneous broadening with respect to the input light spectral width, and using the simplifications

$$\begin{aligned} (\omega' - \chi L/2 + i\gamma_{eg})^{i\zeta/\chi} &\cong (\chi L/2)^{i\zeta/\chi} \exp\{\gamma_{eg}\tau_m - \pi\zeta/\chi\} \exp\{-i\tau_m\omega'\}, \\ (-\chi' L/2 - \omega - i\gamma_{eg})^{-i\zeta/\chi'} &\cong (\chi' L/2)^{-i\zeta/\chi'} \exp\{\gamma_{eg}\tau'_m - \pi\zeta/\chi'\} \exp\{-i\tau'_m\omega\}, \end{aligned}$$

where $\tau'_m = (\zeta/\chi') \frac{(\chi' L/2)}{[(\chi' L/2)^2 + \gamma_{eg}^2]} \cong \frac{(\zeta/\chi')}{(\chi' L/2)}$ and $\tau_m = (\zeta/\chi) \frac{(\chi L/2)}{[(\chi L/2)^2 + \gamma_{eg}^2]} \cong \frac{(\zeta/\chi)}{(\chi L/2)}$, we find, after changing the order of the integrations in Eq. (B6) from ω, z', ω' to ω', z', ω ,

$$\begin{aligned} A_2(\tau, z = -L/2) &= \frac{\zeta}{(2\pi)^2} \exp\{\gamma_{eg}(\tau'_m + \tau_m) \\ &\quad - \pi\zeta(1/\chi' + 1/\chi)\} (\chi' L/2)^{-i\zeta/\chi'} (\chi L/2)^{i\zeta/\chi} \\ &\quad \times \int_{-\infty}^{\infty} d\omega' \exp\{-i\omega'(t_1 + \tau_m)\} \tilde{A}_1(\omega', -L/2) \\ &\quad \times \int_{-L/2}^{L/2} dz' \frac{\exp\{i\delta kz'\}}{(\omega' + \chi z' + i\gamma_{eg})^{1+i\zeta/\chi}} \\ &\quad \times \int_{-\infty}^{\infty} d\omega \frac{\exp\{-i\omega(\tau - t_1 + \tau'_m)\}}{(\chi' z' - \omega - i\gamma_{eg})^{1-i\zeta/\chi'}}. \quad (\text{B7}) \end{aligned}$$

Taking again into account that $\chi L \gg \delta\omega_f$, and using the tabled integral

$$\begin{aligned} (\chi' L/2)^{-i\zeta/\chi'} \int_{-\infty}^{\infty} du \frac{\exp\{iuT\}}{(u - i\gamma_{eg})^{1-i\zeta/\chi'}} &= -\frac{2\pi\eta_\chi(T)}{(\zeta/\chi')\Gamma[-i\zeta/\chi']} (T\chi' L/2)^{-i\zeta/\chi'} \\ &\quad \times \exp\left\{-\gamma_{eg}T + \frac{1}{2}\pi\zeta/\chi'\right\}, \end{aligned}$$

where $\eta_\chi(T \geq 0) = 1$ and $\eta_\chi(T < 0) = 0$ is a Heaviside function, we integrate Eq. (B7) over ω , then over z' and ω' , leading to

$$\begin{aligned} A_2(\tau, z = -L/2) &= -\frac{2\pi\chi' \exp\{-\frac{1}{2}\pi\zeta/\chi' - \frac{1}{2}\pi\zeta/\chi\}}{\zeta \Gamma[-i\zeta/\chi']\Gamma[i\zeta/\chi]} \\ &\quad \times \eta_\chi(\tau - t_1 + \tau_m - \delta k/\chi') \\ &\quad \times [(\chi' L/2)(\tau - t_1 + \tau'_m)]^{-i\zeta/\chi'} \\ &\quad \times [(\chi' L/2)|\tau - t_1 + \tau'_m - \delta k/\chi'|]^{i\zeta/\chi} \\ &\quad \times \exp\{-\gamma_{eg}(1 + \eta)(\tau - t_1) + \gamma_{eg}\eta\tau_z\} \\ &\quad \times A_1\{-\eta[\tau - (1 + 1/\eta)t_1 - \tau_z]\}, \quad (\text{B8}) \end{aligned}$$

where $\eta = \chi'/\chi$ is the compression factor and $\Gamma[\pm ix]$ are Γ functions. Interestingly, as described in Eq. (B8), phase mismatch does not effect to the quantum efficiency of the echo emission but only leads to a temporal shift of the echo emission to $\tau'_{\text{echo}} + \tau_z$ [where $\tau'_{\text{echo}} = (1 + 1/\eta)t_1$, $\tau_z = \delta k/\chi' + \tau_m/\eta - \tau'_m$]. However, this shift results in an additional nonlinear phase shift due to the deviation from perfect temporal reversibility. Furthermore, we find that the echo emission is conditioned on a phase mismatch $\delta k/\chi' > \tau'_m + \delta t - (t_1 + \tau_m)/\eta$. Taking into account that $\tau_m \leq \delta t$ and $\tau'_m \leq \delta t$, we find $\delta k/\chi' > -t_1$.

Finally, taking into account the relations between $A_2(\tilde{\tau}, z)$ and $A_-(\tilde{\tau}, z)$ and those between $A_1(\tilde{\tau}, z)$ and $A_+(\tilde{\tau}, z)$ we find the new carrier frequency $\omega_- = \omega_{eg} + \eta(\omega_{eg} - \omega_+)$ of the echo signal, which coincides with the case of transverse broadening. We also find $A_-(\tilde{\tau}, z) = A_2(\tilde{\tau}, z) \exp\{i\eta(\omega_{eg} - \omega_+)\tilde{\tau}\}$. Using this relation, together with $A_1\{-\eta(\tau - \tau'_{\text{echo}} - \tau_z)\} = A_+\{-\eta(\tau - \tau'_{\text{echo}} - \tau_z)\} \exp\{i\eta(\omega_+ - \omega_{eg})(\tau - \tau'_{\text{echo}} - \tau_z)\}$ in Eq. (B8), we find

the solution for $A_-(\tilde{\tau}, z)$ as a function of A_+ . Putting the obtained solution into Eq. (10), assuming again a Gaussian shape of the photonic wave packets, we ob-

tain the solution for the retrieved photonic time-bin qubit state given in Eq. (43) after a simple algebraic calculation.

- [1] N. Sangouard, C. Simon, H. De Riedmatten, and N. Gisin, e-print [arXiv:0906.2699](https://arxiv.org/abs/0906.2699).
- [2] L.-M. Duan, M. D. Lukin, J. I. Cirac, and P. Zoller, *Nature (London)* **414**, 413 (2001).
- [3] J. McKeever, A. Boca, A. D. Boozer, R. Miller, J. R. Buck, A. Kuzmich, and H. J. Kimble, *Science* **303**, 1992 (2004).
- [4] T. Chanelière, D. N. Matsukevich, S. D. Jenkins, S.-Y. Lan, R. Zhao, T. A. B. Kennedy, and A. Kuzmich, *Phys. Rev. Lett.* **98**, 113602 (2007).
- [5] A. I. Lvovsky, B. C. Sanders, and W. Tittel, *Nat. Photonics* **3**, 706 (2009).
- [6] S. A. Moiseev and S. Kröll, *Phys. Rev. Lett.* **87**, 173601 (2001).
- [7] A. L. Alexander, J. J. Longdell, M. J. Sellars, and N. B. Manson, *Phys. Rev. Lett.* **96**, 043602 (2006).
- [8] M. Nilsson and S. Kröll, *Opt. Commun.* **247**, 393 (2005).
- [9] B. Kraus, W. Tittel, N. Gisin, M. Nilsson, S. Kröll, and J. I. Cirac, *Phys. Rev. A* **73**, 020302(R) (2006).
- [10] W. Tittel, M. Afzelius, T. Chanelière, R. Cone, S. Kröll, S. A. Moiseev, and M. J. Sellars, *Laser Photonics Rev.* **4**, 244 (2010).
- [11] Y. S. Bai and T. W. Mossberg, *Appl. Phys. Lett.* **45**, 1269 (1984).
- [12] Y. S. Bai and T. W. Mossberg, *Opt. Lett.* **11**, 30 (1986).
- [13] X. Wang, M. Afzelius, N. Ohlsson, U. Gustafsson, and S. Kröll, *Opt. Lett.* **25**, 945 (2000).
- [14] J. Ruggiero, J.-L. Le Gouët, C. Simon, and T. Chanelière, *Phys. Rev. A* **79**, 053851 (2009).
- [15] N. Sangouard, C. Simon, J. Minář, M. Afzelius, T. Chanelière, N. Gisin, J.-L. Le Gouët, H. De Riedmatten, and W. Tittel, *Phys. Rev. A* **81**, 062333 (2010).
- [16] S. A. Moiseev and W. Tittel, presentation at *Workshop on the Storage and Manipulation of Quantum Information in Optically-Addressed Solids (SMQIOAS 2008)*, Bozeman, MT, USA, 25–27 January 2008 [http://www.spectrum.montana.edu/Conf/Booklet/SMQIOAS2008Booklet_BW.pdf].
- [17] M. Hosseini, B. M. Sparkes, G. Hétet, J. J. Longdell, P. K. Lam, and B. C. Buchler, *Nature (London)* **461**, 241 (2009).
- [18] B. C. Buchler, M. Hosseini, G. Hétet, B. M. Sparkes, and P. K. Lam, *Opt. Lett.* **35**(7), 1091 (2010).
- [19] W. K. Wootters and W. H. Zurek, *Nature (London)* **299**, 802 (1982).
- [20] D. Dieks, *Phys. Lett. A* **92**, 271 (1982).
- [21] P. W. Shor, *Phys. Rev. A* **52**, 2493(R) (1995).
- [22] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [23] H. Carmichael, *Lecture Notes in Physics* (Springer-Verlag, Berlin/Heidelberg, 1993), Monographs Series, Vol. 18, Chap. 7.
- [24] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [25] W. Tittel and G. Weihs, *Quant. Inf. Comp.* **1**, 3 (2001).
- [26] H. de Riedmatten, I. Marcikic, W. Tittel, H. Zbinden, D. Collins, and N. Gisin, *Phys. Rev. Lett.* **92**, 047904 (2004).
- [27] M. O. Scully and S. M. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, UK, 1997).
- [28] R. Loudon, *The Quantum Theory of Light* (Oxford University Press, Oxford, UK, 2000), 3rd ed.
- [29] M. Hedges, J. J. Longdell, Y. Li, and M. J. Sellars (unpublished).
- [30] S. A. Moiseev, *J. Phys. B* **40**, 3877 (2007).
- [31] N. Gisin, S. A. Moiseev, and C. Simon, *Phys. Rev. A* **76**, 014302 (2007).
- [32] J. R. Klauder and P. W. Anderson, *Phys. Rev.* **125**, 912 (1962).
- [33] K. M. Salikhov, A. G. Semenov, and Y. D. Tsvetkov, *Applications of Electron Spin Echo* (Nauka, Novosibirsk, 1976) [in Russian].
- [34] J.-T. Shen and S. Fan, *Phys. Rev. Lett.* **95**, 213001 (2005); **98**, 153003 (2007).
- [35] S. A. Moiseev and M. I. Noskov, *Laser Phys. Lett.* **1**, 303 (2004).
- [36] N. Sangouard, C. Simon, M. Afzelius, and N. Gisin, *Phys. Rev. A* **75**, 032327 (2007).
- [37] R. M. Macfarlane, *J. Lumin.* **125**, 156 (2007).
- [38] S. R. Hastings-Simon, M. U. Staudt, M. Afzelius, P. Baldi, D. Jaccard, W. Tittel, and N. Gisin, *Opt. Commun.* **266**, 716 (2006).
- [39] S. A. Moiseev and B. S. Ham, *Phys. Rev. A* **70**, 063809 (2004).
- [40] M. U. Staudt, S. R. Hastings-Simon, M. Nilsson, M. Afzelius, V. Scarani, R. Ricken, H. Suche, W. Sohler, W. Tittel, and N. Gisin, *Phys. Rev. Lett.* **98**, 113601 (2007).
- [41] G. Hétet, J. J. Longdell, A. L. Alexander, P. K. Lam, and M. J. Sellars, *Phys. Rev. Lett.* **100**, 023601 (2008).
- [42] S. A. Moiseev and N. M. Arslanov, *Phys. Rev. A* **78**, 023803 (2008).
- [43] Y. C. Sun, in *Spectroscopic Properties of Rare Earths in Optical Materials*, edited by G. Liu and B. Jacquier (Springer, Berlin/Heidelberg/New York, 2005), Chap. 7.
- [44] Y. Sun, C. W. Thiel, R. L. Cone, R. W. Equall, and R. L. Hutcheson, *J. Lumin.* **98**, 281 (2002).
- [45] R. M. Macfarlane, Y. Sun, P. B. Sellin, and R. L. Cone, *Phys. Rev. Lett.* **96**, 033602 (2006).
- [46] M. U. Staudt, S. R. Hastings-Simon, M. Afzelius, D. Jaccard, W. Tittel, and N. Gisin, *Opt. Commun.* **266**, 720 (2006).
- [47] Y. Sun, R. L. Cone, L. Bigot, and B. Jacquier, *Opt. Lett.* **31**, 3453 (2006).
- [48] R. M. Macfarlane, A. Cassanho, and R. S. Meltzer, *Phys. Rev. Lett.* **69**, 542 (1992).
- [49] J. L. Zyskind, E. Desurvire, J. W. Sulhoff, and D. J. Di Giovanni, *IEEE Photonics Technol. Lett.* **2**, 869 (1990).
- [50] G. J. Pryde, M. J. Sellars, and N. B. Manson, *Phys. Rev. Lett.* **84**, 1152 (2000).
- [51] M. Nilsson, L. Rippe, S. Kröll, R. Klieber, and D. Suter, *Phys. Rev. B* **70**, 214116 (2004).
- [52] V. Crozatier, F. de Seze, L. Haals, F. Bretenaker, I. Lorgeré, and J.-L. Le Gouët, *Opt. Commun.* **241**, 203 (2004).
- [53] B. Lauritzen, S. R. Hastings-Simon, H. de Riedmatten, M. Afzelius, and N. Gisin, *Phys. Rev. A* **78**, 043402 (2008).
- [54] F. R. Graf, A. Renn, U. P. Wild, and M. Mitsunaga, *Phys. Rev. B* **55**, 11225 (1997).
- [55] N. Sinclair, E. Saglamyurek, M. George, R. Ricken, C. La Mela, W. Sohler, and W. Tittel, *J. Lumin.* **130**(9), 1586 (2010).
- [56] F. Vewinger, J. Appel, E. Figueroa, and A. L. Lvovsky, *Opt. Lett.* **32** (19), 2771 (2007).