

Teleportation of squeezing: Optimization using non-Gaussian resources

Fabio Dell'Anno,¹ Silvio De Siena,¹ Gerardo Adesso,² and Fabrizio Illuminati^{1,*}

¹*Dipartimento di Matematica e Informatica, Università degli Studi di Salerno, CNR-SPIN, Unità di Salerno, Consorzio Nazionale Italiano di Struttura della Materia (CNISM), Unità di Salerno, and Istituto Nazionale di Fisica Nucleare (INFN), Sezione di Napoli-Gruppo Collegato di Salerno, Via Ponte don Melillo, I-84084 Fisciano (SA), Italy*

²*School of Mathematical Sciences, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom*

(Received 30 September 2010; published 29 December 2010)

We study the continuous-variable quantum teleportation of states, statistical moments of observables, and scale parameters such as squeezing. We investigate the problem both in ideal and imperfect Vaidman-Braunstein-Kimble protocol setups. We show how the teleportation fidelity is maximized and the difference between output and input variances is minimized by using suitably optimized entangled resources. Specifically, we consider the teleportation of coherent squeezed states, exploiting squeezed Bell states as entangled resources. This class of non-Gaussian states, introduced by Illuminati and co-workers [F. Dell'Anno, S. De Siena, L. Albano, and F. Illuminati, *Phys. Rev. A* **76**, 022301 (2007); F. Dell'Anno, S. De Siena, and F. Illuminati, *ibid.* **81**, 012333 (2010)], includes photon-added and photon-subtracted squeezed states as special cases. At variance with the case of entangled Gaussian resources, the use of entangled non-Gaussian squeezed Bell resources allows one to choose different optimization procedures that lead to inequivalent results. Performing two independent optimization procedures, one can either maximize the state teleportation fidelity, or minimize the difference between input and output quadrature variances. The two different procedures are compared depending on the degrees of displacement and squeezing of the input states and on the working conditions in ideal and nonideal setups.

DOI: [10.1103/PhysRevA.82.062329](https://doi.org/10.1103/PhysRevA.82.062329)

PACS number(s): 03.67.Hk, 03.67.Mn, 42.50.Pq

I. INTRODUCTION

Non-Gaussian quantum states, endowed with properly enhanced nonclassical properties, may constitute powerful resources for the efficient implementation of quantum information, communication, computation, and metrology tasks [1–13]. Indeed, it has been shown that, at fixed first and second moments, Gaussian states *minimize* various nonclassical properties [14,15]. Therefore, many theoretical and experimental efforts have been made toward engineering and controlling highly nonclassical, non-Gaussian states of the radiation field (for a review on quantum-state engineering, see, e.g., Ref. [16]). In particular, several proposals for the generation of non-Gaussian states have been presented [17–23], and some successful ground-breaking experimental realizations have been already performed [24–29]. Concerning continuous-variable (CV) quantum teleportation, to date the experimental demonstration of the Vaidman-Braunstein-Kimble (VBK) teleportation protocol [30,31] has been reported both for input coherent states [32–36] and for squeezed vacuum states [37,38]. In particular, Ref. [38] has reported the teleportation of squeezing, and consequently of entanglement, between upper and lower sidebands of the same spatial mode. It is worth noting that the efficient teleportation of squeezing, as well as of entanglement, is a necessary requirement for the realization of a quantum information network based on multistep information processing [39].

In this paper, adopting the VBK protocol, we study in full generality, e.g., including loss mechanisms and nonunity gain regimes, the teleportation of input single-mode coherent squeezed states using as non-Gaussian entangled resources a class of non-Gaussian entangled quantum states, the class of

squeezed Bell states [1,2]. This class includes, for specific choices of the parameters, non-Gaussian photon-added and photon-subtracted squeezed states. In tackling our goal, we use the formalism of the characteristic function introduced in Ref. [40] for an ideal protocol, and extended to the nonideal instance in Ref. [2]. Here, in analogy with the teleportation of coherent states, we first optimize the teleportation fidelity, that is, we look for the maximization of the overlap between the input and the output states. But the presence of squeezing in the unknown input state to be teleported prompts also an alternative procedure, depending on the physical quantities of interest. In fact, if one cares about reproducing in the most faithful way the initial state in phase space, then the fidelity is the natural quantity that needs to be optimized. On the other hand, one can be interested in preserving as much as possible the squeezing degree at the output of the teleportation process, even at the expense of the condition of maximum similarity between input and output states. In this case, one aims at minimizing the difference between the output and input quadrature averages and the quadrature variances. It is important to observe that this distinction makes sense only if one exploits non-Gaussian entangled resources endowed with tunable free parameters, so that enough flexibility is allowed to realize different optimization schemes. Indeed, it is straightforward to verify that this is impossible using Gaussian entangled resources. We will thus show that exploiting non-Gaussian resources, one can identify the best strategies for the optimization of different tasks in quantum teleportation, such as state teleportation versus teleportation of squeezing. A comparison with the same protocols realized by using Gaussian resources will confirm the greater effectiveness of non-Gaussian states versus Gaussian ones as entangled resources in the teleportation of quantum states of CV systems.

The paper is organized as follows. In Sec. II, we introduce the single-mode input states and the two-mode entangled

*illuminati@sa.infn.it

resources, and we recall the basics of both the ideal and the imperfect VKB quantum teleportation protocols. With respect to the instance of Gaussian resources (twin beam), the further free parameters of the non-Gaussian resource (squeezed Bell state) allow one to undertake an optimization procedure to improve the efficiency of the protocols. In Sec. III we investigate the optimization procedure based on the maximization of the teleportation fidelity. We then analyze an alternative optimization procedure, leading to the minimization of the difference between the quadrature variances of the output and input fields. This analysis is carried out in Sec. IV. We show that, unlike Gaussian resources, in the instance of non-Gaussian resources, the two procedures lead to different results and, moreover, always allow one to improve on the optimization procedures that can be implemented with Gaussian resources. Finally, in Sec. V we draw our conclusions and discuss future outlooks.

II. TELEPORTATION OF SQUEEZED COHERENT STATES USING SQUEEZED BELL RESOURCES

In this section, we briefly recall the basics of the ideal and imperfect VBK CV teleportation protocols (for details, see Ref. [2]). The scheme of the (CV) teleportation protocol is the following. Alice wishes to send to Bob, who is at a remote location, a quantum state, drawn from a particular set according to a prior probability distribution. The set of input states and the prior distribution are known to Alice and Bob, however, the specific state to be teleported that is prepared by Alice remains unknown. Alice and Bob share a resource, e.g., a two-mode entangled state. The input state and one of the modes of the resource are available for Alice, while the other mode of the resource is sent to Bob. Alice performs a suitable (homodyne) Bell measurement, and communicates the result to Bob, exploiting a classical communication channel. Then Bob, depending on the result communicated by Alice, performs a local unitary (displacement) transformation, and retrieves the output teleported state. The nonideal (realistic) teleportation protocol includes mechanisms of loss and inefficiency: the photon losses occurring in the realistic Bell measurements, and the noise arising in the propagation of optical fields in noisy channels (fibers) when the second mode of the resource is sent to Bob. The photon losses occurring in the realistic Bell measurements are modeled by placing in front of an ideal detector a fictitious beam splitter with nonunity transmissivity T^2 (and corresponding nonzero reflectivity $R^2 = 1 - T^2$) [41]. The propagation in fiber is modeled by the interaction with a Gaussian bath with an effective photon number n_{th} , yielding a damping process with an inverse-time rate γ [42,43]. Denoting by “in” the input field mode, and by 1 and 2, respectively, the first and the second mode of the entangled resource, the decoherence owing to imperfect photodetection in the homodyne measurement performed by Alice involves the input field mode “in” and one mode of the resource, e.g., mode 1. Throughout, we assume a pure entangled resource. Indeed, it is simple to verify that considering mixed (impure) resources is equivalent to a consider a suitable nonvanishing detection inefficiency R [2]. The degradation owing to propagation in fiber affects the other mode of the resource, e.g., mode 2,

which has to reach Bob’s remote place at the output stage. Denoting now by $\rho_{\text{in}} = |\phi\rangle_{\text{in}}\langle\phi|$ and $\rho_{\text{res}} = |\psi\rangle_{12}\langle\psi|$ the projectors corresponding, respectively, to a generic pure input single-mode state and a generic pure two-mode entangled resource, the characteristic function χ_{out} of the single-mode output field ρ_{out} can be written as [2]

$$\begin{aligned}\chi_{\text{out}}(\alpha) &= \text{Tr}[D_{\text{out}}(\alpha)\rho_{\text{out}}] \\ &= e^{-\Gamma_{\tau,R}|\alpha|^2} \chi_{\text{in}}(gT\alpha) \chi_{\text{res}}(gT, \alpha^*; e^{-\frac{\tau}{2}}\alpha),\end{aligned}\quad (1)$$

where $D_{\text{out}}(\alpha) = e^{\alpha a_{\text{out}}^\dagger + \alpha^* a_{\text{out}}}$ is the Glauber displacement operator, $\chi_{\text{in}}(\alpha) = \text{Tr}[D_{\text{in}}(\alpha)\rho_{\text{in}}]$ is the characteristic function of the input state, $\chi_{\text{res}}(\alpha_1, \alpha_2) = \text{Tr}[D_1(\alpha_1)D_2(\alpha_2)\rho_{\text{res}}]$ is the characteristic function of the resource, g is the gain factor of the protocol [44], $\tau \equiv \gamma t$ is the scaled dimensionless time proportional to the fiber propagation length, and the function $\Gamma_{\tau,R}$ is defined as

$$\Gamma_{\tau,R} = (1 - e^{-\tau})\left(\frac{1}{2} + n_{\text{th}}\right) + g^2 R^2. \quad (2)$$

We assume in principle to have some knowledge about the characteristics of the experimental apparatus: the inefficiency R (or T) of the photodetectors, and the loss parameters τ and n_{th} of the noisy communication channel.

We consider as the input state a single-mode coherent and squeezed (CS) state $|\psi_{\text{CS}}\rangle_{\text{in}}$ with an unknown squeezing parameter ($\varepsilon = s e^{i\phi}$) and an unknown coherent amplitude β . We then consider as a non-Gaussian entangled resource the two-mode squeezed Bell (SB) state $|\psi_{\text{SB}}\rangle_{12}$, defined as [1,2]

$$\begin{aligned}|\psi_{\text{CS}}\rangle_{\text{in}} &= D_{\text{in}}(\beta)S_{\text{in}}(\varepsilon)|0\rangle_{\text{in}}, \\ |\psi_{\text{SB}}\rangle_{12} &= S_{12}(\zeta)\{\cos\delta|0,0\rangle_{12} + e^{i\theta}\sin\delta|1,1\rangle_{12}\}.\end{aligned}\quad (3)$$

Here $D_{\text{in}}(\beta) = e^{\beta a_{\text{in}}^\dagger + \beta^* a_{\text{in}}}$ is, as before, the displacement operator, $S_{\text{in}}(\varepsilon) = e^{-\frac{1}{2}\varepsilon a_{\text{in}}^2 + \frac{1}{2}\varepsilon^* a_{\text{in}}^2}$ is the single-mode squeezing operator, $S_{12}(\zeta) = e^{-\zeta a_1^\dagger a_2^\dagger + \zeta^* a_1 a_2}$ is the two-mode squeezing operator ($\zeta = r e^{i\phi}$), with a_j denoting the annihilation operator for mode j ($j = \text{in}, 1, 2$), $|m, n\rangle_{12} \equiv |m\rangle_1 \otimes |n\rangle_2$ is the two-mode Fock state (of modes 1 and 2) with m photons in the first mode and n photons in the second mode, and θ and δ are two intrinsic free parameters of the resource entangled state, in addition to r and ϕ , which can be exploited for optimization. Note that the particular choices of the angle δ in the class of squeezed Bell states, Eq. (4), allow one to recover different instances of two-mode Gaussian and non-Gaussian entangled states: for $\delta = 0$, the Gaussian twin beam (TWB); for $\delta = \arccos[(\cosh 2r)^{-1/2} \sinh r]$ and $\theta = \phi - \pi$ the two-mode photon-added squeezed (PAS) state $|\psi_{\text{PAS}}\rangle_{12}$; and for $\delta = \arccos[(\cosh 2r)^{-1/2} \cosh r]$ and $\theta = \phi - \pi$ the two-mode photon-subtracted squeezed (PSS) state $|\psi_{\text{PSS}}\rangle_{12}$. The last two non-Gaussian states are defined as

$$|\psi_{\text{PAS}}\rangle_{12} = (\cosh 2r)^{-1/2} a_1^\dagger a_2^\dagger S_{12}(\zeta)|0,0\rangle_{12}, \quad (5)$$

$$|\psi_{\text{PSS}}\rangle_{12} = (\cosh 2r)^{-1/2} a_1 a_2 S_{12}(\zeta)|0,0\rangle_{12}, \quad (6)$$

and are already experimentally realizable with current technology [24–28].

In the following section we study, in comparison with the instance of two-mode Gaussian entangled resources, the performance of the optimized two-mode squeezed Bell states

TABLE I. Summary of the notation employed throughout this work to describe the different parameters that characterize the input coherent squeezed (CS) states [Eq. (3)], the shared entangled two-mode squeezed Bell (SB) resources [Eq. (4)], and the characteristics of nonideal teleportation setups [2]. See text for further details on the role of each parameter.

Object	Parameter	Description
Input state	β	Displacement
	s	Squeezing degree
	φ	Squeezing phase
Two-mode resource state	r	Squeezing degree
	ϕ	Squeezing phase
	δ	Mixing angle
	θ	Mixing phase
	$R = \sqrt{1 - T^2}$	Detection inefficiency
Teleportation apparatus	τ	Fiber loss factor
	n_{th}	Fiber bath temperature
	g	Gain of the protocol

when used as entangled resources for the teleportation of input single-mode coherent squeezed states. For completeness, in the same context, we make also a comparison with the performance, as entangled resources, of the more specific realizations [Eqs. (5) and (6)]. The characteristic functions of states shown in Eqs. (3)–(6) are computed and their explicit expressions are given in Appendix A.

For ease of reference, Table I provides a summary of the parameters associated with the input states, the shared resources, and the sources of noise in the teleportation protocol.

III. OPTIMAL TELEPORTATION FIDELITY

The commonly used measure to quantify the performance of a quantum teleportation protocol is the fidelity of teleportation [45], $\mathcal{F} = \text{Tr}[\rho_{\text{in}}\rho_{\text{out}}]$, which amounts to the overlap between a pure input state ρ_{in} and the (generally mixed) teleported state ρ_{out} . In the formalism of the characteristic function the fidelity reads

$$\mathcal{F} = \frac{1}{\pi} \int d^2\alpha \chi_{\text{in}}(\alpha)\chi_{\text{out}}(-\alpha), \quad (7)$$

where $\chi_{\text{in}}(\alpha)$ is the characteristic function of the single-mode input state $\rho_{\text{in}} = |\psi_{\text{CS}}\rangle_{\text{in}}\langle\psi_{\text{CS}}|$, Eq. (3), and $\chi_{\text{out}}(\alpha)$ is the characteristic function for the output teleported state, Eq. (1). In this section, we will make use of Eq. (7) to analyze the efficiency of the CV teleportation protocol.

In the instance of non-Gaussian squeezed Bell resources [Eq. (4)], at fixed squeezing parameter, the optimization procedure amounts to the maximization of the teleportation fidelity [Eq. (7)] over the free parameters of the entangled resource. It can be shown that the optimal choice for the phases ϕ and θ is $\phi = \pi$ and $\theta = 0$. The analytic expression for the fidelity \mathcal{F}_{CS} of the nonideal quantum teleportation of coherent squeezed states by using squeezed Bell resources reads

$$\mathcal{F}_{\text{CS}} = \frac{4}{\sqrt{\Lambda_1\Lambda_2}} e^{\frac{\omega_1^2}{\Lambda_1} - \frac{\omega_2^2}{\Lambda_2}} \left\{ 1 + e^{-(2r+\tau)} \sin \delta (\Delta_2 \cos \delta - \Delta_1 \sin \delta) \right. \\ \left. \times \left[\frac{1}{\Lambda_1} \left(1 + \frac{2\omega_1^2}{\Lambda_1} \right) + \frac{1}{\Lambda_2} \left(1 - \frac{2\omega_2^2}{\Lambda_2} \right) \right] \right\}, \quad (8)$$

$$+ \frac{1}{4} e^{-2(2r+\tau)} \Delta_2^2 \sin^2 \delta \left[\frac{1}{\Lambda_1^2} \left(3 + \frac{12\omega_1^2}{\Lambda_1} + \frac{4\omega_1^4}{\Lambda_1^2} \right) \right. \\ \left. + \frac{1}{\Lambda_2^2} \left(3 - \frac{12\omega_2^2}{\Lambda_2} + \frac{4\omega_2^4}{\Lambda_2^2} \right) \right. \\ \left. + \frac{2}{\Lambda_1\Lambda_2} \left(1 + \frac{2\omega_1^2}{\Lambda_1} - \frac{2\omega_2^2}{\Lambda_2} - \frac{4\omega_1^2\omega_2^2}{\Lambda_1\Lambda_2} \right) \right] \Bigg\}, \quad (8)$$

where, introducing $\tilde{g} = gT$, the quantities Λ_1 , Λ_2 , Δ_1 , Δ_2 , ω_1 , and ω_2 are defined by the following relations:

$$\Delta_1 = 1 + e^{4r} + 2e^{\frac{r}{2}}(1 - e^{4r})\tilde{g} + e^\tau(1 + e^{4r})\tilde{g}^2, \\ \Delta_2 = 1 - e^{4r} + 2e^{\frac{r}{2}}(1 + e^{4r})\tilde{g} + e^\tau(1 - e^{4r})\tilde{g}^2, \\ \Lambda_1 = e^{-2r-\tau}\Delta_1 + 2e^{2s}(1 + \tilde{g}^2) + 4\Gamma_{\tau,R}, \\ \Lambda_2 = e^{-2r-\tau}\Delta_1 + 2e^{-2s}(1 + \tilde{g}^2) + 4\Gamma_{\tau,R}, \\ \omega_1^2 = (1 - \tilde{g})^2(\beta - \beta^*)^2, \quad \omega_2^2 = (1 - \tilde{g})^2(\beta + \beta^*)^2, \\ \tilde{g} = gT. \quad (9)$$

For different choices of δ in Eq. (9) (see Sec. II), one obtains the teleportation fidelities associated to photon-added and photon-subtracted squeezed resource states. Let us observe that the fidelity in Eq. (9) depends both on the input coherent amplitude β , and on the input single-mode squeezing parameter s , while it is independent of the input squeezing phase φ . Once again, it is worth stressing that, in the teleportation paradigm, the input state is unknown and only partial (probabilistic) knowledge on the alphabet of input states is admitted. It is thus required, in principle, to assume teleportation protocols independent of the input parameters, as it turns out to be the case for the VBK protocol with Gaussian entangled resources and input coherent states. However, in more general cases, one can study the behavior of the so-called one-shot fidelity, that is, the teleportation fidelity at specific values of the input parameters. Suitable averages of the one-shot fidelity over the set of input states and parameters, according to an assigned prior distribution, will then result in the average quantum teleportation fidelity. The latter quantity can then be confronted with so-called classical fidelity thresholds (benchmarks) that correspond to the maximum achievable average fidelity between the input state (measured by Alice in order to achieve an optimal estimation of it) and the output state (prepared by Bob according to Alice's measurement outcomes), without the use of any shared entanglement [45]. While teleportation benchmarks are available for the cases of coherent input states (with completely unknown β) [46], purely squeezed input states (with $\beta = \varphi = 0$ and completely unknown s) [47], as well as for states with known squeezing degree and unknown displacement and phase [48], a benchmark for the case of input states with totally unknown displacement and squeezing has not yet been derived, and stands as a challenging problem in quantum estimation theory.

Henceforth, assuming *a priori* that the input parameters (displacement and squeezing degree) are completely random, we adopt then the following approach to optimize the quantum teleportation fidelity. We exploit a nonunity gain strategy to remove at least the β dependence in the one-shot fidelity; then, we study the behavior of the β -independent one-shot fidelity for specific values of the input squeezing parameter s , in order

to identify an effective, s -independent approximation. Indeed, fixing the gain g at the value $g = 1/T$ ($\tilde{g} = 1$) in Eq. (9) yields the β -independent fidelity \mathcal{F}_S :

$$\mathcal{F}_S = \frac{4}{\sqrt{\Lambda_1 \Lambda_2}} \left[\frac{e^{-2(2r+\tau)}}{4} \Delta_2^2 \sin^2 \delta \left(\frac{3}{\Lambda_1^2} + \frac{3}{\Lambda_2^2} + \frac{2}{\Lambda_1 \Lambda_2} \right) + e^{-(2r+\tau)} \sin \delta (\Delta_2 \cos \delta - \Delta_1 \sin \delta) \left(\frac{1}{\Lambda_1} + \frac{1}{\Lambda_2} \right) + 1 \right], \quad (10)$$

where the quantities $\Lambda_1, \Lambda_2, \Delta_1, \Delta_2, \omega_1$, and ω_2 are defined in Eq. (9). For different choices of δ (see Sec. II), one obtains the teleportation fidelities associated with the use of different Gaussian and non-Gaussian entangled resources: the twin beam, the photon-added squeezed states, and the photon-subtracted squeezed states. For such resources no optimization procedure is possible as δ is a specific function of r . Instead, the optimization of the fidelity [Eq. (10)] with respect to the free non-Gaussian parameter δ identifies the optimal squeezed Bell resource associated to the optimal value:

$$\delta_{\text{opt}} = \frac{1}{2} \arctan \left[\frac{4\Delta_2 \Lambda_1 \Lambda_2 (\Lambda_1 + \Lambda_2)}{4\Delta_1 \Lambda_1 \Lambda_2 (\Lambda_1 + \Lambda_2) - e^{-2r-\tau} \Delta_2^2 (3\Lambda_1^2 + 2\Lambda_1 \Lambda_2 + 3\Lambda_2^2)} \right]. \quad (11)$$

Let us notice that, for $\tau = n_{\text{th}} = R = 0$ (ideal protocol) and $s = 0$ (input coherent states), Eq. (11) reduces to [1]

$$\delta_{\text{opt}} = \frac{1}{2} \arctan(1 + e^{-2r}). \quad (12)$$

The displacement-independent one-shot fidelity \mathcal{F}_S and the optimal angle δ_{opt} are still dependent on s , the input squeezing. Unfortunately, the optimization of the non-Gaussian resource based on the choice [Eq. (11)] as the optimal angle would be practically unfeasible because the input squeezing is not known. In order to circumvent this problem, we introduce a suboptimal angle δ_{subopt} such that

$$\delta_{\text{subopt}} \equiv \delta_{\text{opt}}|_{s=\bar{s}}, \quad (13)$$

where \bar{s} is a fixed effective value of the input squeezing chosen, according to a suitable criterion that will be clarified below, in the range of possible values of the squeezing parameter s .

In the following we will express the squeezing parameters r and s in decibels, according to the relation [49]

$$\kappa \text{ (dB)} = 10 \log_{10} e^{2\kappa}, \quad \kappa = s, r. \quad (14)$$

The practical rationale for introducing a suboptimal characterization in the maximization of the output fidelity is based on the observation that the assumption of a completely random degree of input squeezing s is clearly unrealistic. It is instead very sensible to consider that the range of possible values of s falls in a window $[0, s_{\text{max}}]$ dB. Indeed, to date, the experimentally reachable values of squeezing fall roughly in such a range with $s_{\text{max}} \simeq 10$ dB [50].

We can then study the behavior of \mathcal{F}_S corresponding to the angle δ_{subopt} as a function of the effective input squeezing parameter \bar{s} , at fixed squeezing parameters of the resource and of the input state, respectively r and s , and at fixed loss parameters τ , n_{th} , and R . Figure 1 shows that \mathcal{F}_S is quite insensitive to the value of \bar{s} . By assuming the realistic range $s \in [0, 10]$ dB, the choice of a suboptimal angle such that $\bar{s} = 5$ dB (average value of the interval) leads to a decrease of the optimized fidelity, as compared to the choice of δ_{opt} , of at most 0.3% in ideal conditions, and even smaller in realistic conditions. In other words, the teleportation fidelity

is essentially constant in the considered interval of variability for the angle δ . Therefore, throughout the following, we fix $s = \bar{s} = 5$ dB in Eq. (11) to make it s independent.

In Fig. 2, we plot the teleportation fidelity associated with the various considered resources (Gaussian twin beam, optimized two-mode squeezed Bell-like state, two-mode squeezed photon-subtracted state) both for the ideal protocol (panel I) and for the nonideal protocol (panel II). We see that, at fixed (finite) squeezing r of the resource, the Gaussian twin beam is always outperformed by the optimal non-Gaussian squeezed Bell resource in the ideal protocol. It is worth noting that for very high values of the squeezing r , the advantage of the non-Gaussian resources fades and Gaussian twin beams perform, in practice, equally well for the teleportation of the considered input states. This reflects the well-known fact that, by using the ideal VBK protocol and an ideal




FIG. 1. (Color online) One-shot fidelity \mathcal{F}_S at fixed s , and as a function of the angle δ parametrized by \bar{s} , expressed in dB, i.e., $\delta(\bar{s}) = \delta_{\text{subopt}}$ [see Eq. (13)], both in the instance of the ideal protocol $\tau = n_{\text{th}} = R = 0$ (full lines), and of a nonideal protocol, with $\tau = 0.1$, $n_{\text{th}} = 0$, and $R^2 = 0.05$ (dashed lines). The one-shot fidelities are drawn for three different values of the input squeezing: $s = 0, 5, 10$ dB. The curves are ordered from top to bottom for increasing s .




FIG. 2. (Color online) One-shot fidelity \mathcal{F}_S as a function of the squeezing parameter r of the entangled resource, expressed in dB, for the suboptimal squeezed Bell resource (full black line), for the photon-subtracted squeezed resource (dotted red line) and for the twin-beam resource (dashed blue line), in the instance of the ideal protocol (panel I), $\tau = n_{\text{th}} = R = 0$, and the nonideal protocol (panel II), with $\tau = 0.1$, $n_{\text{th}} = 0$, and $R^2 = 0.05$. The one-shot fidelities are drawn for three different values of the input squeezing: $s = 0, 5$, and 10 dB. In the plots, the fidelity corresponding to the photon-added squeezed resource has been omitted as it is always lower than the ones corresponding to the photon-subtracted squeezed resource and the squeezed Bell resource. The curves are ordered from top to bottom for increasing s .

Einstein-Podolsky-Rosen resource (corresponding, e.g., to a twin beam in the limit $r \rightarrow \infty$), any quantum state can be unconditionally teleported with unit fidelity [30]. All the one-shot fidelities decrease for increasing squeezing s of the input and, interestingly, in the nonideal protocol they achieve a maximum at a finite value r_m of the squeezing r of the resource. The optimal squeezed Bell resource and the twin beam share the same $r_m \simeq 16$ dB and coincide at that point. In Fig. 2 we also plot the one-shot fidelities associated with the two-mode photon-subtracted squeezed states, Eq. (6). The two-mode photon-subtracted squeezed state always outperforms the twin beam in the ideal protocol, and at low and intermediate values of the resource squeezing r in the nonideal case. It is always outperformed by the optimized squeezed Bell resource. We note that, on the other hand, the two-mode photon-added squeezed states always exhibit a performance worse than the two-mode photon-subtracted squeezed states and the squeezed Bell resources (the corresponding fidelities are omitted in the plots for clarity). In a given range of the squeezing r , $|\psi_{\text{PSS}}\rangle_{12}$ and $|\psi_{\text{SB}}\rangle_{12}$ exhibit comparable levels in the fidelity of teleportation. In conclusion, properly optimized non-Gaussian resources maximize the fidelity of teleportation of squeezed coherent states both in the ideal and imperfect VBK protocols, outperforming the corresponding Gaussian resources. In the next section we carry out a similar analysis with the aim of identifying the optimal strategy that maximizes the reproduction at the output of the input squeezing.

IV. TELEPORTATION OF QUADRATURE MOMENTS

In this section, we introduce a different approach to the optimization of the teleportation protocol, aimed at retaining and faithfully reproducing at the output the variances and thus the squeezing of the input state. The strategy is to constrain the first- and second-order moments of the output field to reproduce the ones of the input field, by exploiting the free parameters of the non-Gaussian resources. We introduce the mean values $\langle Z_j \rangle = \text{Tr}[Z_j \rho_j]$, with $Z_j = X_j, P_j$ ($j = \text{in, out}$), and the variances $\langle \Delta Z_j^2 \rangle = \text{Tr}[Z_j^2 \rho_j] - \text{Tr}[Z_j \rho_j]^2$, and $\langle \Delta(X_j P_j)_S \rangle = \text{Tr}[\frac{1}{2}(X_j P_j + P_j X_j) \rho_j] -$

$\text{Tr}[X_j \rho_j] \text{Tr}[P_j \rho_j]$ (the cross-quadrature variance, with S denoting the symmetrization) of the quadrature operators $X_j = \frac{1}{\sqrt{2}}(a_j + a_j^\dagger)$, $P_j = \frac{i}{\sqrt{2}}(a_j^\dagger - a_j)$, associated with the single-mode input state ρ_{in} and the output state ρ_{out} of the teleportation protocol. The explicit expressions for the quantities $\langle Z_j \rangle$, $\langle \Delta Z_j^2 \rangle$, and $\langle \Delta(X_j P_j)_S \rangle$ are reported in the Appendix B.

The quantities measuring the deviation of the output from the input are the differences between the output and input first and second quadrature moments:

$$\begin{aligned} D(X) &\equiv \langle X_{\text{out}} \rangle - \langle X_{\text{in}} \rangle = (\tilde{g} - 1) \langle X_{\text{in}} \rangle, \\ D(P) &\equiv \langle P_{\text{out}} \rangle - \langle P_{\text{in}} \rangle = (\tilde{g} - 1) \langle P_{\text{in}} \rangle, \\ D(\Delta X^2) &\equiv \langle \Delta X_{\text{out}}^2 \rangle - \langle \Delta X_{\text{in}}^2 \rangle = (\tilde{g}^2 - 1) \langle \Delta X_{\text{in}}^2 \rangle + \Sigma, \\ D(\Delta P^2) &\equiv \langle \Delta P_{\text{out}}^2 \rangle - \langle \Delta P_{\text{in}}^2 \rangle = (\tilde{g}^2 - 1) \langle \Delta P_{\text{in}}^2 \rangle + \Sigma, \\ D(\Delta(XP)_S) &\equiv \langle \Delta(X_{\text{out}} P_{\text{out}})_S \rangle - \langle \Delta(X_{\text{in}} P_{\text{in}})_S \rangle \\ &= (\tilde{g}^2 - 1) \langle \Delta(X_{\text{in}} P_{\text{in}})_S \rangle, \end{aligned}$$

with Σ given by Eq. (B11). From the above equations, we see that the assumption $\tilde{g} = 1$ (i.e., $g = 1/T$) yields $D(X) = D(P) = D(\Delta(XP)_S) = 0$ and $D(\Delta X^2) = D(\Delta P^2) = \Sigma|_{\tilde{g}=1}$. Therefore, for $\tilde{g} = 1$, the input and output fields possess equal average position and momentum (equal first moments), and equal cross-quadrature variance; then the optimization procedure reduces to the minimization of the quantity $\Sigma|_{\tilde{g}=1}$ with respect to the free parameters of the non-Gaussian squeezed Bell resource, i.e., $\min_{\delta, \theta, \phi} \Sigma|_{\tilde{g}=1}$. Moreover, as for the optimization procedure discussed in Sec. III, it can be shown that the optimal choice for ϕ and θ is, once again, $\phi = \pi$ and $\theta = 0$. The optimization on the remaining free parameter δ yields the optimal value δ_{optvar} :

$$\delta_{\text{optvar}} = \frac{1}{2} \arctan \left[\frac{(1 + e^{\frac{5}{2}})^2 - e^{4r}(1 - e^{\frac{5}{2}})^2}{(1 + e^{\frac{5}{2}})^2 + e^{4r}(1 - e^{\frac{5}{2}})^2} \right]. \quad (15)$$

The optimal angle δ_{optvar} , corresponding to the minimization of the differences $D(\Delta X^2)$ and $D(\Delta P^2)$ between the output and input quadrature variances, is independent of R , at variance with the optimal value δ_{opt} , Eq. (11), corresponding to the maximization of the teleportation fidelity. It is also important

to note that in this case there are no questions related to a dependence on the input squeezing s . For $\tau = 0$, Eq. (15) reduces to $\delta_{\text{optvar}} = \pi/8$. Such a value is equal to the asymptotic value given by Eq. (12) for $r \rightarrow \infty$, so that in this extreme limit the two optimization procedures become equivalent. In the particular cases of photon-added and photon-subtracted resources, no optimization procedure can be carried out, and the parameter δ is simply a given specific function of r (see Sec. II). We remark that, having automatically zero difference in the cross-quadrature variance at $\tilde{g} = 1$, finding the angles that minimize $D(\Delta X^2)$ and $D(\Delta P^2)$ precisely solves the problem of achieving the optimal teleportation of both the first moments and the full covariance matrix of the input state at once.

In order to compare the performances of the Gaussian and non-Gaussian resources, and to emphasize the improvement of the efficiency of teleportation with squeezed Bell-like states, we consider first the instance of ideal protocol ($\tau = 0$, $n_{\text{th}} = 0$, $R = 0$), and compute, and explicitly report below, the output variances $\langle \Delta Z_{\text{out}}^2 \rangle_f$ of the teleported state associated with non-Gaussian resources [i.e., optimized squeezed Bell-like states ($f = \text{SB}$), photon-added squeezed states ($f = \text{PAS}$), and photon-subtracted squeezed states ($f = \text{PSS}$)], and with Gaussian resources, i.e., twin beams ($f = \text{TwB}$). From Eqs. (B8)–(B13), we get

$$\langle \Delta Z_{\text{out}}^2 \rangle_{\text{TwB}} = \langle \Delta Z_{\text{in}}^2 \rangle + e^{-2r}, \quad (16)$$

$$\langle \Delta Z_{\text{out}}^2 \rangle_{\text{PAS}} = \langle \Delta Z_{\text{in}}^2 \rangle + e^{-2r} \left[1 + \frac{2e^{-2r}(1 + e^{-2r})}{1 + e^{-4r}} \right], \quad (17)$$

$$\langle \Delta Z_{\text{out}}^2 \rangle_{\text{PSS}} = \langle \Delta Z_{\text{in}}^2 \rangle + e^{-2r} \left[1 - \frac{2e^{-2r}(1 - e^{-2r})}{1 + e^{-4r}} \right], \quad (18)$$

$$\langle \Delta Z_{\text{out}}^2 \rangle_{\text{SB}} = \langle \Delta Z_{\text{in}}^2 \rangle + e^{-2r} (2 - \sqrt{2}). \quad (19)$$

Equation (19) is derived by exploiting the optimal angle [Eq. (15)], which reduces to Eq. (12) in the ideal case. Independently of the resource, the teleportation process will in general result in an amplification of the input variance. However, the use of non-Gaussian optimized resources, compared to the Gaussian ones, reduces sensibly the amplification of the variances at the output. Looking at Eq. (16), we see that the teleportation with the twin-beam resource produces an excess, quantified by the exponential term e^{-2r} , of the output variance with respect to the input one. On the other hand, the use of the non-Gaussian squeezed Bell resource [Eq. (19)] yields a reduction in the excess of the output variance with respect to the input one by a factor $(2 - \sqrt{2})$. Let us now analyze the behaviors of the photon-added squeezed resources and of the photon-subtracted squeezed resources, Eqs. (17) and (18), respectively. We observe that, in analogy with the findings of the previous section, the photon-subtracted squeezed resources exhibit an intermediate behavior in the ideal protocol; indeed, for low values of r they perform better than the Gaussian twin beam, but worse than the optimized squeezed Bell states. The photon-added squeezed resources perform worse than both the twin-beam and the other non-Gaussian resources. These considerations follow straightforwardly from a quantitative analysis of the terms associated with the excess of the output variance in Eqs. (17) and (18). Moreover, again in analogy




FIG. 3. (Color online) Output variances $\langle \Delta X_{\text{out}}^2 \rangle$ and $\langle \Delta P_{\text{out}}^2 \rangle$ as a function of the squeezing parameter r of the resource, expressed in dB, for the optimized squeezed Bell resource (full black line), for the photon-subtracted squeezed resource (dotted red line), and for the twin-beam resource (dashed blue line). Panels I and II: Ideal protocol. Panels III and IV: Nonideal protocol. The various curves are to be compared with the given input variances of the input single-mode squeezed coherent state (horizontal solid lines). The squeezing of the input state is fixed at $s = 5$ dB and $\varphi = 0$. In the nonideal protocol, the experimental parameters are fixed at $\tau = 0.1$, $n_{\text{th}} = 0$, and $R^2 = 0.05$. In the plots, the variances associated with the photon-added squeezed resource have been omitted as they are always larger than the ones corresponding to the photon-subtracted squeezed resource and the squeezed Bell resource.

with the analysis of the optimal fidelity, for low values of r , there exists a region in which the performance of photon-subtracted squeezed states and optimized squeezed Bell states are comparable. Finally, again in analogy with the case of the fidelity optimization, the output variance associated with the Gaussian twin beam and with the optimized squeezed Bell states coincide at a specific, large value of r , at which the two resources become identical.

The input variances $\langle \Delta Z_{in}^2 \rangle$ [Eqs. (B3) and (B4)] and the output variances $\langle \Delta X_{out}^2 \rangle$ are plotted in panels I and II of Fig. 3 for the ideal VKB protocol and in panels III and IV of Fig. 3 for the nonideal protocol.

In the instance of realistic protocol, for small-resource squeezing degree r , similar conclusions can be drawn, leading to the same hierarchy among the entangled resources. However, analogously to the behavior of the teleportation fidelity, for high values of r the photon-subtracted squeezed resources are very sensitive to decoherence. In fact, the performance of such resources worsens more than the Gaussian twin beam for r greater than a specific finite threshold value.

Rather than minimizing the differences between output and input quadrature variances, one might be naively tempted to consider minimizing the difference between the ratio of the output variances $\langle \Delta X_{out}^2 \rangle / \langle \Delta P_{out}^2 \rangle$ and the ratio of the input variances $\langle \Delta X_{in}^2 \rangle / \langle \Delta P_{in}^2 \rangle$. This quantity might appear to be of some interest because it is a good measure of how well squeezing is teleported in all those cases in which the input and output quadrature variances are very different, that is, those situations in which the statistical moments are teleported with very low efficiency. However, it is of little use to preserve formally a scale parameter if the noise on the quadrature averages grows out of control. The procedure of minimizing the difference between output and input quadrature statistical moments is the only one that guarantees the simultaneous preservation of the squeezing degree and the reduction of the excess noise on the output averages and statistical moments of the field observables.

V. CONCLUSIONS

We have studied the efficiency of the VBK CV quantum teleportation protocol for the transmission of quantum states and averages of observables by using optimized non-Gaussian entangled resources. We have considered the problem of teleporting Gaussian squeezed and coherent states, i.e., input states with two unknown parameters, the coherent amplitude and the squeezing. The non-Gaussian resources (squeezed Bell states) are endowed with free parameters that can be tuned to maximize the teleportation efficiency either of the state or of physical quantities such as squeezing, quadrature averages, and statistical moments. We have discussed two different optimization procedures: the maximization of the teleportation fidelity of the state, and the optimization of the teleportation of average values and variances of the field quadratures. The first procedure maximizes the similarity in phase space between the teleported and the input state, while the second one maximizes the preservation at the output of the displacement and squeezing contents of the input.

We have shown that optimized non-Gaussian entangled resources such as the squeezed Bell states, as well as other

more conventional non-Gaussian entangled resources, such as the two-mode squeezed photon-subtracted states, outperform, in the realistic intervals of the squeezing parameter r of the entangled resource achievable with the current technology, entangled Gaussian resources both for the maximization of the teleportation fidelity and for the maximal preservation of the input squeezing and statistical moments. These findings are consistent and go in line with previous results on the improvement of various quantum information protocols replacing Gaussian with suitably identified non-Gaussian resources [1,2,6–9]. In the process, we have found that the two optimal values of the resource angle δ associated with the two optimization procedures are different and identified, respectively, by Eqs. (11) and (15). This nonequivalence is connected to the fact that, when using entangled non-Gaussian resources with free parameters that are amenable to optimization, the fidelity is closely related to the form of the different input properties that one wishes to teleport, e.g., quasiprobability distribution in the phase space, squeezing, statistical moments of higher order, and so on. Different quantities correspond to different optimal teleportation strategies.

Finally, regarding the VBK protocol, it is worth mentioning that the maximization of the teleportation fidelity corresponds to the maximization of the squared modulus of the overlap between the input and the output (teleported) state, without taking into account the characteristics of the output with respect to the input state. Therefore, part of the non-Gaussian character of the entangled resource is unavoidably transferred to the output state. The latter then acquires unavoidably a certain degree of non-Gaussianity, even in the presence of pure Gaussian inputs. Moreover, as verified in the case of nonideal protocols, the output state is also strongly affected by decoherence. Thus, in order to recover the purity and the Gaussianity of the teleported state, purification and Gaussification protocols should be implemented serially after transmission through the teleportation channel is completed [51]. If the second (squeezing preserving) procedure is considered instead, the possible deformation of the Gaussian character is not so relevant, because the shape reproduction is not the main goal, while purification procedures are again needed to correct for the extra noise added during teleportation when finite entanglement and realistic conditions are considered.

An important open problem is determining a proper teleportation benchmark for the class of Gaussian input states with unknown displacement and squeezing. Such a benchmark is expected to be certainly smaller than 50% in terms of teleportation fidelity, the latter being the benchmark for purely coherent input states with completely random displacement in phase space [45,46]. Our results indicate that optimized non-Gaussian entangled resources will allow one to beat the classical benchmark, thus achieving unambiguous quantum state transmission via a truly quantum teleportation, with a smaller amount of nonclassical resources, such as squeezing and entanglement, compared to the case of shared Gaussian twin-beam resources. In this context, Fig. 2 provides strong and encouraging evidence that suitable uses of non-Gaussianity in tailored resources, feasible with current technology [24–28], may lead to a genuine demonstration of CV quantum teleportation of displaced squeezed states in realistic conditions of the experimental apparatus. This would constitute a crucial step

forward after the successful recent experimental achievement of the quantum storage of a displaced squeezed thermal state of light into an atomic ensemble memory [52].

ACKNOWLEDGMENT

We acknowledge financial support from the European Union under the FP7 STREP Project HIP (Hybrid Information Processing), Grant Agreement No. 221889.

APPENDIX A: INPUT STATES, ENTANGLED RESOURCES, AND OUTPUT STATES

Here we report the characteristic functions for the single-mode input states and for the two-mode entangled resources. The characteristic function for the coherent squeezed states [Eq. (3)], i.e., $\chi_{\text{in}}(\alpha) = {}_{\text{in}}\langle\psi_{\text{CS}}|D_{\text{in}}(\alpha)|\psi_{\text{CS}}\rangle_{\text{in}}$, reads

$$\chi_{\text{in}}(\alpha) = \exp\left[\frac{1}{2}(\alpha\beta^* - \alpha^*\beta) - \frac{1}{2}(\alpha + \beta)\cosh s + (\alpha^* + \beta^*)e^{i\varphi}\sinh s\right]^2. \quad (\text{A1})$$

The characteristic function for the squeezed Bell-like resource [Eq. (4)], i.e., $\chi_{\text{SB}}(\alpha_1, \alpha_2) = {}_{12}\langle\psi_{\text{SB}}|D_1(\alpha_1)D_2(\alpha_2)|\psi_{\text{SB}}\rangle_{12}$, reads

$$\chi_{\text{SB}}(\alpha_1, \alpha_2) = e^{-\frac{1}{2}(|\xi_1|^2 + |\xi_2|^2)} [1 + \sin\delta \cos\delta (e^{i\theta}\xi_1\xi_2 + e^{-i\theta}\xi_1^*\xi_2^*) + \sin^2\delta (|\xi_1|^2|\xi_2|^2 - |\xi_1|^2 - |\xi_2|^2)], \quad (\text{A2})$$

where the complex variables ξ_k are defined as

$$\xi_k = \alpha_k \cosh r + \alpha_l^* e^{i\phi} \sinh r \quad (k, l = 1, 2; k \neq l). \quad (\text{A3})$$

It is worth noticing that, for $\delta = 0$, Eq. (A2) reduces to the well-known Gaussian characteristic function of the twin beam. Given the characteristic functions for the single-mode input state and for the two-mode entangled resource, Eqs. (A1) and (A2), respectively, it is straightforward to obtain the characteristic function for the single-mode output state of the teleportation protocol by using Eq. (1) and replacing χ_{res} with χ_{SB} .

APPENDIX B: MEAN VALUES AND VARIANCES OF THE QUADRATURES

Here we report the analytical expressions for the mean values $\langle Z_j \rangle = \text{Tr}[Z_j \rho_j]$, with $Z_j = X_j, P_j$ ($j = \text{in}, \text{out}$), and the variances $\langle \Delta Z_j^2 \rangle = \text{Tr}[Z_j^2 \rho_j] - \text{Tr}[Z_j \rho_j]^2$ of the quadrature operators $X_j = \frac{1}{\sqrt{2}}(a_j + a_j^\dagger)$, $P_j = \frac{i}{\sqrt{2}}(a_j^\dagger - a_j)$, associated with the single-mode input state ρ_{in} and the output state ρ_{out} of the teleportation protocol. We also compute the cross-quadrature variance $\langle \Delta(X_j P_j)_S \rangle = \text{Tr}[\frac{1}{2}(X_j P_j + P_j X_j) \rho_j] - \text{Tr}[X_j \rho_j] \text{Tr}[P_j \rho_j]$, associated with the nondiagonal term of the covariance matrix of the density operator,

where the subscript S denotes the symmetrization. The mean values and the variances associated with the input single-mode coherent squeezed state [Eq. (3)] can be computed easily:

$$\langle X_{\text{in}} \rangle = \frac{1}{\sqrt{2}}(\beta + \beta^*), \quad (\text{B1})$$

$$\langle P_{\text{in}} \rangle = \frac{i}{\sqrt{2}}(\beta^* - \beta), \quad (\text{B2})$$

and

$$\langle \Delta X_{\text{in}}^2 \rangle = \frac{1}{2}(\cosh 2s - \cos\varphi \sinh 2s), \quad (\text{B3})$$

$$\langle \Delta P_{\text{in}}^2 \rangle = \frac{1}{2}(\cosh 2s + \cos\varphi \sinh 2s), \quad (\text{B4})$$

$$\langle \Delta(X_{\text{in}} P_{\text{in}})_S \rangle = -\frac{1}{2} \sin\varphi \sinh 2s. \quad (\text{B5})$$

The mean values and the variances associated with the output single-mode teleported state, described by the characteristic function [Eq. (1)] read

$$\langle X_{\text{out}} \rangle = \frac{\tilde{g}}{\sqrt{2}}(\beta + \beta^*), \quad (\text{B6})$$

$$\langle P_{\text{out}} \rangle = \frac{i\tilde{g}}{\sqrt{2}}(\beta^* - \beta), \quad (\text{B7})$$

and

$$\langle \Delta X_{\text{out}}^2 \rangle = \tilde{g}^2 \langle \Delta X_{\text{in}}^2 \rangle + \Sigma, \quad (\text{B8})$$

$$\langle \Delta P_{\text{out}}^2 \rangle = \tilde{g}^2 \langle \Delta P_{\text{in}}^2 \rangle + \Sigma, \quad (\text{B9})$$

$$\langle \Delta(X_{\text{out}} P_{\text{out}})_S \rangle = -\frac{1}{2} \tilde{g}^2 \sin\varphi \sinh 2s, \quad (\text{B10})$$

with

$$\begin{aligned} \Sigma = & \Gamma_{\tau, R} + e^{-\frac{\tau}{2}} \tilde{g} \sin(\theta - \phi) \sin\phi \sin 2\delta \\ & - \frac{1}{4} e^{-2r - \tau} (1 + e^\tau \tilde{g}^2 - 2e^{\frac{\tau}{2}} \tilde{g} \cos\phi) \\ & \times [\cos 2\delta - \cos(\theta - \phi) \sin 2\delta - 2] \\ & - \frac{1}{4} e^{2r - \tau} (1 + e^\tau \tilde{g}^2 + 2e^{\frac{\tau}{2}} \tilde{g} \cos\phi) \\ & \times [\cos 2\delta + \cos(\theta - \phi) \sin 2\delta - 2]. \end{aligned} \quad (\text{B11})$$

For the particular choices $\tilde{g} = 1$, $\phi = \pi$, and $\theta = 0$, Eq. (B11) reduces to

$$\begin{aligned} \Sigma|_{\tilde{g}=1} = & \Gamma_{\tau, R}|_{g=1/T} - \frac{1}{4} e^{-2r - \tau} (1 + e^{\frac{\tau}{2}})^2 [\cos 2\delta + \sin 2\delta - 2] \\ & - \frac{1}{4} e^{2r - \tau} (1 - e^{\frac{\tau}{2}})^2 [\cos 2\delta - \sin 2\delta - 2]. \end{aligned} \quad (\text{B12})$$

In the instance of Gaussian resource ($\delta = 0$), such a quantity simplifies to

$$\Sigma_G = \Gamma_{\tau, R}|_{g=1/T} + \frac{1}{4} e^{-2r - \tau} (1 + e^{\frac{\tau}{2}})^2 + \frac{1}{4} e^{2r - \tau} (1 - e^{\frac{\tau}{2}})^2. \quad (\text{B13})$$

For suitable choices of δ in Eq. (B12) (see Sec. II), one can easily obtain the output variances associated with photon-added and photon-subtracted squeezed states.

[1] F. Dell'Anno, S. De Siena, L. Albano, and F. Illuminati, *Phys. Rev. A* **76**, 022301 (2007).

[2] F. Dell'Anno, S. De Siena, and F. Illuminati, *Phys. Rev. A* **81**, 012333 (2010).

[3] M. S. Kim, W. Son, V. Bužek, and P. L. Knight, *Phys. Rev. A* **65**, 032323 (2002).

[4] V. V. Dodonov and L. A. de Souza, *J. Opt. B* **7**, S490 (2005).

- [5] N. J. Cerf, O. Krüger, P. Navez, R. F. Werner, and M. M. Wolf, *Phys. Rev. Lett.* **95**, 070501 (2005).
- [6] T. Opatrný, G. Kurizki, and D.-G. Welsch, *Phys. Rev. A* **61**, 032302 (2000).
- [7] P. T. Cochrane, T. C. Ralph, and G. J. Milburn, *Phys. Rev. A* **65**, 062306 (2002).
- [8] S. Olivares, M. G. A. Paris, and R. Bonifacio, *Phys. Rev. A* **67**, 032314 (2003).
- [9] A. Kitagawa, M. Takeoka, M. Sasaki, and A. Chefles, *Phys. Rev. A* **73**, 042310 (2006).
- [10] F. Dell'Anno, S. De Siena, L. Albano, and F. Illuminati, *Eur. Phys. J. Special Topics* **160**, 115 (2008).
- [11] Y. Yang and F.-L. Li, *Phys. Rev. A* **80**, 022315 (2009).
- [12] G. Adesso, F. Dell'Anno, S. De Siena, F. Illuminati, and L. A. M. Souza, *Phys. Rev. A* **79**, 040305(R) (2009).
- [13] N. C. Menicucci, P. van Loock, M. Gu, C. Weedbrook, T. C. Ralph, and M. A. Nielsen, *Phys. Rev. Lett.* **97**, 110501 (2006).
- [14] M. M. Wolf, G. Giedke, and J. I. Cirac, *Phys. Rev. Lett.* **96**, 080502 (2006).
- [15] M. G. Genoni and M. G. A. Paris, *Phys. Rev. A* **82**, 052341 (2010).
- [16] F. Dell'Anno, S. De Siena, and F. Illuminati, *Phys. Rep.* **428**, 53 (2006).
- [17] T. Tyc and N. Korolkova, *New J. Phys.* **10**, 023041 (2008).
- [18] G. S. Agarwal and K. Tara, *Phys. Rev. A* **43**, 492 (1991).
- [19] G. Bjork and Y. Yamamoto, *Phys. Rev. A* **37**, 4229 (1988).
- [20] Z. Zhang and H. Fan, *Phys. Lett. A* **165**, 14 (1992).
- [21] M. Dakna, T. Anhut, T. Opatrný, L. Knöll, and D. G. Welsch, *Phys. Rev. A* **55**, 3184 (1997).
- [22] M. S. Kim, E. Park, P. L. Knight, and H. Jeong, *Phys. Rev. A* **71**, 043805 (2005).
- [23] D. Menzies and R. Filip, *Phys. Rev. A* **79**, 012313 (2009).
- [24] A. Zavatta, S. Viciani, and M. Bellini, *Science* **306**, 660 (2004).
- [25] A. I. Lvovsky and S. A. Babichev, *Phys. Rev. A* **66**, 011801 (2002).
- [26] J. Wenger, R. Tualle-Brouiri, and P. Grangier, *Phys. Rev. Lett.* **92**, 153601 (2004).
- [27] A. Ourjoumtsev, A. Dantan, R. Tualle-Brouiri, and P. Grangier, *Phys. Rev. Lett.* **98**, 030502 (2007).
- [28] V. Parigi, A. Zavatta, M. Kim, and M. Bellini, *Science* **317**, 1890 (2007).
- [29] A. Ourjoumtsev, H. Jeong, R. Tualle-Brouiri, and P. Grangier, *Nature (London)* **448**, 784 (2007).
- [30] L. Vaidman, *Phys. Rev. A* **49**, 1473 (1994).
- [31] S. L. Braunstein and H. J. Kimble, *Phys. Rev. Lett.* **80**, 869 (1998).
- [32] A. Furusawa, J. L. Sorensen, S. L. Braunstein, C. A. Fuchs, H. J. Kimble, and E. J. Polzik, *Science* **282**, 706 (1998).
- [33] W. P. Bowen, N. Treps, B. C. Buchler, R. Schnabel, T. C. Ralph, H.-A. Bachor, T. Symul, and P. K. Lam, *Phys. Rev. A* **67**, 032302 (2003).
- [34] T. C. Zhang, K. W. Chou, C. W. Chou, P. Lodahl, and H. J. Kimble, *Phys. Rev. A* **67**, 033802 (2003).
- [35] N. Takei, H. Yonezawa, T. Aoki, and A. Furusawa, *Phys. Rev. Lett.* **94**, 220502 (2005).
- [36] J. F. Sherson, H. Krauter, R. K. Olsson, B. Julsgaard, K. Hammerer, J. I. Cirac, and E. S. Polzik, *Nature (London)* **443**, 557 (2006).
- [37] N. Takei *et al.*, *Phys. Rev. A* **72**, 042304 (2005).
- [38] H. Yonezawa, S. L. Braunstein, and A. Furusawa, *Phys. Rev. Lett.* **99**, 110503 (2007).
- [39] S. L. Braunstein and P. van Loock, *Rev. Mod. Phys.* **77**, 513 (2005).
- [40] P. Marian and T. A. Marian, *Phys. Rev. A* **74**, 042306 (2006).
- [41] U. Leonhardt and H. Paul, *Phys. Rev. A* **48**, 4598 (1993).
- [42] D. Walls and G. Milburn, *Quantum Optics* (Springer, Berlin, 1994).
- [43] A. Serafini, M. G. A. Paris, F. Illuminati, and S. De Siena, *J. Opt. B* **7**, R19 (2005).
- [44] W. P. Bowen, N. Treps, B. C. Buchler, R. Schnabel, T. C. Ralph, T. Symul, and P. K. Lam, *IEEE J. Sel. Top. Quantum* **9**, 1519 (2003).
- [45] S. L. Braunstein, C. A. Fuchs, and H. J. Kimble, *J. Mod. Opt.* **47**, 267 (2000).
- [46] K. Hammerer, M. M. Wolf, E. S. Polzik, and J. I. Cirac, *Phys. Rev. Lett.* **94**, 150503 (2005).
- [47] G. Adesso and G. Chiribella, *Phys. Rev. Lett.* **100**, 170503 (2008).
- [48] M. Owari, M. B. Plenio, E. S. Polzik, A. Serafini, and M. M. Wolf, *New J. Phys.* **10**, 113014 (2008).
- [49] P. Kok and B. W. Lovett, *Introduction to Optical Quantum Information Processing* (Cambridge University Press, Cambridge, 2010).
- [50] S. Suzuki, H. Yonezawa, F. Kannari, M. Sasaki, and A. Furusawa, *Appl. Phys. Lett.* **89**, 061116 (2006); H. Vahlbruch, M. Mehmet, S. Chelkowski, B. Hage, A. Franzen, N. Lastzka, S. Gossler, K. Danzmann, and R. Schnabel, *Phys. Rev. Lett.* **100**, 033602 (2008).
- [51] O. Glöckl, U. L. Andersen, R. Filip, W. P. Bowen, and G. Leuchs, *Phys. Rev. Lett.* **97**, 053601 (2006); J. Heersink, Ch. Marquardt, R. Dong, R. Filip, S. Lorenz, G. Leuchs, and U. L. Andersen, *ibid.* **96**, 253601 (2006); A. Franzen, B. Hage, J. DiGuglielmo, J. Fiurasek, and R. Schnabel, *ibid.* **97**, 150505 (2006); B. Hage, A. Sambrowski, J. DiGuglielmo, A. Franzen, J. Fiurasek, and R. Schnabel, *Nature Phys.* **4**, 915 (2008); R. Dong, M. Lassen, J. Heersink, Ch. Marquardt, R. Filip, G. Leuchs, and U. L. Andersen, *ibid.* **4**, 919 (2008).
- [52] K. Jensen, W. Wasilewski, H. Krauter, T. Fernholz, B. M. Nielsen, A. Serafini, M. Owari, M. B. Plenio, M. M. Wolf, and E. S. Polzik, *Nature Phys.* **7**, 13 (2011).