

Perturbative correction to the ground-state properties of one-dimensional strongly interacting bosons in a harmonic trap

Francis N. C. Paraan* and Vladimir E. Korepin†

C. N. Yang Institute for Theoretical Physics, State University of New York at Stony Brook, New York 11794-3840, USA

(Received 29 November 2010; published 30 December 2010)

We calculate the first-order perturbation correction to the ground-state energy and chemical potential of a harmonically trapped boson gas with contact interactions about the infinite repulsion Tonks-Girardeau limit. With c denoting the interaction strength, we find that, for a large number of particles N , the $1/c$ correction to the ground-state energy increases as $N^{5/2}$, in contrast to the unperturbed Tonks-Girardeau value that is proportional to N^2 . We describe a thermodynamic scaling limit for the trapping frequency that yields an extensive ground-state energy and reproduces the zero temperature thermodynamics obtained by a local-density approximation.

DOI: [10.1103/PhysRevA.82.065603](https://doi.org/10.1103/PhysRevA.82.065603)

PACS number(s): 03.75.Hh, 05.30.Jp, 67.85.-d

I. INTRODUCTION

The realization of quasi-one-dimensional ultracold boson gases with tunable interaction parameters [1–3] and the succeeding advances in atom chip trap technology [4–7] have renewed interest in theoretical models of one-dimensional bosons with short-range interactions. Of particular relevance to workers in this field is the Lieb-Liniger model [8] in which contact interactions are described by Dirac delta functions. The suitability of this model in describing the low-temperature properties of these quasi-one-dimensional bosonic systems has been further strengthened by Olshanii’s analysis of the low-energy scattering of atoms under tight transverse harmonic confinement: The longitudinal s -wave scattering amplitudes are indeed reproduced by a one-dimensional pseudopotential proportional to a Dirac delta function [9]. The resulting demonstration that the magnitude of this effective delta interaction can be explicitly calculated from the three-dimensional atomic scattering length and the dimensions of the confining external trap [10] further strengthens the link between theoretical one-dimensional models and quasi-one-dimensional experiments.

Still, the free Lieb-Liniger model is quite an idealization for actual experiments because the atoms are generally longitudinally confined by an external potential, and thus much effort has been devoted to studying the effects of confinement of interacting bosons [9, 11, 12]. Introducing an external harmonic potential to the free Lieb-Liniger model of spinless bosons leads to the many-particle Schrödinger eigenvalue equation

$$\bar{E}^b \Psi^b = \left[\sum_{i=1}^N -\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x_i^2} + \frac{m\omega^2 x_i^2}{2} \right] \Psi^b - \frac{\hbar^2}{ma} \sum_{i<j} \delta(x_j - x_i) \Psi^b, \quad (1)$$

where m is the mass of each of the N atoms, ω is the angular frequency of the trap, and a is the one-dimensional scattering length. The superscript “b” refers to the bosonic nature of the labeled quantities. Measuring energy in units

of $\hbar\omega$ and length in oscillator units $\ell = \sqrt{\hbar/(m\omega)}$ gives the dimensionless eigenvalue equation

$$E^b \Psi^b = \left[\sum_{i=1}^N -\frac{1}{2} \frac{\partial^2}{\partial x_i^2} + \frac{1}{2} x_i^2 + c \sum_{i<j} \delta(x_j - x_i) \right] \Psi^b, \quad (2)$$

where we have introduced the dimensionless interaction strength $c = -\ell/a$. We consider here the repulsive case $c > 0$ (negative scattering length) to be specific. In the absence of a harmonic potential, the corresponding eigenvalue equation is solvable by the Bethe ansatz and, consequently, much is known about the ground-state and elementary excitations of this system [8], as well as the properties of the various correlation functions at zero and finite temperatures [13]. However, for the important case of harmonic confinement, an exact solution to this problem for general values of the interaction strength c is lacking. The sole exceptions are the two-particle case that is separable in relative and central coordinates [14], and the Tonks-Girardeau (TG) limit of infinite repulsion $c \rightarrow +\infty$ in which the exact N -particle wave functions are absolute values of Slater determinants [15]. These results, especially the latter, are prototypical examples of the fermion-boson duality derived by Girardeau [16] and later generalized by Cheon and Shigehara [17] for one-dimensional systems of particles having contact interactions. For finite values of the interaction parameter c , expressions for the atomic density and collective oscillation frequencies have been calculated using local-density approximations [9, 11, 18–20] and time-dependent density functional theory [20, 21], while formal expressions for the self-consistent Hartree-Fock equations for the single-particle density matrix have been obtained for general trapping potentials [12]. The analogous problem of a system of confined interacting fermions has also been treated by similar methods [22].

In this work, we will use the mentioned fermion-boson relationship to develop perturbative $1/c$ corrections to the ground-state energy and chemical potential of a harmonically confined interacting boson gas about the Tonks-Girardeau solution. The details of the specific fermion-boson mapping we employ here that utilizes a nonlocal pseudopotential [12] are given in Sec. II. In Sec. III we present our perturbation calculations for the general case of an N -particle system obtaining a closed-form analytical result that is calculable for any N . We analyze few-body cases and discuss the

*fparaan@ic.sunysb.edu

†korepin@max2.physics.sunysb.edu

thermodynamic limit $N \rightarrow \infty$ of our solution in Sec. IV. We summarize our results and give concluding remarks in Sec. V.

II. FERMION-BOSON MAPPING

In one dimension it has been demonstrated that a bosonic model with pairwise contact interactions of strength c can be mapped into a fermionic model with pairwise interactions of strength $1/c$ [17]. Specifically, given a fermionic wave function Φ^f that satisfies the eigenvalue equation

$$\left[\sum_{i=1}^N -\frac{1}{2} \frac{\partial^2}{\partial x_i^2} + \frac{1}{2} x_i^2 + \hat{V}^f \right] \Phi^f = E^f \Phi^f, \quad (3)$$

an appropriate choice of a pseudopotential operator \hat{V}^f allows us to make the following correspondence between the bosonic wave function Ψ^b and its fermionic counterpart Φ^f :

$$\Psi^b = \mathcal{A} \Phi^f(x_1, \dots, x_N), \quad (4)$$

where $\mathcal{A} \equiv \prod_{i < j} \text{sgn}(x_j - x_i)$ is a function that is completely antisymmetric under any transposition $x_i \leftrightarrow x_j$. Since the complex squares of the wave functions Ψ^b and Φ^f are identical, the energy eigenvalues E^b and E^f are equal. For our specific problem, a suitable fermionic pseudopotential operator has matrix elements in the coordinate representation given by [12]

$$\langle \varphi^f | \hat{V}^f | \phi^f \rangle = -\frac{4}{c} \sum_{i < j} \int \lim_{r_{ij} \rightarrow 0} \left[\frac{\partial \varphi^{f*}}{\partial r_{ij}} \frac{\partial \phi^f}{\partial r_{ij}} \right] dR_{ij}, \quad (5)$$

where $r_{ij} = x_j - x_i$ and $R_{ij} = \frac{1}{2}(x_j + x_i)$ are relative and central coordinates, respectively, and $\varphi^f(x_1, \dots, x_N)$ and $\phi^f(x_1, \dots, x_N)$ are the coordinate space wave functions corresponding to the fermionic state kets $|\varphi^f\rangle$ and $|\phi^f\rangle$, respectively.

In the infinite repulsion limit, the bosonic eigenvalue equation (2) is solved by the absolute value of the ground-state Slater determinant [15]:

$$\Psi_{\text{TG}}^b = \frac{1}{\sqrt{N!}} |\det \psi_n(x_m)| \equiv \mathcal{A} \Phi_{\text{TG}}^f, \quad (6)$$

where Φ_{TG}^f is the fermionic ground-state wave function, $\{x_m\}$ are the coordinates of the atoms, and $\{\psi_n\}$ are the N lowest energy single-particle harmonic oscillator eigenfunctions $\psi_n(x) = \pi^{-1/4} (2^n n!)^{-1/2} H_n(x) e^{-x^2/2}$. The $H_n(x)$ appearing here are the usual Hermite polynomials. The corresponding energy of this TG ground state (in units of $\hbar\omega$) is

$$E_{\text{TG}}^b = \frac{1}{2} N^2. \quad (7)$$

For a finite and large repulsion strength we may therefore use the quantity $1/c \ll 1$ as a perturbation parameter for the fermionic problem (3) so that ordinary first-order perturbation theory gives the desired correction:

$$E_0^b = E_0^f = \frac{1}{2} N^2 + \langle \Phi_{\text{TG}}^f | \hat{V}^f | \Phi_{\text{TG}}^f \rangle + O(1/c^2). \quad (8)$$

III. GROUND-STATE ENERGY CORRECTION

In this section we work mainly in the fermionic sector and omit the superscripts “f” for brevity. Here, our objective is to explicitly evaluate the leading correction $\delta E \equiv \langle \Phi_{\text{TG}} | \hat{V} | \Phi_{\text{TG}} \rangle$ to the ground-state energy E_0 . Since Φ_{TG} is a Slater determinant and the fermionic interaction operator \hat{V} is a sum of two-body operators \hat{v} having matrix elements

$$v_{klmn} = -\frac{4}{c} \int \lim_{r \rightarrow 0} \left\{ \frac{\partial [\psi_k(x_1) \psi_l(x_2)]^*}{\partial r} \frac{\partial [\psi_m(x_1) \psi_n(x_2)]}{\partial r} \right\} dR, \quad (9)$$

we may calculate the perturbation δE using the Slater-Condon rule $\delta E = \sum_{k < l} (v_{klkl} - v_{kllk})$ [23]. Prior to calculating the derivatives appearing inside the integral (9), we must be careful to write the coordinates $x_1 = R + \frac{1}{2}r$ and $x_2 = R - \frac{1}{2}r$ in terms of the relative and central coordinates r and R . The symmetry of the integrand allows us to write and define $v_{klkl} = -v_{kllk} \equiv \tilde{v}_{kl}$, where

$$\tilde{v}_{kl} = -\frac{4}{c} \int \lim_{r \rightarrow 0} \left\{ \frac{\partial [\psi_k(x_1) \psi_l(x_2)]}{\partial r} \right\}^2 dR. \quad (10)$$

Thus, the leading correction becomes $\delta E = 2 \sum_{k < l} \tilde{v}_{kl}$ and is always negative, as expected. After some manipulation, we obtain a finite series expression for the energy correction that may be evaluated for any number N of atoms:

$$\begin{aligned} \delta E &= \frac{1}{c} \sqrt{\frac{2}{\pi^3}} \sum_{l=1}^{N-1} \frac{\Gamma(l - \frac{1}{2})}{\Gamma(l+1)} \sum_{k=0}^{l-1} \frac{(l-k)^2 \Gamma(k - \frac{1}{2})}{\Gamma(k+1)} \\ &\times {}_3F_2 \left[\begin{matrix} \frac{3}{2}, -k, -l \\ \frac{3}{2} - k, \frac{3}{2} - l \end{matrix}; 1 \right]. \end{aligned} \quad (11)$$

IV. FEW-PARTICLE RESULTS AND THERMODYNAMIC LIMIT

The special case of $N = 2$ particles is separable in relative and central coordinates and the resulting eigenvalue problem for the ground-state energy is exactly solvable. Upon imposing vanishing boundary conditions on the two-body wave function at infinity, we find that the ground-state energy E_0 of the trapped two-boson system satisfies the transcendental equation

$$\frac{2\Gamma[\frac{1}{2}(1 + E_0)] \tan[\frac{1}{2}(1 - E_0)\pi]}{\Gamma[E_0/2]} = -\frac{c}{\sqrt{2}}, \quad (12)$$

where $E_0 \in [1, 2]$. Our perturbative result (11) gives $\delta E = -(2/c)\sqrt{2}/\pi$ and coincides with the leading correction term in the $1/c$ series expansion of this exact solution about the TG ground-state energy $E_{\text{TG}} = 2$. Also, an essentially identical result was obtained by Sen [24] for the reduced single-particle problem using a different local pseudopotential involving the second derivative of the delta function $\propto \delta''(x_2 - x_1)$.¹ A

¹As noted in Ref. [12], Sen’s potential and the pseudopotential used here yield identical interaction matrix elements between states described by continuous fermionic wave functions. Since our calculations involve continuous Slater determinants, the distinction between pseudopotentials is not relevant here.




FIG. 1. (Color online) The first-order perturbation result (dashed line) for the ground-state energy of two delta interacting bosons in a harmonic trap is compared to the exact solution (solid line). The ground-state energy E_{TG} in the infinite repulsion limit is given by the horizontal dotted line.

comparison of our first-order perturbation result and the exact two-particle ground-state energy is shown in Fig. 1 and we observe good agreement between the two results in the strongly interacting regime $c \gg 10$.

Before discussing the situation for more than two particles, let us restore units and rewrite the perturbed ground-state energy as

$$E_0(N) = \frac{1}{2}\hbar\omega N^2[1 + 2\alpha(N)/c] + O(1/c^2), \quad (13)$$

where $\alpha(N)$ is a dimensionless function of N . For values of N up to 10^3 , the magnitude of the scaled first-order correction $-c\delta E(N)/N^2$ is plotted in Fig. 2(a) as a function of N on a double-logarithmic plot. Inspection of this graph suggests a simple power-law scaling for the first-order correction with large N . Leading order asymptotic analysis reveals that the partial sums in Eq. (11) scale as $\sim N^{3/2}$ for large N so that $N^2\alpha(N) \sim N^{5/2}$:

$$E_0(N) \approx \frac{1}{2}\hbar\omega N^2[1 + 2\alpha_0\sqrt{N}/c], \quad N \rightarrow \infty, \quad (14)$$




FIG. 2. (Color online) (a) The scaled perturbation $-c\delta E/N^2$ (dots) grows as a power law \sqrt{N} in the limit $N \rightarrow \infty$. (b) Our large- N result is consistent with numerical calculations (solid line) in the Thomas-Fermi approximation for $\sqrt{N}/c \ll 1$. The dashed lines are regression fits $c\delta E \approx -0.408N^{5/2}$ calculated from values $N \in [100, 1000]$. The prefactor -0.408 is independent of the particular values chosen in this interval.

where α_0 is a constant number. Indeed, for as few as $N \gtrsim 15$ particles the factor $\alpha(N)$ is quite well approximated (within 1%) by the function $\alpha_0\sqrt{N}$ with $\alpha_0 \approx -0.408$. In other words, the correction factor $\alpha(N)$ reaches its asymptotic scaling behavior for systems as small as $N \gg O(10^1)$.

To obtain a thermodynamic limit with an extensive ground-state energy, we observe that, in addition to sending the number of particles to infinity, we must also require the trapping frequency ω to vanish as $1/N$. As we shall see, this condition allows us to reproduce the Thomas-Fermi results near the Tonks-Girardeau limit. This scaling requirement implies that the quantity ℓ/\sqrt{N} approaches a constant in the thermodynamic limit, in contrast to the thermodynamic limit in the case of a Lieb-Liniger model gas confined in a flat-bottomed box, in which the linear dimension of the system is taken to scale proportionally with particle number [25]. Looking back at our asymptotic expression $\alpha(N) \approx \alpha_0\sqrt{N}$, we find that the quantity \sqrt{N}/c approaches a constant value as $N \rightarrow \infty$ in our prescribed thermodynamic limit. This is precisely the condition used by Ma and Yang [11] to obtain the ground-state energy of the trapped interacting boson gas in the Thomas-Fermi formalism, which we reproduce here in Fig. 2(b). We find that our first-order $1/c$ result is reliable for $\sqrt{N}/c \lesssim 0.1$, which means that, for a typical experimental setup with hundreds of atoms, first-order perturbation theory and the Thomas-Fermi result coincide only in the extreme repulsion limit $c \gtrsim 10^2$.

If we now define the Tonks-Girardeau limit chemical potential as $\mu_{\text{TG}} \equiv \lim_{N \rightarrow \infty, \omega \rightarrow 0} \hbar\omega N$ and the scaled interaction parameter as $\gamma \equiv \lim_{N \rightarrow \infty, \omega \rightarrow 0} c/\sqrt{N}$, we obtain the zero temperature chemical potential

$$\mu \approx \mu_{\text{TG}}[1 + \frac{5}{2}\alpha_0/\gamma]. \quad (15)$$

The first term in this expression corresponds to the chemical potential of free fermions in a one-dimensional harmonic trap while the last term gives the reduction in the chemical potential due to the finite repulsion correction and serves as a measure of the departure of the system from the unitarity limit.

V. CONCLUDING REMARKS

In this work we have calculated the first-order finite repulsion correction to the ground-state energy of harmonically trapped bosons having contact interactions for any finite number N of particles. For $N \gg O(10^1)$ we found that, for a fixed interaction strength, this correction scales as a power law $N^{5/2}$, which allowed us to describe a thermodynamic limit that reproduces known results from Thomas-Fermi approaches. This contribution clarifies the smooth transition of the ground-state properties of a harmonically confined interacting boson system as the particle number goes from finite N to infinity near the Tonks-Girardeau limit. We have demonstrated that, in this strongly interacting regime, to at least leading order in $1/c$, finite number effects are negligible in current experimental situations that have $\sim 10^2$ atoms.

A natural extension of this work would involve higher order corrections to the ground-state energy and many-body wave function, as was done recently for a wedge-shaped trapping

potential [26]. If we take the set of all fermionic Slater determinants as an expansion basis for ordinary perturbation theory about the TG limit, we discover that the perturbing pseudopotential couples the ground state to an infinite number of excited states. We therefore expect a complicated analytical result for the second-order energy correction resulting in a numerical problem that may require a truncation of the corresponding Hilbert space. However, on the basis of the agreement between our asymptotic results and the Thomas-Fermi calculation [Fig. 2(b)], we conjecture that the second-order correction scales as $N^3/c^2 > 0$ in the thermodynamic limit.

ACKNOWLEDGMENTS

During the course of this work many insightful discussions were shared with A. G. Abanov, D. Schneble, and F. Franchini. The authors acknowledge financial support by NSF Grant No. DMS-0905744.

-
- [1] A. Görlitz *et al.*, *Phys. Rev. Lett.* **87**, 130402 (2001).
 - [2] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G. V. Shlyapnikov, T. W. Hänsch, and I. Bloch, *Nature (London)* **429**, 277 (2004).
 - [3] T. Kinoshita, T. Wenger, and D. S. Weiss, *Science* **305**, 1125 (2004).
 - [4] M. Horikoshi and K. Nakagawa, *Appl. Phys. B* **82**, 363 (2006).
 - [5] L. Della Pietra, S. Aigner, C. vom Hagen, S. Groth, I. Bar-Joseph, H. J. Lezec, and J. Schmiedmayer, *Phys. Rev. A* **75**, 063604 (2007).
 - [6] A. H. van Amerongen, J. J. P. van Es, P. Wicke, K. V. Kheruntsyan, and N. J. van Druten, *Phys. Rev. Lett.* **100**, 090402 (2008).
 - [7] J. J. P. van Es, P. Wicke, A. H. van Amerongen, C. Rétif, S. Whitlock, and N. J. van Druten, *J. Phys. B* **43**, 155002 (2010).
 - [8] E. H. Lieb and W. Liniger, *Phys. Rev.* **130**, 1605 (1963); E. H. Lieb, *ibid.* **130**, 1616 (1963).
 - [9] V. Dunjko, V. Lorent, and M. Olshanii, *Phys. Rev. Lett.* **86**, 5413 (2001).
 - [10] M. Olshanii, *Phys. Rev. Lett.* **81**, 938 (1998).
 - [11] Z.-Q. Ma and C. N. Yang, *Chin. Phys. Lett.* **26**, 120506 (2009); **27**, 020506 (2010).
 - [12] A. Yu. Cherny and J. Brand, *Phys. Rev. A* **73**, 023612 (2006).
 - [13] V. E. Korepin, N. M. Bogoliubov, and A. G. Azergin, *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge University Press, Cambridge, 1993).
 - [14] Th. Busch, B.-G. Englert, K. Rzążewski, and M. Wilkens, *Found. Phys.* **28**, 549 (1998).
 - [15] M. D. Girardeau, E. M. Wright, and J. M. Triscari, *Phys. Rev. A* **63**, 033601 (2001); M. D. Girardeau and E. M. Wright, *Laser Phys.* **12**, 8 (2002).
 - [16] M. D. Girardeau, *J. Math. Phys.* **1**, 516 (1960).
 - [17] T. Cheon and T. Shigehara, *Phys. Rev. Lett.* **82**, 2536 (1999).
 - [18] E. B. Kolomeisky, T. J. Newman, J. P. Straley, and X. Qi, *Phys. Rev. Lett.* **85**, 1146 (2000).
 - [19] C. Menotti and S. Stringari, *Phys. Rev. A* **66**, 043610 (2002).
 - [20] Y. E. Kim and A. L. Zubarev, *Phys. Rev. A* **67**, 015602 (2003).
 - [21] J. Brand, *J. Phys. B* **37**, S287 (2004).
 - [22] B. E. Granger and D. Blume, *Phys. Rev. Lett.* **92**, 133202 (2004); G. E. Astrakharchik, D. Blume, S. Giorgini, and L. P. Pitaevskii, *ibid.* **93**, 050402 (2004); I. V. Tokatly, *ibid.* **93**, 090405 (2004); R. J. Magyar and K. Burke, *Phys. Rev. A* **70**, 032508 (2004); G. Xianlong, M. Polini, R. Asgari, and M. P. Tosi, *ibid.* **73**, 033609 (2006); Z.-Q. Ma and C. N. Yang, *Chin. Phys. Lett.* **27**, 080501 (2010).
 - [23] J. C. Slater, *Phys. Rev.* **34**, 1293 (1929); E. U. Condon, *ibid.* **36**, 1121 (1930).
 - [24] D. Sen, *Int. J. Mod. Phys. A* **14**, 1789 (1999).
 - [25] M. T. Batchelor, X. W. Guan, N. Oelkers, and C. Lee, *J. Phys. A: Math. Gen.* **38**, 7787 (2005).
 - [26] D. Jukić, S. Galić, R. Pezer, and H. Buljan, *Phys. Rev. A* **82**, 023606 (2010).