

Quantum measurements of atoms using cavity QED

Adetunmise C. Dada,^{1,*} Erika Andersson,¹ Martin L. Jones,² Vivien M. Kendon,² and Mark S. Everitt^{2,3}

¹*SUPA, School of Engineering and Physical Sciences, Heriot-Watt University, Edinburgh EH14 4AS, United Kingdom*

²*School of Physics and Astronomy, University of Leeds, Woodhouse Lane, Leeds LS2 9JT, United Kingdom*

³*National Institute of Informatics, 2-1-2 Hitotsubashi, Chiyoda ku, Tokyo 101-8430, Japan*

(Received 5 November 2010; published 29 April 2011)

Generalized quantum measurements are an important extension of projective or von Neumann measurements in that they can be used to describe any measurement that can be implemented on a quantum system. We describe how to realize two nonstandard quantum measurements using cavity QED. The first measurement optimally and unambiguously distinguishes between two nonorthogonal quantum states. The second example is a measurement that demonstrates superadditive quantum coding gain. The experimental tools used are single-atom unitary operations effected by Ramsey pulses and two-atom Tavis-Cummings interactions. We show how the superadditive quantum coding gain is affected by errors in the field-ionization detection of atoms and that even with rather high levels of experimental imperfections, a reasonable amount of superadditivity can still be seen. To date, these types of measurements have been realized only on photons. It would be of great interest to have realizations using other physical systems. This is for fundamental reasons but also since quantum coding gain in general increases with code word length, and a realization using atoms could be more easily scaled than existing realizations using photons.

DOI: [10.1103/PhysRevA.83.042339](https://doi.org/10.1103/PhysRevA.83.042339)

PACS number(s): 03.67.Hk, 03.67.Lx, 42.50.Pq

I. INTRODUCTION

Generalized quantum measurements or probability operator measures (POMs), also called positive operator valued measures (POVMs), are important mathematical tools for quantum communication and quantum information processing [1]. They are naturally able to describe imperfections and errors in real experimental measurements. In addition, there are also situations where it is advantageous to deliberately engineer a measurement that is not a projective measurement. This is frequently the case when distinguishing between quantum states [1,2]. The simplest such example is when distinguishing between two nonorthogonal states without error [3–5]. In addition, knowledge of optimal measurement strategies may be useful in placing tight bounds on other quantum operations such as quantum cloning [2,6].

In this paper, we describe how to realize two examples of nonstandard quantum measurements using the tools of cavity QED. The methods we describe could, however, be applied also more generally for realizing other generalized quantum measurements. The first measurement is optimal unambiguous discrimination of nonorthogonal quantum states, also known as the Ivanovic-Dieks-Peres (IDP) measurement. This task is relevant for quantum information and communication systems as well as for quantum key distribution (QKD) [7]. The IDP measurement is optimal for the B92 QKD protocol [8], although this was not immediately recognized. To date, all the realizations of the IDP measurement have been optical [9–11]. Nevertheless, generalized quantum measurements could be realized also on ions or atoms using existing experimental techniques [12–14] or using nuclear magnetic resonance [15].

The second example is the measurement required to demonstrate that quantum channel capacities can be superadditive. In this case, at least two uses of a quantum channel, and

a collective measurement of the resulting code block, is required. The quantum coding gain in general grows with the length of the code blocks. Superadditivity has so far only been demonstrated using linear optics [16]. Quantum source coding for message compression is another type of quantum coding scheme that has been optically demonstrated [17], using similar techniques as for the optical demonstration of quantum superadditivity. In both cases, the two uses of the quantum channel were encoded using the path and polarization degrees of freedom of a single photon and the states are manipulated using basic linear optical elements (polarizing beam splitters and wave plates). While this demonstrated the principle of the measurement, extension of the coding to longer code blocks would be impractical due to problems of scalability. Scalability would require effective photon-photon interactions, which are difficult to realize due to prohibitively large overhead costs [18,19].

Generally speaking, some generalized measurements are difficult to realize using linear optics. Therefore it is useful to study how to realize such measurements in other physical systems. For the cavity QED demonstration of superadditivity, we use two atoms and encode each usage of the quantum channel in the state of one atom. This could in principle be scaled to longer codewords using resources which do not scale exponentially, as for the existing optical realizations. Also, other coding schemes, including quantum source coding, or any other realization of collective quantum measurements, could be realized in a cavity QED setting employing similar methods. We also estimate how experimental imperfections would affect the measurement. Cavity QED techniques have indeed been applied extensively in exploring the quantum dynamics of atoms and photons in cavities and has been used, for example, in preparing entangled states of atoms [20], performing phase-gate operations [21], doing quantum nondemolition measurements of cavity fields [22], and in experimental studies of the process of decoherence in quantum measurements [23]. In addition, there are a number of

*acd8@hw.ac.uk

QED-type systems in which a cavity QED based scheme can be easily implemented. These include circuit and photonic crystal based systems [24,25].

The paper is organized as follows. In Sec. II, we describe the IDP measurement and a cavity QED realization. In Sec. III, we then describe the superadditive measurement and a cavity QED scheme to realize it. We consider, in Sec. IV, how experimental imperfections, such as state misidentification, might affect the proposed scheme. Finally, in Sec. V, we give a summary of our conclusions and a few other remarks.

II. DISTINGUISHING BETWEEN TWO NONORTHOGONAL STATES

Generalized quantum measurements are extensions of projective or von Neumann measurements. Just as for projective quantum measurements, probabilities $p(j)$ for measurement outcomes are calculated using the trace rule

$$p(j) = \text{Tr}(\hat{\rho}\hat{\Pi}_j), \quad (1)$$

where ρ is the measured state and $\hat{\Pi}_j$ is the measurement operator corresponding to outcome j . The fact that probabilities are positive means that all eigenvalues of the $\hat{\Pi}_j$ are positive, which is written $\hat{\Pi}_j > 0$, and consequently also that the $\hat{\Pi}_j$ are Hermitian. Also, since the probabilities for all possible outcomes should sum to 1, it follows that

$$\sum_j \hat{\Pi}_j = \hat{\mathbf{I}}, \quad (2)$$

where $\hat{\mathbf{I}}$ is the identity operator. What sets generalized quantum measurements apart from projective measurements is that the measurement operators do not have to be projectors. Also, we can have more measurement outcomes than there are dimensions in the measured quantum system.

The IDP measurement [3–5] is a generalized measurement that distinguishes between two nonorthogonal states without error, in other words, unambiguously. For the measurement to be error free, however, one must accept that it will sometimes be inconclusive. The IDP measurement is optimal in the sense that it minimizes the probability of an inconclusive measurement outcome. Suppose that we wish to distinguish without error between two nonorthogonal quantum states

$$|\psi_1\rangle = \cos\theta|1\rangle - \sin\theta|2\rangle, \quad (3)$$

$$|\psi_2\rangle = \cos\theta|1\rangle + \sin\theta|2\rangle, \quad (4)$$

of a single quantum system such as an atom, where $0 < \theta < \pi/4$. To start, let us note that the optimal measurement will depend on the probabilities for preparing these states, i.e., the prior probabilities. Let us note also that if we make a projective measurement in the basis $\{|\psi_1\rangle, |\psi_1^\perp\rangle\}$, with $|\psi_1^\perp\rangle = \sin\theta|1\rangle + \cos\theta|2\rangle$, then the outcome $|\psi_1^\perp\rangle$ necessarily indicates that the prepared state was $|\psi_2\rangle$. If we obtain $|\psi_1\rangle$, then we cannot be sure which state was prepared, and the outcome is inconclusive. Similarly, if we choose to measure in the basis $\{|\psi_2\rangle, |\psi_2^\perp\rangle\}$, then an outcome $|\psi_2^\perp\rangle$ indicates that the state was certainly $|\psi_1\rangle$ and the outcome $|\psi_2\rangle$ yields an inconclusive result. If we are restricted to standard von Neumann measurements, this is the best we can do.

This procedure above is, however, not always optimal. If the respective probabilities of preparing $|\psi_1\rangle$ and $|\psi_2\rangle$, i.e., the prior probabilities, are similar, then the generalized measurement that gives the lowest possible probability for the inconclusive result has the measurement operators

$$\begin{aligned} \hat{\Pi}_1 &= k|\psi_2^\perp\rangle\langle\psi_2^\perp|, & \hat{\Pi}_2 &= k|\psi_1^\perp\rangle\langle\psi_1^\perp|, \\ \hat{\Pi}_? &= \hat{\mathbf{I}} - \hat{\Pi}_1 - \hat{\Pi}_2, \end{aligned} \quad (5)$$

where k is a positive number which is as large as the positivity of $\hat{\Pi}_?$ will allow, that is, $k = 1/(1 + \langle\psi_1|\psi_2\rangle)$. The minimum probability for the inconclusive result is then given by $p(?) = |\langle\psi_1|\psi_2\rangle|$. Let us denote the prior probabilities as p_1 and $p_2 = 1 - p_1$. It is easy to verify that the generalized measurement is better than the best projective measurement when

$$1 - \sec^2(2\theta) + \sec(2\theta) > p_1 > 2 \sin^2(\theta) \sec^2(2\theta). \quad (6)$$

The measurement described in Eq. (5) can be physically realized as a measurement in a higher-dimensional Hilbert space in an orthonormal basis. This follows from Naimark's theorem which states that any generalized measurement can be realized in this way [1]. We will devise an experimental realization in terms of such a projective measurement in an extended Hilbert space. The IDP measurement can then be realized using the following steps:

(i) Extend the initial 2D Hilbert space into a 3D space by adding an extra state $|3\rangle$ which is orthogonal to both initial states $|\psi_1\rangle$ and $|\psi_2\rangle$, resulting in an orthonormal basis $\{|1\rangle, |2\rangle, |3\rangle\}$.

(ii) Measure in a basis $\{|\Pi_1\rangle, |\Pi_2\rangle, |\Pi_3\rangle\}$, where $|\Pi_1\rangle \perp |\psi_2\rangle$ and $|\Pi_2\rangle \perp |\psi_1\rangle$. This measurement can be implemented in two steps:

(a) Perform a unitary operation \hat{U} given by

$$\hat{U} = |1\rangle\langle\Pi_1| + |2\rangle\langle\Pi_2| + |3\rangle\langle\Pi_3|. \quad (7)$$

(b) Do a standard projective measurement in the $\{|1\rangle, |2\rangle, |3\rangle\}$ basis. A detection in the states $|1\rangle$ or $|2\rangle$ would unambiguously indicate that the unknown state was $|\psi_1\rangle$ or $|\psi_2\rangle$, respectively, while a detection result $|3\rangle$ would make the measurement inconclusive.

The detection probabilities will therefore be

$$\begin{aligned} p(1|1) &= |\langle\Pi_1|\psi_1\rangle|^2 & p(2|2) &= |\langle\Pi_2|\psi_2\rangle|^2 \\ p(1|2) &= p(2|1) = |\langle\Pi_1|\psi_2\rangle|^2 = |\langle\Pi_2|\psi_1\rangle|^2 = 0 & p(?)|1) &= p(?)|2) = |\langle\psi_1|\psi_2\rangle| \end{aligned} \quad (8)$$

Here $p(k|j)$ denotes the probability of obtaining a result k given a state j .

Using the basis state vectors $|1\rangle \equiv [1, 0, 0]^T$, $|2\rangle \equiv [0, 1, 0]^T$, and $|3\rangle \equiv [0, 0, 1]^T$, we work out the unitary operation in (7) for the optimum measurement to be

$$\hat{U} = \frac{1}{\sqrt{2}} \begin{bmatrix} \tan\theta & -1 & -\sqrt{1 - \tan^2\theta} \\ \tan\theta & 1 & -\sqrt{1 - \tan^2\theta} \\ \sqrt{2(1 - \tan^2\theta)} & 0 & \sqrt{2}\tan\theta \end{bmatrix}. \quad (9)$$

For experimental realization, an $n \times n$ discrete unitary operator \hat{U} may be decomposed into a product of unitary

operators, $\hat{T}_{j,k}$, coupling only two levels, j and k , at a time [26]. This involves finding matrices $\hat{T}_{j,k}$ such that

$$\hat{T}_{n-1,n}^\dagger \cdots \hat{T}_{2,n}^\dagger \cdots \hat{T}_{2,4}^\dagger \hat{T}_{2,3}^\dagger \cdots \hat{T}_{1,n}^\dagger \cdots \hat{T}_{1,3}^\dagger \hat{T}_{1,2}^\dagger \hat{U} = I,$$

from which one then deduces that

$$\hat{U} = \hat{T}_{1,2} \hat{T}_{1,3} \cdots \hat{T}_{1,n} \cdots \hat{T}_{2,3} \hat{T}_{2,4} \cdots \hat{T}_{2,n} \cdots \hat{T}_{n-1,n}.$$

In other words, $\hat{T}_{j,k}^\dagger$ are $n \times n$ unitary matrices which successively transform \hat{U} into a diagonal matrix by reducing the off-diagonal elements to zero, one at a time.

Furthermore, especially when there are many outcomes, this decomposition for a generalized quantum measurement may be optimized to use the minimum number of such pairwise operations [27]. In our case, there is only one extra state, and the realization is straightforward,

$$\hat{U} = \hat{T}_{1,2} \hat{T}_{1,3}, \quad (10)$$

where

$$\hat{T}_{1,2} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad (11)$$

$$\hat{T}_{1,3} = \begin{bmatrix} \tan \theta & 0 & -\sqrt{1 - \tan^2 \theta} \\ 0 & 1 & 0 \\ \sqrt{1 - \tan^2 \theta} & 0 & \tan \theta \end{bmatrix}. \quad (12)$$

In summary, one performs $\hat{T}_{1,3}$, followed by $\hat{T}_{1,2}$ on the input state $|\psi\rangle \in \{|\psi_1\rangle, |\psi_2\rangle\}$ to obtain $|\psi'\rangle$, followed by a projective measurement of $|\psi'\rangle$ in the basis $\{|1\rangle, |2\rangle, |3\rangle\}$. This will yield an error probability of zero and a minimum probability of an inconclusive result $p(?) = \cos 2\theta$ (see Fig. 1).

A. Cavity QED implementation

The interaction between an atom and a classical field, resonant or quasiresonant with the atomic transition between




FIG. 1. (Color online) Geometrical representation of the IDP measurement. The initial two-dimensional Hilbert space, where $|\psi_1\rangle$ and $|\psi_2\rangle$ live, is spanned by $\{|1\rangle, |2\rangle\}$. This is extended to three dimensions by adding an extra state $|3\rangle$ which is orthogonal to each of the two initial basis states. The measurement consists of a unitary operation \hat{U} , followed by a standard projective measurement in the $\{|1\rangle, |2\rangle, |3\rangle\}$ basis. The transformed states $\hat{U}|\psi_1\rangle$ and $\hat{U}|\psi_2\rangle$ are orthogonal to $|2\rangle$ and $|1\rangle$, respectively.

two states $|g\rangle$ and $|e\rangle$, can be used to realize the IDP measurement outlined above. The required unitary operations result from the action of the atom-field Hamiltonian, which is [28]

$$\tilde{H} = \frac{\hbar \Delta_r}{2} \sigma_z - i\hbar \frac{\Omega_r}{2} [e^{-i\varphi} \sigma_+ + e^{i\varphi} \sigma_-]. \quad (13)$$

Here $\sigma_z = |g\rangle\langle g| - |e\rangle\langle e|$ is the Pauli-Z operator and σ_\pm , the atomic raising and lowering operators, are defined as $\sigma_+ = |g\rangle\langle e|$ and $\sigma_- = |e\rangle\langle g|$. Ω_r and Δ_r are respectively the classical Rabi frequency and the atom-field detuning, and φ is the phase of the classical field with respect to the atomic transition dipole.

It can be shown from Eq. (13) that an interaction lasting for a time $t = \theta/\Omega_r$ with a resonant field having a phase φ effects the transformations

$$\begin{aligned} |g\rangle &\longrightarrow \cos(\theta/2)|g\rangle + \sin(\theta/2)e^{i\varphi}|e\rangle \\ |e\rangle &\longrightarrow -\sin(\theta/2)e^{-i\varphi}|g\rangle + \cos(\theta/2)|e\rangle. \end{aligned} \quad (14)$$

This transformation corresponds to the operation of $\hat{R}_{g,e}(\theta, \varphi) \equiv \exp(i\tilde{H}t/\hbar)$, which is a unitary operator coupling only two levels at a time, that is, the states denoted by $|e\rangle$ and $|g\rangle$. If we use the notation $|g\rangle \equiv [1, 0]^T$ and $|e\rangle \equiv [0, 1]^T$, then $\hat{R}_{g,e}(\theta, \varphi)$ is given as

$$\hat{R}_{g,e}(\theta, \varphi) = \begin{bmatrix} \cos \frac{\theta}{2} & -e^{-i\varphi} \sin \frac{\theta}{2} \\ e^{i\varphi} \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{bmatrix}. \quad (15)$$

Returning to the definition of the basis $\{|1\rangle, |2\rangle, |3\rangle\}$ above, one may perform the unitary operator \hat{U} for the IDP measurement by setting

$$\hat{T}_{1,2} = R_{1,2}(\pi/2, 0) \quad (16)$$

$$\hat{T}_{1,3} = R_{1,3}(\vartheta, 0), \quad (17)$$

where $\vartheta = \cos^{-1}(\frac{1}{\sqrt{3}})$, and the $R_{g,e}$ is realized using a Ramsey pulse resonant with the transition $|g\rangle \leftrightarrow |e\rangle$.

The physical states representing $|1\rangle$, $|2\rangle$, and $|3\rangle$ will be chosen based on convenience of experimental realization. One needs to bear in mind, for example, that a direct coupling between $|2\rangle$ and $|3\rangle$ will not be necessary and that, at the detection stage, outcome $|3\rangle$ represents an inconclusive result. A possible choice of states could be a ^{85}Rb ladder of Rydberg states, $|e\rangle \equiv 63P$, $|g\rangle \equiv 61D$, and $|i\rangle \equiv 62P$, which are standard micromaser transitions [29]. The ladder of circular Rydberg states with principal quantum numbers $n = 49, 50, 51$ could also be used [23]. A final projective measurement which determines the energy level of the atom would be required. This is commonly done by means of state selective field-ionization detection [30–33], which involves passing the Rydberg atoms through an increasing electric field and measuring the energy at which the atom is ionized.

III. SUPERADDITIVE MEASUREMENT

The maximum amount of information which can be reliably transmitted over a given channel is referred to as its capacity. This is generally determined by the information resources (such as code block length and bandwidth) and the noise characteristics of the channel. Quantum channels may display

superadditivity in classical information capacity [34–36]. This means that

$$C_n > nC_1, \quad (18)$$

where C_1 is the classical information capacity of a single use of the channel, and C_n is the classical information capacity of a combination of n uses of the channel. For classical channels, it holds that

$$C_n = nC_1, \quad (19)$$

meaning that superadditivity is displayed only by quantum channels.

This makes it interesting to experimentally demonstrate the superadditivity of quantum channel capacity. In order to do this, it is necessary to carry out quantum coding followed by an appropriate collective quantum measurement. One possible scheme is outlined below.

A. Trine letter states

Consider a channel coding for sending classical information through a quantum channel with a given ensemble of quantum states representing the letter states. A clear and simple example of an ensemble which can be used to demonstrate superadditivity in classical capacity of a quantum channel is the qubit trine states. Suppose we use the set of ternary symmetric states of a qubit, that is, $\{|\psi_0\rangle, |\psi_1\rangle, |\psi_2\rangle\}$, known as the *qubit trine states*, with

$$\begin{aligned} |\psi_0\rangle &= |0\rangle, & |\psi_1\rangle &= -\frac{1}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle, \\ |\psi_2\rangle &= -\frac{1}{2}|0\rangle + \frac{\sqrt{3}}{2}|1\rangle, \end{aligned} \quad (20)$$

to transmit information, where $\{|0\rangle, |1\rangle\}$ is the orthonormal basis set. Using one quantum state drawn from this ensemble, we can transmit at most $C_1 = 0.6454$ bits. This is achieved by sending any two of the states $|\psi_j\rangle$ with probability $1/2$ each and distinguishing between these with the optimal measurement [16,37,38].

Using two qubits, there are nine possible states. It has been shown [34] that if only three of these are used, namely

$$\begin{aligned} |\psi_{xx}\rangle &= |\psi_x\rangle \otimes |\psi_x\rangle = \frac{1}{2}(1 + \cos \varphi_x)|00\rangle \\ &+ \frac{1}{2} \sin \varphi_x(|01\rangle + |10\rangle) + \frac{1}{2}(1 - \cos \varphi_x)|11\rangle, \end{aligned} \quad (21)$$

with $\varphi_x = 2\pi x/3$, where $x = 0, 1, 2$; then $I_2 = 1.3690$ bits of information can be retrieved if the codeword states are used with equal probabilities. This is larger than $2C_1 (= 1.2908)$. The superadditive quantum coding gain (SQCG), per use of the channel, is

$$(I_2 - 2C_1)/2 = (1.3690 - 1.2908)/2 = 0.0391. \quad (22)$$

The measurement used to decode the codewords is the *square-root measurement* with the measurement basis states $|\Pi_{yy}\rangle$ defined as

$$|\Pi_{yy}\rangle \equiv \left(\sum_x |\psi_{xx}\rangle \langle \psi_{xx}| \right)^{-1/2} |\psi_{yy}\rangle. \quad (23)$$

In explicit form the codeword states are

$$|\psi_{00}\rangle = |00\rangle, \quad |\psi_{11}\rangle = [|00\rangle + \sqrt{3}(|01\rangle + |10\rangle) + 3|00\rangle]/4, \quad (24)$$

$$|\psi_{22}\rangle = [|00\rangle - \sqrt{3}(|01\rangle + |10\rangle) + 3|00\rangle]/4,$$

and the optimal measurement basis is given by

$$\begin{aligned} |\Pi_{00}\rangle &= \cos(\gamma/2)|00\rangle - \sin(\gamma/2)|11\rangle \\ |\Pi_{11}\rangle &= [\sin(\gamma/2)|00\rangle + \cos(\gamma/2)|11\rangle]/\sqrt{2} \\ &+ (|01\rangle + |10\rangle)/2, \end{aligned} \quad (25)$$

$$\begin{aligned} |\Pi_{22}\rangle &= [\sin(\gamma/2)|00\rangle + \cos(\gamma/2)|11\rangle]/\sqrt{2} \\ &- (|01\rangle + |10\rangle)/2, \end{aligned}$$

$$|A\rangle = [|01\rangle - |10\rangle]/\sqrt{2},$$

where

$$\begin{aligned} \cos(\gamma/2) &= (\sqrt{2} + 1)/\sqrt{6} \quad \text{and} \\ \sin(\gamma/2) &= (\sqrt{2} - 1)/\sqrt{6}. \end{aligned} \quad (26)$$

The outcome corresponding to the state $|A\rangle$ will never occur, since all codeword states are orthogonal to $|A\rangle$. This state merely completes the four-dimensional basis.

The states (25) define an entangled measurement basis, and the implementation of the measurement will require similar resources as a Bell measurement, including entangling interactions. The Bell states are the maximally entangled states

$$\begin{aligned} |\Phi^\pm\rangle &= (|00\rangle \pm |11\rangle)/\sqrt{2}, \\ |\Psi^\pm\rangle &= (|01\rangle \pm |10\rangle)/\sqrt{2}, \end{aligned} \quad (27)$$

and a Bell measurement is a projection in this basis. This can be achieved by first performing a unitary transformation \hat{U}_B

$$\hat{U}_B = |00\rangle\langle\Psi^+| + |01\rangle\langle\Phi^+| + |10\rangle\langle\Psi^-| + |11\rangle\langle\Phi^-| \quad (28)$$

on the input Bell state, followed by a projective measurement in the $\{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$ basis, which we refer to as the computational basis. Any other transformation which takes each of the Bell states respectively to any permutation of the computational basis states, up to global phases, would also do.

The superadditive measurement can be realized in a similar fashion by making a unitary transformation \hat{U}_{sa} on the input states, and following this by a projective measurement in the computational basis. From Eq. (25), \hat{U}_{sa} is given as

$$\hat{U}_{sa} = |00\rangle\langle\Pi_{00}| + |01\rangle\langle\Pi_{11}| + |10\rangle\langle A| + |11\rangle\langle\Pi_{22}|. \quad (29)$$

In matrix notation, it takes the form

$$\hat{U}_{sa} = \frac{1}{2} \begin{bmatrix} 2 \cos(\gamma/2) & 0 & 0 & -2 \sin(\gamma/2) \\ \sqrt{2} \sin(\gamma/2) & 1 & 1 & \sqrt{2} \cos(\gamma/2) \\ 0 & \sqrt{2} & -\sqrt{2} & 0 \\ \sqrt{2} \sin(\gamma/2) & -1 & -1 & \sqrt{2} \cos(\gamma/2) \end{bmatrix}. \quad (30)$$

Superadditivity can in general only be achieved when an appropriate *collective* POM is chosen, namely detection in an entangled measurement basis [38]. An SQCG of 0.011 ± 0.003 has been experimentally demonstrated by the previously mentioned optical implementation [35].

B. Cavity QED realization

The unitary operation \hat{U}_{sa} needed to realize this measurement can be decomposed in terms of single-atom operation and entangling interactions. The single-atom rotations correspond to Ramsey pulses $\hat{R}_1(\theta, \varphi)$ and $\hat{R}_2(\theta, \varphi)$. In the four-dimensional Hilbert space spanned by the joint basis states of the two 2-level atoms, the unitary transformation $\hat{R}_1(\theta, \varphi)$ effected by a Ramsey pulse on atom 1 is

$$\hat{R}_1(\theta, \varphi) = \hat{R}(\theta, \varphi) \otimes \hat{I}, \quad (31)$$

while a Ramsey pulse on atom 2 is

$$\hat{R}_2(\theta, \varphi) = \hat{I} \otimes \hat{R}(\theta, \varphi), \quad (32)$$

where \otimes denotes the tensor product operation.

The entangling operations, on the other hand, can be realized using the interactions between atoms and a cavity field governed by the two-atom Tavis-Cummings Hamiltonian in the limit of large detuning [39,40]. This produces an effective Hamiltonian in which the field is removed as a degree of freedom, eliminating atom-field entanglement but allowing virtual excitation of the field to pass excitations between atoms. This ensures that no quantum information is exchanged between the atoms and the cavity, so the cavity merely mediates interactions between the atoms. Each atom is effectively a two-state system, detuned from the cavity resonance by Δ . Let g denote the atom-cavity dipole coupling constant. In the limit of large Δ , the effective Hamiltonian is

$$\tilde{H}_{\text{eff}} = -\frac{\hbar g^2}{\Delta} (\sigma_{eg, eg} + \sigma_{ge, ge} + \sigma_{eg, ge} + \sigma_{ge, eg}). \quad (33)$$

This yields a unitary transformation

$$\hat{T}_{\text{TC}}(t) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & e^{-it\phi} \cos(t\phi) & -ie^{-it\phi} \sin(t\phi) & 0 \\ 0 & -ie^{-it\phi} \sin(t\phi) & e^{-it\phi} \cos(t\phi) & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad (34)$$

where $\phi = g^2/\Delta(s^{-1})$ is the effective coupling constant and we have used the notation $|gg\rangle \equiv [1, 0, 0, 0]^T$, $|ge\rangle \equiv [0, 1, 0, 0]^T$, $|eg\rangle \equiv [0, 0, 1, 0]^T$, and $|ee\rangle \equiv [0, 0, 0, 1]^T$ for the computational basis states.

To elucidate the process of deriving the superadditive measurement using these building blocks, it is instructive to first consider the Bell measurement briefly. The Bell measurement can be performed with a combination of the operations in Eqs. (31), (32), and (34). This will yield a transformation which rotates each of the Bell states into some permutation of the computational basis states up to global phases.

This can be done in only four steps. The principle of the process is to take the entangled states to states which are as close as possible to the separable basis states with each step. The Tavis-Cummings operations are the available two-qubit operations for disentangling the Bell states. However, it turns out that the detuned Tavis-Cummings operation on its own cannot disentangle $|\Psi^\pm\rangle$. It is necessary to precede it with a Ramsey rotation which produces a relative phase shift of $\pi/2$ between $|01\rangle$ and $|10\rangle$. Previous work has achieved the

phase-shifting effect using an extra atomic level $|i\rangle$. This is done by performing a Ramsey operation resonant with the $|e\rangle \leftrightarrow |i\rangle$ transition before and then after the Tavis-Cummings operation [28,39]. Another proposal suggests introducing a slight delay between the passage of the two atoms through the cavity [41]. Here we use another approach.

As a first step, we apply a Ramsey pulse to the atom 2, i.e., $\hat{U}_1 = \hat{R}_2(\pi, \frac{3\pi}{4})$. This gives the following transformation of the Bell states:

$$\begin{aligned} \hat{U}_1|\Phi^\pm\rangle &= (|01\rangle \pm i|10\rangle)/\sqrt{2}, \\ \hat{U}_1|\Psi^\pm\rangle &= (|00\rangle \pm i|11\rangle)/\sqrt{2}. \end{aligned} \quad (35)$$

We then choose the second step as $\hat{U}_2 = \hat{T}_{\text{TC}}(\frac{3\pi}{4\phi})$ to have

$$\begin{aligned} \hat{U}_2\hat{U}_1|\Phi^+\rangle &= (|00\rangle - i|11\rangle)/\sqrt{2}, \\ \hat{U}_2\hat{U}_1|\Psi^+\rangle &= |01\rangle, \\ \hat{U}_2\hat{U}_1|\Phi^-\rangle &= |10\rangle, \\ \hat{U}_2\hat{U}_1|\Psi^-\rangle &= (|00\rangle + i|11\rangle)/\sqrt{2}. \end{aligned} \quad (36)$$

The combination of a preceding Ramsey operation on atom 2 and the detuned Tavis-Cummings operation effectively carries out a transformation which disentangles the $|\Psi^+\rangle$ and $|\Phi^-\rangle$ states.

We now proceed to the third step, which is a Ramsey operation, $\hat{U}_3 = \hat{R}_2(\pi, 0)$. The effect of this operation is simply to interchange $|00\rangle$ with $|01\rangle$ and $|11\rangle$ with $|10\rangle$ simultaneously.

$$\begin{aligned} \hat{U}_2\hat{U}_1|\Phi^+\rangle &= (|01\rangle - i|10\rangle)/\sqrt{2}, \\ \hat{U}_2\hat{U}_1|\Psi^+\rangle &= |00\rangle, \\ \hat{U}_2\hat{U}_1|\Phi^-\rangle &= |11\rangle, \\ \hat{U}_2\hat{U}_1|\Psi^-\rangle &= (|01\rangle + i|10\rangle)/\sqrt{2}. \end{aligned} \quad (37)$$

The resulting entangled states can now be disentangled with a Tavis-Cummings operation in a final step before detection. If we use $\hat{U}_4 = \hat{T}_d(\frac{3\pi}{4\phi})$, then we will have:

$$\begin{aligned} \hat{U}_4\hat{U}_3\hat{U}_2\hat{U}_1|\Phi^+\rangle &= |01\rangle, \\ \hat{U}_4\hat{U}_3\hat{U}_2\hat{U}_1|\Psi^+\rangle &= |00\rangle, \\ \hat{U}_4\hat{U}_3\hat{U}_2\hat{U}_1|\Phi^-\rangle &= |11\rangle, \\ \hat{U}_4\hat{U}_3\hat{U}_2\hat{U}_1|\Psi^-\rangle &= |10\rangle. \end{aligned} \quad (38)$$

Detection results $|00\rangle$, $|01\rangle$, $|10\rangle$, and $|11\rangle$ would indicate that the inputs were the Bell states $|\Psi^+\rangle$, $|\Phi^+\rangle$, $|\Psi^-\rangle$, and $|\Phi^-\rangle$ respectively. This realization is similar in its construction to the realization on atoms considered in Ref. [42].

We now apply a similar method for the superadditive measurement. It is worth noting that since the outcome corresponding to $|A\rangle$ should never occur, it would be sufficient to make a unitary transformation \hat{U}'_{sa} which takes two of the three states $|\Pi_{00}\rangle$, $|\Pi_{11}\rangle$, and $|\Pi_{22}\rangle$ uniquely into two of the four computational basis states in the four-dimensional Hilbert space, say, $|\Pi_{22}\rangle \rightarrow |01\rangle$ and $|\Pi_{11}\rangle \rightarrow |11\rangle$, and the remaining measurement basis state, say, $|\Pi_{00}\rangle$ and $|A\rangle$, each into superpositions of the two other computational basis states, say, $|00\rangle$ and $|10\rangle$. This may somewhat simplify the experimental realization, and we will in fact make use of it.

Obtaining a realization of \hat{U}'_{sa} involves finding a sequence of operations which transforms \hat{U}'_{sa} into a matrix of the form $P_\pi D$, where P_π is a permutation matrix and D is a diagonal matrix. This could be a sequence of unitary operations coupling two basis states at a time [26]. As with most physical settings, not all pairwise coupling operations are available in our case. This is because we are restricted to operations $\hat{T}_{TC}(t)$, which couple the pair of basis states $|01\rangle$ and $|10\rangle$, and single-qubit operations $\hat{R}_1(\theta, \varphi)$ and $\hat{R}_2(\theta, \varphi)$, which each couple two pairs of basis states at the same time. Our strategy for obtaining a realization in terms of these is as follows. Since the operation \hat{T}_{TC} couples basis states $|01\rangle$ and $|10\rangle$, it is natural to first use a Tavis-Cummings interaction to disentangle these components of the measurement states in Eq. (25). In order to do this, it turns out that we need to precede the Tavis-Cummings interaction by two Ramsey pulses. This first pulse sequence then takes the states $|A\rangle = |\Psi^-\rangle$ and $|\Psi^+\rangle$ into the disentangled states $|A'\rangle = |01\rangle$ and $|10\rangle$. Next, in order to use a Tavis-Cummings interaction to disentangle the $|00\rangle$ and $|11\rangle$ components, we need to first change $|0\rangle$ into $|1\rangle$ and vice versa for one of the atoms, which one does not matter, using Ramsey pulses. It turns out that at the end of this process, which thus comprises two Tavis-Cummings interactions and a number of Ramsey pulses, $|A\rangle$ and the measurement basis state $|\Pi_{00}\rangle$ are both mapped to superpositions of $|00\rangle$ and $|10\rangle$, and Ramsey pulses would be needed in order to map these superpositions to $|00\rangle$ and $|10\rangle$. As remarked above, these last Ramsey rotations are not necessarily required.

This leads us to a realization in seven steps. The first step is a Ramsey rotation on the atom 2, $\hat{U}'_1 = \hat{R}_2(\pi, \pi)$. The second step is another Ramsey rotation on atom 2, $\hat{U}'_2 = \hat{R}_2(\pi, 3\pi/4)$. For the third step, we pass the two atoms simultaneously through the first detuned cavity with the effective interaction time $t_1 = 3\pi/(4\phi)$, giving a detuned Tavis-Cummings interaction described by $\hat{U}'_3 = \hat{T}_{TC}(t_1)$. Step four is another Ramsey pulse applied to atom 2 defined as $\hat{U}'_4 = \hat{R}_2(\pi, \pi/2)$. The fifth step is a second detuned Tavis-Cummings type interaction, $\hat{U}'_5 = \hat{T}_d(t_2)$, of duration $t_2 = \frac{\gamma}{2\phi}$. The sixth and seventh steps effectively rotate $|A'\rangle$ into a superposition. In fact, this takes place if we choose $\hat{U}'_6 = \hat{R}_2[\pi, (\gamma - \pi/2)/4]$ as the sixth step and $\hat{U}'_7 = \hat{R}_2(\pi/2, 0)$ as the seventh step.

These steps lead us to the effective unitary operation

$$\hat{U}'_{sa} = \frac{e^{-i\gamma/4}}{2} \begin{bmatrix} -e^{-i\gamma/2}C & -e^{i\gamma/2} & e^{i\gamma/2} & e^{-i\gamma/2}S \\ -e^{-i\gamma/2}C & e^{i\gamma/2} & -e^{i\gamma/2} & e^{-i\gamma/2}S \\ -S & 1 & 1 & -C \\ S & 1 & 1 & C \end{bmatrix}, \quad (39)$$

where $S = \sqrt{2} \sin(\gamma/2)$ and $C = \sqrt{2} \cos(\gamma/2)$. To elucidate the assignment of measurement results, we can also use the alternative form

$$\hat{U}'_{sa} \equiv \frac{1}{\sqrt{2}}(|00\rangle + |10\rangle)\langle\Pi_{00}| + \frac{1}{\sqrt{2}}(|00\rangle - |10\rangle)\langle A| + |01\rangle\langle\Pi_{22}| + |11\rangle\langle\Pi_{11}|. \quad (40)$$

This makes it clear that \hat{U}'_{sa} ideally yields the same value as \hat{U}_{sa} for the mutual information but differs slightly from \hat{U}_{sa} ,

since final detection of $|00\rangle$ and $|10\rangle$ both correspond to $|\Pi_{00}\rangle$. Recall that all three signal states are orthogonal to $|A\rangle$. When \hat{U}_{sa} is used, the final measurement outcome corresponding to $|A\rangle$ should never occur. When experimental imperfections are included, the mutual information and consequently SQCG may differ for \hat{U}_{sa} and \hat{U}'_{sa} . This will be made clear shortly.

After performing the seven steps, a subsequent detection in the computational basis will complete the measurement. The mutual information is given by

$$I_2 = I(X : Y) = \sum_j P(x_j) \sum_k P(y_k|x_j) \log_2 \times \left[\frac{P(y_k|x_j)}{\sum_m P(x_m)P(y_k|x_m)} \right], \quad (41)$$

where X and Y denote the sender and receiver, respectively, and x_j and y_k denote the letter states that were transmitted and received respectively, $k, j, m = 1, 2, 3, \dots$

The channel matrix resulting from applying the derived pulse sequence and a subsequent projective measurement to the input state $\hat{\rho}_x = |\psi_x\rangle\langle\psi_x|$ is

$$P(y|x) = \text{Tr}(\hat{\Pi}'_y \hat{\rho}_x), \quad (42)$$

where $\hat{\Pi}'_y = |\Pi'_{yy}\rangle\langle\Pi'_{yy}|$. Substituting the resulting channel matrix elements into Eq. (41) and using prior probabilities

$$p(x_j) = \frac{1}{3}, \quad (j = 1, 2, 3) \quad (43)$$

gives SQCG of $I_2/2 - C_1 = 0.0391$. A schematic diagram outlining the derived implementation is shown in Fig. 2.

C. Optimality

The question of determining the optimality of a given realization of a POM using certain building blocks in a physical setting is nontrivial. For our realization of the superadditive measurement, we check for the optimality of our proposed realization in terms of the total number of steps. We also try to exclude the more experimentally challenging steps as much as possible. The Tavis-Cummings interaction is clearly more difficult to realize than the Ramsey operations because it involves a synchronous passage of two atoms through a high- Q cavity, which is more experimentally challenging than applying a Ramsey pulse to a single atom.

We supply a short proof by contradiction that at least two detuned Tavis-Cummings interactions is required to realize the superadditive decoding. Using the canonical Cartan decomposition of a two-qubit unitary operator $U \in \text{SU}(4)$ [45], we realize that if a single detuned Tavis-Cummings interaction could be used to implement \hat{U}_{sa} (or \hat{U}'_{sa}), then there would exist $w'_1, w_2, v'_1, v_2, W'_1, W_2, V'_1$, and $V_2 \in \text{SU}(2)$ such that

$$\hat{U}_{sa} = (w'_1 \otimes w_2) T_d(\phi) (v'_1 \otimes v_2) \quad (44)$$

and

$$T_d(\phi) = (W'_1 \otimes W_2) \hat{U}_{sa} (V'_1 \otimes V_2). \quad (45)$$

Equation (45) is a system of 16 equations. It is easily verified that this system of equations has no solution. This concludes the proof and gives evidence of the optimality of our proposed scheme with respect to the number of Tavis-Cummings interactions needed. Jaynes-Cummings interactions through




FIG. 2. (Color online) Schematic diagram of proposed cavity QED implementation of the POM for superadditive decoding. This figure shows the scheme for the unitary operation performed before the final projective measurement by field-ionization detectors. Atom mirrors have been implemented using electric, magnetic, and light-induced forces (see Refs. [43,44] and references therein). If required, it could be realized in this scheme also by using an additional Tavis-Cummings operation to swap the state from one atom to that of a fresh one traveling perpendicularly. The superscripts on the operators indicate the order of their application to the two-atom state. These operations are defined in Eqs. (31), (32), and (34).

sequential passage of the atoms through the cavity can also be used for entangling interactions between atoms. This has proved suitable for preparing specific entangled two-atom states [20,39,41]. However, a main disadvantage of using the Jaynes-Cummings interactions is the leakage of atomic excitation into cavity field modes having more than one excitation, since the field has an infinite number of levels besides $\{|0\rangle, |1\rangle\}$.

IV. EXPERIMENTAL IMPERFECTIONS

Both the IDP measurement and the measurement to demonstrate quantum superadditivity will be affected by experimental imperfections. In particular, when errors are present, error-free or unambiguous state discrimination in general becomes impossible, and we should aim for a maximum confidence measurement strategy instead [46–48]. As for the measurement that demonstrates superadditivity, it is natural to ask how robust the superadditive quantum coding gain is with respect to imperfections. We will now discuss this.

Experimental imperfections that could adversely affect the overall quality of the realizations of the superadditive measurement, and the SQCG, include initial-state preparation fidelity, Ramsey operation fidelity, Tavis-Cummings operation fidelity, and detection errors. The initial-state preparation fidelity would depend largely on the fidelity of the Ramsey operations since they are used to carry out these preparations. In turn, the fidelity of the Ramsey operations depend on the accuracy to which the parameters θ and φ can be set.

Let us consider how the delay between the atoms affect the results and ultimately the SQCG. Zheng and Guo [39] have

estimated the effect of such a delay on the preparation of an maximally entangled pair of the form

$$|\Psi_{\text{EPR}}\rangle = \frac{1}{\sqrt{2}}(|e, g\rangle + i|g, e\rangle), \tag{46}$$

which can be prepared by a single Tavis-Cummings operation. This was done by considering a delay of $t_d = 0.01t$ between the atoms, where t is the time each atom spends in the cavity. In this situation, a fidelity of 0.99 was estimated.

Applying the same idea, we realize that such a delay yields an imperfect Tavis-Cummings operation which affects the coding gain. In Fig. 3, we plot the superadditive coding gain as a function of δ , where δ is the delay as a percentage of the longest cavity interaction time in the sequence, that is, t_2 (s) spent in cavity 2. A delay up to 5% of the longest cavity interaction time in the sequence, which occurs in the second cavity interaction, still gives an SQCG of 0.011 bits.




FIG. 3. (Color online) Plot of the superadditive coding gain as a function of δ , where δ is the delay t_d (s) as a percentage of the time t_2 (s) spent by both atoms in the second cavity. The same delay is applied in both cavities.

In the photonic realization [16], the detection efficiency η , which is the photon count probability, does not degrade the result on its own. This is because the SQCG is calculated using a normalized channel matrix. However, when combined with dark counts which arise from background radiation as well as from carriers generated in a detector even when no photons are incident, the SQCG is degraded since this effectively results in a finite probability of misidentification of states.

In a cavity QED realization, the detection efficiency could even be more of a problem if it depends on the atomic states, for example. Even if the detection efficiency were independent of the atomic state, state misidentification is a usual problem in detection. Consider a detection to determine whether a two-level atom is in a state $|0\rangle$ or $|1\rangle$. In the perfect case, the detection would be an ideal von Neumann measurement which can be described by the two projectors

$$\hat{P}_0 = |0\rangle\langle 0|, \quad \hat{P}_1 = |1\rangle\langle 1|. \quad (47)$$

A nonideal detector, however, might record the wrong state with some probability. This is the case in atomic state detection schemes where projective measurements are carried out using field-ionization detectors, in which the ionization energy of the atoms serves as an indicator of the state. This means that for a two-level atom in the state $|0\rangle$, the measurement will give the result 1 with probability p and the result 0 with probability $1 - p$. In a realistic experimental scenario, the probability of misidentification might not be symmetric. For instance, it might be more likely to misidentify the atomic state $|1\rangle$ as $|0\rangle$ than conversely. Let us denote the probability of misidentifying the $|0\rangle$ and $|1\rangle$ states as $p(0|1) \equiv p$ and $p(1|0) \equiv q$, respectively. Introducing these errors, the effective measurement is a POM with elements defined as the operators

$$\begin{aligned} \hat{\pi}_0 &= (1 - p)\hat{P}_0 + q\hat{P}_1, \\ \hat{\pi}_1 &= (1 - q)\hat{P}_1 + p\hat{P}_0. \end{aligned} \quad (48)$$

To incorporate this into the calculations of the SQCG, we first calculate the resulting single channel capacity C_1 and mutual information for length-two coding I_2 . This is used to obtain the SQCG plotted in Fig. 4 as a function of the




FIG. 4. (Color online) Effects of detection errors when \hat{U}'_{sa} is used to realize the superadditive measurement. (a) Plot of the difference G_0 (bits) between the length-two coding mutual information and the ideal single channel capacity, $G_0 = I_2/2(p,q) - C_1(p=0,q=0)$. (b) Plot of the actual superadditive coding gain G_0 (bits), $G = I_2/2(p,q) - C_1(p,q)$. G_0 and G are plotted as functions of the probabilities, p and q , of misidentifying states $|0\rangle$ and $|1\rangle$ of a two-level atom respectively.




FIG. 5. (Color online) Effects of detection errors when \hat{U}'_{sa} is used to implement the superadditive measurement. (a) Plot of the difference G'_0 (bits) between the length-two coding mutual information and the ideal single channel capacity, $G'_0 = I'_2/2(p,q) - C_1(p=0,q=0)$. (b) Plot of the actual superadditive coding gain G' (bits) for our proposed scheme, where $G' = I'_2/2(p,q) - C_1(p,q)$. G'_0 and G' are plotted as functions of the probabilities, p and q , of misidentifying states $|0\rangle$ and $|1\rangle$ of a two-level atom respectively.

probabilities of misidentification p and q when \hat{U}'_{sa} is realized exactly. The affected channel matrix is given as

$$P(y|x) = \text{Tr}(\hat{U}'_{sa} \hat{\rho}_x \hat{U}'_{sa} \hat{M}_y). \quad (49)$$

Here x and y label the matrix elements. $\hat{M}_1 = \hat{\pi}_0 \otimes \hat{\pi}_0$, $\hat{M}_2 = \hat{\pi}_0 \otimes \hat{\pi}_1$, $\hat{M}_3 = \hat{\pi}_1 \otimes \hat{\pi}_0$, and $\hat{M}_4 = \hat{\pi}_1 \otimes \hat{\pi}_1$ are the elements of the POM describing the imperfect projective measurement in the computational basis. The resulting SQCG plot shows that even with rather high probabilities of misidentification, a reasonable amount of superadditive quantum coding gain can still be accessed. We observe a symmetric trade-off effect between probabilities p and q in Fig. 4.

Let us now use the proposed realization which effects the unitary transformation \hat{U}'_{sa} . The channel matrix elements then become

$$P(y|x) = \text{Tr}(\hat{U}'_{sa} \hat{\rho}_x \hat{U}'_{sa} \hat{M}_y). \quad (50)$$

The corresponding values of mutual information for the double channel and the SQCG are plotted in Fig. 5 as functions of the probabilities of misidentification p and q . As shown in Fig. 5(b), we observe reasonable amounts of SQCG even with rather high levels of detection errors. Since, in our proposed scheme \hat{U}'_{sa} , the SQCG, favors combinations of higher values of p with lower values q , the physical states representing $|0\rangle$ and $|1\rangle$ may need to be chosen to ensure that $p > q$ if there is considerable difference between p and q . Values for the probabilities of state misidentification p (and/or q) of as low as 0.013, for selective detection of some adjacent Rydberg states, have been reported [49]. For a given pair of adjacent Rydberg states, we also note that these error probabilities can be considerably lowered at the expense of the overall detection efficiency.

Finally, it is important to consider feasibility in terms of lifetimes of the atomic states and cavity decay. The earlier mentioned circular Rydberg states, with principal quantum numbers $n = 50, 51$ have radiative lifetimes of $T_r \simeq 30$ ms and coupling constant $g = 2\pi \times 24$ kHz. Using an estimated detuning of $10g$, the atom-atom interaction time will be on the order of $10\pi/g \simeq 0.2$ ms [23]. The time for the entire sequence of operations is on the order of 2 ms, which is a lot less than T_r . It is also worth noting that while in the cavities, the atoms are subject to the Purcell effect [50], which

will further reduce the spontaneous emission rate, thus also enhancing the lifetime. Cavity decay times of up to 130 ms have been achieved [51], allowing for interactions to take place long before the dissipative processes due to the cavity begin to set in. Moreover, since only virtual excitations of the cavity are to be used in our proposed scheme, the effect of cavity decay is reduced much further. With all these taken into consideration, the proposed scheme is realizable within the lifetime of available Rydberg states using current or shortly available technologies.

V. CONCLUSION

In conclusion, we have proposed explicit schemes for experimental realization, using cavity QED, of two generalized quantum measurement strategies. These were unambiguous discrimination of two nonorthogonal quantum states, the so-called IDP measurement, and the measurement to demonstrate superadditive quantum coding using a ternary quantum alphabet. We would like to note that realizations of the minimum-error measurements to distinguish between the trine states in Eq. (20) [1] and between mirror-symmetric states [52] would be very similar to the IDP measurement that we have outlined and that similar methods can be used to implement any generalized quantum measurement using cavity QED.

Our results show that these realizations are feasible using current or shortly available cavity QED technologies. Using a simple proof we have confirmed the optimality of the realization of the measurement that demonstrates quantum

superadditivity in terms of cavity usage. We have also shown how the superadditive quantum coding gain is affected by imperfect detection of the basis states and that even with rather high levels of such experimental imperfections, a reasonable amount of superadditivity can be seen. We have not addressed the fact that in the presence of experimental imperfections, the measurement that one should attempt to implement in order to demonstrate maximum coding gain might change. It is thus conceivable that even with experimental errors, it may be possible to see a somewhat larger quantum coding gain than our estimates indicate. In other words, our estimates are lower bounds on the superadditive quantum coding gain, given the assumed level of errors in the implementation. An example where the optimal quantum measurement changes in the presence of experimental imperfections is when comparing two coherent states [53].

The fact that atoms can interact strongly via cavity fields makes it possible to investigate the implementation of superadditive coding with longer codewords using cavity QED-type systems. It is also interesting to further study realizations of other generalized quantum measurements which are difficult to realize using linear optics.

ACKNOWLEDGMENTS

A.D. is funded by a Scottish Universities Physics Alliance (SUPA) scholarship, M.E. acknowledges support from the Japanese Society for the Promotion of Science, and V.M.K. is funded by a UK Royal Society University Research fellowship.

-
- [1] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976).
- [2] A. Chefles, *Contemp. Phys.* **41**, 401 (2000).
- [3] I. D. Ivanovic, *Phys. Lett. A* **123**, 257 (1987).
- [4] D. Dieks, *Phys. Lett. A* **126**, 303 (1988).
- [5] A. Peres, *Phys. Lett. A* **128**, 19 (1988).
- [6] T. Brougham, E. Andersson, and S. M. Barnett, *Phys. Rev. A* **73**, 062319 (2006).
- [7] N. Gisin, G. Ribordy, W. Tittel, and H. Zbinden, *Rev. Mod. Phys.* **74**, 145 (2002).
- [8] C. H. Bennett, *Phys. Rev. Lett.* **68**, 3121 (1992).
- [9] B. Huttner, A. Muller, J. D. Gautier, H. Zbinden, and N. Gisin, *Phys. Rev. A* **54**, 3783 (1996).
- [10] R. B. M. Clarke, A. Chefles, S. M. Barnett, and E. Riis, *Phys. Rev. A* **63**, 040305 (2001).
- [11] M. Mohseni, A. M. Steinberg, and J. A. Bergou, *Phys. Rev. Lett.* **93**, 200403 (2004).
- [12] S. Franke-Arnold, E. Andersson, S. M. Barnett, and S. Stenholm, *Phys. Rev. A* **63**, 052301 (2001).
- [13] E. Andersson, *Phys. Rev. A* **64**, 032303 (2001).
- [14] L. Roa, J. C. Retamal, and C. Saavedra, *Phys. Rev. A* **66**, 012103 (2002).
- [15] T. Gopinath, R. Das, and A. Kumar, *Phys. Rev. A* **71**, 042307 (2005).
- [16] M. Takeoka, M. Fujiwara, J. Mizuno, and M. Sasaki, *Phys. Rev. A* **69**, 052329 (2004).
- [17] Y. Mitsumori, J. A. Vaccaro, S. M. Barnett, E. Andersson, A. Hasegawa, M. Takeoka, and M. Sasaki, *Phys. Rev. Lett.* **91**, 217902 (2003).
- [18] E. Knill, R. Laflamme, and G. J. Milburn, *Nature* **409**, 46 (2001).
- [19] P. Kok, W. J. Munro, K. Nemoto, T. C. Ralph, J. P. Dowling, and G. J. Milburn, *Rev. Mod. Phys.* **79**, 135 (2007).
- [20] E. Hagley, X. Maître, G. Nogues, C. Wunderlich, M. Brune, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* **79**, 1 (1997).
- [21] A. Rauschenbeutel, G. Nogues, S. Osnaghi, P. Bertet, M. Brune, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* **83**, 5166 (1999).
- [22] C. Guerlin, J. Bernu, S. Deleglise, C. Sayrin, S. Gleyzes, S. Kuhr, M. Brune, J.-M. Raimond, and S. Haroche, *Nature* **448**, 889 (2007).
- [23] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* **77**, 4887 (1996).
- [24] T. Lindström, C. H. Webster, J. E. Healey, M. S. Colclough, C. M. Muirhead, and A. Y. Tzalenchuk, *Supercond. Sci. Technol.* **20**, 814 (2007).
- [25] J. Vuckovic and Y. Yamamoto, *Appl. Phys. Lett.* **82**, 2374 (2003).
- [26] M. Reck, A. Zeilinger, H. J. Bernstein, and P. Bertani, *Phys. Rev. Lett.* **73**, 58 (1994).
- [27] E. Andersson and D. K. L. Oi, *Phys. Rev. A* **77**, 052104 (2008).
- [28] S. Haroche and J.-M. Raimond, *Exploring the Quantum: Atoms, Cavities, and Photons (Oxford Graduate Texts)*, 1st ed. (Oxford University Press, New York, 2006).

- [29] P. Filipovicz, P. Meystre, G. Rempe, and H. Walther, *J. Mod. Opt.* **32**, 1105 (1985).
- [30] H. Walther, B. T. H. Varcoe, B.-G. Englert, and T. Becker, *Rep. Prog. Phys.* **69**, 1325 (2006).
- [31] M. L. Jones, B. Sanguinetti, H. O. Majeed, and B. T. H. Varcoe, *Applied Soft Computing* **11**, 2079 (2011).
- [32] M. Tada, Y. Kishimoto, M. Shibata, K. Kominato, S. Yamada, T. Haseyama, I. Ogawa, H. Funahashi, K. Yamamoto, and S. Matsuki, *Phys. Lett. A* **303**, 285 (2002).
- [33] A. Gurtler and W. J. van der Zande, *Phys. Lett. A* **324**, 315 (2004).
- [34] A. Peres and W. K. Wootters, *Phys. Rev. Lett.* **66**, 1119 (1991).
- [35] M. Sasaki, K. Kato, M. Izutsu, and O. Hirota, *Phys. Rev. A* **58**, 146 (1998).
- [36] J. R. Buck, S. J. van Enk, and C. A. Fuchs, *Phys. Rev. A* **61**, 032309 (2000).
- [37] P. W. Shor, *IBM J. Res. Dev.* **48**, 115 (2004).
- [38] M. Fujiwara, M. Takeoka, J. Mizuno, and M. Sasaki, *Phys. Rev. Lett.* **90**, 167906 (2003).
- [39] S.-B. Zheng and G.-C. Guo, *Phys. Rev. Lett.* **85**, 2392 (2000).
- [40] M. S. Everitt, M. L. Jones, B. T. H. Varcoe, and J. A. Dunningham, *J. Phys. B* **44**, 035504 (2011).
- [41] C. Lazarou and B. M. Garraway, *Phys. Rev. A* **77**, 023818 (2008).
- [42] E. Andersson and S. M. Barnett, *Phys. Rev. A* **62**, 052311 (2000).
- [43] E. Vliegen and F. Merkt, *Phys. Rev. Lett.* **97**, 033002 (2006).
- [44] H. Merimeche, *J. Phys. B* **39**, 3723 (2006).
- [45] J. Zhang, J. Vala, S. Sastry, and K. B. Whaley, *Phys. Rev. A* **67**, 042313 (2003).
- [46] S. Croke, E. Andersson, S. M. Barnett, C. R. Gilson, and J. Jeffers, *Phys. Rev. Lett.* **96**, 070401 (2006).
- [47] S. Croke, E. Andersson, and S. M. Barnett, *Phys. Rev. A* **77**, 012113 (2008).
- [48] P. J. Mosley, S. Croke, I. A. Walmsley, and S. M. Barnett, *Phys. Rev. Lett.* **97**, 193601 (2006).
- [49] S. Osnaghi, P. Bertet, A. Auffeves, P. Maioli, M. Brune, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* **87**, 037902 (2001).
- [50] E. M. Purcell, *Phys. Rev.* **69**, 681 (1946).
- [51] S. Deleglise, I. Dotsenko, C. Sayrin, J. Bernu, M. Brune, J.-M. Raimond, and S. Haroche, *Nature* **455**, 510 (2008).
- [52] E. Andersson, S. M. Barnett, C. R. Gilson, and K. Hunter, *Phys. Rev. A* **65**, 052308 (2002).
- [53] C. S. Hamilton, H. Lavička, E. Andersson, J. Jeffers, and I. Jex, *Phys. Rev. A* **79**, 023808 (2009).