

**Propagation of initially excited states in time-dependent density-functional theory**

Peter Elliott and Neepa T. Maitra

*Department of Physics and Astronomy, Hunter College and the City University of New York, 695 Park Avenue,  
New York, New York 10065, USA*

(Received 29 March 2012; published 22 May 2012)

Many recent applications of time-dependent density functional theory begin in an initially excited state and propagate it using an adiabatic approximation for the exchange-correlation potential. This, however, inserts the excited-state density into a ground-state approximation. By studying a series of model calculations, we highlight the relevance of initial-state dependence of the exact functional when starting in an excited state, discuss different valid choices of the initial Kohn-Sham state, and explore the errors inherent in the adiabatic approximation that neglects this dependence.

DOI: [10.1103/PhysRevA.85.052510](https://doi.org/10.1103/PhysRevA.85.052510)

PACS number(s): 31.15.ee, 31.15.ec, 71.15.Qe, 71.15.Pd

**I. INTRODUCTION**

Time-dependent density functional theory (TDDFT) is an exact reformulation of the time-dependent quantum mechanics of many-electron systems [1,2] that operates by mapping a system of interacting electrons into one of fictitious noninteracting fermions, the Kohn-Sham (KS) system, reproducing the same time-dependent density of the true system. By the theorems of TDDFT, all properties of the true interacting system may be obtained from the KS system, in principle. In practice, approximations are needed for the exchange-correlation effects: in particular, for the exchange-correlation (XC) potential,  $v_{xc}[n; \Psi(0), \Phi(0)](\mathbf{r}, t)$ , a crucial term in the single-particle KS equations determining the evolution of the fictitious noninteracting fermions. This functionally depends on the one-body density  $n(\mathbf{r}, t)$ , the initial many-body state of the interacting system  $\Psi(0)$ , and the initial state of the KS system  $\Phi(0)$ .

From the inception of the theory, subtleties were noted in the functional dependences, which make good approximations for the XC potential more challenging to derive than for its counterpart in the (much older) ground-state density-functional theory [3–5]. One of these is memory: the dependence of the potential at time  $t$  on the density at earlier times and on the initial states  $\Psi(0)$  and  $\Phi(0)$ . Efforts to include memory dependence in functionals [6–14] are not widely used and have all focused on the dependence on the history of the density, neglecting the dependence on the initial states. The vast majority of applications have been in linear response from a nondegenerate ground state, and there this initial-state dependence (ISD) is redundant: by the Hohenberg-Kohn theorem, a nondegenerate ground state is a functional of its own density. In fact, the simple adiabatic approximation, where the instantaneous density is input into a ground-state approximation, neglecting any memory dependence, has shown remarkable success for a great range of excitations and spectra. All commonly available codes with TDDFT capabilities are written assuming the adiabatic approximation. Over the years it has been understood that certain types of excitations cannot be captured by the adiabatic approximation, e.g., double excitations [15] and excitonic series in optical response of semiconductors [16], and there is ongoing research both in developing frequency-

dependent kernels to deal with this as well as in understanding when not to trust the calculated adiabatic TDDFT spectra.

With the results and understanding of recent years lending a level of comfort with calculations of spectra and response, TDDFT has now entered a more mature stage, and with that come more adventurous applications; in particular, for electron dynamics in real time, evolving under strong laser fields, or coupled electron-nuclear dynamics following photoexcitation (e.g., Refs. [17–19]). The role of memory is less well understood in these applications; studies on model systems have shown that sometimes memory dependence is essential [20–24] and at other times it is not important at all [25,26]. Moreover, a new element enters: the initial-state dependence that was conveniently and correctly brushed aside in the linear-response regime now raises its head. In applications such as modeling solar cell processes, the initial photoexcitation of the electronic system is not dynamically modeled; instead the dynamics begins with the electronic system assumed to be in an excited state. The initial state is not the ground state, yet no truly initial-state-dependent functionals are available today; hence we ask how large are the errors in such calculations? Knowing that the true interacting system begins in a certain excited state, is there an optimal choice for the initial KS state when propagating with an adiabatic approximation? How large are the errors due to ISD compared to those due to history dependence?

In this paper, we begin to answer these questions by considering a series of model calculations of two-electron systems. We start by reviewing the underlying theorems of TDDFT and the subtleties of ISD, even in situations where we may not expect it. We show how ISD leads to a nonzero XC potential even for noninteracting electrons. Then, we move to electron dynamics in the model soft-Coulomb helium atom, performing adiabatic TDDFT calculations with different initial states and comparing with exact dynamics. Here we must dissect other sources of error in the usual TDDFT calculations, such as using an initial KS excited state whose density does not equal the true excited-state density, in order to properly ascribe the influence of ISD. Finally, we make the first step towards investigating how ISD can affect an area currently of much interest, namely, coupled electron-ion dynamics, by

performing Ehrenfest calculations for a model LiH system. We work in atomic units throughout ( $e^2 = m_e = \hbar = 1$ ).

## II. INITIAL-STATE DEPENDENCE IN TDDFT

In TDDFT, one evolves a set of single-particle orbitals  $\{\phi_j(\mathbf{r}, t)\}$  with a one-body KS potential:

$$i \frac{\partial}{\partial t} \phi_j(\mathbf{r}, t) = \left( -\frac{1}{2} \nabla^2 + v_s(\mathbf{r}, t) \right) \phi_j(\mathbf{r}, t), \quad (1)$$

$$v_s(\mathbf{r}, t) = v_{\text{ext}}(\mathbf{r}, t) + v_h[n](\mathbf{r}, t) + v_{\text{xc}}[n; \Psi_0, \Phi_0](\mathbf{r}, t), \quad (2)$$

where  $v_h(\mathbf{r}, t) = \int n(\mathbf{r}', t) / |\mathbf{r} - \mathbf{r}'| d^3 r'$  is the usual Hartree potential and  $v_{\text{xc}}(\mathbf{r}, t)$  is the XC potential which depends on the entire history of the density, the interacting initial state  $\Psi_0$ , and the initial KS wave function  $\Phi_0$ .

The origin of the initial-state dependence in the functionals is that the one-to-one Runge-Gross mapping [1] between time-dependent densities and potentials holds for a fixed initial state. Thus, the external potential  $v_{\text{ext}}$  functionally depends on the density and the true initial state  $\Psi(0)$ , and the KS potential  $v_s$  depends on the density and the KS initial state  $\Phi(0)$ . These functional dependences are not directly relevant themselves in a practical calculation, because there only the XC potential  $v_{\text{xc}}$  needs to be approximated. However, they lend their dependences to  $v_{\text{xc}}$  via Eq. (2), which must therefore depend on both the initial states and the density.

There are two known situations in which there is no initial-state dependence. When the initial state is a ground state, then, by the Hohenberg-Kohn theorem of ground-state DFT, it is itself a functional of its own density. ISD is redundant, as the information about the initial state is contained in the initial density. (See also Sec. II A.) The other situation is for one-electron systems: starting in *any* initial state, there is only one potential that can yield a given density evolution [27]. In all other situations, it is assumed that ISD cannot be subsumed into a density dependence; explicit demonstrations for two electrons can be found in Refs. [22,27]. The authors of Ref. [28] derived an exact condition relating the dependence on the history of the density to ISD. This condition is likely violated by any history-dependent functional approximation that has no ISD.

Technically, one may choose any initial KS wave function that has the same  $n$  and  $\dot{n}$  as the true initial state [29]. Usually a single Slater determinant (SSD) of  $N$  spin orbitals  $\phi_i$  is selected, with the required property that

$$\sum_{i=1}^N |\phi_i(\mathbf{r}, 0)|^2 = n(\mathbf{r}, 0) \quad (3)$$

and

$$-\nabla \cdot \text{Im} \sum_{i=1}^N \phi_i^*(\mathbf{r}, 0) \nabla \phi_i(\mathbf{r}, 0) = \dot{n}(\mathbf{r}, 0), \quad (4)$$

where

$$n(\mathbf{r}, 0) = N \sum_{\sigma} \int dx_2 \cdots \int dx_N |\Psi(x, x_2, \dots, x_N, 0)|^2, \quad (5)$$

$$\dot{n}(\mathbf{r}, 0) = -N \nabla \cdot \text{Im} \left\{ \sum_{\sigma} \int dx_2 \cdots \int dx_N \Psi^*(x, x_2, \dots, x_N, 0) \times \nabla \Psi(x, x_2, \dots, x_N, 0) \right\}, \quad (6)$$

where  $x = (\mathbf{r}, \sigma)$  and  $\int dx = \sum_{\sigma} \int d^3 r$ .

In this paper we will consider different choices of  $\Phi(0)$  for two-electron systems that begin in the first excited singlet state of the true system, denoted  $\Psi^*$ . Specifically we will, at various points, investigate the following three forms:

(a) An excited KS singlet state with two occupied orbitals  $\phi_0$  and  $\phi_1$ , whose spatial part has the form

$$\Phi^*(\mathbf{r}_1, \mathbf{r}_2) = \frac{1}{\sqrt{2}} [\phi_0(\mathbf{r}_1) \phi_1(\mathbf{r}_2) + \phi_0(\mathbf{r}_2) \phi_1(\mathbf{r}_1)]. \quad (7)$$

Note that this state is a sum of two Slater determinants.

(b) A ground KS singlet state  $\Phi^{\text{gs}}$ , which for two electrons corresponds to a single doubly occupied orbital

$$\Phi^{\text{gs}}[n^*](\mathbf{r}_1, \mathbf{r}_2) = \phi(\mathbf{r}_1) \phi(\mathbf{r}_2), \quad (8)$$

with

$$\phi(\mathbf{r}) = \sqrt{n^*(\mathbf{r})/2}, \quad (9)$$

where  $n^*(\mathbf{r}) = 2 \int |\Psi^*(\mathbf{r}, \mathbf{r}')|^2 d^3 r'$  is the density of the initial excited state. This is a SSD and is the usual choice when the true initial state is a ground state.

(c) A spin-symmetry-broken excited KS state of the form

$$\Phi_{\text{SB}}^*(\mathbf{r}'\sigma', \mathbf{r}\sigma) = \frac{1}{\sqrt{2}} \begin{vmatrix} \phi_0(\mathbf{r}) \delta_{\sigma\uparrow} & \phi_0(\mathbf{r}') \delta_{\sigma'\uparrow} \\ \phi_1(\mathbf{r}) \delta_{\sigma\downarrow} & \phi_1(\mathbf{r}') \delta_{\sigma'\downarrow} \end{vmatrix}. \quad (10)$$

This state is not a spin eigenstate (although it does have  $\langle \mathbf{S} \rangle = 0$ ), but is a valid choice for an initial KS state, having the same total density as the true system. The up- and down-spin densities are not equal, and we shall evolve them in different spin-up and spin-down KS potentials, but, unlike in spin DFT, we do not consider the spin densities separately meaningful; only their sum is considered as an observable [30]. Instead, (c) will be used to illustrate the effects of orbital-specific functionals, as will be discussed more later.

Now the exact KS potential differs depending on which choice (a), (b), or (c) is made, but any adiabatic approximation is identical for them all. Almost all the calculations being run today, certainly all that are coded in the commonly available codes, utilize such an approximation, and insert the instantaneous density into a ground-state approximation:

$$v_{\text{xc}}^{\text{adia}}[n; \Psi(0), \Phi(0)](\mathbf{r}, t) = v_{\text{xc}}^{\text{gs}}[n(t)](\mathbf{r}). \quad (11)$$

It is not surprising that there are errors inherent in such an approximation, and one question on which we hope to shed light by our investigations here is how much of this error is due to the lack of ISD, rather than the lack of history dependence [dependence on  $n(\mathbf{r}, t' < t)$ ]. Only the latter occurs in the usual calculations that begin in initial ground states, where, as mentioned earlier, ISD may be subsumed into density dependence.

### A. Beginning in the ground state: ISD or not?

Many practical situations start with the system in its ground state; in fact most calculations, except in the most

recent years, have assumed an initial ground state. This is fortunate for TDDFT, since, at least at short times, the adiabatic approximation with its ground-state functionals, which have become increasingly sophisticated over the years, could then be expected to work reasonably well. However, we will show that the subtleties of initial-state dependence can appear even in this case.

A completely legitimate choice for the KS initial state would be the true ground-state wave function as it trivially satisfies the initial conditions Eqs. (3) and (4). The exact KS potential, for this choice, reproduces the density evolution of the exact wave function propagating in the interacting system by propagating the exact initial wave function in a noninteracting system.

Given that at the initial time we are dealing with a ground state, one might expect that the adiabatic approximation would be exact, at least initially; however, this is not the case. The adiabatic approximation returns the XC potential for a ground-state DFT calculation, where the KS wave function is the ground-state KS wave function. If we were to start the KS calculation with this ground-state KS wave function, the adiabatic approximation would then be exact at the initial time. However, the initial interacting wave function is not a ground state of the noninteracting KS system, and so choosing this as the initial KS state means that the adiabatic approximation will be in error from the start.

Thus care must be taken when using the shorthand that there is no initial-state dependence when starting in the ground state. More precisely, what is meant is that (1) the initial state of the true interacting system is the ground state, and the initial KS orbitals are chosen to be those of the KS ground-state wave function of the same density, and (2) the dependence on the initial state for this case can be subsumed into the density via the theorems of ground-state DFT.

### III. NONINTERACTING ELECTRONS

The importance of initial-state dependence is strikingly evident even for the hypothetical case of noninteracting electrons. Due to ISD, the XC potential is *not* always equal to zero, which may be surprising at first sight, given that there is no interaction. Although no-one would perform KS calculations for noninteracting electrons for practical purposes, it is instructive to consider how the KS system behaves in this case. In particular, the studies suggest what is the best choice of KS initial state for a given true initial state when an adiabatic approximation is used. Since the adiabatic approximation is designed for ground states, is the error least if we always choose a KS state that is a ground state?

In the following we consider the “true” system to be noninteracting, i.e., we scale the electron-electron interaction by  $\lambda$  in the limit that  $\lambda \rightarrow 0$  and consider terms zeroth order in  $\lambda$  only, e.g., the Hartree potential vanishes. We then consider the XC potential when the KS system is started in different allowed initial states.

Consider two noninteracting electrons prepared in an excited state  $\Psi^*$  and evolving in some potential  $v_{\text{ext}}(t)$ . To start the KS evolution, any initial state of the same density  $n$  and first time derivative  $\dot{n}$ , zero in this case, may be chosen. We consider the initial-state choices (a) and (b)

introduced in Sec. II, which become here (a)  $\Phi(0) = \Psi^*$  and (b)  $\Phi(0) = \phi_0(\mathbf{r})\phi_0(\mathbf{r}') = \Phi^{\text{gs}}, \phi_0 = \sqrt{n^*/2}$ , where  $n^*$  is the density of  $\Psi^*$ .

For choice (a), the exact XC potential vanishes,  $v_{\text{xc}}[n; \Psi^*, \Phi^*](\mathbf{r}, t) = 0$ . This can be most easily seen by invoking the uniqueness property of the Runge-Gross mapping: for a fixed particle interaction, only one potential  $[v_{\text{ext}}(t)]$  can yield a given density evolution from a given initial state. Since  $\Phi^* = \Psi^*$  for choice (a), we conclude that  $v_{\text{xc}}(t) = 0$ .

Now consider choice (b). In this case,  $v_{\text{xc}}[n; \Psi^*, \Phi^{\text{gs}}](\mathbf{r}, 0)$  is nonzero, which can be seen from the general formula [29]

$$\nabla \cdot \{n(\mathbf{r}, t) \nabla [v_{\text{H}}(\mathbf{r}, t) + v_{\text{xc}}(\mathbf{r}, t)]\} = q(\mathbf{r}, t) - q_{\text{s}}(\mathbf{r}, t), \quad (12)$$

where

$$q(\mathbf{r}, t) = \frac{1}{i\hbar} \nabla \cdot \langle \Psi(t) | \hat{\mathbf{j}}(\mathbf{r}, \hat{T} + \hat{V}_{ee}) | \Psi(t) \rangle, \quad (13)$$

$$q_{\text{s}}(\mathbf{r}, t) = \frac{1}{i\hbar} \nabla \cdot \langle \Phi(t) | \hat{\mathbf{j}}(\mathbf{r}, \hat{T}) | \Phi(t) \rangle, \quad (14)$$

and  $\hat{\mathbf{j}}(\mathbf{r})$  is the current-density operator. At time  $t = 0$ , the right-hand side of Eq. (12) is nonzero, even in the noninteracting limit, due to the difference in the initial wave functions in  $q$  and  $q_{\text{s}}$ . This leads to a nonzero value for  $v_{\text{xc}}$ . This result can also be understood by the fact that  $v_{\text{xc}}$  must be such that  $v_{\text{s}} = v_{\text{ext}} + v_{\text{H}} + v_{\text{xc}}$  evolves the initial  $\Phi^{\text{gs}}$  with the same density for all time as the different initial state  $\Psi^*$  has when evolved in  $v_{\text{ext}}$ . So, ISD leads to a nonzero XC potential, even when the electrons do not interact.

For an explicit demonstration, consider the initial time and realize that  $v_{\text{xc}}(t = 0)$  depends entirely on the initial states, as can be seen from Eq. (12), and not on the choice of the external potential. Figure 1 plots this potential  $v_{\text{xc}}(t = 0)$  for the first excited state of an external harmonic potential ( $\frac{1}{2}x^2$ , on the left), and a soft-Coulomb potential ( $-2/\sqrt{x^2 + 1}$ , on




FIG. 1. (Color online) Top: The first excited-state density  $n^*(x)$  for two noninteracting electrons in a one-dimensional harmonic potential (left) and soft-Coulomb potential (right). Lower: The corresponding correlation potentials  $v_{\text{c}}[n^*, \Psi^*, \Phi^{\text{gs}}[n^*]](x)$  for an initial KS wave function chosen to have ground-state form but yielding the initial excited-state density  $n^*$  [case (b) of the text]. The inset shows the KS potential (solid line) compared with the external potential (dashed line), chosen as harmonic and soft-Coulomb potentials, respectively. When the initial KS state is chosen to be  $\Phi^*$  [case (a) in the text], on the other hand, the correlation potential is zero.

the right) in one dimension. The top panels show the density, and the insets show the external potential in which  $\Psi^*$  is an eigenstate, as well as the the KS potential  $v_{\text{ext}} + v_{\text{c}}$  in which  $\Phi^{\text{gs}}$  is an eigenstate (the ground state, since there are no nodes). Because for two electrons in a spin singlet with one doubly occupied orbital,  $v_{\text{x}} = -v_{\text{H}}/2 = 0$  in the noninteracting limit, this effect appears entirely in the correlation potential and can be interpreted as *static correlation*: due completely to the non-single-determinantal structure of the true initial state. In case (a), this effect vanishes, because the KS initial state is chosen also not to be a SSD.

We now ask what an adiabatic potential would give: approximating  $v_{\text{xc}}[n; \Psi(0), \Phi(0)]$  by a ground-state potential  $v_{\text{xc}}^{\text{gs}}[n]$ . To distinguish errors arising from the choice of the approximate ground-state functional itself, we consider an “adiabatically exact” potential [21,25]  $v_{\text{xc}}^{\text{adia ex}}[n]$ . We shall define this generally in the next section, but for now it suffices to define it such that if both the true and KS wave functions at all times were in fact ground states of some potential, then the exact XC potential would be the exact ground-state one,  $v_{\text{xc}}[n; \Psi^{\text{gs}}, \Phi^{\text{gs}}] = v_{\text{xc}}^{\text{adia ex}}[n]$ . Consider again just the initial time  $t = 0$ . In the noninteracting limit, for both the true and KS wave functions to be ground states that have the density of the excited state  $\Psi^*$ , then  $\Psi(0) = \Phi(0) = \sqrt{n_0(\mathbf{r})}n_0(\mathbf{r}')/2$ , and we rapidly conclude that

$$v_{\text{xc}}[n, \Psi^{\text{gs}}, \Phi^{\text{gs}}](\mathbf{r}, t) = v_{\text{xc}}^{\text{adia ex}}[n](\mathbf{r}, t) = 0 \quad (15)$$

in the noninteracting limit. Applying now this adiabatic approximation to the case of the initial true excited state  $\Psi^*$  considered above, we conclude that *the adiabatic approximation is exact for choice (a)*, when the initial KS state is also chosen excited, as we have argued that there the exact  $v_{\text{xc}}$  also vanishes. It is inaccurate for choice (b), when the initial KS state is chosen to be a ground-state one, where the exact correlation potential is nonzero.

This study suggests that in the general interacting case, errors in adiabatic TDDFT will be least when an initial KS state that most closely resembles the dominant configuration of the true excited state is chosen. This expectation is indeed borne out in the following studies.

#### IV. MODEL TWO-ELECTRON SOFT-COULOMB-INTERACTING SYSTEMS

The case of noninteracting electrons illustrated the importance of ISD, while offering a hopeful diagnosis for the adiabatic approximation; the latter becomes exact if the initial state is chosen appropriately. Of course, electrons do interact, and now we turn to studying the effect of ISD on the dynamics of interacting systems.

When running an approximate TDKS calculation starting in an excited state of the interacting problem, three separate sources of error come into play. First, excited states of the exact ground-state KS potential  $v_{\text{s}}[n_0](\mathbf{r})$  do not have the same densities as interacting excited states of the potential  $v_{\text{ext}}(\mathbf{r})$  whose ground-state density is  $n_0(\mathbf{r})$ . Yet these are usually the ones chosen in practice. We discuss this problem in Sec. IV A. The second source of error is the central one for this paper: the use of the adiabatic approximation, when, from the start, we have an excited state. We consider first

the “adiabatically exact” approximation, mentioned also in Sec. III, but considered now for interacting electrons, in Sec. IV B. This will allow us to separate the errors from the choice of ground-state functional approximation used in the adiabatic approximation, which is the third issue. In Secs. IV D and V, we will study the effect that missing initial-state dependence has in practice on electron dynamics in model two-electron systems when begun in the first singlet excited state of the system.

In our model one-dimensional helium atom, the two electrons interact via a soft-Coulomb electron-electron interaction  $v_{ee}(x_1, x_2) = 1/\sqrt{(x_1 - x_2)^2 + 1}$  and exist in an external soft-Coulomb potential,

$$v_{\text{ext}}(x) = -2/\sqrt{x^2 + 1}. \quad (16)$$

Such a model, straightforward to solve numerically, is popular for understanding and analyzing electron interactions, in both the ground-state and in strong-field dynamics [31], inside and outside the density-functional community. In our model of the diatomic LiH molecule, the soft-Coulomb interaction is also used, while the external potential has an asymmetric soft-Coulomb well structure (Sec. V).

Before proceeding, we make a computational note. The true dynamics are propagated on a real-space grid using the exponential midpoint rule Taylor-expanded to fourth order. Imaginary-time propagation along with Gram-Schmidt orthogonalization is first performed to find the lowest eigenstates. The adiabatic exact-exchange (AEXX) dynamics uses the Crank-Nicolson method with an explicit predictor step for the Hartree-exchange potential. In both cases, a time step of 0.001 is used with a grid spacing of 0.1. When possible these calculations were tested for accuracy against the parallelized code OCTOPUS [32], which was also used for local density approximation (LDA) and self-interaction-corrected LDA (LDASIC) runs.

##### A. The excited-state density

The theorems of TDDFT require that the KS initial state has the same  $n(\mathbf{r}, 0)$  and  $\dot{n}(\mathbf{r}, 0)$  as the true interacting system (Sec. II). In this section we discuss the difficulty of finding an appropriate KS state when the initial state is not a ground state.

When the true initial state is a ground state, the natural and usual choice for the KS initial state is the noninteracting ground state, and this can be found by solving the ground-state KS equations. However, if we want to compute the dynamics of an excited state, we first encounter the problem that there is no DFT scheme to find excited-state densities [33]. Furthermore, even if the true excited state  $\Psi(0)$  can be found using a more computationally expensive higher-level wave-function method, we still must then choose an initial  $\Phi(0)$  in which to start the KS calculation. If we start the interacting system in a first excited state of some  $v_{\text{ext}}(\mathbf{r})$ , the results of the earlier sections suggest that a good choice for adiabatic TDDFT is to start the KS system in a corresponding noninteracting excited state where one electron is excited from the highest occupied to the lowest unoccupied orbital of some  $\tilde{v}_{\text{s}}$  [as in form (a) of Sec. III]. This  $\tilde{v}_{\text{s}}(\mathbf{r})$  however cannot be the ground-state KS potential  $v_{\text{s}}(\mathbf{r})$  corresponding to the interacting  $v_{\text{ext}}$ , because this would not satisfy Eq. (3):  $v_{\text{s}}(\mathbf{r})$  yields the same ground-state






FIG. 2. (Color online) Exact excited-state density (solid line) compared to the exact excited-state density of the exact ground-state KS potential (dashed line) for the soft-Coulomb helium atom.

density in a noninteracting system that  $v_{\text{ext}}(\mathbf{r})$  yields in the interacting one, but the densities of their excited states are different. Yet the usual practice *is* to use the corresponding excited state of the ground-state KS potential [34]. This is partly because the density of the interacting excited state is not often known anyway. In Fig. 2 we compare the exact density of the first excited interacting singlet state in the soft-Coulomb He atom Eq. (16) with the density of the excited state of the corresponding ground-state KS potential. Although matching the general shape, the latter is too narrow and misses some structure.

To be able to separate the error from not quite having the exact initial density from the error from using an adiabatic approximation, we now search for a noninteracting system where the first singlet excited-state density *is* exactly equal to the true excited-state density  $n^*$ , for our model He atom. Several such potentials may exist [33], and in the lower panel of Fig. 3, we plot one such KS potential, denoted  $v_s[n^*, \Phi^*](x)$ , and compare to the KS potential where  $n(x)$  is the exact ground-state density  $v_s[n](x, 0)$  of the model He atom. (The first-excited-state densities of these potentials are precisely those shown in Fig. 2.) We also show  $v_s[n^*](x, 0)$ , i.e., the KS potential for which  $n^*(x)$  is the density of the ground state. This corresponds to choice (b) in Sec. II for the initial KS state.

Once a valid KS state is found satisfying Eqs. (3) and (4), the initial XC potential is completely determined: Eqs. (12) and (13) show that at the initial time,  $v_{\text{xc}}$  is a functional of just the two initial states  $\Psi(0)$  and  $\Phi(0)$ . This can be added to any external potential, prescribed by the physical problem at hand, and Hartree potential, determined by the initial density, to start the evolution. In other words, the choice of the initial KS state fundamentally points to an XC potential rather than to a KS potential, and so in the top panel of Fig. 3 we plot the XC potentials corresponding to choices (a) and (b) of the initial KS state. In the following examples, we take the initial  $v_{\text{ext}}$  to be that of the soft-Coulomb potential Eq. (16); sometimes subsequently an external field is added.

We have therefore now found initial states of the forms (a) and (b) of Sec. II, both yielding the same density as that of the true excited state, meeting the conditions Eqs. (3) and (4). We can now move forward to investigate the time-dependent properties.




FIG. 3. (Color online) Top: The exact XC potentials for the two choices of initial state  $\Phi^*$  (solid line) and  $\Phi^{\text{gs}}$  (dot-dashed line). Bottom: The KS potentials for which the two initial-state choices  $\Phi^*$  (solid line) and  $\Phi^{\text{gs}}$  (dot-dashed line) are eigenstates. Also shown is the exact ground-state KS potential (dashed line) for the soft-Coulomb helium potential (dotted line).

## B. The exact adiabatic approximation

We next move to the error introduced by using the adiabatic approximation. To isolate this error we will calculate the adiabatically exact potential at the initial time, and start in a KS state of the exact same density as the true state, as found in Sec. IV A.

The adiabatically exact potential is defined by [25]

$$v_{\text{xc}}^{\text{adia ex}}[n] = v_s^{\text{adia all}}[n] - v_{\text{ext}}^{\text{adia all}}[n] - v_{\text{H}}[n], \quad (17)$$

where  $v_s^{\text{adia all}}[n](\mathbf{r}, t)$  is the potential for which  $n(\mathbf{r}, t)$  is the noninteracting ground-state density and  $v_{\text{ext}}^{\text{adia all}}[n](\mathbf{r}, t)$  is the potential for which interacting electrons have  $n(\mathbf{r}, t)$  as their ground-state density. If both the true and KS wave functions are actually always in some ground state, then Eq. (17) becomes the exact XC potential. We shall consider this potential for initially excited states, only at the initial time.

The noninteracting potential  $v_s^{\text{adia all}}[n^*](\mathbf{r}, 0)$  may be easily found by inverting the KS equation for a doubly occupied orbital,  $\phi(\mathbf{r}) = \sqrt{n^*(\mathbf{r})}/2$  where  $n^*(\mathbf{r})$  is the density of the initial state  $\Psi^*$ . The interacting potential  $v_{\text{ext}}^{\text{adia all}}[n](\mathbf{r}, 0)$  is solved for using an iterative technique whereby the external potential is updated based on the difference between the ground-state density of the current iteration and the density we are targeting. This is based on the inversion algorithm of Ref. [35], but generalized to the interacting case [25]. We also increase the update to the potential in regions of low density by simply using the inverse of the density (up to a maximum value) as a weighting factor, as in Ref. [30]. When the density converges to the target density, we have found  $v_{\text{ext}}^{\text{adia all}}[n](\mathbf{r}, 0)$ .

The dotted line in the lower panel of Fig. 4 shows at the initial time the full KS potential with the adiabatically exact XC potential, namely,  $v_s^{\text{adia}}(x) = v_{\text{ext}}(x) + v_{\text{H}}[n](x) + v_{\text{xc}}^{\text{adia ex}}[n](x)$ , choosing  $v_{\text{ext}}(t = 0)$  as the soft-Coulomb




FIG. 4. (Color online) Top: The exact XC potentials and the exact adiabatic XC potential. Bottom: The exact and adiabatically exact KS potentials.

potential of Eq. (16), for the first excited singlet state  $\Psi^*$ . Alongside, we compare this to the exact KS potentials found in Sec. IV A (see Fig. 3), for the initial-state choices of cases (a) and (b), respectively. The adiabatically exact KS potential  $v_S^{\text{adia}}(x)$  tracks the shape of the exact KS potential  $v_S[n^*, \Psi^*, \Phi^*]$  although it misses structure. The difference between them is the difference in their XC potentials shown in the upper panel of Fig. 4. The difference is much larger for the comparison with the exact KS potential when the initial KS state is chosen as the ground state (i.e.,  $v_{\text{XC}}^{\text{adia-ex}}[n^*]$  is closer to  $v_{\text{XC}}[n^*, \Psi^*, \Phi^*]$  than it is to  $v_{\text{XC}}[n^*, \Psi^*, \Phi^{\text{GS}}]$ ); in this case, there is a large static-correlation contribution to the exact XC potential. The results here are consistent with the noninteracting case discussed in Sec. III: with interaction, the adiabatic approximation is no longer exact for the choice of  $\Phi^*$ ; however, it is still the better choice over  $\Phi^{\text{GS}}$ .

The fact that the adiabatically exact potential does not equal the true potential, even at the *initial* time, leads to erroneous dynamics. To illustrate this effect we propagate our exact initial wave functions in the adiabatically exact potential and compare with exact dynamics. Since calculating at each time is computationally demanding and delicate, we will simply evolve in time without any additional perturbing potentials, and hold the adiabatic potential fixed to its initial value. The idea is that since the exact evolution is static, the adiabatically exact potential is static and equals its initial value at all times; if this were a good approximation, it would yield density dynamics that were close to static. The deviation of the adiabatic propagation from the exact is a measure of its error. Note that such a calculation does not treat the adiabatic potential self-consistently; later calculations with approximate adiabatic functionals suggest that having self-consistency decreases the error somewhat.

In Fig. 5, we plot snapshots of the density at 5 a.u. intervals until  $T = 20$  a.u. for the two choices of exact initial states discussed above, evolving in the fixed adiabatically exact potential plotted in Fig. 4. As each time we show the exact




FIG. 5. (Color online) The density at times 5, 10, 15, and 20 a.u. for initial states  $\Phi^*$  (solid line) and  $\Phi^{\text{GS}}$  (dashed line) propagating in the static exact adiabatic initial potential. Also shown is the exact excited-state density (dotted line) which is the exact solution at each time.

static excited-state density. While both choices show density “leaking” out, the  $\Phi^*$  choice is more successful in preventing this, as might be expected from the above discussions, but cannot stop the density melting away in its outer regions. The  $\Phi^{\text{GS}}[n^*]$  choice is poor from the start, jettisoning density outward and becoming too narrow.

This error in the adiabatically exact evolution is caused *entirely* by initial-state dependence: for the analogous calculation beginning in the interacting ground state and starting with the ground-state KS wave function, the adiabatic approximation would be exact at all times when there are no perturbing fields.

### C. Approximate ground-state functionals

Finally we look at the third source of error, namely, use of an approximate ground-state XC functional instead of the exact one. This error contributes both at the ground-state level in obtaining the initial orbitals and in the adiabatic functional when computing dynamics. We first turn our attention to the latter and investigate the adiabatic exact exchange functional AEXX, which, for a two-electron system, is simply  $v_{\text{XC}}^{\text{AEXX}}[n](\mathbf{r}, t) = -v_{\text{h}}[n](\mathbf{r}, t)/2$ . In Fig. 6 we compare the adiabatically exact XC potential found in the previous section to the AEXX




FIG. 6. (Color online) The exact adiabatic XC (solid) and AEXX (dashed) potentials evaluated on the true excited-state density  $n^*$ . (Note that the solid line here is the same as the dotted line in the top panel of Fig. 4.)




FIG. 7. (Color online) The ground and first excited orbitals from ground-state EXX and LDA calculations, compared to those from the exact ground-state (g.s.) KS potential. Also shown are those orbitals found in Sec. IV A of this configuration that yield the exact excited-state density.

potential, at the initial time. The undulations of the exact adiabatic XC potential that are missing in the AEXX potential represent correlation; they are relatively small in this particular case once added to the external potential, so we expect that AEXX dynamics fairly approximates the adiabatically exact dynamics in this situation at least for short times.

We now return to the error of not having the correct initial density (Sec. IV A). In the usual TDDFT calculation, a ground-state DFT calculation is first performed and the orbitals from this are used to create the initial wave function. So we would start with the ground-state KS density of Fig. 2 except here there is a further error that will be made, as an approximate ground-state XC functional is used instead of the exact one.

In Fig. 7 we plot the ground and first excited KS orbitals from ground-state DFT calculations using EXX and LDA results for the soft-Coulomb helium atom, and compare to the exact KS orbitals. We also plot, for completeness, the pair of orbitals found for this configuration that yield the exact *interacting* excited-state density, i.e., these are the exact orbitals which, when singly occupied, make up the densities shown in Fig. 2.

If we first compare the approximate orbitals to the exact KS orbitals of the soft-Coulomb potential, we find that both the LDA and EXX methods perform very well, particularly for the ground-state orbital. Their ground-state KS potentials are, however, quite different: it is well known that the LDA potential is much too shallow, with a much too rapid exponential decay away from the nucleus, while the EXX potential is closer to the exact, with the correct asymptotic decay. This is reflected in the first excited orbital in the LDA, which is more diffuse than that of EXX and the first excited orbital of the exact soft-Coulomb KS potential. The asymptotic behavior will be important especially when we turn on an electric field, so we will (mostly) use the EXX orbitals to build our initial KS wave functions.

In the next section we will add the spin-symmetry-broken initial state (c) introduced in Sec. II to our investigations; while no longer a spin eigenstate as in the exact case or  $\Phi^*$  and  $\Phi^{gs}$ , this still has the same total density as the exact system. We will simply construct this from the same EXX orbitals composing  $\Phi^*$ , but will evolve the two orbitals using *different* XC potentials  $v_{xc}^\uparrow$  and  $v_{xc}^\downarrow$ . For exact exchange, we have

$v_{\text{HXC}}^\uparrow[n](x) = v_{\text{H}}[n^\uparrow](x)$  for two electrons. We are however *not* performing spin DFT since the individual spin densities are not physical ones; we only ever consider their sum as observable. (Indeed, even the initial spin densities are wrong.) The point of considering such an initial state and dynamics is that, for two electrons, it is an example of dynamics in *orbital-specific* potentials, as would occur in generalized KS approaches [36–38]. Orbital-dependent functionals require an optimized effective potential (OEP) approach to find a single potential in which all the KS orbitals can exist. This procedure is numerically very intensive, so often a generalized KS approach is used, that relaxes the condition that all orbitals evolve under the same potential; moreover, the latter has been shown to have certain advantages over the OEP in certain cases, e.g., better band gaps [39]. The interest in orbital-dependent functionals is that they can work less hard to capture both spatially nonlocal and time nonlocal density dependence since the orbitals themselves are nonlocal functionals of the density.

We note nevertheless that it can be shown there is no ISD for two-electron dynamics with spin TDDFT; this follows straightforwardly by generalizing the arguments of Ref. [27] to each spin density. This is not the case for our spin-broken case, however, as here we are *not* attempting to reproduce the exact spin densities but rather the total density evolution; only  $n(\mathbf{r}) = n_\uparrow + n_\downarrow$  will be considered meaningful.

Having delineated and explored the three possible sources of error in an adiabatic TDDFT calculation of excited-state dynamics, we are now ready to run a typical adiabatic TDDFT calculation on our model systems, starting with our different choices of initial KS states, and compare with the exact result.

#### D. Propagation in an electric field

We simulate electric-field-driven dynamics in the soft-Coulomb helium model, using adiabatic TDDFT. We apply a relatively weak oscillating electric field of amplitude 0.01 a.u. with an off-resonant frequency of 0.2 a.u. We run for six cycles including a trapezoidal envelope consisting of a two-cycle linear switch-on and a two-cycle switch-off.

As a point of reference, we first look at the performance of TDDFT when starting in the *ground state*. We propagate the doubly occupied EXX ground-state orbital in the electric field, using AEXX, and compare to the exact dynamics of the initial interacting ground state. The resulting dipole moment is very accurate as can be seen in Fig. 8. In fact, almost exact dynamics can be achieved in this case with an adiabatic approximation: the same figure shows the result of using the adiabatic LDASIC to propagate the LDASIC ground-state orbital. The dipole moment lies practically on top of the exact curve. Including correlation in a self-interaction-free functional therefore improves the result over exact exchange, but an adiabatic approximation is certainly adequate in this case. We also show for comparison the LDA result.

The adiabatic approximation is, however, not so good, as we shall see, in the case of an initially excited state. The initial interacting state is the first singlet excited state ( $E = -1.705$  a.u.).

The typical TDDFT calculation will begin in an excited state determined by the orbitals of the corresponding ground-state KS system, but, as discussed in Sec. IV A, this




FIG. 8. (Color online) The exact dipole moment and those from TDDFT calculations starting in the self-consistent ground-state KS wave function and propagated with the corresponding adiabatic approximation, all under the influence of the electric field described in the text.

does not have quite the right density to start with. Since in our model system we found an initial KS state of the correct density, we first check the error made by using the AEEXX orbitals. We plot in Fig. 9 the dipole moments for AEEXX calculations starting in the  $\Phi^*$  initial state with exact orbitals and with the ground-state EXX orbitals (i.e., using the solid and dotted orbitals of Fig. 7, respectively). As previously anticipated, the two calculations perform similarly, especially at shorter times.

At last we are ready to study the excited-state electron dynamics, using, as in practical calculations, an approximate ground-state KS potential (in this case EXX) to generate the orbitals composing the various initial state choices. In Fig. 10, we plot the dipole moment for each choice of initial state. The spin-broken-symmetry case gives the best dynamics; as mentioned earlier, functionals expressed directly in terms of instantaneous orbitals automatically contain time-nonlocal and spatially nonlocal density information. In this particular case the orbital-specific EXX potentials, obtained from spin scaling the (spin-restricted) AEEXX potential, are self-interaction-free, while the latter is not for the case of the *excited* two-orbital singlet state. Unlike in the propagation of the ground-state, adding correlation to the adiabatic potential does not significantly improve the results, as can be seen in Fig. 11; the symmetry-broken LSDASIC and EXX calculations are both




FIG. 9. (Color online) The dipole moments when beginning in  $\Phi^*$  composed of EXX orbitals (dashed line) and the exact orbitals of Sec. IV A (solid line) in the electric field (see text).




FIG. 10. (Color online) The dipole moments for dynamics in the given electric field starting in the  $\Phi^*$ ,  $\Phi_{SB}^*$ ,  $\Phi_{SS}^*$  initial KS wave functions and using AEEXX, compared to the exact result.

off by roughly the same amount. Figure 11 also shows the local spin density approximation (LSDA) result, clearly worse than the others by comparison.

The  $\Phi^*$  initial state has the correct spin symmetry and is closest in character to the true excited state, being a double Slater determinant. It gives better dynamics than the  $\Phi_{SS}^*[n^*]$  initial state, whose dynamics are completely incorrect. This is consistent with the results in the previous sections. We can say with confidence that it is ISD and its underlying static correlation that is causing poor dynamics for this doubly-occupied-singlet case.

We can understand the poor result for the initial state of the Eq. (8) form also by looking at the exact adiabatic XC potential and  $v_{xc}[n^*, \Psi^*, \Phi_{SS}^*[n^*]](x)$ , both shown in Fig. 4. This form treats the density as if it were a ground state and uses one doubly occupied orbital; inversion of this orbital yields  $v_s^{\text{dia all}}[n](x)$ , i.e., this is the potential in which the wave function is static at the initial time. However, in a calculation using the adiabatic approximation, it will fall into a much deeper potential, drastically affecting its dynamics. Without the initial-state dependence of the XC potential giving a bump similar to that in Fig. 1, it will perform poorly.

## V. MODEL LiH DYNAMICS

Our final example is perhaps the most topical one, where the initial state is an excited one: coupled electron-ion dynamics




FIG. 11. (Color online) Dynamics for the spin-broken case,  $\Phi_{SB}^*$ , for different orbital-specific adiabatic functionals EXX, LSDASIC, and LSDA, compared to the exact result.



after photoexcitation. The initial excitation is assumed to place the initial nuclear wave packet on an excited potential energy surface, vertically up from its ground-state equilibrium. Field-free dynamics on the excited surface ensues; usually the nuclei are treated classically, coupled in either an Ehrenfest or a surface-hopping scheme to the quantum electron dynamics. For Ehrenfest dynamics in the TDDFT framework [34] and the simplest type of surface hopping [40], the electrons start in the KS excited state obtained simply from promoting an electron from the highest occupied orbital in the ground-state KS configuration to a virtual orbital. This initial state is then evolved in the time-dependent external potential caused by the moving ions. The dynamics of the ions is given by simple Newtonian mechanics where the electronic system provides a force  $\int n(\mathbf{r}, t) \nabla_R v_{\text{ext}}(\mathbf{r}; \{\mathbf{R}\})$ , where  $\{\mathbf{R}\}$  represent the nuclear coordinates. All the three errors mentioned earlier therefore raise their heads: the initial density is not that of the true excited state (even if the exact XC potential was used), an adiabatic approximation is used to propagate it, and this involves an approximate ground-state functional. It should be noted that in the more accurate surface-hopping method of Ref. [41], where correct TDDFT surfaces are used to provide the nuclear forces [42], the first problem does not arise; the method in fact circumvents the necessity to define the electronic state.

We will use the parameters found in Ref. [43] for a one-dimensional two-electron lithium hydride model, where

$$v_{\text{ext}}(x) = -1/\sqrt{x^2 + 0.7} - 1/\sqrt{x^2 + 2.25} \quad (18)$$

and  $v_{nn}(R) = -1.0/\sqrt{R^2 + 1.95}$ . We integrate Newton's equations of motion using the leapfrog algorithm, and for simplicity, we use the same time step for both electrons and ions.

The Born-Oppenheimer potential energy surfaces are shown in Fig. 12 for both the exact system and the bare KS system calculated within the spin-restricted EXX method. The ground-state surface has a minimum at  $R = 1.55$ , whereas the EXX surface has the g.s. equilibrium at  $R = 1.45$ . Near equilibrium the surfaces are somewhat similar. The spin-restricted EXX surfaces, however, encounter the well-known fractional charge problem as  $R$  increases, with the calculations becoming extremely difficult to converge and the lowest two surfaces collapsing onto each other. In the exact surface,




FIG. 12. (Color online) The exact Born-Oppenheimer potential energy surfaces (solid lines) and the EXX KS Born-Oppenheimer surfaces (dashed lines). Note that due to degeneracy between the lowest two surfaces, these calculations cannot be converged for large values of  $R$ , the interatomic separation.




FIG. 13. (Color online) The interatomic distance,  $R(t)$ , for Ehrenfest dynamics starting in the first excited-state and propagated freely. The various KS initial wavefunctions ( $\Phi^*$ ,  $\Phi_{\text{SB}}^*$ ,  $\Phi^{\text{gs}}$ ) are propagated with the adiabatic exact exchange approximation. Inset: short-time dynamics for the various initial states.

an avoided crossing can be seen between the ground and first excited states around  $R = 4$ , where the latter establishes ionic character, eventually decaying as  $-0.896 - 1/x$ , while the neutral dissociation curve flattens out. In fact there are multiple avoided crossings at larger distances, the next being around  $R = 10$ . Unrestricted EXX calculations break the spin symmetry of the true ground state at a critical separation, but describe dissociation and the shape of the potential energy surfaces at larger distances qualitatively [44,45].

We vertically excite the system at the ground-state equilibrium geometry into the first excited singlet state and then begin the propagation with the initial nuclear momentum chosen as zero. In Fig. 13, we show the internuclear separation as a function of time for our three choices of initial states, calculated within the EXX method. For  $\Phi^*$  we make the usual practice of taking  $\phi_0$  in Eqs. (7) and (10) as the highest occupied molecular orbital and  $\phi_1$  as the lowest unoccupied molecular orbital; likewise for the spin-symmetry-broken version  $\Phi_{\text{SB}}^*$ . The density of this state  $n^*$  is then used to define the doubly occupied orbital in the SSD state  $\Phi^{\text{gs}}[n^*]$  [Eq. (8)].

For this particular system, the exact system remains on the first excited surface; the nuclear coordinate oscillates between the turning points at  $R = 1.5$  and  $R = 11.165$ . The ground state  $\Phi^{\text{gs}}$  completely fails to capture this behavior, making only very small oscillations around its initial position. The excited states  $\Phi^*$  and  $\Phi_{\text{SB}}^*$  show reasonable agreement for short times, with the spin-broken state once again edging out the other. For longer times, however, the spin-broken state is dramatically wrong, yielding nuclei moving away from each other with constant speed. The excited state  $\Phi^*$  shows behavior more qualitatively exact but, because of the deviation of its potential energy curve compared to the exact one, the oscillation period is quite wrong.

The conclusions regarding ISD in this example, at least at short times, are consistent with the results throughout this paper and support the idea that when using an adiabatic approximation the choice of a KS initial state with a configuration close to that of the exact interacting initial state yields the better result. However, the behavior at longer times in this problem highlights, above all, the need for accurate ground-state functionals: EXX produces poor nuclear dynamics when  $\Phi^*$  is chosen as the KS state primarily because the shape of its

potential energy surfaces is wrong—the ground-state as well as the excited-state ones. Any adiabatic approximation that does not have strong correlation will not perform well. Despite giving improved potential energy surfaces, the spin-broken EXX approach fails for dynamics for longer times. We believe this is because, as the molecule dissociates, symmetry breaking localizes each electron on one nucleus or the other, with each evolving according to a different XC potential; each atom experiences the other as just a neutral atom, leading to a net zero potential and a constant velocity (somewhat as if the molecule were dissociating on the ground-state surface). It should be noted that the spin-broken solution is not evolving on the spin-broken potential energy surface, e.g., even at the initial time, at the equilibrium geometry there is no symmetry breaking in a spin-unrestricted approach, so the spin-broken surface is on top of the spin-restricted surface shown in the figure. However the spin-broken wave function (c) is evolved using different potentials for each orbital, unlike the evolution dictated by either the spin-restricted or the unrestricted EXX surface (the symmetry-breaking point occurs around  $R = 2.5$ ).

## VI. CONCLUSIONS AND OUTLOOK

Until relatively recently, ISD fell in the realm of a theoretical curiosity, but due to an increasing number of topical applications beginning with the system not in its ground state, ISD is now also of much practical concern. Almost all calculations today use adiabatic functionals that neglect the ISD that the exact functionals are known to have.

By considering several choices of initial KS states when the initial interacting state is excited, we explored the effect of ISD on dynamics in several exactly solvable model systems, and the performance of the adiabatic approximation. We noted that there are three sources of error: excited KS initial states not having the same density as the corresponding true excited states even when the exact functional is used, the use of the adiabatic approximation to propagate the initial state, and the ground-state functional generating the adiabatic approximation itself being approximate. By separating these errors from each other, we were able to properly assign how badly they impacted the dynamics in model systems, allowing us to see the effect of ISD more clearly.

When the initial KS state has a configuration that is significantly different from that of the true state, the adiabatic

approximation fails severely—even for noninteracting electrons, where there is a significant error that can be interpreted as a static correlation effect. The optimal choice for the KS initial state is one whose configuration is most similar to the true excited state; this is in fact the usual practice in recent calculations (e.g., Ref. [34]). The error in using an adiabatic XC functional to propagate such a state, however, is significantly larger than the error the same functional produces when describing the propagation of an initial true ground state, as demonstrated by the model soft-Coulomb helium atom example. For two electrons, the spin-broken initial KS state, with the two orbitals evolved under different spin-decomposed potentials (depending on the instantaneous spin densities), often appeared to be the best choice, suggesting that orbital-specific potentials (allowed by the generalized KS scheme) could be a useful future avenue of research for TDDFT. However, how to obtain such potentials in a general  $N$ -electron case is not obvious (at least without the need for empirical parameters). Generally, orbital-dependent functionals treated within the OEP may be promising for these problems where memory (of the density) dependence, including ISD at the KS level only, is naturally captured by the KS orbitals; again, devising suitable orbital-dependent functionals with the appropriate ISD is a direction for future research. The example of coupled electron-nuclear dynamics in the model LiH molecule, although supporting the earlier results of the optimal choice of a KS initial state, above all, highlighted the need for more accurate ground-state functional approximations for dissociation.

We do not provide in this work an approximation that includes ISD, of both the true interacting state and the KS state; this is likely a difficult task. But what is clear from the results in this paper, and the stage of the field, is that now is the right time to address this issue.

## ACKNOWLEDGMENTS

Financial support from the National Science Foundation (Grant No. CHE-1152784) and a grant of computer time from the CUNY High Performance Computing Center under NSF Grants No. CNS-0855217 and No. CNS-0958379 are gratefully acknowledged. Thanks also to Johanna Fuks for comments on the manuscript.

- 
- [1] E. Runge and E. K. U. Gross, *Phys. Rev. Lett.* **52**, 997 (1984).
  - [2] *Fundamentals of Time-Dependent Density Functional Theory*, edited by M. A. L. Marques, N. T. Maitra, F. Nogueira, E. K. U. Gross, and A. Rubio, Lecture Notes in Physics, Vol. 837 (Springer-Verlag, Berlin, 2012).
  - [3] P. Hohenberg and W. Kohn, *Phys. Rev. B* **136**, 864 (1964).
  - [4] W. Kohn and L. J. Sham, *Phys. Rev.* **140**, A1133 (1965).
  - [5] W. Kohn, *Rev. Mod. Phys.* **71**, 1253 (1999).
  - [6] E. K. U. Gross and W. Kohn, *Phys. Rev. Lett.* **55**, 2850 (1985); **57**, 923(E) (1986).
  - [7] J. F. Dobson, *Phys. Rev. Lett.* **73**, 2244 (1994).
  - [8] G. Vignale and W. Kohn, *Phys. Rev. Lett.* **77**, 2037 (1996).
  - [9] J. F. Dobson, M. J. Büchner, and E. K. U. Gross, *Phys. Rev. Lett.* **79**, 1905 (1997).
  - [10] G. Vignale, C. A. Ullrich, and S. Conti, *Phys. Rev. Lett.* **79**, 4878 (1997).
  - [11] I. V. Tokatly and O. Pankratov, *Phys. Rev. B* **67**, 201103 (2003).
  - [12] Y. Kurzweil and R. Baer, *J. Chem. Phys.* **121**, 8731 (2004).
  - [13] H. O. Wijewardane and C. A. Ullrich, *Phys. Rev. Lett.* **100**, 056404 (2008).
  - [14] I. V. Tokatly, *Fundamentals of Time-Dependent Density Functional Theory* (Ref. [2]).
  - [15] N. T. Maitra, F. Zhang, R. J. Cave, and K. Burke, *J. Chem. Phys.* **120**, 5932 (2004).

- [16] M. Gatti, V. Olevano, L. Reining, and I. V. Tokatly, *Phys. Rev. Lett.* **99**, 057401 (2007).
- [17] E. Penka Fowe and A. D. Bandrauk, *Phys. Rev. A* **84**, 035402 (2011).
- [18] E. Tapavicza, I. Tavernelli, U. Rothlisberger, C. Filippi, and M. E. Casida, *J. Chem. Phys.* **129**, 124108 (2008).
- [19] J. Gavnholt, A. Rubio, T. Olsen, K. S. Thygesen, and J. Schiøtz, *Phys. Rev. B* **79**, 195405 (2009).
- [20] I. D'Amico and G. Vignale, *Phys. Rev. B* **59**, 7876 (1999).
- [21] P. Hessler, N. T. Maitra, and K. Burke, *J. Chem. Phys.* **117**, 72 (2002).
- [22] N. T. Maitra and K. Burke, *Chem. Phys. Lett.* **359**, 237 (2002).
- [23] C. A. Ullrich, *J. Chem. Phys.* **125**, 234108 (2006).
- [24] J. I. Fuks, N. Helbig, I. V. Tokatly, and A. Rubio, *Phys. Rev. B* **84**, 075107 (2011).
- [25] M. Thiele, E. K. U. Gross, and S. Kummel, *Phys. Rev. Lett.* **100**, 153004 (2008).
- [26] R. Baer, *J. Mol. Struct.: THEOCHEM* **914**, 19 (2009).
- [27] N. T. Maitra and K. Burke, *Phys. Rev. A* **63**, 042501 (2001); **64**, 039901(E) (2001).
- [28] N. T. Maitra, K. Burke, and C. Woodward, *Phys. Rev. Lett.* **89**, 023002 (2002).
- [29] R. van Leeuwen, *Phys. Rev. Lett.* **82**, 3863 (1999).
- [30] I. Dreissigacker and M. Lein, *Chem. Phys.* **391**, 143 (2011).
- [31] J. Javanainen, J. H. Eberly, and Q. Su, *Phys. Rev. A* **38**, 3430 (1988); W.-C. Liu, J. H. Eberly, S. L. Haan, and R. Grobe, *Phys. Rev. Lett.* **83**, 520 (1999); D. M. Villeneuve, M. Y. Ivanov, and P. B. Corkum, *Phys. Rev. A* **54**, 736 (1996); A. D. Bandrauk and N. H. Shon, *ibid.* **66**, 031401(R) (2002); M. Lein, E. K. U. Gross, and V. Engel, *Phys. Rev. Lett.* **85**, 4707 (2000).
- [32] A. Castro, H. Appel, M. Oliveira, C. A. Rozzi, X. Andrade, F. Lorenzen, M. A. L. Marques, E. K. U. Gross, and A. Rubio, *Phys. Status Solidi B* **243**, 2465 (2006).
- [33] R. Gaudoin and K. Burke, *Phys. Rev. Lett.* **93**, 173001 (2004).
- [34] I. Tavernelli, U. F. Röhrig, and U. Rothlisberger, *Mol. Phys.* **103**, 963 (2005).
- [35] K. Peirs, D. Van Neck, and M. Waroquier, *Phys. Rev. A* **67**, 012505 (2003).
- [36] A. Seidl, A. Görling, P. Vogl, J. A. Majewski, and M. Levy, *Phys. Rev. B* **53**, 3764 (1996).
- [37] S. Kümmel, *Fundamentals of Time-Dependent Density Functional Theory* (Ref. [2]).
- [38] A. Görling and M. Levy, *J. Chem. Phys.* **106**, 2675 (1997).
- [39] L. Kronik, T. Stein, S. Refaely-Abramson, and R. Baer, *J. Chem. Theory Comput.* **8**, 1515 (2012).
- [40] C. F. Craig, W. R. Duncan, and O. V. Prezhdo, *Phys. Rev. Lett.* **95**, 163001 (2005).
- [41] E. Tapavicza, I. Tavernelli, and U. Rothlisberger, *Phys. Rev. Lett.* **98**, 023001 (2007).
- [42] N. T. Maitra, *J. Chem. Phys.* **125**, 014110 (2006).
- [43] D. G. Tempel, T. J. Martínez, and N. T. Maitra, *J. Chem. Theory Comput.* **5**, 770 (2009).
- [44] M. E. Casida, F. Gutierrez, J. Guan, F.-X. Gadea, D. Salahub, and J.-P. Daudey, *J. Chem. Phys.* **113**, 7062 (2000).
- [45] J. I. Fuks, A. Rubio, and N. T. Maitra, *Phys. Rev. A* **83**, 042501 (2011).