

Formation of ultracold fermionic NaLi Feshbach molecules

Myoung-Sun Heo,¹ Tout T. Wang,^{1,2} Caleb A. Christensen,¹ Timur M. Rvachov,¹ Dylan A. Cotta,^{1,3} Jae-Hoon Choi,¹ Ye-Ryoung Lee,¹ and Wolfgang Ketterle¹

¹MIT-Harvard Center for Ultracold Atoms, Research Laboratory of Electronics, Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

²Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

³Département de Physique, École Normale Supérieure de Cachan, 94235 Cachan, France

(Received 23 May 2012; published 6 August 2012)

We describe the formation of fermionic NaLi Feshbach molecules from an ultracold mixture of bosonic ²³Na and fermionic ⁶Li. Precise magnetic field sweeps across a narrow Feshbach resonance at 745 G result in a molecule conversion fraction of 5% for our experimental densities and temperatures, corresponding to a molecule number of 5×10^4 . The observed molecular decay lifetime is 1.3 ms after removing free Li and Na atoms from the trap. Due to its extremely low reactivity, NaLi molecules in the ground state will have a much longer lifetime than KRb.

DOI: 10.1103/PhysRevA.86.021602

PACS number(s): 67.85.-d, 05.30.Fk, 34.50.-s

The preparation and control of ultracold atoms has led to major advances in precision measurements and many-body physics. One current frontier is to extend this to diatomic molecules. Early experiments focused on homonuclear molecules, where highlights included the study of fermion pairs across the BEC-BCS crossover [1]. The preparation of heteronuclear molecules is more challenging because it requires a controlled reaction between two distinct atomic species. However, heteronuclear molecules can have a strong electric dipole moment, which leads to a range of new scientific directions [2], including precision measurements, such as of the electron electric dipole moment [3], quantum computation mediated by dipolar coupling between molecular qubits [4] or in a hybrid system of molecules coupled to superconducting waveguides [5], many-body physics with anisotropic long-range interactions [6,7], and ultracold chemistry [8].

A number of experiments have explored molecule formation in ultracold atoms using photoassociation and Feshbach resonances [2]. Due to the lower abundance of fermionic alkali-metal isotopes, only one heteronuclear fermionic molecule ⁴⁰K⁸⁷Rb has been produced at ultracold temperatures [9]. Fermionic molecules are appealing due to Pauli suppression of *s*-wave collisions between identical fermions [10], as well as prospects for preparing fermions with long-range interactions as a model system for electrons with Coulomb interactions [6]. In this Rapid Communication, we report the formation of a fermionic heteronuclear molecule ²³Na⁶Li.

NaLi has at least three unique features due to its constituents being the two smallest alkali-metal atoms. First, its small reduced mass gives it a large rotational constant, which suppresses inelastic molecule-molecule collisions that occur via coupling between rotational levels [11]. Second, NaLi is reactive in its singlet $X^1\Sigma^+$ ground state, meaning that the reaction $\text{NaLi} + \text{NaLi} \rightarrow \text{Na}_2 + \text{Li}_2$ is energetically allowed [12], but with an unusually small predicted rate constant of 10^{-13} cm³/s that is by far the lowest among all reactive heteronuclear alkali-metal molecules [13,14] and should allow lifetimes >1 s even without dipolar suppression [15]. This is related to NaLi having the smallest van der Waals C_6 coefficient of all heteronuclear alkali-metal atom pairs [14], which results in weak scattering by the long-range potential.

Finally, this slow collision rate, together with weak spin-orbit coupling in diatomic molecules with small atomic numbers *Z* of its constituents [16], may allow a long-lived triplet $a^3\Sigma^+$ ground state in NaLi. This state has nonzero electric and magnetic dipole moments, opening up the possibility of exploring physics ranging from magnetic [17] and electric field control [18] of molecule-molecule collisions to realizing novel lattice spin Hamiltonians with coupling mediated by the electric dipole moment [19]. In addition to these three features, we note that NaLi has only a moderate dipole moment of 0.5 D [20] in its $X^1\Sigma^+$ ground state, comparable to KRb. Larger electric fields are required to align this dipole moment due to the large rotational constant $B = 0.4$ cm⁻¹ [21].

An earlier effort to produce NaLi Feshbach molecules was unsuccessful [22] due to incorrect assignments of interspecies Feshbach resonances between ²³Na and ⁶Li [23,24], which predicted many resonances to be much stronger than in the recently revised assignments [25]. For molecule formation the relevant figure of merit of a Feshbach resonance is its energetic width E_0 [1]. The more familiar width ΔB characterizes the visibility of scattering length modification relative to the background scattering length a_{bg} and is not directly relevant for molecule formation. E_0 and ΔB are related by $\Delta\mu\Delta B = \sqrt{E_0\hbar^2/ma_{\text{bg}}^2}$, where $\Delta\mu$ is the differential magnetic moment between the atomic and molecular states and m is the reduced mass of the two-particle system. Previously, NaLi resonances observed below 1000 G were assigned to $l = 0$ molecular bound states, where l is the rotational angular momentum of the molecule. The strongest of these resonances had a predicted width $E_0/h = 6$ kHz [24]. The revised interpretation [25] assigns these resonances to $l = 2$ molecular bound states coupled to the $l = 0$ open channel via magnetic dipole-dipole interactions, with $E_0/h = 5$ Hz for the strongest resonance, which is three orders of magnitude smaller.

This small E_0 results in a uniquely challenging situation for molecule formation. The Feshbach molecule has an open-channel character only for magnetic fields B around resonance B_0 such that $B - B_0 < E_0/\Delta\mu = 2$ μG [26]. This is impossible to resolve experimentally, and precludes molecule formation via radio-frequency (rf) association [27]

or modulation of the magnetic field at a frequency resonant with the molecular binding energy [28], both of which require a large open-channel contribution for good wave-function overlap between atomic and molecular states. This leaves magnetic field sweeps across resonance as the only feasible approach.

Such sweeps can be described as a Landau-Zener level crossing of the molecular state and the unbound pair of atoms. In this simplified picture the molecule conversion fraction $f_{\text{mol}} = 1 - e^{-2\pi\delta}$, with a Landau-Zener parameter δ given by [1,29]

$$\delta = \frac{(\hbar\Omega)^2}{\hbar\Delta\mu\dot{B}} = \frac{4\pi\hbar n}{2m} \frac{\Delta B a_{\text{bg}}}{\dot{B}} = \frac{1}{3\pi} \frac{(\tilde{E}_F/E_0)^{3/2}}{\hbar\Delta\mu\dot{B}/E_0^2}, \quad (1)$$

where $\Omega = \frac{1}{\hbar} \sqrt{\tilde{E}_F^{3/2} E_0^{1/2}} / 3\pi$ is the Rabi frequency for atom-molecule coupling, with an effective Fermi energy $\tilde{E}_F = (6\pi^2 n)^{2/3} \hbar^2 / m$ which expresses the atomic density n in energy units for bosons as well as for fermions. Ω corresponds to the width of the level crossing, and for efficient molecule formation $\delta > 1$, the magnetic field B must be varied across a region of width $\hbar\Omega/\Delta\mu$ around resonance in a time $> 1/\Omega$. In our experiment we use the strongest Feshbach resonance from the revised assignments [25], located at $B_0 = 745$ G, with $\Delta B = 10$ mG, $a_{\text{bg}} = -70a_0$, $\Delta\mu = 2\mu_B$, $E_0/\Delta\mu = 2$ μG as mentioned earlier, and $\hbar\Omega/\Delta\mu = 2$ mG for the experimental densities specified below. Working with such a narrow resonance at 745 G requires careful stabilization of large magnetic fields. Molecule formation experiments with other comparably narrow resonances have been done, but at much lower fields, for example, Cs_2 [30] and $^6\text{Li}^{40}\text{K}$ [31].

Experimentally we achieve < 10 mG root-mean-square (rms) magnetic field noise. The power supply produces current fluctuations corresponding to 200 mG of noise at 745 G. Active feedback stabilization of the current reduces the noise to < 10 mG rms after synchronizing experimental sequences with the 60-Hz line frequency.

We produce a near-degenerate mixture of ^{23}Na atoms in the $^2S_{1/2} |F, m_F\rangle = |2, 2\rangle$ state and ^6Li atoms in the $^2S_{1/2} |3/2, 3/2\rangle$ state in a cloverleaf magnetic trap, using an apparatus similar to the one described in Ref. [32]. The mixture is then transferred into a 5-W single-beam optical dipole trap at 1064 nm. In the optical trap we spin flip both species with simultaneous Landau-Zener rf sweeps at a small bias field of 15 G to the hyperfine ground states Na $|1, 1\rangle$ and Li $|1/2, 1/2\rangle$.

From low magnetic field we first jump far above the resonance to 752 G, and after waiting 200 ms the mixture is evaporatively cooled, leaving 1×10^6 each of Na and Li at a temperature of 1.2 μK . Here, the weak magnetic field curvature along the optical dipole trap provides additional axial confinement, giving radial and axial trap frequencies (ν_r, ν_z) of (920 Hz, 13 Hz) for Na, and (2.0 kHz, 26 Hz) for Li. The temperature of the mixture is near the onset of condensation for the bosonic Na, with $T/T_c = 1.2$, and $T/T_F = 0.3$, where for our trap parameters the condensation temperature for Na is $T_c = 1.0$ μK and the Fermi temperature for Li is $T_F = 4.1$ μK . The peak in-trap densities are $2.5 \times 10^{13} \text{ cm}^{-3}$ for Na and $1.4 \times 10^{13} \text{ cm}^{-3}$ for Li.




FIG. 1. Determination of Feshbach resonance location. The remaining atom number as a function of the end point of a downward magnetic field sweep across resonance is shown. Results are fitted to a hyperbolic tangent function (solid line) with center 745.38 G and width 40 mG. The normalized atom number along the vertical axis is a sum of Na and Li atom numbers, each normalized to 1.

The precise location of the 745-G resonance is determined by jumping down to 745.9 G, waiting 15 ms for the field to stabilize, and then sweeping the magnetic field down at a constant rate of 50 mG/ms but with a variable end point. The total atom number remaining after the sweep drops sharply as the end point crosses the Feshbach resonance (Fig. 1). This measurement determines the location of the Feshbach resonance to be 745.38 ± 0.04 G, providing a precise target field for molecule formation sweeps. Note the 40-mG width from Fig. 1 is not directly related to E_0 , Ω , or ΔB .

A typical molecule formation sequence is shown in Fig. 2. The start and end points of the magnetic field sweep are chosen to be as close to resonance as possible while still being outside the region of atom-molecule coupling defined by Ω . This maximizes the formation efficiency, because molecules form and begin to decay immediately after crossing the coupling region around resonance.




FIG. 2. Experimental sequence for molecule formation. After waiting for the magnetic field B to stabilize at 745.68 G, we sweep the field across resonance to $B = 745.19$ G in 800 μs , at a constant rate of 600 mG/ms. At the end of the sweep, simultaneous pulses of resonant light remove Na and Li atoms from the trap within 100 μs while leaving molecules unaffected. After a variable hold time for lifetime measurements, molecules are detected by jumping back above resonance and imaging dissociated atoms. The magnetic field can also be kept below resonance to confirm that bound molecules are invisible to the imaging light.




FIG. 3. Absorption imaging of molecular clouds using dissociated Na (a), (b) and Li atoms (c), (d). The lower images in each pair (b), (d) are taken after switching above resonance and dissociating molecules into free atoms. The upper images (a), (c) are reference images with the field below resonance, where free atoms have been removed from the trap and the molecules are invisible to the imaging light.

At the end of the sweep, laser pulses resonant with the 589-nm $^2S_{1/2}|-1/2, 3/2\rangle \rightarrow ^2P_{3/2}|-3/2, 3/2\rangle$ transition in Na and the 671-nm $^2S_{1/2}|-1/2, 1\rangle \rightarrow ^2P_{3/2}|-3/2, 1\rangle$ transition in Li remove $>90\%$ of the remaining free atoms from the trap (states written in high-field basis $|m_J, m_I\rangle$). For closed-channel Feshbach molecules, the excitation spectrum is sufficiently different from free atoms, so NaLi molecules are mostly unaffected if the laser pulses are not too long. After a variable hold time for molecule lifetime measurements, the field is switched above resonance, where bound molecules rapidly dissociate [33]. Imaging the free atoms gives us a measure of the molecule number, which for our optimized sweep parameters gives a formation fraction of 5%. This corresponds to a molecule number of 5×10^4 .

To confirm that the detected atomic signal is indeed from dissociated molecules, we check that imaging while keeping the magnetic field below resonance gives a negligible signal (Fig. 3). Below resonance, bound molecules have a vanishing absorption cross section σ and are invisible to the imaging light, while for free atoms σ is nearly identical to the one above resonance because the Zeeman shift produced by magnetic field changes of 500 mG is much smaller than the atomic linewidth. A second, independent confirmation of the molecular signal comes from releasing the trapped mixture in a magnetic field gradient, and demonstrating that the expanding atomic and molecular clouds separate from one another (Fig. 4)




FIG. 4. Separation of atoms and molecules in a magnetic field gradient. Absorption images of (a) free Li atoms, (b) separated clouds of atoms and dissociated molecules, and (c) molecules with free atoms removed, taken 3 ms after release from the crossed optical dipole trap in a 6-G/cm gradient.




FIG. 5. Lifetime of trapped NaLi molecules after removing the remaining free Na and Li atoms from the trap (solid squares). Fitting to an exponential gives a decay time constant of 1.3 ms. For comparison, the lifetime without removing the remaining free atoms from the trap is 270 μ s (open circles). The normalized molecule number along the vertical axis is a sum of Na and Li atom numbers, each normalized to 1 at zero hold time, detected from dissociating bound NaLi molecules.

because of their differential magnetic moment $\Delta\mu$. To observe this separation, we added additional optical confinement along the magnetic field gradient direction, using a second beam to form a crossed-beam optical dipole trap.

We measure the lifetime of trapped NaLi molecules (Fig. 5) by varying the hold time before switching the magnetic field above resonance for dissociation and imaging. With $>90\%$ of both species of the remaining atoms removed, the molecular lifetime is 1.3 ms. This lifetime appears to be limited by collisions with other molecules or leftover atoms rather than by photon scattering, since it does not increase significantly with a reduced intensity of the trapping laser. It can be enhanced by suppressing molecule-molecule collisions in a three-dimensional (3D) optical lattice [34] and fully removing residual free atoms from the trap. The molecular lifetime drops to 270 μ s when keeping free Na and Li atoms trapped with the molecules. Finally, if only Li atoms are removed before the hold time and not Na, the lifetime is 550 μ s. The presence of free atoms increases the molecular decay rate because of inelastic collisions with molecules. Our lifetime measurements show that Na and Li each give comparable contributions to this increased decay, which is consistent with the constituent atoms in the closed-channel NaLi molecule being distinguishable from free Na and Li atoms, meaning that quantum statistics does not play a role in collisions. In contrast, experiments with open-channel KRb molecules [35] showed a sharp dependence of lifetime on the quantum statistics of the atomic collisional partner. The lifetimes above can also be reported as two-body loss-rate constants $\beta_{\text{NaLi+Na}} \sim 1 \times 10^{-10}$ cm^3/s and $\beta_{\text{NaLi+Li}} \sim 4 \times 10^{-10}$ cm^3/s .

The Landau-Zener parameter in our experiment is estimated to be $\delta = 0.13$ by choosing n in Eq. (1) to be the larger of n_{Na} and n_{Li} . This corresponds to $f_{\text{mol}} = 56\%$. The discrepancy with the observed molecule fraction of 5% has several explanations. The simplified Landau-Zener picture above assumes a full phase-space overlap between the atoms involved in molecule formation, which is only true in a Bose-Einstein condensate at $T = 0$ [36,37]. In a Bose-Fermi mixture such as NaLi, the phase-space overlap is lower, reaching a maximum

around the boson condensation temperature T_c because at lower T the Na condensate begins to have less spatial overlap with the fermionic Li [27]. The phase-space overlap between Na and Li is further reduced because of their different trapping potentials [38]. Finally, the optimized sweep time of 800 μ s is much longer than the 270- μ s lifetime of NaLi molecules when trapped with free atoms, meaning that 80% of molecules formed are lost, assuming that molecule formation occurs at the midpoint of the sweep.

In summary, we have succeeded in forming fermionic $^{23}\text{Na}^6\text{Li}$ molecules around a narrow, closed-channel-dominated Feshbach resonance at 745 G. Optimized magnetic field sweeps across resonance result in a molecule conversion fraction of 5% for our experimental densities and temperatures, corresponding to a molecule number of 5×10^4 . The observed

molecular decay lifetime of 1.3 ms is enough to begin exploring routes for reaching the rovibrational ground state of the molecule by stimulated Raman transitions [9], and to explore the unique features of this smallest heteronuclear alkali-metal molecule.

This work was supported an AFOSR MURI on Ultracold Molecules, by the NSF and the ONR, and by ARO Grant No. W911NF-07-1-0493 with funds from the DARPA Optical Lattice Emulator program. T.T.W. acknowledges additional support from NSERC. We are grateful to Jook Walraven, Tobias Tiecke, Roman Krems, John Doyle, and Dave Pritchard for valuable discussions, and also Colin Kennedy and Gregory Lau for experimental assistance.

-
- [1] W. Ketterle and M. W. Zwierlein, in *Ultracold Fermi Gases*, Proceedings of the International School of Physics “Enrico Fermi,” Course CLXIV, Varenna, 2006, edited by M. Inguscio, W. Ketterle, and C. Salomon (IOS, Amsterdam, 2007).
- [2] *Cold Molecules: Theory, Experiment, Applications*, edited by R. V. Krems, W. C. Stwalley, and B. Friedrich (CRC, Boca Raton, FL, 2009).
- [3] E. A. Hinds, *Phys. Scr.*, **T 70**, 34 (1997).
- [4] D. DeMille, *Phys. Rev. Lett.* **88**, 067901 (2002).
- [5] A. André, D. DeMille, J. M. Doyle, M. D. Lukin, S. E. Maxwell, P. Rabl, R. J. Schoelkopf, and P. Zoller, *Nat. Phys.* **2**, 636 (2006).
- [6] M. A. Baranov, *Phys. Rep.* **464**, 71 (2008).
- [7] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, *Rep. Prog. Phys.* **72**, 126401 (2009).
- [8] R. V. Krems, *Phys. Chem. Chem. Phys.* **10**, 4079 (2008).
- [9] K.-K. Ni, S. Ospelkaus, M. H. G. de Miranda, A. Pe’er, B. Neyenhuis, J. J. Zirbel, S. Kotochigova, P. S. Julienne, D. S. Jin, and J. Ye, *Science* **322**, 231 (2008).
- [10] S. Ospelkaus, K.-K. Ni, D. Wang, M. H. G. de Miranda, B. Neyenhuis, G. Quémener, P. S. Julienne, J. L. Bohn, D. S. Jin, and J. Ye, *Science* **327**, 853 (2010).
- [11] R. V. Krems and A. Dalgarno, *J. Chem. Phys.* **120**, 2296 (2004).
- [12] P. S. Żuchowski and J. M. Hutson, *Phys. Rev. A* **81**, 060703 (2010).
- [13] P. S. Julienne, T. M. Hanna, and Z. Idziaszek, *Phys. Chem. Chem. Phys.* **13**, 19114 (2011).
- [14] G. Quémener, J. L. Bohn, A. Petrov, and S. Kotochigova, *Phys. Rev. A* **84**, 062703 (2011).
- [15] M. H. G. de Miranda, A. Chotia, B. Neyenhuis, D. Wang, G. Quémener, S. Ospelkaus, J. L. Bohn, J. Ye, and D. S. Jin, *Nat. Phys.* **7**, 502 (2011).
- [16] P. F. Bernath, *Spectra of Atoms and Molecules*, 2nd ed. (Oxford University Press, New York, 2005).
- [17] T. V. Tscherbul, Y. V. Suleimanov, V. Aquilanti, and R. V. Krems, *New J. Phys.* **11**, 055021 (2009).
- [18] T. V. Tscherbul and R. V. Krems, *Phys. Rev. Lett.* **97**, 083201 (2006).
- [19] A. Micheli, G. K. Brennen, and P. Zoller, *Nat. Phys.* **2**, 341 (2006).
- [20] M. Aymar and O. Dulieu, *J. Chem. Phys.* **122**, 204302 (2005).
- [21] M. M. Hessel, *Phys. Rev. Lett.* **26**, 215 (1971).
- [22] C. A. Christensen, Ph.D. thesis, Massachusetts Institute of Technology, 2011.
- [23] C. A. Stan, M. W. Zwierlein, C. H. Schunck, S. M. F. Raupach, and W. Ketterle, *Phys. Rev. Lett.* **93**, 143001 (2004).
- [24] M. Gacesa, P. Pellegrini, and R. Côté, *Phys. Rev. A* **78**, 010701 (2008).
- [25] T. Schuster, R. Scelle, A. Trautmann, S. Knoop, M. K. Oberthaler, M. M. Haverhals, M. R. Goosen, S. J. J. M. F. Kokkelmans, and E. Tiemann, *Phys. Rev. A* **85**, 042721 (2012).
- [26] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, *Rev. Mod. Phys.* **82**, 1225 (2010).
- [27] J. J. Zirbel, K.-K. Ni, S. Ospelkaus, T. L. Nicholson, M. L. Olsen, P. S. Julienne, C. E. Wieman, J. Ye, and D. S. Jin, *Phys. Rev. A* **78**, 013416 (2008).
- [28] S. T. Thompson, E. Hodby, and C. E. Wieman, *Phys. Rev. Lett.* **95**, 190404 (2005).
- [29] T. Köhler, K. Góral, and P. S. Julienne, *Rev. Mod. Phys.* **78**, 1311 (2006).
- [30] J. Herbig, T. Kraemer, M. Mark, T. Weber, C. Chin, H.-C. Nägerl, and R. Grimm, *Science* **301**, 1510 (2003).
- [31] F. M. Spiegelhalter, A. Trenkwalder, D. Naik, G. Kerner, E. Wille, G. Hendl, F. Schreck, and R. Grimm, *Phys. Rev. A* **81**, 043637 (2010).
- [32] Z. Hadzibabic, S. Gupta, C. A. Stan, C. H. Schunck, M. W. Zwierlein, K. Dieckmann, and W. Ketterle, *Phys. Rev. Lett.* **91**, 160401 (2003).
- [33] T. Mukaiyama, J. R. Abo-Shaeer, K. Xu, J. K. Chin, and W. Ketterle, *Phys. Rev. Lett.* **92**, 180402 (2004).
- [34] A. Chotia, B. Neyenhuis, S. A. Moses, B. Yan, J. P. Covey, M. Foss-Feig, A. M. Rey, D. S. Jin, and J. Ye, *Phys. Rev. Lett.* **108**, 080405 (2012).
- [35] J. J. Zirbel, K.-K. Ni, S. Ospelkaus, J. P. D’Incao, C. E. Wieman, J. Ye, and D. S. Jin, *Phys. Rev. Lett.* **100**, 143201 (2008).
- [36] E. Hodby, S. T. Thompson, C. A. Regal, M. Greiner, A. C. Wilson, D. S. Jin, E. A. Cornell, and C. E. Wieman, *Phys. Rev. Lett.* **94**, 120402 (2005).
- [37] J. E. Williams, N. Nygaard, and C. W. Clark, *New J. Phys.* **8**, 150 (2006).
- [38] F. Deuretzbacher, K. Plassmeier, D. Pfannkuche, F. Werner, C. Ospelkaus, S. Ospelkaus, K. Sengstock, and K. Bongs, *Phys. Rev. A* **77**, 032726 (2008).