

Photonic band gap in an imperfect atomic diamond lattice: Penetration depth and effects of finite size and vacancies

Mauro Antezza^{1,2} and Yvan Castin³

¹*Université Montpellier 2, Laboratoire Charles Coulomb UMR 5221–F-34095, Montpellier, France*

²*CNRS, Laboratoire Charles Coulomb UMR 5221–F-34095, Montpellier, France*

³*Laboratoire Kastler Brossel, ENS, UPMC and CNRS, 24 rue Lhomond, 75231 Paris Cedex 05, France*

(Received 26 April 2013; published 25 September 2013)

We study the effects of finite size and of vacancies on the photonic band gap recently predicted for an atomic diamond lattice. Close to a $J_g = 0 \rightarrow J_e = 1$ atomic transition, and for atomic lattices containing up to $N \approx 3 \times 10^4$ atoms, we show how the density of states can be affected by both the shape of the system and the possible presence of a fraction of unoccupied lattice sites. We numerically predict and theoretically explain the presence of shape-induced border states and of vacancy-induced localized states appearing in the gap. We also investigate the penetration depth of the electromagnetic field which we compare to the case of an infinite system.

DOI: [10.1103/PhysRevA.88.033844](https://doi.org/10.1103/PhysRevA.88.033844)

PACS number(s): 42.70.Qs, 42.50.Ct

I. INTRODUCTION

Wave propagation in periodic potentials constitutes a problem shared by several domains of classical and quantum physics, ranging from the study of electron motion in metals [1], to that of x- and γ -ray scattering by crystals [1–3], and of light by photonic crystals and metamaterials [4]. Periodicity leads to an organization of modes according to bands, and to the possible presence of band gaps, i.e., energy intervals where modes are absent. By periodicity here we really refer to an infinitely extended three-dimensional periodic system, not even to a system on a torus, hence it is strictly speaking not physical. Nonetheless, predictions made on the basis of infinite systems can become really satisfactory for systems large enough, as in solid-state physics, and present the advantage of benefiting from the Bloch theorem, and of being solved in the reciprocal space avoiding typical real-space oscillating functions. Models based on infinite systems may, however, present some subtleties related to the way in which the infinite limiting process is performed, often requiring ad hoc Ewald's summations type strategies.

The recent experimental realization of a Mott phase with ultracold-atomic gases [5,6], i.e., of artificial crystals made by single atoms trapped at the nodes of laser optical lattices, leads to the necessity of understanding the features of the band structure of light interacting with such systems. The peculiarity of this new system is that it presents several remarkable features: incident light scatters on pointlike elementary quantum objects with an internal energy level structure, and a quantum delocalized position in space [7], the lattice periodicity is of the order of the incident light wavelength, allowing the exploration of the entire Brillouin zone and hence of possible band gaps [8]; experiments reached a remarkable accuracy and control, permitting the realization of ultraprecise atomic clocks [9–11]; the long-range dipolar coupling among the scattering objects induces a strongly cooperative response that can be used to tailor the light amplitude on a subwavelength scale [12]. First attempts toward the description of such a system overlooked divergence problems, resulting in noncorrect prediction of band gaps [13,14], or were based on an ad hoc ultraviolet regularization procedure allowing us to explore only a particular class of lattice geometries not presenting any band gap [15].

Photonic band gaps of one-dimensional (1D) cold-atomic vapors have been realized [16] and exploited to generate optical parametric oscillation with distributed feedback [17]. Scattered photons have been suggested as a signature of the Mott insulator and superfluid quantum states [18] and studied in the framework of polaritons [19], excitons, and cavity polaritons [20]. Recently, by exploiting a microscopic theory of light-atom interaction [21] (see also [22–25]), and by explicitly introducing the presence of the unavoidable atomic quantum motion, it was possible to naturally regularize the divergences in a way independent of the lattice geometry, and at the same time to study the quantitative effects of the quantum atomic motion on the band structure [7]. The explicit dependence of the photonic band structure on quantum features, as the atomic internal energy levels and the external atomic quantum motion, allows us to consider this artificial structure as an example of quantum metamaterial [26]. In the framework of the Fano-Hopfield self-consistent quadratic theory [3,19,27], it was also possible to find an exact solution valid for the full Brillouin zone and for arbitrary Bravais and non-Bravais lattices, allowing the prediction of the diamond as the first three-dimensional (3D) atomic lattice geometry presenting a complete photonic band gap [8]. Although an optical diamond lattice was to our knowledge not obtained yet in the laboratory, the technique to realize it, elaborating on the ideas of [28], is perfectly known and consists in superimposing four plane-wave laser beams of well-chosen directions and polarizations [8,29]. Further investigations suggested to add external magnetic fields to open band gaps in other geometric structures [30].

In cold-atom realizations of 3D optical lattices, the atomic Mott state extends over 10–20 lattice sites in diameter, so a natural question regards the features of the band gap in this finite-size system. A further question concerns the effects of an imperfect finite portion of a lattice containing site defects, i.e., a fraction of vacancies resulting in a not complete filling of the lattice.¹ The experimental interest of these issues is related to the fact that both the finite-size and vacancy effects,

¹The $n = 1$ ground-state Mott phase for a nonzero tunneling-to-interaction ratio t/U has indeed vacant and multiply occupied sites.

separately, could in principle drastically affect the presence and the experimental visibility of the band gap due to the appearance of states in the gap. The main questions we address in this paper are as follows: What does happen to the band gap for systems of realistic sizes and of different shapes? What is the fraction of vacancies which still permits us to have a reasonable band-gap visibility? What is the value of the penetration depth of an electromagnetic wave in the atomic diamond lattice for finite and infinite systems, i.e., how is it affected by finite-size effects? Even if we discuss in detail the case of a diamond lattice, we will present a general formulation and will discuss the main features which will remain valid for other lattice geometries with a photonic band gap (which excludes the simple cubic lattice).

The paper is organized as follows. In Sec. II, we illustrate the model we use, and the resulting main equations we solve. In Sec. III, we present and discuss a numerical study on the density of states and on the penetration depth in a finite-size system, possibly in presence of imperfections due to vacant sites in the lattice. In Sec. IV, we provide an analytical analysis to support and illustrate some of the main features of the numerical findings. We conclude in Sec. V.

II. MODEL

We consider a system made by a collection of N identical atoms having fixed positions and an optical dipolar transition between a $J_g = 0$ electronic ground state and a $J_e = 1$ electronic excited state.² Such a transition is available in appropriate atomic species, such as strontium [32]. Having a spinless electronic ground state of course considerably simplifies the physical description, as the atomic density operator has no off-diagonal matrix elements in the ground state and there can not be ground-state-changing photon scattering, that is, optical-pumping processes [33]; in the low-intensity regime, the scattering of a photon by a ground-state atom is then purely elastic, which is a crucial advantage to study coherent multiple scattering of light in an atomic ensemble as experimentally done in [32], and only the optical coherences of the density operator, that give the mean atomic dipoles, are relevant in the theoretical description.

In our model, which applies in the regime of low-atomic excitation, the atomic dipoles are thus the only dynamical variables, and they are coupled by the electromagnetic field they radiate. The resulting eigenmodes of the mean atomic dipoles are given by the solutions of the eigenvalue

We may here treat the multiply occupied sites as effectively vacant sites, either because the resonance frequency of two or more atoms confined in the Lamb-Dicke regime differs from the single atom one ω_0 by much more than the natural width Γ , or because there are techniques to selectively empty the multiply occupied sites [31].

²It is worth stressing that the often-used scalar model for light has the evident advantage of drastically reducing the numerical effort, but also the disadvantage of providing a qualitatively and quantitatively wrong description of the physical system. For instance, it is possible to show that already for a simple cubic atomic lattice, the scalar model, in contradiction with the vectorial one, predicts the presence of a band gap.

problem [13,21]

$$\begin{aligned} \left(\hbar\omega_0 - i\frac{\hbar\Gamma}{2}\right)d_{i,\alpha} + \sum_{\substack{j=1 \\ j \neq i}}^N \sum_{\beta=x,y,z} g_{\alpha\beta}(\mathbf{r}_i - \mathbf{r}_j)d_{j,\beta} \\ = \hbar(\omega - i\gamma)d_{i,\alpha}. \end{aligned} \quad (1)$$

Here, $d_{i,\alpha}$ is the component along the direction $\alpha = x, y, z$ of the mean dipole carried by the atom i , $\omega - i\gamma$ is the mode eigenfrequency (it is complex in general with $\gamma > 0$, and may be measured as suggested in [7]), ω_0 and Γ are the single-atom resonance frequency and spontaneous emission rate. The tensor $g_{\alpha\beta}(\mathbf{r})$ gives the α component of the electric field at the position \mathbf{r} radiated by a dipole oscillating along the direction β at the origin of coordinates $E_\alpha(\mathbf{r}) = -g_{\alpha\beta}(\mathbf{r})d_\beta/d^2$, d being the atomic dipole moment such that $\Gamma = d^2\omega_0^3/(3\pi\epsilon_0\hbar c^3)$. Here, we consider the case where $\omega - i\gamma$ is very close to ω_0 , so that $g_{\alpha\beta}$ can be evaluated for a dipole oscillating at the resonance frequency; introducing the vacuum wave number

$$k_0 = \frac{\omega_0}{c}, \quad (2)$$

we take

$$\begin{aligned} g_{\alpha\beta}(\mathbf{r}) &= -\frac{3\hbar\Gamma}{4k_0^3} \left[(k_0^2\delta_{\alpha\beta} + \partial_{r_\alpha}\partial_{r_\beta}) \frac{e^{ik_0r}}{r} + 4\pi\delta_{\alpha\beta}\delta(\mathbf{r}) \right] \\ &\stackrel{r>0}{=} \frac{3}{4}\hbar\Gamma \frac{e^{ik_0r}}{k_0r} \left[\left(-1 - \frac{i}{k_0r} + \frac{1}{(k_0r)^2} \right) \delta_{\alpha\beta} \right. \\ &\quad \left. + \left(1 + \frac{3i}{k_0r} - \frac{3}{(k_0r)^2} \right) \frac{r_\alpha r_\beta}{r^2} \right]. \end{aligned} \quad (3)$$

Our first expression in (3) for $g_{\alpha\beta}(\mathbf{r})$ differs by a scalar $\delta(\mathbf{r})$ contribution from the usual expression for the electric field radiated by a dipole [see Eqs. (4.20) and (9.18) of [34]]; this ensures compatibility with our previous works and it is of course irrelevant here since atoms are never at the same position.³ The first expression is particularly useful to directly extract its Fourier transform, needed in the Bloch description of infinite systems (see Sec. IV), while the second one [which differs from the first one by another scalar $\delta(\mathbf{r})$ contribution] has the well-known dipole-dipole interaction form, and will be used in numerical calculations on finite-size systems in Sec. III.

Equation (1) allows one to determine the density of states of the system. In case an infinite number of atoms are periodically arranged at the nodes of a diamond lattice, it has been shown that the system may exhibit an omnidirectional photonic band gap [8]. Here, by numerical solution of (1), we investigate the fate of such a gap, in situations close to realistic experimental ones, where the number of atoms is finite and/or there are vacancies in the lattice. A further interesting quantity related to the occurrence of a gap is the so-called ‘‘penetration depth’’ ξ : an incident electromagnetic wave at a frequency in the band gap can not penetrate the medium, and its amplitude will decay exponentially over a characteristic distance ξ . In order to calculate such a length, we consider a pointlike dipolar source immersed in the atomic medium, and we extract ξ from the total field and the induced dipole spatial profiles: we fix at the

³Our convention amounts to omitting the $\delta(\mathbf{r})$ term in Eq. (3) of [21].

position \mathbf{r}_s a forced dipole $d_\alpha^s = \check{d}_\alpha^s e^{-i\omega_s t}$; the atomic dipoles at the positions \mathbf{r}_i will reach a steady state $d_{i,\alpha} = \check{d}_{i,\alpha} e^{-i\omega_s t}$ given by the linear system

$$-\left[\hbar(\omega_s - \omega_0) + i\frac{\hbar\Gamma}{2}\right]\check{d}_{i,\alpha} + \sum_{\substack{j=1 \\ j \neq i}}^N \sum_{\beta=x,y,z} g_{\alpha\beta}(\mathbf{r}_i - \mathbf{r}_j)\check{d}_{j,\beta} = -\sum_{\beta=x,y,z} g_{\alpha\beta}(\mathbf{r}_i - \mathbf{r}_s)\check{d}_\beta^s. \quad (4)$$

III. NUMERICAL RESULTS FOR A FINITE-SIZE SYSTEM

In this section, we study a system of N atoms at the nodes of a diamond lattice. We recall that the diamond lattice is formed by the superposition of two copies of the same Bravais lattice: the fcc lattice of lattice constant a , generated by the three basis vectors

$$\mathbf{e}_1 = (0, a/2, a/2), \quad \mathbf{e}_2 = (a/2, 0, a/2), \quad \mathbf{e}_3 = (a/2, a/2, 0), \quad (5)$$

and a second fcc lattice obtained by translating the first lattice by the vector $(a/4, a/4, a/4)$. The corresponding basis of the reciprocal lattice is

$$\tilde{\mathbf{e}}_1 = (-2\pi/a, 2\pi/a, 2\pi/a), \quad \tilde{\mathbf{e}}_2 = (2\pi/a, -2\pi/a, 2\pi/a), \quad \tilde{\mathbf{e}}_3 = (2\pi/a, 2\pi/a, -2\pi/a). \quad (6)$$

In our simulations, the atoms occupy a finite region in space, which can be a ball or a cube centered at the origin of the coordinates, each geometry corresponding to the usually used harmonic traps [5,6] or to the recently realized flat-bottom cuboid traps [35]. From the numerical solution of (1) we extract the density of states for the case of a unit filling factor (Sec. III A) and for the case with a low concentration of vacancies (Sec. III B). Finally, we analyze the penetration depth in Sec. III C solving (4).

A. Finite-size effects on the density of states

In this section, we discuss the density of states obtained by solving Eq. (1) for a finite-size diamond lattice, in the absence of vacancies. In particular, in Fig. 1 we show the density of states $\rho(\omega)$ for a number of atoms corresponding to typical experimental realizations $N \approx 2.5 \times 10^4$. Here, $\rho(\omega)$ is defined as the distribution of the real part of the complex spectrum of Eq. (1), normalized as $\int \rho(\omega)d\omega = 6/\mathcal{V}_L$, where $\mathcal{V}_L = a^3/4$ is the volume of the direct lattice primitive cell, in order to facilitate the comparison with the infinite-system results of [8], plotted as a bar histogram in the figure. If the atoms occupy a ball (see the black solid line), we observe partial filling of the spectral gap, most pronounced in the upper region. On the contrary, the region close to the lower border of the gap remains relatively weakly affected by the finite size of the system, considering the sharp rise of $\rho(\omega)$ to the left of this border. The remaining part of the density of states remains very close to the one of the infinite system. If the atoms occupy a cube (see the red solid line), the finite-size effects are quite different. Two peaks appear: a very pronounced one in the middle of the gap [at $(\omega - \omega_0)/\Gamma \approx -3.2$], and a second one




FIG. 1. (Color online) Density of the real part of the eigenfrequencies $\rho(\omega)$ obtained from Eq. (1), in the absence of vacancies and for $k_0 a = 2$ where a is the fcc lattice constant. Red solid lines: finite system with a cubic shape of side length $14a$, and $N = 2.7 \times 10^4$. Black solid lines: finite system with a spherical shape of diameter $18a$, and $N = 2.4 \times 10^4$. The histogram provides the same quantity for an infinite system [8]. Each of the three curves is composed of 250 bins. \mathcal{V}_L is the volume of the direct lattice primitive cell. The inset is a magnification.

at $(\omega - \omega_0)/\Gamma \approx 0.5$. We investigated the nature of the states belonging to the peak in the gap, by looking at 10 successive eigenstates of (1), finding that they are “border states”: they reach their maxima in a spherical shell of radius $\approx 5a$, and decay exponentially towards the center of the cube with a law

$$|d_i|^2 \equiv \sum_{\alpha=x,y,z} |d_{i,\alpha}|^2 \lesssim e^{-22+4.6r_i/a}, \quad (7)$$

where the dipole eigenvectors are normalized to the maximum value of their modulus equal to unity. This suggests a value of the penetration depth of the order of $0.5a$, in agreement with the calculation done in Sec. III C.

In Fig. 2, we show the distribution of the eigenvalues of Eq. (1) in the complex plane, restricted to small values of




FIG. 2. (Color online) Complex eigenvalues $\omega - i\gamma$ obtained from Eq. (1). The system is of finite size, in the absence of vacancies, for $k_0 a = 2$, with a spherical shape of diameter $18a$, and $N = 2.4 \times 10^4$ atoms. The two vertical red dashed lines give the borders of the band gap of the infinite periodic system.

γ/Γ . In this region, the figure shows that the band gap is not filled, apart from two narrow intervals of values of ω , in the center and close to the upper border of the gap. Then, in the finite-size system, the partial filling of the gap is mostly due to eigenvalues with larger values of γ/Γ . The smallest values of γ/Γ we obtained are $\approx 2 \times 10^{-5}$. The real part of the corresponding eigenvalues are located on the borders of the band gap for the infinite system, marked in the figure by vertical dashed lines, and on the upper bound of the values of ω represented in Fig. 2 for the finite system, i.e., around $(\omega - \omega_0)/\Gamma = 9.5$.

B. Effects of vacancies on the density of states

In this section, we address the case where the finite-size diamond lattice is not perfectly filled, presenting a concentration $1 - p$ of defects made by the presence of a random uniform distribution of nonoccupied lattice sites. For each value of the parameters and of the system geometry, the numerical results are obtained for a *single* realization of the disorder, randomly chosen by the computer. The fact that such a random realization is *typical* was checked by repeating the simulations with other random arrangements, which produced similar results (not shown); this could also have been expected from the fact that the densities of states reported here have a smooth variation with the concentration $1 - p$ of vacancies, as the reader shall see. From an experimental point of view, this is an interesting piece of information, indicating that the relevant physics (such as peaks in the density of states) can be accessed without averaging over the disorder.

In Fig. 3, we show the density of states $\rho(\omega)$ obtained solving Eq. (1) for atoms occupying a ball, as a function of the lattice filling factor p . The figure, and its inset, show that already a small concentration of vacancies equal to $1 - p = 0.01$ (red solid line) produces a remarkable signature in the density of states manifested by the appearance of a pronounced peak in the middle of the band gap, at $(\omega - \omega_0)/\Gamma \approx -3.08$. We explain the nature of the peak with the presence of single-vacancy states localized at the vacancy position. Since the vacancy concentration is small, most frequent vacancy states have a single-site nature. In Sec. IV C, we theoretically calculate the value of the single-vacancy state frequency, signaled in the inset by a black vertical dotted line, which seems to coincide quite satisfactorily with that of the numerically observed peak. By increasing the vacancy concentration, Fig. 3 shows for $1 - p = 0.05$ the occurrence of a clear second peak in the gap, which seems to match quite well the frequency of a two-vacancy in-gap state calculated in Sec. IV C [see the red vertical dotted line at $(\omega - \omega_0)/\Gamma \simeq -4$]. Peaks corresponding to other two-vacancy states predicted in Sec. IV C are less visible (see the other vertical dotted lines in the inset of Fig. 3). Further increase of the concentration of vacancies produces a gradual filling of the band gap, whose visibility completely deteriorates for a vacancy concentration around $1 - p = 0.2$.

In Fig. 4, for exactly the same spherical system with a vacancy concentration of $1 - p = 0.2$, we show the distribution of the eigenvalues of Eq. (1) in the complex plane, restricted to small values of γ/Γ . The figure shows that the band gap is completely filled. The states filling the gap, for such a large




FIG. 3. (Color online) Density of the real part of the eigenfrequencies $\rho(\omega)$ obtained from Eq. (1), for different concentrations of vacancies, that is, for various filling factors p , and for $k_0a = 2$. The finite system has a spherical shape of diameter $18a$, and $N = 2.4 \times 10^4$ for $p = 1$. The histogram provides the same quantity for an infinite system with no vacancies [8]. V_L is the volume of the direct lattice primitive cell. The inset is a magnification, where the vertical dotted lines correspond to frequencies of the single-vacancy in-gap state (black, central) and to two-vacancy in-gap states ($\check{R}_2 - \check{R}_1 = \mathbf{e}_1$, $\check{\mu}_1 = \check{\mu}_2 = 1$ in red, outer; $\check{R}_2 - \check{R}_1 = a\mathbf{e}_x$, $\check{\mu}_1 = 2, \check{\mu}_2 = 1$ in blue, inner; these quantities are defined in Appendix B) theoretically predicted in Sec. IV C. Decreasing values of p correspond to increasing values of $\rho(\omega)$ in the band gap (see the color code and, in the inset, the values of p linked with the curves by arrows).

vacancy concentration, are completely delocalized over the entire system size, and have a spectral imaginary part mostly concentrated around $\gamma/\Gamma \approx 10^{-2}$, with $\gamma/\Gamma \geq 10^{-3}$.

In Fig. 5, we study the effect of vacancies on a system of cubic shape. The figure shows that for a concentration of vacancies $1 - p = 0.01$ (red solid line), two peaks are present in the band gap. They have a different origin: the first one, that at smallest frequency, is nothing but the peak related to shape-induced states, already present in the absence of vacancies (see




FIG. 4. (Color online) Complex eigenvalues $\omega - i\gamma$ obtained from Eq. (1). The system is of finite size, in presence of vacancies, that is with a filling factor $p = 0.8$, for $k_0a = 2$, with a spherical shape of diameter $18a$, and $N = 1.9 \times 10^4$ atoms. The two vertical red dashed lines give the borders of the band gap of the infinite periodic system.




FIG. 5. (Color online) Density of the real part of the eigenfrequencies $\rho(\omega)$ obtained from Eq. (1), for two concentrations of vacancies, that is, for the filling factors $p = 1$ (black) and $p = 0.99$ (red), and for $k_0a = 2$. The finite system has a cubic shape of side $14a$, and $N = 2.7 \times 10^4$ for $p = 1$. The histogram provides the same quantity for an infinite system with no vacancies [8]. \mathcal{V}_L is the volume of the direct lattice primitive cell. We note the double-peak structure for $p = 0.99$ (see text). The vertical dotted line corresponds to the frequency of the single-vacancy in-gap state theoretically predicted in Sec. IV C.

black solid line, and the discussion in Sec. III A). The second peak is instead the signature of single-vacancy localized states, and its central frequency is the same as that shown in Fig. 3 for a spherical shape at the same vacancy concentration.

C. Penetration depth

To numerically calculate the penetration depth ξ for a diamond finite-size atomic lattice, we numerically solve the forced dipole equation (4) for a pointlike dipolar oscillating source $\check{d}_\alpha^s = \check{d}_\alpha^s e^{-i\omega_s t}$ at the position \mathbf{r}_s (at the center of the system within a distance a , where a is the lattice constant), and with ω_s in the band gap. Solutions of Eq. (4) provide the induced atomic dipoles amplitudes $\check{d}_{i,\alpha}$ at the lattice positions \mathbf{r}_i . We point out that this geometry is quite favorable to extract the penetration depth ξ for *small* systems, whereas the usual geometry, that sends an electromagnetic plane wave on the medium, may lead here to complicated border effects, with light propagating around our spherical medium, making the precise extraction of ξ difficult, even numerically. Experimentally, the source inside the medium may be realized by driving one or a few atomic dipoles close to the center of the medium, with the two-photon transition already considered in [7]; the two focused laser beams, whose joint action shall excite the two-photon transition, spatially overlap at the center of the medium only, and individually have a frequency far from single-photon resonance frequencies so that they freely propagate in the medium.

An interesting question is to know if the penetration depth ξ is the same for the textbook geometry used to define the penetration depth (an infinite half-space medium excited by a plane wave) and for our geometry (a finite spherical medium with a source at its center). The analytical result of Sec. IV B for ξ [see Eqs. (33), (35), and (38)] strongly suggests that this is the case since it only involves the dispersion relation for light




FIG. 6. (Color online) Penetration depth ξ in (a) and its inverse κ in (b), as functions of the dipole source frequency ω_s . Symbols (the lines are a guide to the eye) correspond to the numerical solution of Eq. (4) for a finite system of spherical shape, diameter $18a$, filling factor $p = 1$, $k_0a = 2$, and containing $N = 2.4 \times 10^4$ atoms. Red squares and black circles correspond to values obtained using the methods of Eqs. (8) and (10), respectively. The vertical dashed lines correspond to the borders (13) of the band gap for the infinite periodic system.

in the bulk. Mathematically, one may try to obtain an explicit proof of that property by extending the analytics of Sec. IV B to a medium filling a half-space; this is, however, beyond the scope of this work and it is reserved for a future work.

From our numerical simulations, we extract ξ according to different methods. The first method is based on the direct analysis of the induced dipoles, and consists in averaging the norm $\sqrt{\sum_\alpha |\check{d}_{i,\alpha}|^2}$ on spherical shells of radius $\approx u = \|\mathbf{r} - \mathbf{r}_s\|$ centered at the source position. We then obtain an average real dipole function $\mathcal{D}(u)$ that we fit in a certain range of u [where the behavior of $\mathcal{D}(u)$ is clearly exponential over several decades] as

$$\mathcal{D}(u) = C \frac{e^{-u/\xi}}{u}, \quad (8)$$

where ξ and C are the two fitting parameters. The factor $1/u$ in (8) is introduced to take into account the direct effect of the source which is dominant at small distances, allowing us to fit the function on a larger range. This method provides the results presented by red squares in Fig. 6. Its specialization to the analysis of the penetration depth along some given direction (without averaging over spherical shells) is straightforward, and leads to the filled diamonds and circles in Figs. 7(a) and 7(b), respectively.

The second method is based on the calculation of the total electric field amplitude generated by the source and induced




FIG. 7. Inverse penetration depth $\kappa = 1/\xi$ along direction \mathbf{e}_1 in (a) and direction \mathbf{e}_x in (b), as a function of the source frequency ω_s expressed through a change of variable mapping the band gap $[\omega_{\text{inf}}, \omega_{\text{sup}}]$ onto \mathbb{R}^+ . Same physical parameters as in Fig. 6. Filled diamonds in (a) and filled circles in (b): finite-size system with first extraction method; empty circles in (b): finite-size system with second extraction method; for those data, ω_{inf} and ω_{sup} were obtained by linear extrapolation of κ^2 to zero. Stars: for the infinite medium from a numerical evaluation of Eq. (30). Dashed lines: analytical predictions, close to the band borders, deduced from Eq. (42) (see text). Note that the x and y axes are in \log_{10} and \log_2 scale.

dipoles obtained by (4):

$$\check{E}_\alpha(\mathbf{r}) = - \sum_{\beta} g_{\alpha\beta}(\mathbf{r} - \mathbf{r}_s) \frac{\check{d}_\beta^s}{d^2} - \sum_{i=1}^N \sum_{\beta} g_{\alpha\beta}(\mathbf{r} - \mathbf{r}_i) \frac{\check{d}_{i,\beta}}{d^2}. \quad (9)$$

We evaluate $\check{E}_\alpha(\mathbf{r})$ on three lines, parallel to the Cartesian axes and passing through the source position \mathbf{r}_s . We first average the norm $\sqrt{\sum_{\alpha} |\check{E}_\alpha(\mathbf{r})|^2}$ on the two directions (\pm) of the three axes α , then we obtain and fit the six corresponding average real electric functions $\mathcal{E}_\alpha^{(\pm)}(u)$ as

$$\mathcal{E}_\alpha^{(\pm)}(u) = K_\alpha^{(\pm)} \frac{e^{-u/\xi_\alpha^{(\pm)}}}{u}, \quad (10)$$

obtaining six values of $\xi_\alpha^{(\pm)}$, whose average is presented by empty black circles in Figs. 6 and 7(b).

In Fig. 6, it is apparent that the extractions of the penetration depth from Eqs. (8) and (10) give different values. This shows that ξ is not isotropic; it depends on the considered direction of space, a property that will be recovered analytically in Sec. IV B. Whereas use of Eq. (10) is expected to give the penetration depth along the x axis, the first method, when it involves a directional average as in Eq. (8), is expected to pull out the maximal penetration depth (maximized over the directions of space). A second property, apparent

in Fig. 6(a), is the divergence of ξ at the borders of the infinite-medium forbidden gap (represented by vertical dashed lines at frequencies $\omega_{\text{inf}}, \omega_{\text{sup}}$). Figure 6(b) even suggests that κ vanishes there with a vertical slope. We indeed find that κ^2 vanishes linearly with ω_s (not shown), as also predicted analytically in Sec. IV B. By a linear extrapolation of κ^2 as a function of ω_s , we get for the borders of the forbidden bands

$$\left(\frac{\omega_{\text{inf}} - \omega_0}{\Gamma}, \frac{\omega_{\text{sup}} - \omega_0}{\Gamma} \right) \stackrel{\text{Eq. (10)}}{\simeq} (-4.748, -1.962) \quad (11)$$

$$\stackrel{\text{Eq. (8)}}{\simeq} (-4.747, -1.948), \quad (12)$$

which are indeed quite close to the infinite-medium results [8]

$$\left(\frac{\omega_{\text{inf}} - \omega_0}{\Gamma}, \frac{\omega_{\text{sup}} - \omega_0}{\Gamma} \right) \simeq (-4.743, -1.962). \quad (13)$$

To better put in evidence the vanishing of κ at the band edges, and to more easily compare the various methods, we show κ as a function of $(\omega_s - \omega_{\text{inf}})/(\omega_{\text{sup}} - \omega_s)$ in Fig. 7, with the band edges ω_{inf} and ω_{sup} deduced for the finite-size simulations by linear extrapolation of κ^2 .⁴ This change of variable on ω_s has the advantage of mapping the band edges to 0 and $+\infty$, respectively, which is then combined with a log-scale representation on both figure axes. This figure was produced for two particular directions of penetration, along the direct lattice basis vector \mathbf{e}_1 in Fig. 7(a), and along the Cartesian axis direction \mathbf{e}_x in Fig. 7(b). First, in Fig. 7(b), it appears that the two extraction methods for the penetration depth in the finite-size system (the first method from the dipoles, see the filled circles; the second method from the electric field, see the empty circles) give compatible results if they are applied along the *same* direction (here \mathbf{e}_x , which is equivalent to \mathbf{e}_y or \mathbf{e}_z due to symmetry of the diamond lattice). Second, in Figs. 7(a) and 7(b), the results of the finite-size systems are compatible with the ones (stars) for the infinite system in Sec. IV B, and even if they do not cover a as large range for κ , they do nicely follow the analytical prediction (dashed lines) for the vanishing of κ close to the band edges.

IV. THEORY FOR THE INFINITE SYSTEM

We show in this section that several features of the numerical simulations, such as the sharp rise of the penetration depth ξ close to the band-gap borders, and some peaks induced by vacancies in the density $\rho(\omega)$ of the real part of the eigenfrequencies, can be interpreted analytically for an infinite system. In this case, a reformulation of Eqs. (1) and (4) in Fourier space is more appropriate. It is known, however, that the resulting series over the reciprocal lattice presents subtle convergence issues [15] that were overlooked in [13, 14]. These issues were solved in [8] by coupling each atomic dipole to a spatially smoothed version $\check{\mathbf{E}}_\perp(\mathbf{r}) = \int d^3u \mathbf{E}_\perp(\mathbf{r} - \mathbf{u}) \chi(\mathbf{u})$ of the transverse electromagnetic field operator $\mathbf{E}_\perp(\mathbf{r})$, where the smoothing function $\chi(\mathbf{u})$ may be taken as a positive

⁴Note that, according to Eqs. (19) and (20), this rational fraction of the source frequency is the same for the original model and the Gaussian spatially smoothed model $(\bar{\omega}_s - \bar{\omega}_{\text{inf}})/(\bar{\omega}_{\text{sup}} - \bar{\omega}_s) \simeq (\omega_s - \omega_{\text{inf}})/(\omega_{\text{sup}} - \omega_s)$, within an exponentially small error in $1/b^2$.

rotationally invariant function of unit integral and of small width b . This cuts off the dipolar coupling at high-wave-number field modes and regularizes the theory for the infinite system.

One then finds that two changes have to be applied to Eqs. (1) and (4). First, the function $g_{\alpha\beta}(\mathbf{r})$ has to be replaced by the smoothed function $\bar{g}_{\alpha\beta}(\mathbf{r})$ such that

$$\bar{g}_{\alpha\beta}(\mathbf{r}_i - \mathbf{r}_j) = \int d^3 u_i \int d^3 u_j g_{\alpha\beta}(\mathbf{r}_i + \mathbf{u}_i - \mathbf{r}_j - \mathbf{u}_j) \chi(\mathbf{u}_i) \chi(\mathbf{u}_j). \quad (14)$$

In Fourier space, the convolution products take a simple form so that

$$\bar{\tilde{g}}_{\alpha\beta}(\mathbf{k}) = \frac{3\pi\hbar\Gamma}{k_0^3} \frac{k^2\delta_{\alpha\beta} - k_\alpha k_\beta}{k^2 - k^2 + i0^+} \tilde{\chi}^2(\mathbf{k}), \quad (15)$$

where $\tilde{\tilde{g}}_{\alpha\beta}(\mathbf{k}) = \int d^3 r e^{-i\mathbf{k}\cdot\mathbf{r}} \bar{g}_{\alpha\beta}(\mathbf{r})$ is the Fourier transform of $\bar{g}_{\alpha\beta}$ and $\tilde{\chi}(\mathbf{k})$ is the one of $\chi(\mathbf{r})$. Second, the spontaneous emission rate Γ in Eqs. (1) and (4) has to be replaced by

$$\bar{\Gamma} = \Gamma \tilde{\chi}^2(\mathbf{k}_0), \quad (16)$$

where \mathbf{k}_0 is a vector of modulus equal to k_0 and of arbitrary direction. If one would treat the atomic motion quantum mechanically, as in [7], for atoms trapped at the nodes of an optical lattice, $\chi(\mathbf{u}) = \phi^2(\mathbf{u})$ would be the probability distribution of the fluctuations \mathbf{u} of the atomic position around a node \mathbf{r}_i , where ϕ is the underlying atomic center-of-mass wave function. Then, Eq. (14) would have a straightforward physical interpretation. Also, $\bar{\Gamma}$ would simply be the elastic spontaneous emission rate, where the atomic center-of-mass after decay to the electronic ground state remained in the wave function ϕ . In practice, a Gaussian choice for χ is convenient, which corresponds to

$$\tilde{\chi}(\mathbf{k}) = e^{-k^2 b^2/2}. \quad (17)$$

It is useful to know to which extent the results from the spatially smoothed model differ from the original model. For the Gaussian smoothing function, one then has the remarkable result that, when the width b is much smaller than all interatomic distances $|\mathbf{r}_i - \mathbf{r}_j|$, one has the approximate relation

$$\bar{g}_{\alpha\beta}(\mathbf{r}_i - \mathbf{r}_j) \simeq e^{-k_0^2 b^2} g_{\alpha\beta}(\mathbf{r}_i - \mathbf{r}_j) \quad (18)$$

with an exponentially small error in $1/b^2$ [7], that is, one has the same Gaussian factor as for $\bar{\Gamma}$. For the eigenvalue problem (1), this shows that the eigenvalues $\bar{\omega} - i\bar{\gamma}$ of the spatially smoothed model may be related to the ones $\omega - i\gamma$ of the original model by

$$\bar{\omega} - \omega_0 - i\bar{\gamma} \simeq e^{-k_0^2 b^2} (\omega - \omega_0 - i\gamma) \quad (19)$$

within an exponentially small error in $1/b^2$. For the steady-state problem (4), it is found that the forced dipoles of the spatially smoothed model will (within an exponentially small error) coincide with the ones of the original model if one takes in the smoothed model the modified source frequency such that

$$\bar{\omega}_s - \omega_0 = e^{-k_0^2 b^2} (\omega_s - \omega_0). \quad (20)$$

A. Density of states for the infinite periodic system

In this section, we show how to recover Fourier space results of [8] for the density of states in the infinite periodic system, starting from the smoothed version of the real-space equation (1).

According to Bloch theorem, solutions of (1) can be taken of the form $d_{i,\alpha} = d_\alpha^{(\mu)} e^{i\mathbf{q}\cdot\mathbf{R}}$, where \mathbf{q} is the Bloch vector, \mathbf{R} is a vector of the Bravais lattice, the index μ labels the sublattices (for the diamond lattice, given by the combination of two shifted fcc Bravais lattices, μ assumes two values), so that all atomic positions can be written as $\mathbf{r}_i = \mathbf{R} + \mathbf{r}^{(\mu)}$, where $\mathbf{r}^{(\mu)}$ is the position with respect to the Bravais lattice vector \mathbf{R} . Injecting this ansatz in Eq. (1) modified according to Eqs. (14) and (16) gives the eigenvalue problem

$$\sum_{\beta,\nu} \bar{\mathbb{P}}_{\alpha\mu,\beta\nu}(\mathbf{q}) \bar{d}_\beta^{(\nu)} = \hbar(\bar{\omega} - \omega_0 - i\bar{\gamma}) \bar{d}_\alpha^{(\mu)} \quad (21)$$

with

$$\bar{\mathbb{P}}_{\alpha\mu,\beta\nu}(\mathbf{q}) = - \left[\bar{g}_{\alpha\beta}(\mathbf{0}) + i \frac{\hbar\bar{\Gamma}}{2} \delta_{\alpha\beta} \right] \delta_{\mu\nu} + \sum_{\mathbf{R} \in \mathcal{L}} \bar{g}_{\alpha\beta}(\mathbf{R} + \mathbf{r}^{(\mu)} - \mathbf{r}^{(\nu)}) e^{-i\mathbf{q}\cdot\mathbf{R}}. \quad (22)$$

Here, indices α, β and μ, ν label the spatial direction and the sublattices, respectively, and eigenvalues $\bar{\omega} - i\bar{\gamma}$ and eigenvectors $\bar{d}_\beta^{(\nu)}$ depend on the choice of the cutoff smooth function $\chi(\mathbf{u})$, hence, for the Gaussian choice as in Eq. (17), they depend on the value of b . By considering the first contribution of Eq. (22), inside the square brackets, it is found from the inverse Fourier transform of (15) that the tensor $\bar{g}_{\alpha\beta}(\mathbf{0})$ is scalar (it is proportional to $\delta_{\alpha\beta}$); further, using $1/(X + i0^+) = \mathcal{P} \frac{1}{X} - i\pi\delta(X)$ and (16), one finds that the imaginary part of $\bar{g}_{\alpha\beta}(\mathbf{0})$ exactly cancels with the $\bar{\Gamma}$ term. The second contribution, that is the sum over the Bravais lattice in (22), can be transformed with the Poisson summation formula. For the Gaussian smoothing function (17), the real part of $\bar{g}_{\alpha\beta}(\mathbf{0})$ can be calculated explicitly; one obtains as in [8]

$$\bar{\mathbb{P}}_{\alpha\mu,\beta\nu}(\mathbf{q}) = \frac{\hbar\Gamma}{2} \delta_{\alpha\beta} \delta_{\mu\nu} \left[\frac{1 + 2(k_0 b)^2}{2\pi^{1/2} (k_0 b)^3} - \text{Erfi}(k_0 b) e^{-k_0^2 b^2} \right] + \frac{1}{\mathcal{V}_L} \sum_{\mathbf{K} \in \text{RL}} e^{i(\mathbf{K}+\mathbf{q})\cdot(\mathbf{r}^{(\mu)} - \mathbf{r}^{(\nu)})} \bar{\tilde{g}}_{\alpha\beta}(\mathbf{K} + \mathbf{q}), \quad (23)$$

where the wave vectors \mathbf{K} run over the reciprocal lattice of the Bravais lattice, and Erfi is the imaginary error function. As expected for an infinite system, the matrix $\bar{\mathbb{P}}$ is Hermitian, so that $\bar{\gamma} = 0$.

Turning back to the original problem (1), that is, in the absence of any smoothing function, we conclude for the infinite periodic system that the spectrum is real ($\gamma = 0$) and that $\hbar\omega - \hbar\omega_0$ is any of the eigenvalues of the matrix

$$\mathbb{P}(\mathbf{q}) = \lim_{b \rightarrow 0} \bar{\mathbb{P}}(\mathbf{q}), \quad (24)$$

as in the perturbative limit of [8] (that is, for the eigenfrequencies close to ω_0 when $\omega_p^2/\omega_0^2 \rightarrow 0$, ω_p being the plasma frequency). The resulting density of states is

$$\rho(\omega) = \sum_n \int_{\mathcal{D}} \frac{d^3 q}{(2\pi)^3} \delta(\omega - \omega_{\mathbf{q},n}), \quad (25)$$

where the integral over \mathbf{q} is taken in the unit cell $\mathcal{D} = \{\sum_{i=1}^3 Q_i \tilde{\mathbf{e}}_i, -\frac{1}{2} \leq Q_i < \frac{1}{2}\}$ of the reciprocal lattice of basis $(\tilde{\mathbf{e}}_i)_{1 \leq i \leq 3}$, the sum over n runs over the all the eigenvectors of $\mathbb{P}(\mathbf{q})$, and $\omega_{\mathbf{q},n}$ is the corresponding eigenfrequency.

For the Gaussian smoothing function, the limit of the band structure for $b \rightarrow 0$ is computed in practice from the relation

$$\tilde{\mathbb{P}}_{\alpha\mu,\beta\nu}(\mathbf{q}) \simeq \mathbb{P}_{\alpha\mu,\beta\nu}(\mathbf{q})e^{-k_0^2 b^2}, \quad (26)$$

which holds within an exponentially small error in $(d_{\min}/b)^2 \gg 1$ where d_{\min} is the minimal interatomic distance [7,8]. Note that this relation, obtained for the particular case of a periodic system, is consistent with the general result (19), and implies that the eigenvectors of $\tilde{\mathbb{P}}(\mathbf{q})$ essentially coincide with those of $\mathbb{P}(\mathbf{q})$. For the diamond, $d_{\min} = a\sqrt{3}/4$. We used typically $b = 0.05a$, to which we applied the above $b \rightarrow 0$ extrapolation formula to obtain the histogram in Figs. 1, 3, and 5.

B. Penetration depth for the infinite periodic system

In this section we wish to derive, for an infinite system, the value of the penetration depth ξ and to confirm that it depends on the considered direction of the direct space and that it diverges at the band edges, both properties having already been observed for a finite-size system in Sec. III C.

Hence, we have to solve Eq. (4) in the presence of a forcing source dipole $d_{\alpha}^s = \check{d}_{\alpha}^s e^{-i\omega_s t}$ placed in \mathbf{r}_s . The solutions we look for are the steady-state dipole amplitudes $\check{d}_{i,\alpha} = \check{d}_{\mathbf{R},\alpha}^{(\mu)}$ on each diamond lattice site of position $\mathbf{r}_i = \mathbf{R} + \mathbf{r}^{(\mu)}$, where \mathbf{R} belongs to the Bravais direct lattice. Since the scope is to determine the penetration depth ξ , we restrict ourselves to the case where the source frequency ω_s is in the band gap. Then, the dipole amplitudes are expected to decay exponentially at large distances, and one may introduce the Fourier transform

$$\check{d}_{\mathbf{q},\alpha}^{(\mu)} = \sum_{\mathbf{R} \in \mathbb{L}} \check{d}_{\mathbf{R},\alpha}^{(\mu)} e^{-i\mathbf{q} \cdot \mathbf{R}}. \quad (27)$$

One applies this Fourier transform to the spatially smoothed version of Eq. (4); for a Gaussian smoothing function, the source frequency is actually chosen to be $\bar{\omega}_s$ given by Eq. (20), which ensures that the forced dipole amplitudes are essentially unaffected by the smoothing. In what follows, we can thus omit the bar (indicating the spatial smoothing) over the dipoles and the penetration depth. After calculations that closely resemble those of Sec. IV A,

$$\begin{aligned} & -\hbar(\bar{\omega}_s - \omega_0) \check{d}_{\mathbf{q},\alpha}^{(\mu)} + \sum_{\beta,\nu} \tilde{\mathbb{P}}_{\alpha\mu,\beta\nu}(\mathbf{q}) \check{d}_{\mathbf{q},\beta}^{(\nu)} \\ &= -\frac{1}{\mathcal{V}_{\mathbb{L}}} \sum_{\mathbf{K} \in \mathbb{RL}} e^{i(\mathbf{K}+\mathbf{q}) \cdot (\mathbf{r}^{(\mu)} - \mathbf{r}_s)} \sum_{\beta} \tilde{\tilde{g}}_{\alpha\beta}(\mathbf{K} + \mathbf{q}) \check{d}_{\beta}^s. \end{aligned} \quad (28)$$

One writes the formal solution of this linear system in terms of the inverse of the matrix $\tilde{\mathbb{P}}(\mathbf{q}) - \hbar(\bar{\omega}_s - \omega_0)\mathbb{1}$, where $\mathbb{1}$ is the identity; this inverse exists for all \mathbf{q} since $\bar{\omega}_s$ is in the band gap of the spatially smoothed model. Then, applying the inverse Fourier transform

$$d_{\mathbf{R},\alpha}^{(\mu)} = \int_{\mathcal{D}} \frac{d^3 q}{\mathcal{V}_{\mathbb{RL}}} d_{\mathbf{q},\alpha}^{(\mu)} e^{i\mathbf{q} \cdot \mathbf{R}}, \quad (29)$$

and using $\mathbf{K} \cdot \mathbf{R} = 0$ modulo 2π , one obtains the forced dipole amplitude on each lattice site:

$$\begin{aligned} d_{\mathbf{R},\alpha}^{(\mu)} &= - \sum_{\beta,\gamma,\nu} \sum_{\mathbf{K} \in \mathbb{RL}} \int_{\mathcal{D}} \frac{d^3 q}{(2\pi)^3} e^{i(\mathbf{K}+\mathbf{q}) \cdot (\mathbf{R} + \mathbf{r}^{(\nu)} - \mathbf{r}_s)} \\ &\quad \times \{[\tilde{\mathbb{P}}(\mathbf{q}) - \hbar(\bar{\omega}_s - \omega_0)\mathbb{1}]^{-1}\}_{\alpha\mu,\beta\nu} \tilde{\tilde{g}}_{\beta\gamma}(\mathbf{K} + \mathbf{q}) \check{d}_{\gamma}^s. \end{aligned} \quad (30)$$

A first application of Eq. (30) is to evaluate the dipole amplitudes from a numerical integration over \mathbf{q} and, fitting them in a region of large values of R in some direction \mathbf{u} , to extract the penetration depth in that direction. Using up to 256^3 points in the numerical integration over \mathbf{q} leads to the stars in Fig. 7, that compare well to the penetration depth extracted from the simulations on a finite-size system in Sec. III C. Furthermore, this approach is numerically more efficient close to the borders of the band gap, where the penetration depth diverges and the finite-size effects of the simulations become stronger.

A second strategy to obtain the penetration depth from Eq. (30) is to use the residue theorem. Since $\mathbf{K} + \mathbf{q}$ spans \mathbb{R}^3 when \mathbf{K} spans the reciprocal lattice and \mathbf{q} spans its unit cell \mathcal{D} , and since $\tilde{\mathbb{P}}(\mathbf{q}) = \tilde{\mathbb{P}}(\mathbf{q} + \mathbf{K})$, Eq. (30) can be rewritten as

$$\begin{aligned} d_{\mathbf{R},\alpha}^{(\mu)} &= - \sum_{\beta,\gamma,\nu} \int_{\mathbb{R}^3} \frac{d^3 k}{(2\pi)^3} e^{i\mathbf{k} \cdot (\mathbf{R} + \mathbf{r}^{(\nu)} - \mathbf{r}_s)} \\ &\quad \times \{[\tilde{\mathbb{P}}(\mathbf{k}) - \hbar(\bar{\omega}_s - \omega_0)\mathbb{1}]^{-1}\}_{\alpha\mu,\beta\nu} \tilde{\tilde{g}}_{\beta\gamma}(\mathbf{k}) \check{d}_{\gamma}^s. \end{aligned} \quad (31)$$

To take the large- \mathbf{R} limit in the direction \mathbf{u} , we set

$$\mathbf{R} = r\mathbf{u} + O(1) \quad \text{with} \quad r > 0, r \rightarrow \infty. \quad (32)$$

We split the integration over \mathbf{k} into an integral over the component k_{\parallel} of \mathbf{k} along \mathbf{u} and over the transverse components \mathbf{k}_{\perp} of \mathbf{k} . Then, $(\mathbf{k}_{\perp}/k) \cdot \mathbf{R}$ remains bounded, whereas $\mathbf{u} \cdot \mathbf{R}$ is divergent.

First, we consider the integral over $k_{\parallel} \in \mathbb{R}$ for a fixed \mathbf{k}_{\perp} . The integrand involves the exponential factor $e^{ik_{\parallel}r}$; since $r > 0$ we close the integration contour with a half-circle (of diverging radius) in the upper complex plane.⁵ Whereas the equation for k_{\parallel} ,

$$\bar{\omega}_{k_{\parallel}\mathbf{u}+\mathbf{k}_{\perp},n} = \bar{\omega}_s, \quad (33)$$

where $\bar{\omega}_{\mathbf{k},n}$ is the dispersion relation of the n th band of eigenfrequencies for the spatially smoothed periodic system, has for sure no real solution since $\bar{\omega}_s$ is in the band gap, it may have complex solutions $k_{\parallel}^{(0)}$ with a positive imaginary part. Due to the occurrence of the inverse matrix involving $\tilde{\mathbb{P}}(\mathbf{k})$ in the integrand, such complex solutions provide poles in the half upper plane, which according to the residue theorem lead to the damped exponential $\exp(ik_{\parallel}^{(0)}r)$. If (33) admits several roots, or roots for various band index n , one has to keep the value n_0 of n and the root $k_{\parallel}^{(0)}$ leading to the smallest imaginary part, that provides the leading contribution in the large- r limit.

⁵To this end, the Gaussian smoothing function χ is not appropriate. One can rather take $\chi(\mathbf{r}) \propto e^{-r/b}/r$, whose Fourier transform is a Lorentzian.

Then, one has to remember that there is still an integral over \mathbf{k}_\perp , and that $k_\parallel^{(0)}$ depends on \mathbf{k}_\perp . We thus face an integral of the form

$$\check{d}_{\mathbf{R},\alpha}^{(\mu)} = \int_{\mathbb{R}^2} \frac{d^2 k_\perp}{(2\pi)^2} e^{ik_\parallel^{(0)}(\mathbf{k}_\perp)r} \frac{f(\mathbf{k}_\perp)}{\partial_{k_\parallel} \bar{\omega}_{k_\parallel^{(0)}(\mathbf{k}_\perp)\mathbf{u}+\mathbf{k}_\perp, n_0}}, \quad (34)$$

where the derivative of the band dispersion relation in the denominator originates from the residue of the pole in $k_\parallel^{(0)}(\mathbf{k}_\perp)$ and the r -independent function f in the numerator is easily reconstructed from Eq. (31). To obtain an asymptotic equivalent of the integral (34) in the large- r limit, we use the saddle-point method: Eq. (34) is dominated by the contribution of the vicinity of the stationary point of the ‘‘phase,’’ that is, $\mathbf{k}_\perp^{(0)}$ such that⁶

$$\partial_{\mathbf{k}_\perp} k_\parallel^{(0)}(\mathbf{k}_\perp^{(0)}) = \mathbf{0}. \quad (35)$$

As we shall see, in general $\mathbf{k}_\perp^{(0)}$ has complex coordinates (in the plane orthogonal to \mathbf{u}) and one has to deform the integration domain of (34) to let the integration go through the stationary point.⁷ Then, one quadratizes the variation of the pole around the stationary point:

$$k_\parallel^{(0)}(\mathbf{k}_\perp^{(0)} + \delta\mathbf{k}_\perp) = k_\parallel^{(0)}(\mathbf{k}_\perp^{(0)}) + \delta\mathbf{k}_\perp \cdot B \delta\mathbf{k}_\perp + O(\delta k_\perp^3), \quad (36)$$

where the relevant values of the deviation δk_\perp of \mathbf{k}_\perp from the stationary point scale as $1/r^{1/2}$. One finally gets the equivalent

$$\check{d}_{\mathbf{R},\alpha}^{(\mu)} \underset{r \rightarrow \infty}{\sim} \frac{e^{ik_\parallel^{(0)}(\mathbf{k}_\perp^{(0)})r} f(\mathbf{k}_\perp^{(0)})}{\partial_{k_\parallel} \bar{\omega}_{k_\parallel^{(0)}(\mathbf{k}_\perp^{(0)})\mathbf{u}+\mathbf{k}_\perp^{(0)}, n_0}} \int_{\mathbb{R}^2} \frac{d^2 \delta k_\perp}{(2\pi)^2} e^{ir\delta\mathbf{k}_\perp \cdot B \delta\mathbf{k}_\perp}, \quad (37)$$

where the Gaussian integral provides a factor $1/r$. The inverse of the penetration depth in direction \mathbf{u} is thus

$$\kappa(\mathbf{u}) \equiv \frac{1}{\xi(\mathbf{u})} = \text{Im} [k_\parallel^{(0)}(\mathbf{k}_\perp^{(0)})]. \quad (38)$$

In general, this procedure is however difficult to use, even numerically, as one has to look for poles of the dispersion relation for a wave vector $\mathbf{k}^{(0)}$ with three complex coordinates. An important and manageable limiting case is for a source frequency $\bar{\omega}_s$ very close to the lower border $\bar{\omega}_{\text{inf}}$ or the upper border $\bar{\omega}_{\text{sup}}$ of the band gap. The penetration depth is then expected to diverge, so that the imaginary components of the wave vector are small and its real components are close to

⁶If there are several stationary points, one has to keep the one leading to the smallest imaginary part of $k_\parallel^{(0)}$.

⁷In the limiting case to come, close to a border of the band gap and for a given direction \mathbf{u} , the naive minimization of the imaginary part of $k_\parallel^{(0)}(\mathbf{k}_\perp)$ over real component \mathbf{k}_\perp leads to $\mathbf{k}_\perp = \mathbf{0}$ and gives an expression for the inverse penetration depth that in general differs from the correct one (42). Surprisingly, this naive minimization gives an *upper* bound on the penetration depth in the direction \mathbf{u} , i.e., it *underestimates* the damping of the field. The clue to this paradox is the fact that the value $\mathbf{k}_\perp = \mathbf{0}$, although it minimizes the imaginary part of $k_\parallel^{(0)}(\mathbf{k}_\perp)$, is *not* a stationary point of the real part of $k_\parallel^{(0)}(\mathbf{k}_\perp)$. The integration of the oscillating function $\exp\{i \text{Re} [k_\parallel^{(0)}(\mathbf{k}_\perp)]r\}$ over \mathbf{k}_\perp then induces an additional blurring of the dipole amplitudes, that *reduces* the penetration depth as compared to the naive prediction $1/\text{Im}[k_\parallel^{(0)}(\mathbf{0})]$.

the location \mathbf{q}_0 in the Bloch vector space of the band-gap border (such that $\bar{\omega}_{\mathbf{q}_0, n_0}$ is equal to $\bar{\omega}_{\text{inf}}$ or $\bar{\omega}_{\text{sup}}$). One can then quadratize the dispersion relation around the location of the border:

$$\bar{\omega}_{\mathbf{q}_0+\delta\mathbf{q}, n_0} = \bar{\omega}_{\mathbf{q}_0} + \delta\mathbf{q} \cdot \bar{A} \delta\mathbf{q} + O(\delta q^3), \quad (39)$$

where \bar{A} (resp. $-\bar{A}$) is a positive-definite matrix for the upper (resp. lower) border of the band gap. In practice, to determine the matrix \bar{A} , we diagonalize the matrix $\bar{\mathbb{P}}(\mathbf{q}_0)$ and we analytically expand the matrix $\bar{\mathbb{P}}(\mathbf{q}_0 + \delta\mathbf{q})$ up to second order in $\delta\mathbf{q}$, where the matrix $\bar{\mathbb{P}}(\mathbf{q})$ is given as a function of \mathbf{q} by Eq. (23). To obtain the resulting change of the eigenvalues of $\bar{\mathbb{P}}(\mathbf{q}_0)$ up to order δq^2 included, we treat the deviation of $\bar{\mathbb{P}}(\mathbf{q}_0 + \delta\mathbf{q})$ from $\bar{\mathbb{P}}(\mathbf{q}_0)$ linear in $\delta\mathbf{q}$ up to second order in the usual perturbation theory, and the deviation of $\bar{\mathbb{P}}(\mathbf{q}_0 + \delta\mathbf{q})$ from $\bar{\mathbb{P}}(\mathbf{q}_0)$ quadratic in $\delta\mathbf{q}$ up to first order in perturbation theory. Note that, according to Eq. (26), \bar{A} is related to its zero- b limit A , that is, to the matrix A of the original model, by

$$\bar{A} \simeq e^{-k_0^2 b^2} A \quad (40)$$

within an exponentially small error in $1/b^2$. Then, the solution of (33) obeying the stationarity condition (35) can be obtained analytically:

$$\mathbf{k}^{(0)} \equiv k_\parallel^{(0)}(\mathbf{k}_\perp^{(0)})\mathbf{u} + \mathbf{k}_\perp^{(0)} \simeq \mathbf{q}_0 + i\kappa(\mathbf{u}) \frac{\bar{A}^{-1}\mathbf{u}}{\mathbf{u} \cdot \bar{A}^{-1}\mathbf{u}} \quad (41)$$

with the expression for the inverse penetration depth

$$\kappa(\mathbf{u}) = [(\bar{\omega}_{\mathbf{q}_0} - \bar{\omega}_s)\mathbf{u} \cdot \bar{A}^{-1}\mathbf{u}]^{1/2}, \quad (42)$$

where \bar{A}^{-1} is the inverse of the matrix \bar{A} . In practice, one may find that the band-gap border is obtained for several values of \mathbf{q}_0 , due to symmetry properties (as it shall be the case for the diamond lattice). At fixed direction \mathbf{u} , one then has to select the value of \mathbf{q}_0 leading to the minimal value of $\kappa(\mathbf{u})$ in Eq. (42). Equations (41) and (42) are derived in Appendix A, where the complete resulting expression for $\check{d}_{\mathbf{R},\alpha}^{(\mu)}$ is also given.

A simple consequence of (42) is the asymptotic expression for the *maximal* penetration depth at a given frequency $\bar{\omega}_s$, i.e., maximized over the direction \mathbf{u} , close to a band-gap border:

$$\xi_{\text{max}} \underset{\bar{\omega}_s \rightarrow \bar{\omega}_{\text{bord}}}{\sim} \left(\frac{\bar{A}_{\text{max}}}{\bar{\omega}_{\text{bord}} - \bar{\omega}_s} \right)^{1/2}, \quad (43)$$

where \bar{A}_{max} is the eigenvalue of the matrix \bar{A} of maximal modulus.

We have explicitly evaluated the prediction (42) in the vicinity of the upper border of the band gap. Irrespective of the value of $k_0 a$, we find that the frequency $\bar{\omega}_{\text{sup}}$ of this upper border is reached on the so-called L point of the first Brillouin zone of the lattice, corresponding to $\mathbf{q}_0 = (\tilde{\mathbf{e}}_1 + \tilde{\mathbf{e}}_2 + \tilde{\mathbf{e}}_3)/2 = (\pi/a)(\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)$ [see Eq. (6) for the values of the $\tilde{\mathbf{e}}_i$], as it was already suspected in [8]. This point is so symmetric that all the six components of the corresponding eigenvector of the

matrix $\bar{\mathbb{P}}$ are equal, which leads to the quite explicit expression

$$\begin{aligned} \bar{\omega}_{\text{sup}} - \omega_0 &= \frac{\Gamma}{2} \left[\frac{1 + 2(k_0 b)^2}{2\pi^{1/2}(k_0 b)^3} - \text{Erfi}(k_0 b) e^{-k_0^2 b^2} \right] \\ &+ \frac{2\pi\Gamma}{k_0^3} \sum_{\mathbf{K} \in \text{RL}} \sum_v \cos[\mathbf{K}' \cdot (\mathbf{r}^{(\mu)} - \mathbf{r}^{(v)})] \\ &\times \frac{K'^2 e^{-K'^2 b^2}}{k_0^2 - K'^2}, \end{aligned} \quad (44)$$

where $\mathbf{K}' = \mathbf{K} + \mathbf{q}_0$. However, this frequency is also exactly reached for 13 other values of \mathbf{q}_0 , so that

$$\bar{\omega}_{\mathbf{q}_0} = \bar{\omega}_{\text{sup}} \text{ for } 2\mathbf{q}_0 \in \{\pm\tilde{\mathbf{e}}_1 \pm \tilde{\mathbf{e}}_2 \pm \tilde{\mathbf{e}}_3, \pm\tilde{\mathbf{e}}_1, \pm\tilde{\mathbf{e}}_2, \pm\tilde{\mathbf{e}}_3\}. \quad (45)$$

For a given direction \mathbf{u} , one thus calculates the 14 corresponding matrices \bar{A} , which are all *similar*, and one keeps the one giving the smallest contribution to Eq. (42). For $\mathbf{u} = \mathbf{e}_1$ and $\mathbf{u} = \mathbf{e}_x$, this leads to the dashed line in the right part of Figs. 7(a) and 7(b), respectively, in excellent agreement with the numerical evaluation of (30) and in good agreement with the finite-size simulations. Furthermore, for $k_0 a = 2$ as in the simulations, the direction \mathbf{e}_1 corresponds to the twice-degenerate, maximal modulus eigenvalue \bar{A}_{max} of some of the 14 matrices \bar{A} (the ones associated to $\mathbf{q}_0 = \pm\frac{1}{2}\tilde{\mathbf{e}}_2$ and $\mathbf{q}_0 = \pm\frac{1}{2}\tilde{\mathbf{e}}_3$) so that the maximal penetration depth ξ_{max} is obtained in that direction \mathbf{e}_1 . Remarkably, for $k_0 a$ large enough (but smaller than the value $k_0 a \simeq 5.14$ leading to a closure of the gap), we find that the conclusion changes, and that the maximal penetration depth is now obtained in the direction $(\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)/\sqrt{3}$. This change suggests that there exists a magic value of $k_0 a$ such that the matrix \bar{A} is scalar and, close to the upper border of the band gap, the penetration depth is isotropic, which is confirmed by the diagonalization of \bar{A} that leads to⁸

$$(k_0 a)_{\text{iso}}^{\text{sup}} \simeq 2.8632. \quad (46)$$

We have also explicitly evaluated the prediction of Eq. (42) in the vicinity of the lower border of the band gap. We have found that the frequency $\bar{\omega}_{\text{inf}}$ of this lower border is obtained in 12 values \mathbf{q}_0 of the Bloch vector, that weakly depend on $k_0 a$ and that can be parametrized in terms of a single positive dimensionless unknown quantity σ :

$$\begin{aligned} \bar{\omega}_{\mathbf{q}_0} &= \bar{\omega}_{\text{inf}} \text{ for } \mathbf{q}_0 \in \{\pm\sigma(\tilde{\mathbf{e}}_1 - \tilde{\mathbf{e}}_2), \pm\sigma(\tilde{\mathbf{e}}_1 - \tilde{\mathbf{e}}_3), \\ &\pm\sigma(\tilde{\mathbf{e}}_2 - \tilde{\mathbf{e}}_3), \pm[\sigma(\tilde{\mathbf{e}}_1 + \tilde{\mathbf{e}}_2) + (2\sigma - 1)\tilde{\mathbf{e}}_3], \\ &\pm[\sigma(\tilde{\mathbf{e}}_1 + \tilde{\mathbf{e}}_3) + (2\sigma - 1)\tilde{\mathbf{e}}_2], \\ &\pm[\sigma(\tilde{\mathbf{e}}_2 + \tilde{\mathbf{e}}_3) + (2\sigma - 1)\tilde{\mathbf{e}}_1]\}, \end{aligned} \quad (47)$$

where the basis vectors of the reciprocal of the fcc lattice are given by Eq. (6). Note that the last six elements of (47) have a σ -independent component $\pm 2\pi/a$ in the Cartesian basis,

⁸For $k_0 a$ below that value, \bar{A}_{max} is twice degenerate and the corresponding eigenspace is the plane orthogonal to $(\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)/\sqrt{3}$ for $\mathbf{q}_0 = (\tilde{\mathbf{e}}_1 + \tilde{\mathbf{e}}_2 + \tilde{\mathbf{e}}_3)/2$ and the plane orthogonal to $(-\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)/\sqrt{3}$ for $\mathbf{q}_0 = \tilde{\mathbf{e}}_1/2$. For $k_0 a$ above that value, \bar{A}_{max} is not degenerate and the corresponding eigenvector is $(\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)/\sqrt{3}$ for $\mathbf{q}_0 = (\tilde{\mathbf{e}}_1 + \tilde{\mathbf{e}}_2 + \tilde{\mathbf{e}}_3)/2$, and $(-\mathbf{e}_x + \mathbf{e}_y + \mathbf{e}_z)/\sqrt{3}$ for $\mathbf{q}_0 = \tilde{\mathbf{e}}_1/2$.

along \mathbf{e}_z , \mathbf{e}_y , \mathbf{e}_x , respectively, and their components along the other two Cartesian axes are equal; these six elements are thus located on the straight line XU , where X and U are standard remarkable points of the first Brillouin zone of the diamond lattice. For the value $k_0 a = 2$ taken in the figures, we numerically obtained $\sigma \simeq 0.330346$. For those 12 values of \mathbf{q}_0 , we have determined the 12 similar matrices \bar{A} describing the local quadratization of $\bar{\omega}_{\mathbf{q}}$ and we have kept, for a given \mathbf{u} equal to \mathbf{e}_1 or \mathbf{e}_x , the one giving the smallest contribution to Eq. (42). This has led to the dashed line in the left part of Figs. 7(a) and 7(b), respectively, again in excellent agreement with the numerical evaluation of (30) and in good agreement with the finite-size simulations. For $k_0 a = 2$, it is also found that \mathbf{e}_1 is the eigenvector of two of the 12 similar \bar{A} matrices [the ones corresponding to the last two elements of (47)] with the nondegenerate, largest modulus eigenvalue \bar{A}_{max} , so that the maximal penetration depth ξ_{max} is actually achieved in that direction, close to the lower border of the band gap. For larger values of $k_0 a$, the situation can change to a maximal penetration depth obtained along direction \mathbf{e}_x . This change occurs for the magic value

$$(k_0 a)_{\text{change}}^{\text{inf}} \simeq 2.9412, \quad (48)$$

where $\sigma \simeq 0.353740$ and the maximal modulus eigenvalue \bar{A}_{max} of the matrices \bar{A} is twice degenerate.

C. States in the gap due to vacancies

We now create a single vacancy in the infinite periodic system still using the spatially smoothed version by removing the atom at the location $\mathbf{r}_{i_0} = \mathbf{R}_0 + \mathbf{r}^{(\mu_0)}$, that is, at the lattice site \mathbf{R}_0 on the sublattice μ_0 . The eigenspectrum of the spatially smoothed version of (1) is expected to remain real ($\bar{\gamma} = 0$) but there may now be eigenvalues with $\bar{\omega}$ in the band gap of the periodic system, corresponding to states exponentially localized around the vacancy. As we will see, the corresponding $\bar{\omega}$ are given by Eq. (53).

To look for such in-gap states, we use the following trick: Starting from a periodic system in presence of a source dipole in \mathbf{r}_s (of imposed frequency $\bar{\omega}_s$ and amplitudes \check{d}_α^s), we imagine that the vacancy on site \mathbf{r}_{i_0} results from the coalescence of the corresponding forced dipole $\check{d}_{i_0, \alpha}$ with the source dipole in the limit where the source location tends to the location of the vacancy:

$$\lim_{\mathbf{r}_s \rightarrow \mathbf{r}_{i_0}} \check{d}_{i_0, \alpha} = -\check{d}_\alpha^s, \forall \alpha. \quad (49)$$

In this case, the total dipole carried by the vacancy site vanishes, as if there was indeed a vacancy there. Obviously, condition (49) can be satisfied only for specific values of $\bar{\omega}_s$ in the band gap of the spatially smoothed model, that we now determine.

Writing Eq. (30) for $\mathbf{R} = \mathbf{R}_0$, $\mu = \mu_0$, $\mathbf{r}_s = \mathbf{R}_0 + \mathbf{r}^{(\mu_0)}$, and replacing $\check{d}_{\mathbf{R}_0, \alpha}^{(\mu_0)}$ with $-\check{d}_\alpha^s$, we obtain the homogeneous linear system

$$\begin{aligned} \check{d}_\alpha^s &= \sum_{\beta, \gamma, \nu} \int_D \frac{d^3 q}{V_{\text{RL}}} \{[\bar{\mathbb{P}}(\mathbf{q}) - \hbar(\bar{\omega}_s - \omega_0)\mathbb{1}]\}_{\alpha\mu_0, \beta\nu} \\ &\times \bar{\mathbb{Q}}_{\beta\nu, \gamma\mu_0}(\mathbf{q}) \check{d}_\gamma^s, \end{aligned} \quad (50)$$

where we used $(2\pi)^3 = \mathcal{V}_{\text{RL}}\mathcal{V}_{\text{L}}$ and we called $\tilde{\mathbb{Q}}_{\alpha\mu,\beta\nu}(\mathbf{q})$ the nonscalar contribution to $\tilde{\mathbb{P}}_{\alpha\mu,\beta\nu}(\mathbf{q})$, which is the second contribution in the right-hand side of Eq. (23). In terms of matrices,

$$\tilde{\mathbb{P}}(\mathbf{q}) = \Lambda \mathbb{1} + \tilde{\mathbb{Q}}(\mathbf{q}), \quad (51)$$

where Λ is the coefficient of the scalar contribution, that is, of the first term in Eq. (23), $\Lambda = -[\bar{g}_{\alpha\alpha}(\mathbf{0}) + i\hbar\Gamma/2]$ (this is independent of the direction α). We recognize a matrix product in Eq. (50), related to the sum over ν and β ; we then use

$$[\tilde{\mathbb{P}}(\mathbf{q}) - \hbar\bar{\delta}\mathbb{1}]^{-1}\tilde{\mathbb{Q}}(\mathbf{q}) = \mathbb{1} + (\hbar\bar{\delta} - \Lambda)[\tilde{\mathbb{P}}(\mathbf{q}) - \hbar\bar{\delta}\mathbb{1}]^{-1}, \quad (52)$$

with $\bar{\delta} = \bar{\omega}_s - \omega_0$. The contribution to (50) of $\mathbb{1}$ in that expression exactly reproduces the term \check{d}_α^s of the left-hand side of (50) since the integral over \mathbf{q} on the primitive cell \mathcal{D} of the reciprocal lattice is equal to \mathcal{V}_{RL} . Simplifying the remaining contribution by the factor $\hbar\bar{\delta} - \Lambda$, it remains

$$0 = \sum_{\gamma} \left\{ \int_{\mathcal{D}} \frac{d^3q}{\mathcal{V}_{\text{RL}}} [\tilde{\mathbb{P}}(\mathbf{q}) - \hbar(\bar{\omega}_s - \omega_0)\mathbb{1}]^{-1} \right\}_{\alpha\mu_0,\gamma\mu_0} \check{d}_\gamma^s, \quad \forall \alpha. \quad (53)$$

This must have a nonzero solution for the source dipole, which is equivalent to requiring that the $\bar{\omega}_s$ -dependent 3×3 Hermitian matrix in-between curly brackets in (53) has a zero eigenvalue. To show that the condition (53) is not only sufficient, but also necessary, we have performed an alternative calculation, presented in Appendix B, that has also the advantage of including the case of several vacancies.

For the diamond lattice, we have evaluated numerically the integral over the Bloch vector \mathbf{q} in Eq. (53). We then find that the resulting 3×3 Hermitian matrix (in-between curly brackets) is scalar. As the eigenvalues of that matrix are increasing functions of $\bar{\omega}_s$, as can be shown with the Hellmann-Feynman theorem, this implies that there is at most one solution for $\bar{\omega}_s$ in the band gap. Numerically, we find that there is a solution, whose value [after extrapolation to $b \rightarrow 0$ using Eq. (19)] for $k_0a = 2$ is indicated by a vertical dotted line in Fig. 3, in agreement with a peak location in the density of states in the numerical simulations.

In the case of several vacancies, we can extend our analysis as described in Appendix B. By numerical solution of Eq. (B7), we have then investigated the in-gap states for two vacancies on sites separated by $\check{R}_2 - \check{R}_1 = \mathbf{0}, \mathbf{e}_1$ or $a\mathbf{e}_x$, being either on the same sublattice ($\check{\mu}_1 = \check{\mu}_2$) or on different sublattices ($\check{\mu}_1 \neq \check{\mu}_2$). In most cases, we have found allowed frequencies close to the one of the single-vacancy state, within the width of the central peak in the inset of Fig. 3; those states can not be resolved in that figure and we have not indicated them. For the two geometries specified in the caption of Fig. 3, we have found frequencies of two-vacancy states that are clearly out of the central peak (see the red and blue vertical dotted lines); in particular, the prediction with $(\omega - \omega_0)/\Gamma \simeq -4$ seems to match quite well the very clear secondary peak that emerges in the figure for increasing concentration of vacancies.

V. CONCLUSION

Three-dimensional infinite periodic arrangements of extended scattering objects leading to an omnidirectional band gap for light have been known since the 1990s, starting from the diamond lattice configuration of dielectric microspheres of [36]. In the case of a periodic ensemble of *pointlike* scatterers, the technical issues affecting the calculation of the band structure of light have been solved only recently [7,8,15], which has allowed us to show that the diamond lattice can also lead to a photonic band gap in the pointlike case [8].

With cold-atom experiments, a diamondlike ensemble of pointlike scatterers is in principle realizable, provided that one produces, in the appropriate optical lattice geometry [8,29], a high-quality Mott phase of atoms [5,6] having an optical transition between a spin-zero ground state and a spin-one electronic excited state [37]. In practical realizations, there will be of course unavoidable deviations from the ideal infinite periodic case, which we have quantified in this work with numerical solutions of linearly coupled dipoles equations with about 3×10^4 particles.

A first issue is due to effects of the finite size of the atomic medium. Rather than having a band structure, light has a continuous spectrum of scattering states; by analytic continuation to the lower half of the complex plane, however, it is more physical to consider, as we have done, the discrete complex eigenfrequencies $\omega - i\gamma$ of the *resonances* of the system. In the distribution function of ω , the forbidden gap remains visible in our simulations. It remains actually quite visible if one restricts to the resonances with a half decay rate γ much smaller than the free-space single-atom spontaneous emission rate Γ ; such a filtering of the resonances could be realized experimentally by performing a frequency measurement after an adjustable time delay, during which the short-lived resonances decay and are suppressed. Amusingly, a narrow peak in the distribution function of ω was observed close to the center of the infinite system band gap, when the finite-size atomic medium has a cubic shape; such a peak, absent when the medium has a spherical shape, is a very clear finite-size effect.

A second issue is due to vacancies inside the atomic medium. For a concentration of a few percent of vacancies, narrow peaks emerge in the distribution function of ω inside the gap. We were able to identify several of these peaks as corresponding to the frequencies of localized states around one or two close vacancies in an otherwise infinite periodic medium. At higher concentrations of vacancies, e.g., 20%, with no filtering on γ , the gap disappears.

From our finite-size sample, we have shown that one can quite accurately extract the penetration depth ξ of the light in the medium, and that the obtained values compare well with independent calculations in a periodic medium. Away from the borders of the band gap, ξ as a function of the imposed field frequency ω_s exhibits a plateau at a remarkably low value, between $0.5a$ and a , where a is the lattice constant of the underlying fcc lattice. Close to the borders ω_{bord} of the band gap, one can even directly observe, in our finite-size system, the onset of the divergence of ξ as $1/|\omega_s - \omega_{\text{bord}}|^{1/2}$, with a prefactor close to our analytical predictions. We have also observed from the simulations that ξ is anisotropic (it depends

on the direction of space), in agreement with our theoretical analysis, and that this anisotropy becomes quite pronounced close to the lower border of the band gap.

ACKNOWLEDGMENTS

We acknowledge a discussion with D. Delande at an early stage of this project. M.A. is member of the LabEx NUMEV.

APPENDIX A: PENETRATION DEPTH

In this appendix, for the spatially smoothed model, we derive the results (41) and (42) for the penetration depth in the direction \mathbf{u} at a frequency $\bar{\omega}_s$ close to a border of the band gap, which justifies the use of the quadratized dispersion relation (39) around the Bloch vector \mathbf{q}_0 , and we give the large-distance equivalent of the forced dipole amplitude, as obtained from the saddle-point method.

As shorthand notations, we introduce $z = k_{\parallel} - q_{0\parallel}$ and $\mathbf{x} = \mathbf{k}_{\perp} - \mathbf{q}_{0\perp}$ as the components along \mathbf{u} and in the plane orthogonal to \mathbf{u} of the vector $\mathbf{k} - \mathbf{q}_0$. We also introduce the frequency deviation from the nearest band border $\bar{\Delta} \equiv \bar{\omega}_{\mathbf{q}_0} - \bar{\omega}_s$. Then, Eq. (33) reduces to a degree-two equation for z :

$$z^2 \mathbf{u} \cdot \bar{A} \mathbf{u} + 2z \mathbf{x} \cdot \bar{A} \mathbf{u} + \mathbf{x} \cdot \bar{A} \mathbf{x} + \bar{\Delta} = 0. \quad (\text{A1})$$

Furthermore, z has to be stationary with respect to a variation of \mathbf{k}_{\perp} [see Eq. (35)]. Differentiating the trinomial (A1) with respect to \mathbf{x} , and using $\partial_{\mathbf{x}} z = \mathbf{0}$, one obtains the vectorial equation $z Q \bar{A} \mathbf{u} + Q \bar{A} Q \mathbf{x} = \mathbf{0}$ where Q projects orthogonally to \mathbf{u} . The solution is

$$\mathbf{x} = -z(Q \bar{A} Q)^{-1} \bar{A} \mathbf{u}, \quad (\text{A2})$$

where the matrix inverse is intended within the vectorial plane orthogonal to \mathbf{u} . Inserting this solution into Eq. (A1) and using $[P \bar{A} P - P \bar{A} Q (Q \bar{A} Q)^{-1} Q \bar{A} P] P \bar{A}^{-1} P = P$ where $P = 1 - Q$ is the orthogonal projector on \mathbf{u} [see relation (B.23) of Sec. III B2 in [38]], one obtains

$$z = i\kappa(\mathbf{u}) \text{ with } \kappa(\mathbf{u}) \text{ given by Eq. (42)}. \quad (\text{A3})$$

Similarly, we can derive the chain of identities $\bar{A}[\mathbf{u} - (Q \bar{A} Q)^{-1} \bar{A} \mathbf{u}] = [P \bar{A} P - P \bar{A} Q (Q \bar{A} Q)^{-1} Q \bar{A} P] \mathbf{u} = (P \bar{A}^{-1} P)^{-1} \mathbf{u} = \mathbf{u} / (\mathbf{u} \cdot \bar{A}^{-1} \mathbf{u})$, simply injecting the closure relation $P + Q = \mathbb{1}$ on both sides of the first factor \bar{A} . This gives, as in Eq. (41),

$$\frac{z \mathbf{u} + \mathbf{x}}{i\kappa(\mathbf{u})} = \mathbf{u} - (Q \bar{A} Q)^{-1} \bar{A} \mathbf{u} = \frac{\bar{A}^{-1} \mathbf{u}}{\mathbf{u} \cdot \bar{A}^{-1} \mathbf{u}}. \quad (\text{A4})$$

To determine the residue appearing in (37), one takes the derivative of the trinomial (A1) with respect to z for a fixed \mathbf{x} . Using the previous relations, one obtains

$$\partial_{k_{\parallel}} \bar{\omega}_{\mathbf{k}^{(0)}, n_0} = 2z \mathbf{u} \cdot \bar{A} \mathbf{u} + 2\mathbf{x} \cdot \bar{A} \mathbf{u} = \frac{2i\kappa(\mathbf{u})}{(\mathbf{u} \cdot \bar{A}^{-1} \mathbf{u})}. \quad (\text{A5})$$

Next, we determine the matrix B in Eq. (37) originating from the quadratization of z around the stationary point \mathbf{x} . Since \mathbf{x} is a stationary point, a first-order variation $\delta \mathbf{x}$ induces a variation δz of z that is second order in $\delta \mathbf{x}$. Performing these variations in Eq. (A1) up to second order in $\delta \mathbf{x}$, and thus up to first order

in δz , and using the previous relations, we obtain

$$B = \frac{i}{2\kappa(\mathbf{u})} (\mathbf{u} \cdot \bar{A}^{-1} \mathbf{u}) Q \bar{A} Q. \quad (\text{A6})$$

We conclude that the matrix iB appearing in the Gaussian integral (37) is negative, which justifies the fact that the saddle point is approached along the real-axis direction as in (37). If one performs the Gaussian integral, Eq. (37) reduces to

$$\check{d}_{\mathbf{R}, \alpha}^{(\mu)} \underset{r \rightarrow \infty}{\sim} \frac{e^{-\kappa(\mathbf{u})r} f(\mathbf{k}_{\perp}^{(0)})}{4i\pi r} [\det(Q \bar{A} Q)]^{-1/2}. \quad (\text{A7})$$

The determinant in that expression is conveniently transformed as $\det(Q \bar{A} Q) = (\mathbf{u} \cdot \bar{A}^{-1} \mathbf{u}) \det \bar{A}$ using the expression of the matrix of \bar{A}^{-1} in terms of the comatrix of \bar{A} (in an orthonormal basis containing the direction \mathbf{u}).

To obtain our final asymptotic form for the forced dipole amplitude, we note that, for any acceptable vector $\mathbf{k}^{(0)}$ of the pole plus saddle-point analysis, $\mathbf{k}^{(0)} + \mathbf{K}$ is again acceptable, where \mathbf{K} is any vector of the reciprocal lattice; this is due to the periodicity of the dispersion relation $\bar{\omega}_{\mathbf{k}, n_0}$. We also include a sum over possibly degenerate Bloch vector \mathbf{q}_0 leading to the same value $\bar{\omega}_{\mathbf{q}_0, n_0}$ (as discussed in the main text). We also note that, when $R \rightarrow +\infty$,

$$\begin{aligned} & |(\bar{A}^{-1} \bar{\Delta})^{1/2} (\mathbf{R} + \mathbf{r}^{(v)} - \mathbf{r}_s)| \\ &= -i(\mathbf{k}^{(0)} - \mathbf{q}_0) \cdot (\mathbf{R} + \mathbf{r}^{(v)} - \mathbf{r}_s) + o(1), \end{aligned} \quad (\text{A8})$$

which gives a simple physical interpretation to the expression (41) of $\mathbf{k}^{(0)}$: The apparently obscure correction to \mathbf{q}_0 in (41) simply originates from the fact that what more precisely matters in the asymptotic behavior of the dipole amplitudes is not $r\mathbf{u}$ but really the vectorial distance $\mathbf{R} + \mathbf{r}^{(v)} - \mathbf{r}_s$ between the considered lattice site and the source. Finally, we obtain, for $\bar{\omega}_s$ close to a border of the band gap, the asymptotic equivalent for $R \rightarrow \infty$:

$$\begin{aligned} \check{d}_{\mathbf{R}, \alpha}^{(\mu)} \sim & - \sum_{\mathbf{q}_0} \left(\frac{\bar{\Delta}}{\det \bar{A}} \right)^{1/2} \frac{e^{i\mathbf{q}_0 \cdot \mathbf{R}} e^{-|(\bar{A}^{-1} \bar{\Delta})^{1/2} (\mathbf{R} + \mathbf{r}^{(v)} - \mathbf{r}_s)|}}{4\pi \hbar |(\bar{A}^{-1} \bar{\Delta})^{1/2} (\mathbf{R} + \mathbf{r}^{(v)} - \mathbf{r}_s)|} \\ & \times \sum_{\beta, \mu, \nu} \sum_{\mathbf{K} \in \text{RL}} e^{i(\mathbf{q}_0 + \mathbf{K}) \cdot (\mathbf{r}^{(v)} - \mathbf{r}_s)} \phi_{\alpha\mu}^{(n_0)} \phi_{\beta\nu}^{(n_0)*} \tilde{g}_{\beta\gamma}(\mathbf{q}_0 + \mathbf{K}) \check{d}_{\gamma}^s, \end{aligned} \quad (\text{A9})$$

where $\phi_{\alpha\mu}^{(n_0)}$ are the components of the normalized eigenvector of $\bar{\mathbb{P}}(\mathbf{q}_0)$ of eigenvalue $\bar{\omega}_{\mathbf{q}_0, n_0}$, we approximated $\mathbf{k}^{(0)}$ by \mathbf{q}_0 in the argument of $\tilde{g}_{\beta\gamma}$, and the square root $(\bar{A}^{-1} \bar{\Delta})^{1/2}$ of the matrix $\bar{A}^{-1} \bar{\Delta}$ is well defined since this matrix is positive. Note that the second line of (A9) does not depend on \mathbf{R} .

APPENDIX B: A GENERAL VACANCY CALCULATION

We consider here the infinite periodic system, with a finite number of vacancies at nodes $(\bar{\mathbf{R}}_i, \check{\mu}_i)$, $1 \leq i \leq n$, where we recall that \mathbf{R} belongs to the Bravais lattice and μ labels the sublattices. The scope is to determine the frequencies $\bar{\omega}$ of the localized states that can exist, due to the presence of the vacancies, in the band gap of the periodic system, in the spatially smoothed version of the model.

The idea is to formally introduce, in the coupled equations for the dipoles, fictitious dipoles carried by the vacancies. Among the physical dipoles, the spatially smoothed version of

Eq. (1) holds:

$$0 = (\Lambda - \hbar\bar{\delta})d_{\mathbf{R}}^{(\mu)} + \sum_{\mathbf{R}',\mu'} \bar{g}(\mathbf{R} + \mathbf{r}^{(\mu)} - \mathbf{R}' - \mathbf{r}^{(\mu')})d_{\mathbf{R}'}^{(\mu')}. \quad (\text{B1})$$

Here, the prime over the summation symbol means that the sum is restricted to the physical dipoles, $\bar{\delta} = \bar{\omega} - \omega_0$ is the detuning from the atomic resonance, Λ is defined below Eq. (51), and we have used for conciseness an implicit vectorial notation for the dipoles and an implicit matrix notation for \bar{g} . For the fictitious dipoles, the equation is that they are equal to zero:

$$0 = (\Lambda - \hbar\bar{\delta})d_{\check{\mathbf{R}}_i}^{(\check{\mu}_i)}, \quad \forall i \in \{1, \dots, n\}. \quad (\text{B2})$$

This allows us to formally extend the sum in Eq. (B1) to the fictitious dipoles, that is, one can remove the prime over the summation symbol. One can then merge the two series of equations using the usual plus-minus trick: for *all* \mathbf{R} in the Bravais lattice and for *all* sublattices μ , one requires that

$$0 = (\Lambda - \hbar\bar{\delta})d_{\mathbf{R}}^{(\mu)} + \sum_{\mathbf{R}',\mu'} \bar{g}(\mathbf{R} + \mathbf{r}^{(\mu)} - \mathbf{R}' - \mathbf{r}^{(\mu')})d_{\mathbf{R}'}^{(\mu')} - \sum_{i=1}^n \delta_{\mathbf{R},\check{\mathbf{R}}_i} \delta_{\mu,\check{\mu}_i} \bar{s}_i, \quad (\text{B3})$$

where δ is the Kronecker symbol and we have introduced the auxiliary unknowns

$$\bar{s}_i \equiv \sum_{\mathbf{R}',\mu'} \bar{g}(\check{\mathbf{R}}_i + \mathbf{r}^{(\check{\mu}_i)} - \mathbf{R}' - \mathbf{r}^{(\mu')})d_{\mathbf{R}'}^{(\mu')}. \quad (\text{B4})$$

Then, taking the Fourier transform (27) of Eq. (B3) and using (22),

$$0 = \sum_{\mu'} [\bar{\mathbb{P}}(\mathbf{q}) - \hbar\bar{\delta}\mathbb{1}]_{\mu\mu'} d_{\mathbf{q}}^{(\mu')} - \sum_{i=1}^n e^{-i\mathbf{q}\cdot\check{\mathbf{R}}_i} \delta_{\mu,\check{\mu}_i} \bar{s}_i. \quad (\text{B5})$$

Since the frequency $\bar{\omega}$ is in the gap, the matrix is invertible, and taking the inverse Fourier transform, one obtains

$$d_{\mathbf{R}}^{(\mu)} = \sum_{i=1}^n \int_{\mathcal{D}} \frac{d^3q}{\mathcal{V}_{\text{RL}}} e^{i\mathbf{q}\cdot(\mathbf{R}-\check{\mathbf{R}}_i)} \{[\bar{\mathbb{P}}(\mathbf{q}) - \hbar\bar{\delta}\mathbb{1}]^{-1}\}_{\mu\check{\mu}_i} \bar{s}_i. \quad (\text{B6})$$

Expressing the fact that the fictitious dipoles are all equal to zero, we find the homogeneous system of equations:

$$\sum_{i=1}^n \left\{ \int_{\mathcal{D}} \frac{d^3q}{\mathcal{V}_{\text{RL}}} e^{i\mathbf{q}\cdot(\check{\mathbf{R}}_j - \check{\mathbf{R}}_i)} [\bar{\mathbb{P}}(\mathbf{q}) - \hbar\bar{\delta}\mathbb{1}]^{-1} \right\}_{\check{\mu}_j \check{\mu}_i} \bar{s}_i = 0, \quad (\text{B7})$$

to be satisfied $\forall j \in \{1, \dots, n\}$. The acceptable in-gap frequencies are such that the system admits a nonidentically zero solution $(\bar{s}_i)_{1 \leq i \leq n}$, that is the determinant of the corresponding $3n \times 3n$ matrix (the one in-between curly brackets) must vanish. In the case of a single vacancy, this reproduces Eq. (53).

Finally, we have performed the consistency check that, if one replaces in Eq. (B4) the dipoles in terms of the auxiliary unknowns \bar{s}_j , as given by (B6), one recovers exactly the same system as (B7), using Eqs. (22) and (51) and the fact that the integral over \mathbf{q} on the primitive cell \mathcal{D} of the reciprocal lattice is equal to its volume \mathcal{V}_{RL} .

-
- [1] G. Grosso and G. Pastori-Parravicini, *Solid State Physics* (Academic, New York, 2000).
- [2] A. M. Afanas'ev and Yu. Kagan, *Zh. Eksp. Teor. Fiz.* **52**, 191 (1966) [*Sov. Phys.-JETP* **25**, 124 (1967)]; G. B. Smirnov and Y. V. Shvydko, *Pis'ma Zh. Eksp. Teor. Fiz.* **34**, 409 (1982) [*JETP Lett.* **35**, 505 (1982)].
- [3] J. J. Hopfield, *Phys. Rev.* **112**, 1555 (1958); V. Agranovich, *Zh. Eksp. Teor. Fiz.* **10**, 216 (1960) [*Sov. Phys.-JETP* **10**, 307 (1960)].
- [4] J. D. Joannopoulos, S. G. Johnson, J. N. Winn, and R. D. Meade, *Photonic Crystals: Molding the Flow of Light*, 2nd ed. (Princeton University Press, Princeton, NJ, 2008); F. Zolla, G. Renversez, A. Nicolet, B. Kuhlmeiy, S. Guenneau, D. Felbacq, A. Argyros, and S. Leon-Saval, *Foundations of Photonic Crystal Fibres*, 2nd ed. (Imperial College Press, London, 2012).
- [5] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, *Nature (London)* **415**, 39 (2002).
- [6] M. Anderlini, P. J. Lee, B. L. Brown, J. Sebby-Strabley, W. D. Phillips, and J. V. Porto, *Nature (London)* **448**, 452 (2007).
- [7] M. Antezza and Y. Castin, *Phys. Rev. Lett.* **103**, 123903 (2009).
- [8] M. Antezza and Y. Castin, *Phys. Rev. A* **80**, 013816 (2009).
- [9] Masao Takamoto, Feng-Lei Hong, Ryoichi Higashi, and Hidetoshi Katori, *Nature (London)* **435**, 321 (2005).
- [10] T. L. Nicholson, M. J. Martin, J. R. Williams, B. J. Bloom, M. Bishof, M. D. Swallows, S. L. Campbell, and J. Ye, *Phys. Rev. Lett.* **109**, 230801 (2012), and references therein.
- [11] A. D. Ludlow, T. Zelevinsky, G. K. Campbell, S. Blatt, M. M. Boyd, M. H. G. de Miranda, M. J. Martin, J. W. Thomsen, S. M. Foreman, Jun Ye, T. M. Fortier, J. E. Stalnaker, S. A. Diddams, Y. Le Coq, Z. W. Barber, N. Poli, N. D. Lemke, K. M. Beck, and C. W. Oates, *Science* **139**, 1805 (2008).
- [12] S. D. Jenkins and J. Ruostekoski, *Phys. Rev. A* **86**, 031602(R) (2012).
- [13] D. V. van Coevorden, R. Sprik, A. Tip, and A. Lagendijk, *Phys. Rev. Lett.* **77**, 2412 (1996).
- [14] P. de Vries, D. V. van Coevorden, and A. Lagendijk, *Rev. Mod. Phys.* **70**, 447 (1998).
- [15] J. A. Klugkist, M. Mostovoy, and J. Knoester, *Phys. Rev. Lett.* **96**, 163903 (2006).
- [16] A. Schilke, C. Zimmermann, P. W. Courteille, and W. Guerin, *Phys. Rev. Lett.* **106**, 223903 (2011); A. Schilke, C. Zimmermann, and W. Guerin, *Phys. Rev. A* **86**, 023809 (2012).
- [17] A. Schilke, C. Zimmermann, P. W. Courteille, and W. Guerin, *Nat. Photonics* **6**, 101 (2011).
- [18] S. Rist, C. Menotti, and G. Morigi, *Phys. Rev. A* **81**, 013404 (2010).
- [19] I. Carusotto, M. Antezza, F. Bariani, S. De Liberato, and C. Ciuti, *Phys. Rev. A* **77**, 063621 (2008).
- [20] H. Zoubi and H. Ritsch, *Phys. Rev. A* **76**, 013817 (2007).
- [21] O. Morice, Y. Castin, and J. Dalibard, *Phys. Rev. A* **51**, 3896 (1995).
- [22] W. Zhang and D. F. Walls, *Phys. Rev. A* **49**, 3799 (1994).

- [23] J. Ruostekoski and J. Javanainen, *Phys. Rev. A* **55**, 513 (1997).
- [24] H. Wallis, *Phys. Rev. A* **56**, 2060 (1997).
- [25] K. V. Krutitsky, F. Burgbacher, and J. Audretsch, *Phys. Rev. A* **59**, 1517 (1999).
- [26] D. Felbacq and M. Antezza, SPIE Newsroom (2012), doi: [10.1117/2.1201206.004296](https://doi.org/10.1117/2.1201206.004296), and references therein.
- [27] U. Fano, *Phys. Rev.* **103**, 1202 (1956).
- [28] A. Chelnokov, S. Rowson, J.-M. Lourtioz, V. Berger, and J.-Y. Courtois, *J. Opt. A: Pure Appl. Opt.* **1**, L3 (1999).
- [29] O. Toader, T. Y. M. Chan, and S. John, *Phys. Rev. Lett.* **92**, 043905 (2004).
- [30] D. Yu, *Phys. Rev. A* **84**, 043833 (2011).
- [31] C. Weitenberg, M. Endres, J. F. Sherson, M. Cheneau, P. Schauß, T. Fukuhara, I. Bloch, and S. Kuhr, *Nature (London)* **471**, 319 (2011).
- [32] Y. Bidel, B. Klappauf, J. C. Bernard, D. Delande, G. Labeyrie, C. Miniatura, D. Wilkowski, and R. Kaiser, *Phys. Rev. Lett.* **88**, 203902 (2002).
- [33] O. Morice, Ph.D. thesis, Université Pierre et Marie Curie, 1995, <http://tel.archives-ouvertes.fr/tel-00011902>.
- [34] J. D. Jackson, *Classical Electrodynamics*, 2nd ed. (Wiley, New York, 1975).
- [35] A. L. Gaunt, T. F. Schmidutz, I. Gotlibovych, R. P. Smith, and Z. Hadzibabic, *Phys. Rev. Lett.* **110**, 200406 (2013).
- [36] K. M. Ho, C. T. Chan, and C. M. Soukoulis, *Phys. Rev. Lett.* **65**, 3152 (1990).
- [37] T. Fukuhara, S. Sugawa, M. Sugimoto, S. Taie, and Y. Takahashi, *Phys. Rev. A* **79**, 041604 (2009).
- [38] C. Cohen-Tannoudji, J. Dupont-Roc, and G. Grynberg, *Processus d'interaction entre photons et atomes* (InterEditions/ Editions du CNRS, Paris, 1988).