

# Role of the phase-matching condition in nondegenerate four-wave mixing in hot vapors for the generation of squeezed states of light

M. T. Turnbull,<sup>1</sup> P. G. Petrov,<sup>1</sup> C. S. Embrey,<sup>1</sup> A. M. Marino,<sup>2</sup> and V. Boyer<sup>1</sup>

<sup>1</sup>*Midlands Ultracold Atom Research Centre, School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom*

<sup>2</sup>*Homer L. Dodge Department of Physics and Astronomy, The University of Oklahoma, 440 W. Brooks St., Norman, Oklahoma 73019, USA*

(Received 28 March 2013; published 26 September 2013)

Nondegenerate forward four-wave mixing in hot atomic vapors has been shown to produce strong quantum correlations between twin beams of light [McCormick *et al.*, *Opt. Lett.* **32**, 178 (2007)], in a configuration which minimizes losses by absorption. In this paper, we look at the role of the phase-matching condition in the trade-off that occurs between the efficiency of the nonlinear process and the absorption of the twin beams. To this effect, we develop a semiclassical model by deriving the atomic susceptibilities in the relevant double- $\Lambda$  configuration and by solving the classical propagation of the twin-beam fields for parameters close to those found in typical experiments. These theoretical results are confirmed by a simple experimental study of the nonlinear gain experienced by the twin beams as a function of the phase mismatch. The model shows that the amount of phase mismatch is key to the realization of the physical conditions in which the absorption of the twin beams is minimized while the cross coupling between the twin beams is maintained at the level required for the generation of strong quantum correlations. The optimum is reached when the four-wave mixing process is not phase matched for fully resonant four-wave mixing.

DOI: [10.1103/PhysRevA.88.033845](https://doi.org/10.1103/PhysRevA.88.033845)

PACS number(s): 42.50.Gy, 42.65.Yj, 42.65.Hw

## I. INTRODUCTION

Continuous-variable entanglement can be generated deterministically with a phase-insensitive optical amplifier. For a gain larger than 1, the system produces a two-mode squeezed state where the signal and the idler (here referred to as probe and conjugate, respectively) display Einstein-Podolsky-Rosen-type entanglement [1]. Such an amplifier can be realized using a nonlinear optical process such as parametric down-conversion [2,3] or four-wave mixing (4WM) [4]. In real physical systems, the presence of absorption reduces the amount of quantum correlations which can be generated, and although 4WM in atomic vapors can lead to large gains, resonant atomic processes are also responsible for losses, which limit the amount of observable squeezing. Recently, a configuration in 4WM was found which reduces absorption [5,6], and generates large degrees of squeezing [7–10].

There have been a number of reasons put forward to explain this success. The main one is the nature of the nonlinearity, which is based on coherence effects between the hyperfine electronic ground states rather than on the saturation of a transition of a two-level atom [6,11]. Indeed, avoiding a large atomic population in the excited state is key to the reduction of the noise associated with spontaneous emission. More specifically, it was pointed out that the  $D_1$  line of alkali-metal atoms is particularly amenable to the establishment of a ground state coherence [12]. Phenomenological [7] and sophisticated microscopic [13] models have also been able to reproduce these results. We show here that the production of squeezing is also due in great part to a judicious choice of the parameters that most greatly influence the phase-matching condition of the nonlinear process, specifically the relative frequencies of the beams and the angle between them. Maybe surprisingly, the highest levels of squeezing are achieved when the system is not phase matched for resonant 4WM.

The paper is divided as follows. In Sec. II, we theoretically study 4WM in an atomic vapor in a double- $\Lambda$  configuration where both pumps are detuned from the atomic resonance. From the atomic susceptibilities we evaluate the impact of the phase-matching condition on the gain and the absorption in the forward-4WM configuration. In Sec. III, we report on a systematic experimental study of the phase-matching condition which confirms the findings of Sec. II. In Sec. IV, we extend our model to take into account the Doppler effect due to the thermal motion of the atoms. Finally in Sec. V we discuss the impact of our theoretical and experimental findings on the possibility of generating strong two-mode squeezing with 4WM in a hot atomic vapor. From the model we deduce the best parameters in terms of beam geometry and beam detunings, and compare them to the recent squeezing experiments of Refs. [7–10].

## II. THEORETICAL MODEL OF NONDEGENERATE FOUR-WAVE MIXING

When describing nonlinear media in the presence of off-resonant fields, it is common to separate the response of the system into a linear part and a nonlinear part [14]. The linear contribution leads to an index of refraction which modifies the linear dispersion relation for each of the individual light fields in the medium. The nonlinear part acts as a perturbative source term in the propagation equations. It enables energy transfer between light fields for those configurations where the phase-matching condition is fulfilled, that is to say, when the total wave vector of the waves giving up energy equals the total wave vector of the waves receiving the energy. The relevant wave vectors are those in the medium. They are equal to the wave vectors in vacuum times their corresponding indices of refraction.




FIG. 1. (Color online) Double- $\Lambda$  scheme on the  $D_1$  line of  $^{85}\text{Rb}$ . The hyperfine splitting of the excited state is not resolved due to Doppler broadening. The  $\Lambda$  transitions, detuned by  $\Delta_1$  and  $\Delta_2$ , are driven by pump fields with resonant Rabi frequencies  $\Omega_1$  and  $\Omega_2$ , respectively. Note that the pumps can actually be a single laser beam.

In resonant media, such as atomic vapors excited close to atomic transitions, the medium is strongly perturbed by the presence of light. For instance, optical pumping by a strong pump beam can affect the atomic populations in the hyperfine levels, leading to a strong change in the index of refraction seen by a weaker beam. In this case, the usual expansion separating the linear and the nonlinear responses may not be appropriate. Instead we consider an expansion of the nonlinear polarization to first order in the electric fields of the weak beams and to all orders in the electric fields of the pump beams [6].

We consider the double- $\Lambda$  configuration, shown in Fig. 1, which was used to demonstrate intensity-difference squeezing and quadrature entanglement in a vapor of  $^{85}\text{Rb}$  [15,16]. A nondegenerate 4WM parametric process drives an atom from one of the hyperfine ground states to the other hyperfine ground state and back to the initial state. In the process, two pump photons are converted into two twin photons, called probe and conjugate, with wave vectors  $\mathbf{k}_p$  and  $\mathbf{k}_c$  in vacuum, and frequencies  $\omega_p$  and  $\omega_c$ . The nonlinearity originates in a strong coupling between the probe and the conjugate fields mediated by the coherence of the electronic ground states [11]. Following the usual experimental configuration, we further assume that the two pump photons come from a single field, of wave vector  $\mathbf{k}_0$  in vacuum and frequency  $\omega_0$ . We denote  $\delta$  the two-photon detuning of the pump and the probe:  $\delta = \omega_0 - \omega_p - \omega_{\text{HF}}$ , where  $\omega_{\text{HF}}$  is the hyperfine splitting of the ground state. The natural linewidth of the excited state is  $\gamma = 2\pi \times 6$  MHz. In the rest of the paper, we vary  $\delta$  by changing the frequency of the probe. The 4WM resonance occurs roughly at  $\delta = 0$  (more on this below).

For simplicity, we assume that the coupling strengths of the pump to both transitions are equal, corresponding to a resonant Rabi frequency  $\Omega = \Omega_1 = \Omega_2$ . Following Ref. [6], we calculate the atomic susceptibilities at the probe and conjugate frequencies in the limit of weak probe and conjugate fields. The detailed calculation is developed in the Appendix. The susceptibilities are derived by calculating the steady-state value of the density matrix of a four-level system interacting with the four fields of the double  $\Lambda$ . The atomic polarization at the frequencies of the probe and the conjugate is proportional

to the average oscillating atomic electric dipole at those frequencies and therefore to the off-diagonal components of the density matrix corresponding to these transitions. To first order in the probe and conjugate fields, the atomic polarization is described by two direct susceptibilities,  $\chi_{pp}$  and  $\chi_{cc}$ , and two cross susceptibilities,  $\chi_{pc}$  and  $\chi_{cp} = \chi_{pc}^*$ , given by Eqs. (A13)–(A16). The cross susceptibilities are responsible for the 4WM.

The propagation equations for the slowly varying envelopes of the probe and conjugate fields  $\mathcal{E}_p$  and  $\mathcal{E}_c$ , using the polarization expressions (A11) and (A12), are given in steady state by

$$\frac{\partial}{\partial z} \mathcal{E}_p = \frac{ik_p}{2\epsilon_0} P(\omega_p) e^{-ik_p \cdot \mathbf{r}}, \quad (1)$$

$$\frac{\partial}{\partial z} \mathcal{E}_c = \frac{ik_c}{2\epsilon_0} P(\omega_c) e^{-ik_c \cdot \mathbf{r}}. \quad (2)$$

Furthermore, if we consider the case of copropagating, or nearly copropagating, beams along the  $z$  axis, these equations read

$$\frac{\partial}{\partial z} \mathcal{E}_p = \frac{ik_p}{2} \chi_{pp}(\omega_p) \mathcal{E}_p + \frac{ik_p}{2} \chi_{pc}(\omega_p) e^{i\Delta k_z z} \mathcal{E}_c^*, \quad (3)$$

$$\frac{\partial}{\partial z} \mathcal{E}_c = \frac{ik_c}{2} \chi_{cc}(\omega_c) \mathcal{E}_c + \frac{ik_c}{2} \chi_{cp}(\omega_c) e^{i\Delta k_z z} \mathcal{E}_p^*, \quad (4)$$

where  $\Delta k_z$  is the projection of the geometric phase mismatch  $\Delta \mathbf{k} = 2\mathbf{k}_0 - \mathbf{k}_p - \mathbf{k}_c$  on the  $z$  axis, and the conservation of energy imposes the condition  $\omega_p + \omega_c = 2\omega_0$ .

In the low pump depletion limit, which is usually experimentally the case, the pump Rabi frequency  $\Omega$  is constant along the vapor cell and these equations are simply first order coupled linear differential equations. When the dynamics is dominated by the cross terms and no conjugate field is injected, the probe and conjugate fields grow asymptotically exponentially and the system behaves like a phase-insensitive amplifier for a probe input field [14,17]. At the quantum level, these cross terms are responsible for the creation of quantum correlations between the output probe and conjugate fields, leading to the production of a two-mode squeezed state [1,16]. The larger the gain of the amplifier, the greater the amount of squeezing.

The full dynamics is more complicated than pure 4WM because of the presence of the direct terms  $\chi_{pp}$  and  $\chi_{cc}$ . However, the general form of Eqs. (3) and (4) offers us a straightforward interpretation of  $\chi_{pp}$  and  $\chi_{cc}$  in terms of effective linear susceptibilities for the probe and the conjugate fields. Note that unlike the usual linear susceptibilities, these effective susceptibilities depend, nonlinearly, on the pump field. Therefore they give rise to a pump-dependent complex index of refraction for the probe and the conjugate. The real part influences the phase matching of the process, as discussed below. The imaginary part translates into absorption.

In the original proposal by Lukin *et al.* [6], the pump beams are resonant with an atomic transition ( $\Delta_1 = \Delta_2 = 0$ ). This results in a remarkable situation where the pumps and the twin beams fulfill the two-photon Raman resonance and enter an electromagnetically induced transparency (EIT) condition [18]. As a result, they see a perfectly transparent medium, as witnessed by a vanishing imaginary part of  $\chi_{pp}$  and  $\chi_{cc}$ . At the same time, the cross susceptibilities are enhanced by the coherence of the hyperfine electronic ground states. In

theory, it should lead to very efficient 4WM and virtually no absorption, even for a weak pump. In practice, the EIT effect in hot atomic vapors is limited because the Doppler effect, the presence of multiple excited hyperfine levels, and the finite transit time of the atoms in the laser beams all act to increase the decoherence rate between the ground states. This causes residual absorption of the probe field and as a result, only low levels of squeezing have been observed in this configuration [19].

In contrast, most recent squeezing experiments in hot atomic vapors operate at large detuning  $\Delta_1/2\pi$ , typically 0.5–1 GHz, and at larger pump power [8–10,15]. In these conditions, the susceptibilities take a different form from the resonant case. This off-resonant form is depicted in Fig. 2, for typical experimental parameters.

The first important property of these susceptibilities is that the 4WM resonance is shifted from the bare two-photon resonance ( $\delta = 0$ ) by the light shift created by the more resonant pump, on the  $5S_{1/2}(F = 2) \rightarrow 5P_{1/2}$  transition. For typical experimental parameters, the shift is of the order of  $-5\gamma$ . In the rest of the paper, the “blue side” of the 4WM resonance refers to the range of detunings  $\delta$  for which the frequency of the probe is above the 4WM resonance. The other side of the 4WM resonance is the “red side” (see Fig. 2).

The second property is that the imaginary part of  $\chi_{pp}$  [ $\text{Im}(\chi_{pp})$ , dashed line] is maximum at the 4WM resonance, due to Raman absorption. It is therefore not a good place to observe 4WM because the medium is essentially opaque for the probe at this detuning. On the other hand,  $\text{Im}(\chi_{pp})$  decays much faster than the magnitude of  $\chi_{pc}$  when moving away from resonance, therefore there is a range of  $\delta$  on each side of the 4WM resonance where  $\chi_{pc}$  still exhibits a substantial magnitude, while  $\text{Im}(\chi_{pp})$  has almost completely vanished (see Fig. 3). These regions of the two-photon detuning are better places to observe quantum effects.

The third important property is the behavior of the real part of  $\chi_{pp}$  [ $\text{Re}(\chi_{pp})$ , solid line], which is effectively responsible for the index of refraction for the probe. Around the 4WM resonance,  $\text{Re}(\chi_{pp})$  is the sum of a dispersive feature resulting from the 4WM coupling and the off-resonance negative susceptibility resulting from the one-photon transition between ground state and excited state. These competing terms lead to a cancellation of  $\text{Re}(\chi_{pp})$  around the bare two-photon resonance ( $\delta = 0$ ).

It is legitimate to wonder if it would not be possible to take advantage of the coherence between the ground states to reduce the probe absorption through the EIT phenomenon, as envisioned in the original proposal. Theoretically, for a very low value of the ground state decoherence rate  $\gamma_c$ , it is indeed possible to observe the transparency window at the bare two-photon resonance, as shown in Fig. 3. In practice, such low decoherence rates may not be achieved due to imperfections associated with hot atomic vapors. Experimental results reported below and in Refs. [13,20] are compatible with values of  $\gamma_c$  higher than  $\gamma/20$ , for which there is no marked transparency window. In spite of the lack of efficient EIT, the off-resonant value of  $\text{Im}(\chi_{pp})$  is small enough compared to the cross coupling to ensure efficient 4WM. In fact, the cross coupling is large enough that experiments set to demonstrate large levels of squeezing do not require special precautions




FIG. 2. (Color online) The direct and cross susceptibilities for the probe and conjugate fields as a function of the two-photon detuning  $\delta$ , varied by changing the probe frequency, expressed in units of the excited state decay rate  $\gamma = 2\pi \times 6$  MHz. The solid black lines are the real parts; the dashed red lines are the imaginary parts. The units on the y axes are arbitrary but identical for all four susceptibilities. The resonant Rabi frequency of the pump is  $\Omega = 60\gamma$ , the detuning of the pump is  $\Delta_1 = 140\gamma$ , and the decoherence rate of the excited state is taken here to be  $\gamma_c = 0.2\gamma$ . The feature at  $\delta = 150\gamma$ , visible on  $\chi_{pp}$  in the inset, corresponds to the one-photon resonance of the probe from the  $F = 2$  hyperfine ground state. The region of positive  $\text{Re}(\chi_{pp})$  around the 4WM resonance is indicated with a thicker line.




FIG. 3. (Color online) 4WM coupling  $|\chi_{pc}|$  (solid black line) and probe absorption  $\text{Im}(\chi_{pp})$  (dashed red line) as a function of the two-photon detuning  $\delta$ , for  $\gamma_c = 0.2\gamma$ . The probe absorption is also shown for  $\gamma_c = \gamma$  (dashed-dotted line) and for  $\gamma_c = 0.02\gamma$  (dotted line). In the latter case, the transparency window around  $\delta = 0$  starts to become visible.

with regard to decoherence sources such as stray magnetic fields [13].

Note that these considerations do not apply to the conjugate field, which is much further detuned from resonance than the probe. The direct susceptibility of the conjugate  $\chi_{cc}$  is substantially smaller than the other susceptibilities and can be regarded as zero in practice.

Now that adequate ranges of  $\delta$  have been identified for which probe losses are negligible compared to 4WM amplification, the question is whether there is a geometric configuration of the light fields for which 4WM is phase matched. To answer this question, we recall from Ref. [6] the solutions to the propagation equations (3) and (4), for a seed  $\mathcal{E}_s$  on the input probe and no input conjugate:

$$\mathcal{E}_p = \mathcal{E}_s \exp(\delta a L) \left[ \cosh(\xi L) + \frac{a}{\xi} \sinh(\xi L) \right], \quad (5)$$

$$\mathcal{E}_c^* = \mathcal{E}_s \exp(\delta a L) \frac{a_{cp}}{\xi} \sinh(\xi L), \quad (6)$$

where  $L$  is the length of the medium,  $a_{pj} = ik_p \chi_{pj}/2$ ,  $a_{cj} = ik_c \chi_{cj}^*/2$ ,  $\delta a = (a_{pp} - a_{cc} + i\Delta k_z)/2$ ,  $a = (a_{pp} + a_{cc} - i\Delta k_z)/2$ ,  $\xi = \sqrt{-a_{pc}a_{cp} + a^2}$ , and  $j = p, c$ . We define the probe and conjugate intensity gains  $g_p$  and  $g_c$  as  $|\mathcal{E}_p|^2 = g_p |\mathcal{E}_s|^2$  and  $|\mathcal{E}_c|^2 = g_c |\mathcal{E}_s|^2$ .

From these expressions, we plot in Fig. 4 the probe and conjugate gains as a function of  $\delta$  and the geometric phase mismatch  $\Delta k_z$ . One can see that for  $\Delta k_z \simeq 0$ , the maximum gain is obtained around  $\delta \simeq 0$ , for both the probe and the conjugate. When  $\Delta k_z$  increases, the gain on both the probe and the conjugate increases, while the gain resonance moves towards the 4WM resonance (shown with the vertical dashed line). At larger  $\Delta k_z$ , the gain resonance comes asymptotically within  $\gamma$  of the 4WM resonance, and the probe intensity drops while the conjugate intensity keeps increasing. Finally, at large  $\Delta k_z$ , the conjugate intensity also drops.




FIG. 4. (Color online) Theoretical output probe and conjugate gains  $g_p$  and  $g_c$  as a function of the two-photon detuning and the geometrical phase mismatch. Here the decoherence rate is  $\gamma_c = 0.5\gamma$ , the atom density is  $\mathcal{N} = 3 \times 10^{12} \text{ cm}^{-3}$ , the length of the medium is  $L = 12.5 \text{ mm}$  and the pump Rabi frequency is  $\Omega = 60\gamma$ . The dashed lines indicate the position of the 4WM resonance.

The position of the gain resonance is the result of the 4WM phase matching (or absence thereof), and is influenced by the effective index of refraction seen by the probe and the conjugate as follows. Since  $\text{Re}(\chi_{cc})$  is much smaller than the other susceptibilities, the conjugate effectively propagates in a medium of index 1. The situation is different for the probe, for which the index of refraction changes sign at the 4WM resonance and at  $\delta \simeq 0$ , as indicated in Fig. 2. The change in sign of  $\text{Re}(\chi_{pp})$  at  $\delta \simeq 0$  means that the probe experiences an effective index of refraction,  $n_p$ , smaller than 1 for  $\delta \gtrsim 0$  and larger than 1 between  $\delta \lesssim 0$  and the 4WM resonance. We assume that the index of refraction experienced by the pump,  $n_0$ , is close to 1 since the pump tends to optically pump the atoms towards the ground state of the off-resonant transition  $5S_{1/2}(F=3) \rightarrow 5P_{1/2}$ . This assumption will be refined later.

The geometric phase-matching condition  $\Delta k_z = 0$  is the phase-matching condition in free space. It is fulfilled only when the beams are rigorously copropagating, as shown in Fig. 5(a). For the process to be efficient, the effective phase-matching condition must be fulfilled:

$$2\mathbf{k}_0 - n_p \mathbf{k}_p - \mathbf{k}_c = 0, \quad (7)$$

as shown in Fig. 5(b). This condition is identical to the geometric phase-matching condition ( $\Delta k_z = 0$ ) only when  $n_p = \sqrt{1 + \text{Re}(\chi_{pp})} = 1$ , which occurs around  $\delta = 0$  (Fig. 2). When  $n_p > 1$ , the effective phase-matching condition (7) imposes  $\Delta k_z > 0$ , which corresponds to having a finite angle  $\theta$  between the pump and the probe and conjugate [21]. This occurs on the red side of the 4WM resonance, for  $\delta \lesssim 0$  (see Fig. 2). As  $\theta$  increases, the gain resonance is shifted towards higher values of  $\text{Re}(\chi_{pp})$  and therefore towards the 4WM resonance. A negative geometric phase mismatch  $\Delta k_z$  cannot be fulfilled. For this reason, no effective phase matching can happen on the blue side of the 4WM resonance, where  $n_p < 1$ .

Close to the 4WM resonance, the increase in  $\text{Im}(\chi_{pp})$  accounts for the reduction in probe power with respect to the




FIG. 5. (Color online) (a) Configuration where the geometric phase-matching condition is fulfilled ( $\Delta k_z = 0$ ); (b) configuration where the effective phase-matching condition for an effective index of refraction of the probe  $n_p \gtrsim 1$  is fulfilled. In this case, there is a necessary geometric phase mismatch ( $\Delta k_z > 0$ ). The wave vectors in vacuum  $\mathbf{k}_0$ ,  $\mathbf{k}_p$ , and  $\mathbf{k}_c$ , for the pump, probe, and conjugate, respectively, all have nearly the same magnitude. Energy conservation ensures that  $k_p + k_c = 2k_0$ .

conjugate power, seen in Fig. 4. In this region, a high level of probe absorption coupled to a large 4WM gain still produces a strong conjugate output. At larger angle  $\theta$ , the effective phase-matching condition requires a value of  $\delta$  so close to the 4WM resonance that the high probe loss prevents the 4WM from happening at all.

### III. EXPERIMENTAL VERIFICATION

In order to verify the theoretical predictions, a test was performed, as shown in Fig. 6 in which a 750 mW pump laser of beam waist 0.9 mm drives the  $D_1$  line at 795 nm in a 12-mm-long cell of  $^{85}\text{Rb}$  vapor, heated and temperature stabilized at  $\sim 110^\circ\text{C}$ . A seed beam at the probe frequency is produced by diverting a fraction of the pump through an acousto-optic modulator (AOM) operating at  $\omega_{\text{HF}}/2 \simeq 2\pi \times 1.5$  GHz in a double-pass arrangement. This seed beam, of power 10–20  $\mu\text{W}$  and waist 0.4 mm, then intersects with the pump inside the cell at a small angle  $\theta$ . The probe and conjugate beams are perpendicularly polarized with respect to the pump which is rejected at the output with a polarizing beam splitter. From




FIG. 6. (Color online) Experimental setup. The Rb cell is pumped with a bright pump beam, in blue, and seeded at an angle  $\theta$  with a probe of power  $P_s$ , in red. Emitted is an amplified probe of power  $P_p$ , also red, and a conjugate of power  $P_c$ , in yellow. The three powers,  $P_s$ ,  $P_p$ , and  $P_c$ , are monitored by photodiodes while the probe frequency is scanned. PBS: polarizing beam splitter.




FIG. 7. (Color online) Measurement of the probe gain  $g_p$  (a) and conjugate gain  $g_c$  (b) as a function of the two-photon detuning and the probe-pump angle. The region below the dashed line is used to fit the model to the data (see Fig. 8).

the measured input seed power  $P_s$ , output probe power  $P_p$ , and output conjugate power  $P_c$ , the probe and conjugate gains ( $g_p = P_p/P_s$  and  $g_c = P_c/P_s$ , respectively) are obtained.

A dichroic-atomic-vapor laser lock [22] is in place to regulate the pump frequency thus maintaining a constant  $\Delta_1$ . The angle of intersection  $\theta$  between the pump and probe is set by manually adjusting a pair of input mirrors, and  $\delta$  is adjusted by changing the AOM drive frequency. For a selection of values of  $\theta$ ,  $\delta$  is swept over a range of typically  $2\pi \times 60$  MHz, where noticeable gains  $g_p$  and  $g_c$  are observed. The angular range used, from  $0^\circ$  to  $1^\circ$ , satisfies the condition that there must be a full overlap of the beams over the cell length (this is achievable for  $\theta$  up to  $5^\circ$ ). Figure 7 shows the contour plots of  $g_p$  and  $g_c$  as a function of  $\delta$  and  $\theta$ .

The general features of Fig. 4 are reproduced, including the shift of the gain peaks towards the 4WM resonance when  $\theta$  is increased, as well as the crossover between probe power and conjugate power. The crossover is the value of  $\theta$  for which the peak conjugate power is equal to the peak probe power. The main discrepancy between the experimental data and the theoretical prediction is the measured drop in probe and conjugate power for  $\theta > 0.6^\circ$ . This leads to peak gains for both the probe and the conjugate which are well below the theoretical prediction. There are two main reasons for this behavior. First, for a finite  $\theta$ , the Doppler effect due to the thermal motion of the atoms does not cancel between the pump and the twin beams. For  $\theta = 1^\circ$ , the residual Doppler effect on  $\delta$  reaches  $2\gamma$ , which is roughly the width of the gain peak itself. This results in a broadening of the gain resonance at larger angles  $\theta$ , when compared to the resonance given by the theoretical model.

Secondly, and perhaps more importantly, when  $\theta$  is increased past  $0.5^\circ$ , the probe beam is subject to a strong effective cross-Kerr interaction with the pump around the gain resonance. This is due to the fact that as the gain peak moves closer to the 4WM resonance the effective index of the probe is resonantly enhanced, as shown by the behavior of  $\text{Re}(\chi_{pp})$  in Fig. 2. As a result, the transverse intensity variation of the pump realizes a strong lens for the probe, causing it to emerge from the cell with an angle of divergence comparable to or larger than  $\theta$  itself [23].

The good agreement between the theoretical model and the measurements at angles where the probe focusing is negligible gives us the opportunity to extract the values

of those parameters which are not readily measurable, in particular the pump Rabi frequency  $\Omega$ , the atom density  $\mathcal{N}$ , and most importantly the decoherence rate  $\gamma_c$ . Moreover, as explained in the Appendix, we can also introduce the effect of the index of refraction of the pump on the phase-matching condition by replacing the geometric phase mismatch with  $\Delta k_z = 2n_0k_0 - k_p \cos \theta - k_c \cos \theta$ , where  $n_0 = 1 - \varepsilon$  is the pump index of refraction. This index is less than 1 for the pump on the blue side of the atomic resonance. Although all these parameters enter the model in an intricate way, their respective influences on the gain map as represented in Fig. 7 follow simple trends:  $\Omega$  controls both the magnitude of the gain and the position of the gain feature along the  $\delta$  axis,  $\mathcal{N}$  controls the overall magnitude of the gain,  $\gamma_c$  controls the ratio of the probe gain over the conjugate gain, and  $\varepsilon$  controls the position of the gain feature along the  $\theta$  axis.

By fitting the model to the data for the values of  $\theta$  smaller than  $0.5^\circ$ , as shown in Fig. 8, we find that  $\Omega = 58\gamma$ ,  $\mathcal{N} = 2.6 \times 10^{12} \text{ cm}^{-3}$ , and  $\varepsilon = 8 \times 10^{-6}$ . Unlike these parameters,




FIG. 8. (Color online) Fit of the model to the experimental data of Fig. 7 at small angles, both for the probe gain  $g_p$  (a) and the conjugate gain  $g_c$  (b). The thick black lines show the theoretical model; the thin red lines show the data. The fitting procedure assumes the same set of parameters for all the curves except for the decoherence rate  $\gamma_c$  which is allowed to vary from angle to angle and is plotted in Fig. 9. The fit is of good quality except for the smallest angle, where the probe and conjugate power measurements are polluted by pump leakage through the output polarizer. For  $\theta > 0.5^\circ$ , the model predicts gains much larger than those measured.




FIG. 9. (Color online) Fitted decoherence rate  $\gamma_c$  as a function of the probe-pump angle  $\theta$ . The other parameters ( $\Omega$ ,  $\mathcal{N}$ , and  $\varepsilon$ ) have fixed values, given in the text. The uncertainties on  $\gamma_c$  are estimated from the uncertainties on these parameters.

the decoherence rate  $\gamma_c$  is allowed to depend on  $\theta$  and is plotted in Fig. 9. For this data set, the pump detuning was determined to be  $\Delta_1 = 140\gamma$  by calibrating the position of the 4WM gain against a Rb spectroscopy spectrum. We estimate the uncertainties on the values of the fitting parameters by observing how they fluctuate when fitting for the different values of  $\theta$  independently. The full relative spreads are found to be  $\pm 3\%$  for  $\Omega$ ,  $\pm 5\%$  for  $\mathcal{N}$ , and  $\pm 20\%$  for  $\varepsilon$ .

It can be seen in Fig. 9 that  $\gamma_c$  increases with  $\theta$ , under the influence of the residual Doppler broadening [24], the Kerr focusing of the probe, or possibly the transverse inhomogeneity of the pump intensity. The value of  $\gamma_c$  at small angles, below  $0.05\gamma$ , gives an indication of the intrinsic decoherence rate of the process. In our experiment the relatively low temperature leads to a low atomic density and therefore to negligible collisional decoherence. The measurable decoherence rate is most likely limited by the finite transit time of the atoms through the beams. The theoretical estimate of 500 kHz reported in Ref. [13] is compatible with our inferred  $\gamma_c$ , given that our beams are 50% wider. Note that the predominance of finite transit time over atomic collisions in the reduction of ground state coherence depends on the specific experimental conditions, in particular, the temperature. In typical squeezing experiments, e.g., in Ref. [7], the gain and the squeezing drop for temperatures larger than  $120^\circ\text{C}$ . This is compatible with the observation by Siddons *et al.* [25] that collision-driven decoherence dominates at those temperatures.

One can check that the parameters extracted from the fit are broadly consistent with the estimated experimental conditions. Our pump beam parameters lead to a peak intensity of  $60 \text{ W cm}^{-2}$ , which results in a resonant Rabi frequency  $\Omega = 80\gamma$  for a mean electric dipole  $d = 1.47 \times 10^{-29} \text{ C m}$  [26]. From the vapor pressure data summarized in Ref. [26], the number density of  $^{85}\text{Rb}$  at  $100^\circ\text{C}$  is  $\mathcal{N} \simeq 4 \times 10^{12} \text{ cm}^{-3}$ . The estimate on  $\mathcal{N}$  is relatively poor because unlike in more direct and accurate methods [25], we do not have a good knowledge of the average electric dipole moments involved in the 4WM. Finally the index of refraction for the pump, evaluated for ground state populations of 6% in the lower hyperfine state and 94% in the upper hyperfine state, as given




FIG. 10. (Color online) Probe gain as a function of the two-photon detuning at low angle  $\theta$  and fit to the data including the Doppler broadening in the model. The discrepancy in the width of the Doppler-broadened single-photon absorption dip is caused by the presence of two excited hyperfine levels separated by  $60\gamma$ . The model only considers two excited states which are degenerate in energy (Fig. 11).

by Eqs. (A17)–(A20), and for the electric dipole value given above, is  $n_0 = 1 - 1.6 \times 10^{-5}$ .

#### IV. DOPPLER BROADENING

The model developed above does not directly take into account the Doppler broadening caused by the thermal motion of the atoms in the cell. The good agreement between the model and the experimental data suggests that considering only average values of the single-photon detunings  $\Delta_1$  and  $\Delta_2$  captures most of the physics at play. However, considering that the probe is typically tuned to the edge of the Doppler profile, it is legitimate to wonder what level of absorption this causes. It turns out that although the EIT has a limited impact, the pump field is highly saturating at resonance and causes a wide Autler-Townes splitting for those atoms resonant with the probe field. This renders even the resonant part of the atomic vapor highly transparent for the probe as long as the optical depth is not too large. In practice, noticeable levels of squeezing can be observed for  $\Delta_1/2\pi$  as low as 500 MHz [7], which is well inside the Doppler profile at our operational temperature.

In order to verify these assumptions, we extended the above model to include the full Doppler distribution of detunings. The result, fitted to the gain curve of the probe as a function of  $\delta$ , is shown in Fig. 10. The small discrepancy in the width of the single-photon resonance is due to the fact that the model does not include the hyperfine structure of the excited state, whose main effect is to broaden the apparent Doppler profile.

#### V. OPTIMIZING FOR QUANTUM NOISE REDUCTION

The observation of large levels of intensity-difference quantum noise reduction requires a near-perfect phase-insensitive amplifier with a gain of at least a few units. In the case of 4WM in a hot atomic vapor, this means that the absorption of the twin beams must be kept to a minimum while ensuring

that efficient 4WM can take place. As pointed out previously, the two-photon detuning which fulfills those two conditions is  $\delta \simeq 0$ , and not the 4WM resonance in itself. Firstly, this is because away from the 4WM resonance the probe susceptibility responsible for the absorption drops faster than the cross susceptibility responsible for the 4WM gain, and secondly because the ground state coherence reduces the probe absorption at that detuning [27].

From the previous discussion, at a detuning  $\delta \simeq 0$ , effective phase matching of the 4WM process requires geometric phase matching  $\Delta k_z = 0$ . When corrected for the index of refraction for the pump, this condition corresponds to the introduction of a small angle  $\theta \ll 1^\circ$  between the pump and the probe beams. This is indeed how the best levels of squeezing have been experimentally observed [7–10]. This also justifies the implicit geometric phase-matching condition used in previous theoretical studies [13]. The production of squeezed light by nondegenerate 4WM in a hot vapor therefore benefits from two favorable circumstances. First, the optimum angle  $\theta$  is small enough that the residual Doppler effect acting on the nearly copropagating beams is much smaller than the width of the gain peak. Second, the same angle is large enough that the beams participating in the 4WM can be spatially separated at the output of the vapor cell.

It is worth noting that in certain conditions, a small amount of loss on the probe beam can be beneficial. For instance, for a probe seed containing a large amount of classical noise, it is useful to have twin beams of equal powers in order to ensure proper rejection of the classical noise in the balanced detection [7,28]. In particular, the existence of points in the  $(\delta, \theta)$  parameter space where the probe and conjugate powers are perfectly balanced has allowed the detection of 8 dB of intensity-difference squeezing at frequencies as low as 2.5 kHz, despite the presence of substantial technical noise on the input probe [7].

#### VI. CONCLUSION

We have shown that it is possible to phase match 4WM in a hot atomic vapor so that absorption is reduced to a level where quantum effects could manifest themselves. Most reports of large squeezing generated by nondegenerate forward 4WM in hot vapors to date have indeed used a similar arrangement as the one presented here, with nearly identical atomic and beam parameters. Furthermore, the model we have developed appears very accurate for small angles between beams, even in its simple form neglecting both the Doppler effect and the Zeeman substructure. It allows the robust extraction of parameters which would be difficult to determine by direct measurement, such as the decoherence rate of the ground states.

By providing further insight into the mechanism of the process, the present results may lead to the realization of different configurations or different regimes amenable to the production of interesting quantum states of light, such as multi-spatial-mode phase-sensitive amplifiers [29].

#### ACKNOWLEDGMENTS

We acknowledge useful discussions with Colin McCormick, Ennio Arimondo, and Paul Lett, and we thank

Etienne Pertreux for his help in the early stages of this project. This research was supported by the Engineering and Physical Sciences Research Council Grants No. EP/E036473/1 and No. EP/I001743/1.

## APPENDIX: EXPRESSIONS FOR THE SUSCEPTIBILITIES

### 1. Derivation of the susceptibilities for the probe and the conjugate

In this Appendix we derive the dynamics of the double- $\Lambda$  configuration described in the paper. The response of an atomic system to an optical field is determined by the polarization of the medium, which acts as the driving term in the wave equation. For a medium that consists of noninteracting particles, such as a dilute atomic vapor, the polarization of the medium is of the form [14]

$$\mathbf{P} = \mathcal{N} \langle \hat{\mathbf{d}} \rangle,$$

where  $\mathcal{N}$  is the number density of the atomic medium and  $\hat{\mathbf{d}}$  is the atomic dipole moment operator. The polarization of the medium can be written in terms of the atomic eigenstates, such that it takes the form

$$\mathbf{P} = \mathcal{N} \sum_{n,m} \mathbf{d}_{mn} \sigma_{nm} e^{-i\omega_{f,nm}t},$$

where the sum is over all the involved atomic transitions,  $\omega_{f,nm}$  is the frequency of the field that couples the transition between levels  $n$  and  $m$ ,  $\sigma_{nm}$  is the density matrix element between levels  $n$  and  $m$  in a rotating frame at frequency  $\omega_{f,nm}$ , and  $\mathbf{d}_{mn}$  is the dipole matrix element between levels  $n$  and  $m$ . For an isotropic medium, the polarization of the atomic medium at a particular frequency is given by

$$\mathbf{P}(\omega_{f,nm}) = \mathcal{N} \mathbf{d}_{mn} \sigma_{nm}.$$

Thus, the response is completely determined by the atomic coherence of the corresponding transition.

The equations of motion for the density matrix elements in the rotating frame can be shown to be of the form [14]

$$\begin{aligned} \dot{\sigma}_{nm} &= i(\Delta_{nm} - \gamma_{nm})\sigma_{nm} \\ &+ \frac{i}{\hbar} \sum_v [\mathbf{d}_{nv} \cdot \mathbf{E}(\mathbf{r},t) \sigma_{vm} e^{-i(\omega_{f,vm} - \omega_{f,nm})t} \\ &- \sigma_{nv} \mathbf{d}_{vm} \cdot \mathbf{E}(\mathbf{r},t) e^{-i(\omega_{f,nv} - \omega_{f,nm})t}] \quad \text{for } n \neq m, \\ \dot{\sigma}_{nn} &= \frac{i}{\hbar} \sum_v [\mathbf{d}_{nv} \cdot \mathbf{E}(\mathbf{r},t) \sigma_{vn} e^{-i\omega_{f,nn}t} \\ &- \sigma_{nv} \mathbf{d}_{vn} \cdot \mathbf{E}(\mathbf{r},t) e^{-i\omega_{f,nn}t}] \\ &+ \sum_{E_m > E_n} \Gamma_{nm} \sigma_{mm} - \sum_{E_m < E_n} \Gamma_{mn} \sigma_{nn}, \end{aligned}$$

where  $\Delta_{nm}$  is the detuning of the field at frequency  $\omega_{f,nm}$  from the transition between levels  $n$  and  $m$ ,  $\Gamma_{mn}$  is the population decay rate from level  $n$  to level  $m$ ,  $\gamma_{nm} = (\Gamma_n + \Gamma_m)/2 + \gamma_{nm}^c$  is the dipole dephasing rate,  $\Gamma_n$  is the total decay rate out of level  $n$ , and  $\gamma_{nm}^c$  is the dipole dephasing rate due to any other source of decoherence.

We now specialize the above equations to our 4WM process in the double- $\Lambda$  configuration, shown in Figs. 1 and 11, with a single pump field,  $\mathcal{E}_0$ . For this case the total electric field is




FIG. 11. (Color online) Double- $\Lambda$  scheme with a single pump field  $\mathcal{E}_0$ . States |3> and |4> are orthogonal linear combinations of magnetic states of the excited hyperfine levels.

of the form

$$\begin{aligned} \mathbf{E}(\mathbf{r},t) &= \mathcal{E}_0 e^{i(\mathbf{k}_0 \cdot \mathbf{r} - \omega_0 t)} \boldsymbol{\epsilon}_0 + \mathcal{E}_c e^{i(\mathbf{k}_c \cdot \mathbf{r} - \omega_c t)} \boldsymbol{\epsilon}_c \\ &+ \mathcal{E}_p e^{i(\mathbf{k}_p \cdot \mathbf{r} - \omega_p t)} \boldsymbol{\epsilon}_p + \text{c.c.}, \end{aligned}$$

where  $\mathcal{E}_0$ ,  $\mathcal{E}_p$ , and  $\mathcal{E}_c$  are the field amplitudes for the pump, the probe, and the conjugate, respectively, and  $\boldsymbol{\epsilon}_i$  are unit vectors describing the polarization of the fields. We assume that the pump couples the transitions  $|1\rangle \rightarrow |3\rangle$  and  $|2\rangle \rightarrow |4\rangle$ , the probe couples transition  $|2\rangle \rightarrow |3\rangle$ , the conjugate couples transition  $|1\rangle \rightarrow |4\rangle$ , and that the transitions  $|1\rangle \rightarrow |2\rangle$  and  $|3\rangle \rightarrow |4\rangle$  are not dipole allowed. With these assumptions and using the rotating-wave approximation, the equations of motion for the density matrix elements take the form

$$\begin{aligned} \dot{\sigma}_{11} &= \frac{i}{2} (\Omega_1^* e^{-i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{31} + \Omega_p^* e^{-i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{41} - \Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{13} \\ &- \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{14}) + \Gamma_{13} \sigma_{33} + \Gamma_{14} \sigma_{44}, \end{aligned} \quad (\text{A1})$$

$$\begin{aligned} \dot{\sigma}_{22} &= \frac{i}{2} (\Omega_c^* e^{-i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{32} + \Omega_2^* e^{-i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{42} - \Omega_c e^{i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{23} \\ &- \Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{24}) + \Gamma_{23} \sigma_{33} + \Gamma_{24} \sigma_{44}, \end{aligned} \quad (\text{A2})$$

$$\begin{aligned} \dot{\sigma}_{33} &= \frac{i}{2} (\Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{13} + \Omega_c e^{i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{23} - \Omega_1^* e^{-i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{31} \\ &- \Omega_c^* e^{-i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{32}) - \Gamma_3 \sigma_{33}, \end{aligned} \quad (\text{A3})$$

$$\begin{aligned} \dot{\sigma}_{44} &= \frac{i}{2} (\Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{14} + \Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{24} - \Omega_p^* e^{-i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{41} \\ &- \Omega_2^* e^{-i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{42}) - \Gamma_4 \sigma_{44}, \end{aligned} \quad (\text{A4})$$

$$\begin{aligned} \dot{\sigma}_{43} &= \frac{i}{2} (\Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{23} + \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{13} - \Omega_1^* e^{-i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{41} \\ &- \Omega_c^* e^{-i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{42}) + (i\Delta_2 - i\Delta_1 - \gamma_{43}) \sigma_{43}, \end{aligned} \quad (\text{A5})$$

$$\begin{aligned} \dot{\sigma}_{42} &= \frac{i}{2} (\Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{22} + \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{12} - \Omega_c e^{i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{43} \\ &- \Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{44}) + (i\Delta_2 - i\delta - \gamma_{42}) \sigma_{42}, \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} \dot{\sigma}_{41} &= \frac{i}{2} (\Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{21} + \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{11} - \Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{43} \\ &- \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{44}) + (i\Delta_2 - \gamma_{43}) \sigma_{41}, \end{aligned} \quad (\text{A7})$$



$$\dot{\sigma}_{32} = \frac{i}{2}(\Omega_c e^{i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{22} + \Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{12} - \Omega_c e^{i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{32} - \Omega_2 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{34}) + (i\Delta_1 - i\delta - \gamma_{32})\sigma_{32}, \quad (\text{A8})$$

$$\dot{\sigma}_{31} = \frac{i}{2}(\Omega_c e^{i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{21} + \Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{11} - \Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{33} - \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{34}) + (i\Delta_1 - \gamma_{31})\sigma_{31}, \quad (\text{A9})$$

$$\dot{\sigma}_{21} = \frac{i}{2}(\Omega_c^* e^{-i\mathbf{k}_c \cdot \mathbf{r}} \sigma_{31} + \Omega_2^* e^{-i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{41} - \Omega_1 e^{i\mathbf{k}_0 \cdot \mathbf{r}} \sigma_{23} - \Omega_p e^{i\mathbf{k}_p \cdot \mathbf{r}} \sigma_{24}) + (i\delta - \gamma_{21})\sigma_{21}, \quad (\text{A10})$$

where  $\Omega_1$  and  $\Omega_2$  give the Rabi frequencies for the two transitions coupled by the single pump,  $\Omega_p$  gives the Rabi frequency for the transition coupled by the probe, and  $\Omega_c$  gives the Rabi frequency for the transition coupled by the conjugate. In order to completely eliminate the explicit time dependence when doing the rotating-wave approximation, the frequencies of the pump, probe, and conjugate fields need to satisfy the relation

$$2\omega_0 = \omega_p + \omega_c,$$

which is just the energy conservation condition for the 4WM process.

In order to solve the above equations and obtain analytical expressions, we assume that the probe and conjugate fields are weak fields, such that we only keep terms to first order in  $\Omega_p$  and  $\Omega_c$ . In this case, the polarization of the medium at frequency  $\omega_i$  (where  $i$  indicates probe or conjugate frequency) can be divided into two different terms, one that is proportional to the field at frequency  $\omega_i$  and one that is proportional to the field at frequency  $2\omega_0 - \omega_i$ , such that

$$P(\omega_p) = \epsilon_0 \chi_{pp}(\omega_p) \mathcal{E}_p e^{i\mathbf{k}_p \cdot \mathbf{r}} + \epsilon_0 \chi_{pc}(\omega_p) \mathcal{E}_c^* e^{i(2\mathbf{k}_0 - \mathbf{k}_c) \cdot \mathbf{r}}, \quad (\text{A11})$$

$$P(\omega_c) = \epsilon_0 \chi_{cc}(\omega_c) \mathcal{E}_c e^{i\mathbf{k}_c \cdot \mathbf{r}} + \epsilon_0 \chi_{cp}(\omega_c) \mathcal{E}_p^* e^{i(2\mathbf{k}_0 - \mathbf{k}_p) \cdot \mathbf{r}}. \quad (\text{A12})$$

In doing this, we have introduced the susceptibility of the atomic medium  $\chi_{ij}$ , which completely characterizes the response of the atomic system for a given field. The direct susceptibilities  $\chi_{pp,cc}$  act as the effective linear susceptibilities for the probe and conjugate, respectively; the cross susceptibilities  $\chi_{pc,cp}$  are responsible for the 4WM process.

In addition to the approximations mentioned above, we assume the dipole moments for the two pump transitions to be equal, such that  $\Omega_1 = \Omega_2 \equiv \Omega$ . Under these approximations we can solve the density matrix equations (A1)–(A10) to all orders in the pump field ( $\Omega$ ) and in steady-state condition for  $\sigma_{41}$  and  $\sigma_{32}$  to find that

$$\chi_{pp} = \frac{i\mathcal{N}|d_{23}|^2 \xi_{41}^*}{\epsilon_0 \hbar D^*} \left[ \frac{\xi_{21}^*}{\xi_{42}^*} \sigma_{22,44} + \frac{\xi_{43}^*}{\xi_{31}^*} \sigma_{11,33} - \left( \frac{\xi_{21}^* + \xi_{43}^*}{\xi_{41}^*} + \frac{\xi_{21}^* \xi_{43}^*}{|\Omega|^2/4} \right) \sigma_{22,33} \right], \quad (\text{A13})$$

$$\chi_{cc} = \frac{i\mathcal{N}|d_{14}|^2 \xi_{32}^*}{\epsilon_0 \hbar D} \left[ \frac{\xi_{43}}{\xi_{42}^*} \sigma_{22,44} + \frac{\xi_{21}}{\xi_{31}^*} \sigma_{11,33} - \left( \frac{\xi_{21} + \xi_{43}}{\xi_{32}^*} + \frac{\xi_{21} \xi_{43}}{|\Omega|^2/4} \right) \sigma_{11,44} \right], \quad (\text{A14})$$

$$\chi_{pc} = \frac{i\mathcal{N}d_{14}d_{23}\xi_{41}^*\Omega^2}{\epsilon_0 \hbar D^* |\Omega|^2} \left[ \frac{\xi_{21}^*}{\xi_{31}} \sigma_{11,33} + \frac{\xi_{43}^*}{\xi_{42}} \sigma_{22,44} + \left( \frac{\xi_{21}^* + \xi_{43}^*}{\xi_{41}^*} \right) \sigma_{11,44} \right], \quad (\text{A15})$$

$$\chi_{cp} = \frac{i\mathcal{N}d_{14}d_{23}\xi_{32}^*\Omega^2}{\epsilon_0 \hbar D |\Omega|^2} \left[ \frac{\xi_{43}}{\xi_{31}} \sigma_{11,33} + \frac{\xi_{21}}{\xi_{42}} \sigma_{22,44} + \left( \frac{\xi_{21} + \xi_{43}}{\xi_{32}^*} \right) \sigma_{22,33} \right], \quad (\text{A16})$$

where we have defined

$$D = (\xi_{43} + \xi_{21})(\xi_{32}^* + \xi_{41}) + \frac{\xi_{32}^* \xi_{41} \xi_{43} \xi_{21}}{|\Omega|^2/4},$$

the population differences

$$\sigma_{11,33} \equiv \sigma_{11} - \sigma_{33} = \frac{|\xi_{31}|^2}{|\Omega|^2 + |\xi_{31}|^2 + |\xi_{42}|^2}, \quad (\text{A17})$$

$$\sigma_{11,44} \equiv \sigma_{11} - \sigma_{44} = \frac{|\xi_{31}|^2}{|\Omega|^2 + |\xi_{31}|^2 + |\xi_{42}|^2}, \quad (\text{A18})$$

$$\sigma_{22,33} \equiv \sigma_{22} - \sigma_{33} = \frac{|\xi_{42}|^2}{|\Omega|^2 + |\xi_{31}|^2 + |\xi_{42}|^2}, \quad (\text{A19})$$

$$\sigma_{22,44} \equiv \sigma_{22} - \sigma_{44} = \frac{|\xi_{42}|^2}{|\Omega|^2 + |\xi_{31}|^2 + |\xi_{42}|^2}, \quad (\text{A20})$$

and the complex decay rates

$$\xi_{43} = i(\Delta_2 - \Delta_1) - \gamma, \quad \xi_{42} = i(\Delta_2 - \delta) - \frac{\gamma}{2},$$

$$\xi_{41} = i\Delta_2 - \frac{\gamma}{2}, \quad \xi_{32} = i(\Delta_1 - \delta) - \frac{\gamma}{2},$$

$$\xi_{31} = i\Delta_1 - \frac{\gamma}{2}, \quad \xi_{21} = i\delta - \gamma_c.$$

In deriving these equations, we have assumed that the total decay rate out of the excited states are the same and with equal branching ratios to the two ground states; that is,  $\Gamma_4 = \Gamma_3 \equiv \gamma$  and  $\Gamma_{14} = \Gamma_{24} = \Gamma_{13} = \Gamma_{14} = \gamma/2$ . In addition, we assume that the additional dipole dephasing term  $\gamma_{ij}^c$  is only significant for the ground state coherence, such that  $\gamma_{ij \neq 12}^c = 0$  and  $\gamma_{12}^c \equiv \gamma_c$ .

It is notable that within the approximation of first-order expansion of the susceptibilities in  $\Omega_p$  and  $\Omega_c$ , the atomic populations in Eqs. (A17)–(A20) only depend on the pump Rabi frequency. In other words, the atomic dynamics is dominated by the pumping mechanism and is independent from the level of seeding. Experimentally we have verified that gain and squeezing levels are indeed independent from the input probe power up to 200  $\mu\text{W}$ , only limited by the amount of seed optical power that was experimentally available.

## 2. Index of refraction for the pump

In the above derivation of the susceptibilities for the probe and the conjugate, we have implicitly set  $\mathcal{E}_0$ , the amplitude for the pump electric field, to be a constant throughout the medium. This assumes that the pump is neither dephased nor

absorbed. In practice, the pump detuning is large enough to neglect absorption, but even a small index of refraction may have a substantial impact on the phase-matching condition of the 4WM process. The effect of refraction on the pump is to multiply  $\mathcal{E}_0$  (and therefore  $\Omega$ ) by a running phase factor as the pump propagates. This can be taken into account by simply replacing  $\mathbf{k}_0$  in  $\Delta\mathbf{k}$  by  $n_0\mathbf{k}_0$ , where  $n_0$  is the index of refraction. The index of refraction created by the population  $\mathcal{N}_i$  of the ground state

$i$  can be estimated to be  $n_0 = \sqrt{1 + \chi} \simeq 1 + \chi/2$  with

$$\chi = -\frac{\mathcal{N}_i d^2}{\varepsilon_0 \hbar} \frac{\Delta_i}{\Delta_i^2 + \gamma^2/4},$$

where  $d$  is the dipole matrix element of the transition for large detunings and  $\Delta_i$  is the detuning, which is taken to be much larger than the hyperfine splitting of the excited state.

- 
- [1] M. D. Reid, *Phys. Rev. A* **40**, 913 (1989).
- [2] A. Heidmann, R. J. Horowicz, S. Reynaud, E. Giacobino, C. Fabre, and G. Camy, *Phys. Rev. Lett.* **59**, 2555 (1987).
- [3] Z. Y. Ou, S. F. Pereira, H. J. Kimble, and K. C. Peng, *Phys. Rev. Lett.* **68**, 3663 (1992).
- [4] C. Silberhorn, P. K. Lam, O. Weiß, F. König, N. Korolkova, and G. Leuchs, *Phys. Rev. Lett.* **86**, 4267 (2001).
- [5] P. R. Hemmer, D. P. Katz, J. Donoghue, M. Cronin-Golomb, M. S. Shahriar, and P. Kumar, *Opt. Lett.* **20**, 982 (1995).
- [6] M. D. Lukin, P. R. Hemmer, M. Löffler, and M. O. Scully, *Phys. Rev. Lett.* **81**, 2675 (1998); M. D. Lukin, P. R. Hemmer, and M. O. Scully, in *Advances in Atomic Molecular, and Optical Physics*, edited by B. Bederson and H. Walther (Elsevier Academic Press Inc., San Diego, 2000), Vol. 42, pp. 347–386.
- [7] C. F. McCormick, A. M. Marino, V. Boyer, and P. D. Lett, *Phys. Rev. A* **78**, 043816 (2008).
- [8] Q. Glorieux, L. Guidoni, S. Guibal, J.-P. Likforman, and T. Coudreau, *Phys. Rev. A* **84**, 053826 (2011).
- [9] M. Jasperse, L. D. Turner, and R. E. Scholten, *Opt. Express* **19**, 3765 (2011).
- [10] Z. Qin, J. Jing, J. Zhou, C. Liu, R. C. Pooser, Z. Zhou, and W. Zhang, *Opt. Lett.* **37**, 3141 (2012).
- [11] M. D. Reid, D. F. Walls, and B. J. Dalton, *Phys. Rev. Lett.* **55**, 1288 (1985).
- [12] R. C. Pooser, A. M. Marino, V. Boyer, K. M. Jones, and P. D. Lett, *Opt. Express* **17**, 16722 (2009).
- [13] Q. Glorieux, R. Dubessy, S. Guibal, L. Guidoni, J.-P. Likforman, T. Coudreau, and E. Arimondo, *Phys. Rev. A* **82**, 033819 (2010).
- [14] R. W. Boyd, *Nonlinear Optics* (Academic, New York, 2008).
- [15] C. F. McCormick, V. Boyer, E. Arimondo, and P. D. Lett, *Opt. Lett.* **32**, 178 (2007).
- [16] V. Boyer, A. M. Marino, R. C. Pooser, and P. D. Lett, *Science* **321**, 544 (2008).
- [17] R. C. Pooser, A. M. Marino, V. Boyer, K. M. Jones, and P. D. Lett, *Phys. Rev. Lett.* **103**, 010501 (2009).
- [18] K.-J. Boller, A. Imamoglu, and S. E. Harris, *Phys. Rev. Lett.* **66**, 2593 (1991).
- [19] C. H. van der Wal, M. D. Eisaman, A. André, R. L. Walsworth, D. F. Phillips, A. S. Zibrov, and M. D. Lukin, *Science* **301**, 196 (2003).
- [20] V. Boyer, C. F. McCormick, E. Arimondo, and P. D. Lett, *Phys. Rev. Lett.* **99**, 143601 (2007).
- [21] As the angle between the pump and the probe,  $\theta$ , is small and  $n_p$  is close to 1, the angle between the pump and the conjugate is very close to  $\theta$ .
- [22] K. L. Corwin, Z.-T. Lu, C. F. Hand, R. J. Epstein, and C. E. Wieman, *Appl. Opt.* **37**, 3295 (1998).
- [23] The full Kerr action of the pump on the probe is the compound effect of the change of magnitude of the index of refraction at resonance and the frequency shift of the resonance due to the light shift caused by the pump.
- [24] P. R. S. Carvalho, L. E. E. de Araujo, and J. W. R. Tabosa, *Phys. Rev. A* **70**, 063818 (2004).
- [25] P. Siddons, C. S. Adams, C. Ge, and I. G. Hughes, *J. Phys. B* **41**, 155004 (2008).
- [26] D. A. Steck, Rubidium 85 D Line Data, <http://steck.us/alkalidata> (revision 2.1.5).
- [27] It is not exactly  $\delta = 0$  due to the light shift created by the pump beam on the  $5S_{1/2}(F = 3) \rightarrow 5P_{1/2}$  transition.
- [28] We consider here only the case where the detector is perfectly balanced, with equal electronic gains on both inputs.
- [29] N. Corzo, A. M. Marino, K. M. Jones, and P. D. Lett, *Opt. Express* **19**, 21358 (2011); N. V. Corzo, A. M. Marino, K. M. Jones, and P. D. Lett, *Phys. Rev. Lett.* **109**, 043602 (2012).