

Angular and polarization analysis for two-photon decay of $2s$ hyperfine states of hydrogenlike uranium

L. Safari,^{1,*} P. Amaro,² J. P. Santos,² and F. Fratini^{1,3,4}¹*Department of Physics, University of Oulu, Fin-90014 Oulu, Finland*²*Centro de Física Atômica, Departamento de Física, Faculdade de Ciências e Tecnologia, Fundação para a Ciência e a Tecnologia, Universidade Nova de Lisboa, P-2829-516 Caparica, Portugal*³*Departamento de Física, Universidade Federal de Minas Gerais, 31270-901 Belo Horizonte, MG, Brazil*⁴*Institut Néel, Centre National de la Recherche Scientifique, BP 166, 25 rue des Martyrs, 38042 Grenoble Cedex 9, France*

(Received 8 April 2014; published 29 July 2014)

The amplitude of two-photon transitions between hyperfine states in hydrogenlike ions is derived based on the relativistic Dirac equation and second-order perturbation theory. We study angular and linear polarization properties of the photon pair emitted in the decay of $2s$ states, where spin-flip and non-spin-flip transitions are highlighted. We pay particular attention to hydrogenlike uranium, since it is an ideal candidate for investigating relativistic and high-multipole effects, such as spin-flip transitions. Two types of emission patterns are identified: (i) non-spin-flip transitions are found to be characterized by an angular distribution of the type $W(\theta) \sim 1 + \cos^2 \theta$ while the polarizations of the emitted photons are parallel; and (ii) spin-flip transitions have somewhat smaller decay rates and are found to be characterized by an angular distribution of the type $W(\theta) \sim 1 - 1/3 \cos^2 \theta$ while the polarizations of the emitted photons are orthogonal, where θ is the angle between photons directions. Deviations due to nondipole and relativistic contributions are evaluated for both types of transitions. This work is the first step toward exploring the effect of the nucleus over the angular and polarization properties of the photon pairs emitted by two-photon transitions.

DOI: [10.1103/PhysRevA.90.014502](https://doi.org/10.1103/PhysRevA.90.014502)

PACS number(s): 32.70.Cs, 32.70.Jz, 32.80.Wr, 31.10.+z

I. INTRODUCTION

Two-photon decay in atoms and ions was introduced by Max Born's Ph.D. student Goepfert-Mayer in 1931 [1]. Since then, many aspects of such a process, like the total decay rate and the spectral distribution, have been extensively investigated in the context of few-electron atoms and ions, both in theory and experiments [2–12]. Recently, some interest has been also devoted to the relativistic effects on angular and polarization properties of the two emitted photons [13–19] and to electron-electron interaction effects on the total decay rate [20]. Apart from fundamental interest, two-photon transitions revealed themselves as a useful tool for investigation of different physical areas and applied science. Already in 1940, for instance, Breit and Teller derived that the double photon emission was the principal cause of the decay of interstellar hydrogen atoms from their metastable $2s$ state [21], while, more recently, polarization properties of the emitted photons have been employed to successfully explore quantum entanglement [22–24]. Furthermore, two-photon transitions have been proposed as a tool to measure weak-interaction properties [25,26].

In this Brief Report, the angular and polarization properties of the photon pair emitted by the two-photon decay of $2s$ hyperfine states in hydrogenlike ions are presented. Particular attention is paid to hydrogenlike uranium (U^{91+}). In fact, due to its strong electromagnetic field, high multipoles contributions that lead to spin-flip transitions are enhanced in this system. This makes hydrogenlike uranium an ideal candidate for our studies. The angular and polarization analysis of the emitted light is carried out within the independent particle approximation (IPA), i.e., by coupling the spin-angular

momenta of electron and nucleus and by neglecting any hyperfine interaction between electron and nucleus. We derive the analytical expression for the transitions amplitude within IPA. This work is a first step toward exploring the effect of the nuclear angular momentum (spin) on the angular and polarization properties of the emitted photons. Our analysis may pave the way for a new route to get information on the direction and the magnitude of the spin distribution inside the nucleus (which is still quite an unraveled problem) by using two-photon angular and polarization correlations.

II. THEORETICAL BACKGROUND

A. Construction of the overall set of states

The presence of the nuclear spin has a twofold effect on the states of hydrogenlike systems. First, the energies of the atomic metastable states are slightly shifted, mainly due to the magnetic dipole interaction that the nucleus and electron experience. This energy correction can be described by using first-order perturbation theory with additional contributions, such as the relativistic, Bohr-Weisskopf, Breit-Rosenthal, and QED contributions [27,28]. Since this energy correction does not influence the angular and polarization properties of the emitted radiation, it will be totally neglected in the following. Second, the atomic states acquire a new quantum number, usually denoted by F , that represents the total angular momentum of the overall–nucleus plus electron–system.

The overall atomic state can be described by coupling the nucleus and electron angular momenta (referred to as IPA), i.e., by

$$|n, \beta; F, I, \kappa, m_F\rangle = \sum_{m_I, m_J} \langle j, m_J, I, m_I | F, m_F \rangle |n; \kappa, m_J\rangle |\beta; I, m_I\rangle, \quad (1)$$

*laleh.safari@oulu.fi

where n , κ , and j are the (Bohr) principal, the Dirac, and the angular momentum quantum numbers of the electron, respectively, while I represents the nuclear spin. On the other hand, m_I , m_j , and m_F are the projections of the nuclear, electronic, and total (nucleus plus electron) angular momenta onto the quantization axis, respectively. Finally, β is a collective label that denotes any other quantum number needed to specify the nuclear state apart from I and m_I . Using standard notation, $\langle j, m_j, I, m_I | F, m_F \rangle$ are Clebsch-Gordan coefficients.

To further proceed, we suppose that the nucleus does not interact with the radiation field. In the language of quantum mechanics, this equates to considering that the interaction Hamiltonian couples only electron fields through the photon emission, while it does not act on the quantum space of nuclear states. This hypothesis holds for decays which involve bound states of neutral atoms, since the energy released in such decays is far lower than the nuclear excitation energies (there are few exceptions to this, like the nucleus of Th^{229} , where the first metastable excited state is $\lesssim 10$ eV above the ground state). For highly charged ions, on the other hand, nuclear excitations are of the order of $\sim \text{MeV}$ while photon energies can take values up to hundreds of keV. Equation (1) represents, therefore, only a rough approximation for such systems [29]. As a result of this assumption, we shall find in the next subsection that the radial part of

the decay amplitude is characterized by only electron state components. On the other hand, we shall see that the angular part of the decay amplitude is characterized by both electron and nucleus states components, due to the coupling of their angular momenta. We shall see that the value for the total spin quantum number will directly determine the shape of the angular and polarization distributions in the atomic transitions.

B. Second-order transition amplitude

The theory of two-photon decay is based on the second-order transition amplitude and has been discussed in a number of recent papers [17,18,26]. One of the characteristic features of such amplitude is that it contains a summation over the intermediate atomic states which runs through the whole atomic spectrum, including a summation over the discrete part as well as an integration over the (positive and negative) continuum. For the problem under consideration, such a summation splits up into summations over (i) the principal quantum number n_ν , (ii) the Dirac quantum number κ_ν , (iii) the total angular momentum F_ν , and (iv) its projection onto the quantization axis m_{F_ν} .

By using Eq. (1) and by taking into account the orthonormality of the nuclear states, the amplitude for two-photon transitions between hyperfine states takes the form

$$\begin{aligned} \mathcal{M}^{\lambda_1 \lambda_2}(i \rightarrow f) = & -(2\pi) \sum_{T T'} \sum_{\substack{\kappa_\nu \\ m_I m_{j\nu}}} \sum_{\substack{L_1 L_2 \\ M_1 M_2}} \sum_{p_1} \sum_{p_2 \Delta_1 \Delta_2} (\lambda_1)^{p_1} (\lambda_2)^{p_2} [L_1, L_2]^{1/2} i^{-L_1 - L_2 - p_1 - p_2} \xi_{L_1 \Lambda_1}^{p_1} \xi_{L_2 \Lambda_2}^{p_2} P^T P^{T'} \\ & \times D_{M_2 \lambda_2}^{L_2*}(\varphi_2, \theta_2, 0) D_{M_1 \lambda_1}^{L_1*}(\varphi_1, \theta_1, 0) [U_{\Lambda_1 \Lambda_2}^{T T'} \chi_{m_I m_{j\nu}}^{f T v T'} \chi_{m_I m_{j\nu}}^{v T' i T'} + (1 \leftrightarrow 2)], \end{aligned} \quad (2)$$

where λ_j and \mathbf{k}_j are the helicity and wave vector of the j th photon. The term $D_{M_j \lambda_j}^{L_j}(\theta_j, \varphi_j)$ stands for the Wigner rotation matrices of order L_j with angle coordinates (θ_j, φ_j) . The notation $[L]$ stands for $2L + 1$ and Λ_j runs from $L_j - 1$ to $L_j + 1$. $T, T' = L, S$ denote the large (L) and small (S) components of the electron Dirac spinor, for which the factor P^T is defined as $P^L = 1$ and $P^S = -1$. Furthermore, $p_{1,2} = 0, 1$ and the function $\xi_{L \Lambda}^p$ is given by

$$\xi_{L \Lambda}^0 = \delta_{L, \Lambda}, \quad \xi_{L \Lambda}^1 = \begin{cases} \sqrt{\frac{L+1}{2L+1}} & \text{for } \Lambda = L - 1 \\ -\sqrt{\frac{L}{2L+1}} & \text{for } \Lambda = L + 1 \\ 0 & \text{otherwise} \end{cases} \quad (3)$$

The radial part of the amplitude in Eq. (2) is represented by the integral $U_{\Lambda_1 \Lambda_2}^{T T'}$, which reads

$$U_{\Lambda_1 \Lambda_2}^{T T'} = \int dr dr' r'^2 r'^2 j_{\Lambda_1}(k_1 r') j_{\Lambda_2}(k_2 r) g_f^{\bar{T} *} g_{E_i + \omega_1}^{T \bar{T}'} g_i^{T'}, \quad (4)$$

where $g_{f,i}^T$ are the small and large radial components of the final and initial electron state, while

$$g_{E_i + \omega_1}^{T \bar{T}'} = \sum_{n_\nu} \frac{g_\nu^T g_\nu^{\bar{T}' *}}{E_\nu - E_i - \omega_1} \quad (5)$$

is the radial Green's function of the process. Here $E_{i,\nu}$ are the energies of the initial and intermediate atomic states, while \bar{T} refers to the reverse radial component of T , i.e., $\bar{T} = L$ for $T = S$ and vice versa.

The integral in Eq. (4) involves only electron state components. However, its evaluation is not an easy task due to the (infinite) summation over the principal quantum number n_ν contained in the radial Green's function. In the present work, such an integral has been computed by using the Green's Function Library [30]. Other computational techniques could be also used, such as the B spline finite basis set method [31,32]. The angular part of the amplitude in Eq. (2) is represented by the elements $\chi_{m_I m_{j\nu}}^{f T v T'}$ and $\chi_{m_I m_{j\nu}}^{v T' i T'}$ therein contained and can be computed analytically:

$$\begin{aligned} \chi_{m_I m_{j\nu}}^{f T v T'} &= \sum_{m_{j_f}} \langle j_f, m_{j_f}, I, m_I | F_f, m_{F_f} \rangle \\ &\quad \times \langle \kappa_f, l_f^T, m_{j_f} | \boldsymbol{\sigma} \cdot \mathbf{T}_{L_2 \Lambda_2}^* | \kappa_\nu, l_\nu^{\bar{T}}, m_{j_\nu} \rangle, \\ \chi_{m_I m_{j\nu}}^{v T' i T'} &= \sum_{m_{j_i}} \langle j_i, m_{j_i}, I, m_I | F_i, m_{F_i} \rangle \\ &\quad \times \langle \kappa_\nu, l_\nu^{T'}, m_{j_\nu} | \boldsymbol{\sigma} \cdot \mathbf{T}_{L_1 \Lambda_1 M_1}^* | \kappa_i, l_i^{\bar{T}'}, m_{j_i} \rangle, \end{aligned} \quad (6)$$




FIG. 1. (Color online) Geometry considered for the two-photon emission. The propagation direction of the first photon is adopted as the z direction. x is chosen such that xz is the reaction plane (plane spanned by the photons directions). θ is the angle between the photons directions, while angles $\chi_{1,2}$ define the linear polarizations of the first and second photon, respectively, with respect to their respective polarization planes. The polarization plane of the first (second) photon is denoted by A (B) and represents the plane orthogonal to the photon direction.

where σ are Pauli matrices. The elements $\langle \kappa_f, l_f^T, m_j | \sigma \cdot \mathbf{T}_{L_2 \Lambda_2 M_2}^* | \kappa_v, l_v^T, m_{j_v} \rangle$ and $\langle \kappa_v, l_v^T, m_{j_v} | \sigma \cdot \mathbf{T}_{L_1 \Lambda_1 M_1}^* | \kappa_i, l_i^T, m_{j_i} \rangle$ have been already discussed elsewhere [14,33] and will not be here recalled.

The initial and final states involved in the two-photon transitions which we shall analyze below are *unpolarized*. It has been shown that, for this case, we may arbitrarily adopt the quantization axis (\hat{z}) along the momentum of the “first” photon: $\hat{z} \parallel \hat{k}_1$ [14–18]. We furthermore adopt \hat{x} such that the xz plane is the reaction plane (plane spanned by the photons directions). Figure 1 sketches the geometry we consider for the decay. Within this geometry, the Wigner matrices in Eq. (2) simplify to $D_{M_1 \lambda_1}^{L_1 *}(\varphi_1, \theta_1, 0) = \delta_{M_1, \lambda_1}$ and $D_{M_2 \lambda_2}^{L_2 *}(\varphi_2, \theta_2, 0) = d_{M_2 \lambda_2}^{L_2}(\theta)$, where $d_{M \lambda}^L(\theta)$ is the reduced Wigner matrix and $\theta \equiv \theta_2$ is the polar angle of the second photon, which coincides, in the chosen geometry, with the angle between the photons directions (opening angle). Hence, the relative photons directions are uniquely determined by assigning the opening angle θ , which will be the independent variable for plotting the angular distributions.

Since part of this work is devoted to analyze photons linear polarizations, further details concerning the detection geometry must be provided before proceeding with the analysis. In Fig. 1, we show how the photon polarizations may be defined in a case experiment. The polarization of each photon is measured in the “polarization plane,” which is the plane orthogonal to the photon direction. In Fig. 1, the polarization planes of the first and second photon are denoted by A and B , respectively. Each detector is supposed to have a transmission axis, along which the linear polarization of the photon is measured. Such a transmission axis is rotated by an angle χ with respect to the reaction plane shown in Fig. 1 by the red dashed lines. Finally, each detector is supposed to work as a filter: Whenever a photon hits it, the detector either gives or does not give off a “click,” which would, respectively,

indicate that the photon has been measured as having its linear polarization along the direction χ or $\chi + 90^\circ$.

C. Definition of angular and polarization correlations

Within IPA, Eq. (2) represents the relativistic transition amplitude for the two-photon decay between hyperfine states in hydrogenlike ions. It contains the complete information on the emitted radiation. Assuming that the ion is initially unpolarized and that the polarization of the final atomic state remains unobserved, taking into account the axes geometry chosen for the two-photon emission, and using the well-known relations between linear and circular polarization bases [34], we can write the polarization-dependent differential decay rate as a function of the opening angle θ [3,35]:

$$W^{\chi_1 \chi_2}(\theta) \equiv \frac{dw^{\chi_1 \chi_2}}{d \cos \theta} = \frac{8\pi^2}{2F_i + 1} \sum_{m_{F_i} m_{F_f}} \sum_{\substack{\lambda_1 \lambda_2 \\ \lambda'_1 \lambda'_2}} \int d\omega_1 \frac{\omega_1 \omega_2}{4(2\pi)^3 c^2} e^{i(\lambda_1 - \lambda'_1)\chi_1} \times e^{i(\lambda_2 - \lambda'_2)\chi_2} \mathcal{M}^{\lambda_1 \lambda_2} \mathcal{M}^{\lambda'_1 \lambda'_2 *}. \quad (7)$$

In this Brief Report, the integration over the photon energies is numerically carried out by using the trapezoidal rule method. The number of points we used for the numerical integration has been checked to provide a precision of 1%. Hereafter, the function $W^{\chi_1 \chi_2}(\theta)$ shall be called “polarization correlation.” It represents the probability density of detecting the emitted photons at the opening angle θ with defined linear polarizations χ_1 and χ_2 .

Finally, by summing over the photons polarizations, we define the “angular correlation” as

$$W(\theta) \equiv \frac{dw}{d \cos \theta} = \frac{8\pi^2}{2F_i + 1} \int d\omega_1 \frac{\omega_1 \omega_2}{(2\pi)^3 c^2} \sum_{m_{F_i} m_{F_f}} \sum_{\lambda_1 \lambda_2} |\mathcal{M}^{\lambda_1 \lambda_2}(i \rightarrow f)|^2, \quad (8)$$

which represents the probability density of detecting the emitted photons at the opening angle θ , irrespective of their polarizations.

III. RESULTS AND DISCUSSIONS

Here we analyze the angular and polarization correlations defined in Eqs. (8) and (7), respectively, for decays of hyperfine $2s$ states in hydrogenlike ions, with special attention to the hydrogenlike ${}_{92}^{235}\text{U}$ ion, whose nuclear spin is $I = 7/2$ [36]. The function $W(\theta)$ obtained for $2s_{1/2} (F = 4, 3) \rightarrow 1s_{1/2} (F = 4, 3)$ transitions in the hydrogenlike ${}_{92}^{235}\text{U}$ ion is displayed in Fig. 2. The full multipoles and the electric dipole (E1E1) contributions are separately displayed. Within the dipole approximation, the angular correlation for non-spin-flip transitions can be well described by the familiar shape $W_{E1E1}(\theta) \sim 1 + \cos^2 \theta$ [10]. However, the full multipole calculation shows some asymmetric deviations from such a shape. This effect is already known from the past literature,




FIG. 2. (Color online) Angular correlations in the hydrogenlike $^{235}_{92}\text{U}$ ion. The function $W(\theta)$ is shown for the transitions $2s_{1/2} (F = 4, 3) \rightarrow 1s_{1/2} (F = 4, 3)$. The dashed-red curve refers to the electric dipole approximation while the solid-black curve refers to the full multipoles contribution.

where it has been shown that high multipoles contribute with terms of the type $\sim \cos \theta$ to the angular correlation in $2s_{1/2} \rightarrow 1s_{1/2}$ transitions in highly charged ions [13].

On the other hand, for spin-flip transitions, the angular correlation within the dipole approximation is well described by the function $W_{E1E1}(\theta) \sim 1 - 1/3 \cos^2 \theta$. This emission pattern is typical for two-photon transitions of the type $J_{\text{TOT}} = 1 (0) \rightarrow J_{\text{TOT}} = 0 (1)$, where J_{TOT} is the total angular momentum of the system which undergoes the decay. The two-photon decay $(1s 2s)^3 S_{J=1} \rightarrow (1s 1s)^1 S_{J=0}$ in heliumlike ions, where J is the total angular momentum of the two-electron system, shows approximately the same behavior [17]. As in the previous case, the full multipole calculation shows remarkable asymmetric deviations from the symmetric shape. Quantitatively, the ratio $W(\pi)/W(0)$ is $\simeq 1.09$ for non-spin-flip transitions and $\simeq 1.35$ for spin-flip transitions.

For low-charged ions, we find that the angular correlation is fully described by the functions $\sim 1 + \cos^2 \theta$ and $\sim 1 - 1/3 \cos^2 \theta$ for non-spin-flip and spin-flip transitions, respectively (i.e., the full multipole calculations coincide with the calculations performed within the dipole approximation).

From the figures, we also notice that spin-flip transitions are overall suppressed with respect to the non-spin-flip transition, which is to be expected in view of the fact that the electric dipole, which is the leading multipole here, conserves the electron spin if evaluated nonrelativistically. This entails that the curves we obtained for spin-flip transitions are fully determined by relativistic and high multipole contributions.

We now turn to analyze the polarization correlations for some of the hyperfine transitions considered in Fig. 2. In Fig. 3, we plot the function $W^{\chi_1 \chi_2}(\theta)$ as obtained for the transitions $2s_{1/2} (F = 3) \rightarrow 1s_{1/2} (F = 3)$ and $2s_{1/2} (F = 3) \rightarrow 1s_{1/2} (F = 4)$. We see that, in general, photons coming from spin-flip transitions and non-spin-flip transitions have mainly orthogonal and parallel linear polarizations, respectively. We find that this polarization scheme holds




FIG. 3. (Color online) Polarization correlations in the hydrogenlike $^{235}_{92}\text{U}$ ion. The function $W^{\chi_1 \chi_2}(\theta)$ is shown for the transitions $2s_{1/2} (F = 3) \rightarrow 1s_{1/2} (F = 3)$ (solid-black curve) and $2s_{1/2} (F = 3) \rightarrow 1s_{1/2} (F = 4)$ (dashed-green curve). The four polarization configurations $(\chi_1, \chi_2 = 0^\circ, 90^\circ)$ are displayed.

perfectly (i.e., without deviations) for low- Z hydrogenlike ions. However, for hydrogenlike uranium, as well as for any highly charged ions, sizable deviations are evident, as can be seen in Fig. 3.

Hydrogenlike heavy ions can be nowadays efficiently produced in storage rings [37]. The energy of the emitted radiation in hydrogenlike uranium is in the range of hard x rays. An experimental polarization analysis of such energetic photons would be nowadays possible through the use of Compton polarimeters [38–42]. By analyzing the decay spectrum, a conventional photon-photon coincidence measurement enables one to distinguish two-photon decay events from the dominant single-photon M1 decay channel [10,43]. Therefore, information on the polarization state of two photons can be achieved by selecting events which have been recorded in coincidence by two polarimeters and which have the desired scattering angle [26,44–48].

IV. SUMMARY AND PERSPECTIVES

In summary, the amplitude for two-photon transitions between hyperfine states in hydrogenlike atoms has been calculated. By using such amplitude, the angular and linear polarization properties of the photon pair emitted in two-photon decays of $2s$ hyperfine states have been investigated within second-order perturbation theory and the Dirac relativistic framework. Special attention has been paid to the hydrogenlike $^{235}_{92}\text{U}$ ion. Results have been shown for the transitions $2s_{1/2} (F = 4, 3) \rightarrow 1s_{1/2} (F = 4, 3)$. It has been possible to identify two emission patterns: (i) two-photon non-spin-flip transitions are found to be characterized by an angular distribution approximately of the type $W(\theta) \sim 1 + \cos^2 \theta$ and by photon polarizations approximately parallel to one another; and (ii) two-photon spin-flip transitions have somewhat smaller decay rates and are found to be characterized by an angular distribution

approximately of the type $W(\theta) \sim 1 - 1/3 \cos^2 \theta$ as well as by photon polarizations approximately orthogonal to one another. Deviations to this pattern come from high multipoles and relativistic contributions, are negligible for low-charged ions, and are of size 1 to 25% in the hydrogenlike $^{235}_{92}\text{U}$ ion.

This Brief Report is the first step toward exploring the nuclear-spin effect on the angular and polarization properties of the photon pair emitted in two-photon decays. This study might pave the way for a new route to get information on the direction and the magnitude of the spin distribution inside the nucleus, which is still quite an unraveled problem [49].

ACKNOWLEDGMENTS

F.F. acknowledges support by Fundação de Amparo à Pesquisa do estado de Minas Gerais and Conselho Nacional de Desenvolvimento Científico e Tecnológico. F.F. and L.L. acknowledge support by the Research Council for Natural Sciences and Engineering of the Academy of Finland. J.P.S. and P.A. acknowledge support by Fundação para a Ciência e a Tecnologia (Portugal), through the Projects No. PEstOE/FIS/UI0303/2011 and No. PTDC/FIS/117606/2010, financed by the European Community Fund through the Competitiveness Factors Operational Programme.

-
- [1] M. Goepfert-Mayer, *Ann. Phys. (Leipzig)* **9**, 273 (1931).
 [2] J. Shapiro and G. Breit, *Phys. Rev.* **113**, 179 (1959).
 [3] S. P. Goldman and G. W. F. Drake, *Phys. Rev. A* **24**, 183 (1981).
 [4] G. W. F. Drake, *Phys. Rev. A* **34**, 2871 (1986).
 [5] A. Derevianko and W. R. Johnson, *Phys. Rev. A* **56**, 1288 (1997).
 [6] J. P. Santos, F. Parente, and P. Indelicato, *Eur. Phys. J. D* **3**, 43 (1998).
 [7] R. W. Dunford, H. G. Berry, S. Cheng, E. P. Kanter, C. Kurtz, B. J. Zabransky, A. E. Livingston, and L. J. Curtis, *Phys. Rev. A* **48**, 1929 (1993).
 [8] R. Ali, I. Ahmad, R. W. Dunford, D. S. Gemmell, M. Jung, E. P. Kanter, P. H. Mokler, H. G. Berry, A. E. Livingston, S. Cheng, and L. J. Curtis, *Phys. Rev. A* **55**, 994 (1997).
 [9] H. W. Schäffer, P. H. Mokler, R. W. Dunford, C. Kozhuharov, A. Krämer, A. E. Livingston, T. Ludziejewski, H. T. Prinz, P. Rymuza, L. Sarkadi, Z. Stachura, Th. Stöhlker, P. Swiat, and A. Warczak, *Phys. Lett. A* **260**, 489 (1999).
 [10] P. H. Mokler and R. W. Dunford, *Phys. Scr.* **69**, C1 (2004).
 [11] A. Kumar, S. Trotsenko, A. V. Volotka, D. Banaś, H. F. Beyer, H. Bräuning, A. Gumberidze, S. Hagmann, S. Hess, C. Kozhuharov, R. Reuschl, U. Spillmann, M. Trassinelli, G. Weber, and Th. Stöhlker, *Eur. Phys. J. Special Topics* **169**, 19 (2009).
 [12] S. Trotsenko, A. Kumar, A. V. Volotka *et al.*, *Phys. Rev. Lett.* **104**, 033001 (2010).
 [13] C. K. Au, *Phys. Rev. A* **14**, 531 (1976).
 [14] A. Surzhykov, P. Koval, and S. Fritzsche, *Phys. Rev. A* **71**, 022509 (2005).
 [15] L. Borowska, A. Surzhykov, Th. Stöhlker, and S. Fritzsche, *Phys. Rev. A* **74**, 062516 (2006).
 [16] A. Surzhykov, T. Radtke, P. Indelicato, and S. Fritzsche, *Eur. J. Phys. Special Topics* **169**, 29 (2009).
 [17] A. Surzhykov, A. Volotka, F. Fratini, J. P. Santos, P. Indelicato, G. Plunien, Th. Stöhlker, and S. Fritzsche, *Phys. Rev. A* **81**, 042510 (2010).
 [18] F. Fratini and A. Surzhykov, *Hyp. Int.* **199**, 85 (2011).
 [19] P. Amaro, F. Fratini, S. Fritzsche, P. Indelicato, J. P. Santos, and A. Surzhykov, *Phys. Rev. A* **86**, 042509 (2012).
 [20] A. V. Volotka, A. Surzhykov, V. M. Shabaev, and G. Plunien, *Phys. Rev. A* **83**, 062508 (2011).
 [21] G. Breit and E. Teller, *Astrophys. J.* **91**, 215 (1940).
 [22] W. Perrie, A. J. Duncan, H. J. Beyer, and H. Kleinpoppen, *Phys. Rev. Lett.* **54**, 1790 (1985).
 [23] H. Kleinpoppen, A. J. Duncan, H. J. Beyer, and Z. A. Sheikh, *Phys. Scr. T* **72**, 7 (1997).
 [24] F. Fratini, M. C. Tichy, Th. Jahrsetz, A. Buchleitner, S. Fritzsche, and A. Surzhykov, *Phys. Rev. A* **83**, 032506 (2011).
 [25] A. Schäfer, G. Soff, P. Indelicato, B. Müller, and W. Greiner, *Phys. Rev. A* **40**, 7362 (1989).
 [26] F. Fratini, S. Trotsenko, S. Tashenov, Th. Stöhlker, and A. Surzhykov, *Phys. Rev. A* **83**, 052505 (2011).
 [27] V. M. Shabaev, M. Tomaselli, T. Kühl, A. N. Artemyev, and V. A. Yerokhin, *Phys. Rev. A* **56**, 252 (1997).
 [28] S. G. Karshenboim and V. G. Ivanov, *Phys. Lett. B* **524**, 259 (2002).
 [29] A. Pálffy, *Cont. Phys.* **51**, 471 (2010).
 [30] P. Koval and S. Fritzsche, *Comp. Phys. Commun.* **152**, 191 (2003).
 [31] L. Safari, P. Amaro, S. Fritzsche, J. P. Santos, and F. Fratini, *Phys. Rev. A* **85**, 043406 (2012).
 [32] L. Safari, P. Amaro, S. Fritzsche, J. P. Santos, S. Tashenov, and F. Fratini, *Phys. Rev. A* **86**, 043405 (2012).
 [33] A. Surzhykov, S. Fritzsche and Th. Stöhlker, *J. Phys. B* **35**, 3713 (2002).
 [34] M. E. Rose, *Elementary Theory of Angular Momentum* (Wiley, New York, 1953).
 [35] F. Fratini and A. G. Hayrapetyan, *Phys. Scr.* **84**, 035008 (2011).
 [36] N. J. Stone, *At. Data Nucl. Data Tables* **90**, 75 (2005).
 [37] A. Gumberidze *et al.*, *Phys. Rev. Lett.* **110**, 213201 (2013).
 [38] U. Spillmann, H. Bräuning, S. Hess, H. Beyer, Th. Stöhlker, J. C. Dousse, D. Protic, and T. Krings, *Rev. Sci. Instrum.* **79**, 083101 (2008).
 [39] G. Weber, H. Bräuning, S. Hess, R. Martin, U. Spillmann, and Th. Stöhlker, *Journal of Instrumentation* **5**, C07010 (2010).
 [40] P. S. Shaw, U. Arp, A. Henins, and S. Southworth, *Rev. Sci. Instrum.* **67**, 3362 (1996).
 [41] D. Protic, E. L. Hull, T. Krings, and K. Vetter, *IEEE Trans. Nucl. Sci.* **52**, 3181 (2005).
 [42] D. Protic, Th. Stöhlker, T. Krings, I. Mohos, and U. Spillmann, *IEEE Trans. Nucl. Sci.* **52**, 3194 (2005).
 [43] A. Simionovici *et al.*, *Phys. Rev. A* **48**, 1695 (1993).
 [44] S. Tashenov, Th. Stöhlker *et al.*, *Phys. Rev. Lett.* **97**, 223202 (2006).
 [45] M. H. L. Pryce and J. C. Ward, *Nature (London)* **160**, 435 (1947).
 [46] H. S. Snyder, S. Pasternack, and J. Hornbostel, *Phys. Rev.* **73**, 440 (1948).
 [47] C. S. Wu and I. Shaknov, *Phys. Rev.* **77**, 136 (1950).
 [48] E. Bleuler and H. L. Brandt, *Phys. Rev.* **73**, 1398 (1948).
 [49] R. L. Jaffe, *Phys. Today* **48**, 24 (1995).