

# Lowest-order relativistic corrections to the fundamental limits of nonlinear-optical coefficients

Nathan J. Dawson\*

Department of Physics, Case Western Reserve University, Cleveland, Ohio 44106, USA

(Received 20 August 2014; published 22 January 2015)

The effects of small relativistic corrections to the off-resonant polarizability, hyperpolarizability, and second hyperpolarizability are investigated. Corrections to linear- and nonlinear-optical coefficients are demonstrated in the three-level ansatz, which includes corrections to the Kuzyk limits when scaled to semirelativistic energies. It is also shown that the maximum value of the hyperpolarizability is more sensitive than the maximum polarizability or second hyperpolarizability to lowest-order relativistic corrections. These corrections illustrate how the intrinsic nonlinear-optical response is affected at semirelativistic energies.

DOI: [10.1103/PhysRevA.91.013832](https://doi.org/10.1103/PhysRevA.91.013832)

PACS number(s): 42.65.An, 42.70.Mp, 31.30.jx, 31.30.jc

## I. INTRODUCTION

Over a decade ago, Kuzyk [1] showed that there are fundamental limits to the off-resonant, electronic, nonlinear-optical response. This was discovered by manipulating both the on- and off-diagonal elements of the Thomas-Reiche-Kuhn (TRK) sum rule [2–4], which limits the oscillator strengths of a quantum system with respect to fundamental constants in the nonrelativistic regime. The oscillator strength is limited by the nonrelativistic kinetic energy of a free particle, where field interactions from a four-potential do not contribute to the maximum oscillator strength. The intrinsic values of the hyperpolarizability and second hyperpolarizability in the nonrelativistic limit have been studied in great detail [5–23], where there is a looming gap between the measured and calculated values and the fundamental limits in the nonrelativistic regime.

There have been several approaches to reduce this gap using optimization routines on one-dimensional potentials, which have resulted in the confirmation of the apparent gap [24–32]. Another approach to breach the gap involves a systematic search for new classes of organic nonlinear-optical molecules with multipolar charge-density analysis from crystallographic data [33–35]. New abstract methods of calculating large nonlinear responses have also been studied for low-dimensional quantum graphs [36–38]. All of these approaches focus on breaching the gap between the fundamental limit and the largest calculated (or directly measured) intrinsic values.

Although a four-potential does not contribute to the nonrelativistic TRK sum rule, there may be other ways to change the limiting value on the oscillator strength, and thereby adjust the fundamental limits of the nonlinear-optical response. Instead of focusing on optimizing the intrinsic value based on a specific potential, I will discuss the changes in the limiting constant of the TRK sum rules for a specific type of quantum system that is not properly represented by a closed Schrodinger equation. Specifically, a relativistic system is examined which no longer has a simple  $p^2/2m$  kinetic energy approximation, and therefore, the energy-momentum relationship directly affects the fundamental limits. Thus, this paper is dedicated to the study of the fundamental limits of the hyperpolarizability and second hyperpolarizability for systems that have non-negligible relativistic energies.

## II. THEORY

In the far off-resonant limit (frequency approaches zero), the respective one-dimensional polarizability, hyperpolarizability, and second hyperpolarizability are [39]

$$\alpha = 2e^2 \sum_n' \frac{x_{0n}x_{n0}}{E_{n0}}, \quad (1)$$

$$\beta = 3e^3 \sum_{n,l}' \frac{x_{0n}\bar{x}_{nl}x_{l0}}{E_{n0}E_{l0}}, \quad (2)$$

and

$$\gamma = 4e^4 \left( \sum_{n,l,k}' \frac{x_{0n}\bar{x}_{nl}\bar{x}_{lk}x_{k0}}{E_{n0}E_{l0}E_{k0}} - \sum_{n,l}' \frac{x_{0n}x_{n0}x_{0l}x_{l0}}{E_{n0}^2E_{l0}} \right), \quad (3)$$

where  $x$  is the position operator in one dimension,  $e$  is the magnitude of an electron's charge,  $E_i$  is the  $i$ th energy eigenstate, and the prime restricts the summation by excluding the ground state. The shorthand notation,  $x_{ij} = \langle i | x | j \rangle$  and  $E_{ij} = E_i - E_j$ , was introduced in Eqs. (1)–(3). Note that the barred operator presented in the expressions for the nonlinear coefficients is the origin-specific expectation value, which is given as  $\bar{x}_{ii} = x_{iii} - x_{00}$  when the indices are matched.

The TRK sum rules for the Dirac equation, gives the well-known result of all states summing to zero, where the zero value is due to the sum of the positive and corresponding negative energy states [40]. We wish to only observe the positive energy states from an electron in an atom or molecule, and therefore, we must project out the positive energy states. For a single electron system, the *positive energy* TRK sum rules to lowest-relativistic order (ordered in  $1/c$ ) have previously been derived [41–43] using a Foldy-Wouthuysen (FW) transformation [44],

$$\begin{aligned} & \sum_{l=0}^{\infty} \langle k | \mathbf{r} | l \rangle \langle l | \mathbf{r} | n \rangle \left[ E_l - \frac{1}{2} (E_k + E_n) \right] \\ & = \langle k' | \left( \frac{3\hbar^2}{2m} + \frac{5\hbar^4}{4m^3c^2} \nabla^2 \right) | n' \rangle, \end{aligned} \quad (4)$$

where  $|n'\rangle = e^{i\mathcal{S}}|n\rangle$  with  $\mathcal{S}$  being a unitary operator. Equation (4) differs slightly from the results of Ref. [42], where we have derived the relation with arbitrary eigenstates because both the on- and off-diagonal components of the sum rules

\*dawsphys@hotmail.com

are essential in determining the off-resonant, nonlinear-optical responses [1]. In the FW approach  $p = p'$ , where an operator  $A$  in the FW approximation is defined as  $A' = e^{i\mathcal{S}} A e^{-i\mathcal{S}}$ . Thus, the momentum operator commutes with  $e^{i\mathcal{S}}$ , and therefore  $\langle k'|p^2|n'\rangle = \langle k|e^{-i\mathcal{S}} p^2 e^{i\mathcal{S}}|n\rangle = \langle k|p^2|n\rangle$ . Note that while transforming the Hamiltonian for an electron interacting with fields,  $\mathcal{S}$  is chosen at every iteration to remove all odd operators.

Inasmuch as the  $\nabla$  operator is related to the momentum operator, and that there is an equivalence between the momentum operator in the Schrodinger equation and the momentum in the FW transformation, the right-hand side (RHS) of Eq. (4) may be written as

$$\langle k|\left(\frac{3\hbar^2}{2m} - \frac{5\hbar^2 p^2}{4m^3 c^2}\right)|n\rangle.$$

Thus, to the lowest-order relativistic correction, the RHS of the TRK sum rules given in Eq. (4) decreases for any real value of the momentum.

The lowest-order relativistic approximation to the Hamiltonian (for an electron in the presence of a scalar potential only) is given as

$$H = H_0 - \frac{p^4}{8m^3 c^2} + \frac{1}{4m^2 c^2} (\boldsymbol{\sigma} \cdot \mathbf{p}) V(\mathbf{r}) (\boldsymbol{\sigma} \cdot \mathbf{p}) - \frac{1}{8m^2 c^2} (p^2 V(\mathbf{r}) + V(\mathbf{r}) p^2), \quad (5)$$

where

$$H_0 = \frac{p^2}{2m} + V(\mathbf{r}), \quad (6)$$

with  $V(\mathbf{r})$  denoting a spatially dependent scalar potential and  $\boldsymbol{\sigma}$  representing the Pauli spin matrices. We may rewrite Eq. (5) as the well-known result [45]:

$$H = \frac{p^2}{2m} - \frac{p^4}{8m^3 c^2} + V(\mathbf{r}) + \frac{\hbar}{4m^2 c^2} \boldsymbol{\sigma} \cdot \{[\nabla V(\mathbf{r})] \times \mathbf{p}\} + \frac{\hbar^2}{8m^2 c^2} \nabla^2 V(\mathbf{r}). \quad (7)$$

The Hamiltonian with lowest-order relativistic corrections is a quartic equation with respect to momentum. For central potentials, the spin-orbit term may be recast in terms of the angular momentum operator, and thereby reduces the Hamiltonian to a quadratic equation in  $p^2$ .

There is an alternative method of reducing Eq. (7) to a quadratic that does not require one to collapse the parameter space to the centrosymmetric limit, which is observed when limiting the system to one dimension. In one dimension, there is no orbital angular momentum,  $\nabla V \times \mathbf{p} = 0$ , and therefore, the spin-orbit term vanishes. Thus, Eq. (7) reduces to a simplified quadratic equation in  $p_x^2$ , where

$$H = \frac{p_x^2}{2m} - \frac{p_x^4}{8m^3 c^2} + V(x) + \frac{\hbar^2}{8m^2 c^2} \nabla^2 V(x). \quad (8)$$

Note that the Darwin still term survives the one-dimensional approximation. Although this approach simplifies the study of generalized semirelativistic interactions while maintaining a noncentrosymmetric parameter space, one should note that many recent advances in quantum chemistry have been

introduced for numerically approximating specified relativistic systems. Most notably are the electrostatic-potential-ordered Douglass-Kroll-Hess method [46–49], the ordered regular approximations [50–53], and others based on exact decoupling methods [54–57].

By restricting ourselves to one dimension, we may write

$$\langle k|V + \frac{p_x^2}{2m} - \frac{p_x^4}{8m^3 c^2} + \frac{\hbar^2}{8m^2 c^2} \frac{\partial^2 V}{\partial x^2} = E_n|n\rangle. \quad (9)$$

Solving Eq. (9) for  $p_x^2$  gives

$$\langle k|p_x^2|n\rangle = 2m^2 c^2 \delta_{k,n} - 2m^2 c^2 \times \langle k|\sqrt{1 - \frac{2(E_n - V)}{mc^2} + \frac{\hbar^2}{4m^3 c^4} \frac{\partial^2 V}{\partial x^2}}|n\rangle, \quad (10)$$

where  $\delta$  is the Kronecker delta function, and the negative root was chosen which reduces Eq. (10) to the nonrelativistic TRK sum rules as  $1/c \rightarrow 0$ . Using Eq. (10), the one-dimensional TRK sum rule with lowest-order relativistic corrections,

$$\sum_{l=0}^{\infty} \langle k|x|l\rangle \langle l|x|n\rangle \left[ E_l - \frac{1}{2}(E_k + E_n) \right] = \langle k|\left(\frac{\hbar^2}{2m} - \frac{3\hbar^2 p_x^2}{4m^3 c^2}\right)|n\rangle, \quad (11)$$

may be rewritten as

$$\sum_{l=0}^{\infty} \langle k|x|l\rangle \langle l|x|n\rangle \left[ E_l - \frac{1}{2}(E_k + E_n) \right] = \frac{\hbar^2}{m} \left( \frac{3}{2} \lambda_{kn} - \delta_{k,n} \right), \quad (12)$$

where

$$\lambda_{kn} = \langle k|\sqrt{1 - \frac{2}{mc^2}(E_n - V) + \frac{\hbar^2}{4m^3 c^4} \frac{\partial^2 V}{\partial x^2}}|n\rangle. \quad (13)$$

Under the current set of approximations, we take the element ( $k = 0, n = 0$ ), or (0,0), which gives

$$|x_{10}|^2 E_{10} = \frac{\hbar^2}{m} \left( \frac{3}{2} \lambda_{00} - 1 \right) - \sum_{l=2}^{\infty} |x_{l0}|^2 E_{l0}. \quad (14)$$

Considering the diagonal components and neglecting the Darwin term, there are two regimes that adjust the fundamental limit. If  $E_n > V_{n,n}(x)$ , then the electron is moving inside a potential and  $\lambda_{nn}$  is real. This causes a decrease in the maximum oscillator strength. If the electron is expected to be outside a potential such that  $E_n < V_{n,n}(x)$ , then  $\lambda_{nn}$  becomes imaginary, which cannot occur for bound states with positive energies. Therefore, the oscillator strength of a one-dimensional semirelativistic system decreases with respect to the nonrelativistic approximation; however, a competing parameter may increase the final numerical value of some systems (not the intrinsic value) of the off-resonant response because the relativistic corrections reduce the transition energies with respect to those mapped from the nonrelativistic Hamiltonian.

In prior studies that began with a Hamiltonian in the nonrelativistic limit, it was shown that the largest nonlinear-optical responses occur when all other transition energies become much larger than  $E_{10}$ . In other words, the sum-over-state (SOS) expressions are dominated by the first excited state transition. This is also true for relativistically corrected systems which is obvious from Eq. (14). Thus, we adopt the same method as Kuzyk [1] and assume a three-level model. Then, Eq. (14) reduces to

$$|x_{10}|^2 E_{10} + |x_{20}|^2 E_{20} = \frac{\hbar^2}{m} \left( \frac{3}{2} \lambda_{00} - 1 \right). \quad (15)$$

Likewise, (1,1) produces the resultant equation,

$$|x_{12}|^2 E_{21} - |x_{10}|^2 E_{10} = \frac{\hbar^2}{m} \left( \frac{3}{2} \lambda_{11} - 1 \right). \quad (16)$$

In the same manner as Eqs. (15) and (16) we take (1,0), which gives

$$x_{10} \bar{x}_{11} E_{10} + x_{12} x_{20} (E_{21} + E_{20}) = \frac{3\hbar^2}{2m} \lambda_{10}. \quad (17)$$

Note that the left-hand side of Eq. (17) is identical for (1,0) and (0,1) when we assume real transition moments, i.e.,  $x_{ij} = x_{ji}$  [6]. Thus, it is of no surprise that the corresponding  $\lambda$  parameter must also possess the property  $\lambda_{10} = \lambda_{01}$ . Finally, we set the matrix elements corresponding to (2,0), or (0,2), which gives

$$x_{20} \bar{x}_{22} E_{20} + x_{10} x_{12} (E_{10} - E_{21}) = \frac{3\hbar^2}{2m} \lambda_{20}. \quad (18)$$

Note that Eqs. (17) and (18) contain off-diagonal components that are real and positive for well-behaved systems.

Solving Eqs. (15)–(18) for the transition dipole moments, we find

$$|x_{10}| = \frac{\hbar}{\sqrt{mE_{10}}} X \sqrt{\frac{3}{2} \lambda_{00} - 1}, \quad (19)$$

$$|x_{12}| = \frac{\hbar}{\sqrt{mE_{10}}} \sqrt{\frac{E}{1-E}} G_\lambda(X), \quad (20)$$

$$\alpha'_{3L} = \frac{2e^2 \hbar^2}{mE_{10}^2} [X^2 + E^2(1-X^2)] H_\lambda^2, \quad (27)$$

$$\beta'_{3L} = \frac{6e^3 \hbar^3}{\sqrt{m^3 E_{10}^7}} H_\lambda \left[ \sqrt{1-X^2} X (1-E)^{3/2} \left( 1 + \frac{3}{2} E + E^2 \right) H_\lambda G_\lambda(X) - 3X \lambda_{10} - 3\sqrt{1-X^2} E^{7/2} \lambda_{20} \right], \quad (28)$$

and

$$\begin{aligned} \gamma'_{3L} = & \frac{e^4 \hbar^4}{m^2 E_{10}^5} \left\{ 4[4 - (1 + 2X^2 + 5X^4)E^5 - (1 - 2X^2 - 5X^4)E^3 - (3 - 5X^4)E^2 - 5X^4] \right. \\ & - 9[(1 - 2X^2 + 5X^4)E^5 + (2X^2 - 5X^4)E^3 + (4X^2 - 5X^4)E^2 + 5X^4 - 4X^2] \lambda_{00}^2 \\ & - 6[4 - 4X^2 + (4X^2 - 3)E^2 + (4X^2 - 1)E^3 - 4X^2 E^5] \lambda_{11} \\ & + 6[(2 + 10X^4)E^5 + (1 - 10X^4)E^3 + (3 + 4X^2 - 10X^4)E^2 + 10X^4 - 4X^2 - 4] \lambda_{00} \\ & - 9[4X^2 E^5 (1 - 4X^2) E^3 + (3 - 4X^2) E^2 + 4X^2 - 4] \lambda_{00} \lambda_{11} + 9(E^5 \lambda_{20}^2 + \lambda_{10}^2) \\ & \left. - 12H_\lambda G_\lambda(X) [\lambda_{10}(2 + E)\sqrt{1-E}\sqrt{1-X^2} - \lambda_{20} X \sqrt{E(1-E)}(E^3 + 2E^4)] \right\}, \quad (29) \end{aligned}$$

$$\begin{aligned} \bar{x}_{11} = & \frac{\hbar}{\sqrt{mE_{10}}} \left[ \frac{E-2}{\sqrt{1-E}} \frac{\sqrt{1-X^2}}{X} G_\lambda(X) \right. \\ & \left. + \frac{3\lambda_{10}}{2X\sqrt{\frac{3}{2}\lambda_{00}-1}} \right], \quad (21) \end{aligned}$$

$$\begin{aligned} \bar{x}_{22} = & \frac{\hbar}{\sqrt{mE_{10}}} \left[ \frac{1-2E}{\sqrt{1-E}} \frac{X}{\sqrt{1-X^2}} G_\lambda(X) \right. \\ & \left. + \frac{3\sqrt{E}\lambda_{20}}{2\sqrt{1-X^2}\sqrt{\frac{3}{2}\lambda_{00}-1}} \right], \quad (22) \end{aligned}$$

and

$$|x_{20}| = \frac{\hbar}{\sqrt{mE_{10}}} \sqrt{E} \sqrt{1-X^2} \sqrt{\frac{3}{2}\lambda_{00}-1}, \quad (23)$$

where

$$G_\lambda(X) = \sqrt{X^2 \left( \frac{3}{2} \lambda_{00} - 1 \right) + \frac{3}{2} \lambda_{11} - 1}. \quad (24)$$

Here, we used the notation inline with previous expressions for the nonlinear-optical limits of nonrelativistic systems such that

$$X = \frac{|x_{10}|}{|x_{10}^{\max}|} \quad \text{and} \quad E = \frac{E_{10}}{E_{20}}, \quad (25)$$

where we can see that the maximum value for the  $x_{10}$  transition moment is

$$|x_{10}^{\max}| = \frac{\hbar}{\sqrt{mE_{10}}} \sqrt{\frac{3}{2}\lambda_{00}-1}. \quad (26)$$

To find an expression for the off-resonant polarizability, hyperpolarizability, and second hyperpolarizability of a three-level system, we substitute Eqs. (19)–(26) into Eqs. (1)–(3). The three-level polarizability, hyperpolarizability, and second hyperpolarizability reduce to

where

$$H_\lambda = \sqrt{\frac{3}{2}\lambda_{00} - 1}. \quad (30)$$

The primed coefficients in Eqs. (27)–(29) denote relativistic corrections to the TRK sum rules. Note that the energies in these primed equations for the nonlinear-optical coefficients are also relativistically corrected.

In the nonrelativistic limit, i.e., when  $c \rightarrow \infty$ , Eqs. (27)–(29) reduce to the off-resonant, three-level model calculated from the nonrelativistic TRK sum rules [17,19]. The polarizability, hyperpolarizability, and second hyperpolarizability in the nonrelativistic limit are given by

$$\alpha_{3L} = \frac{e^2 \hbar^2}{m E_{10}^2} [X^2 + E^2(1 - X^2)], \quad (31)$$

$$\beta_{3L} = \frac{3e^3 \hbar^3}{2\sqrt{2}m^3 E_{10}^7} X \sqrt{1 - X^4} \times (1 - E)^{3/2} \left(1 + \frac{3}{2}E + E^2\right), \quad (32)$$

and

$$\gamma_{3L} = \frac{e^4 \hbar^4}{m^2 E_{10}^5} [4 - 2(E^2 - 1)E^3 X^2 - 5(E - 1)^2(E + 1)(E^2 + E + 1)X^4 - (E^3 + E + 3)E^2]. \quad (33)$$

### III. DISCUSSION

Transition moments (and expectation values of many types) in addition to diagonal energy or potential differences can appear in the relativistically corrected equation via the  $\lambda_{ij}$  terms. If the values of  $\lambda_{ij}$  are known for a specific potential, then the second hyperpolarizability can be approximated by Eq. (29). In other words, the inclusion of the momentum term in the TRK sum rules no longer gives a simple relationship between the transition moments and energies.

It is clear that the linear polarizability for all  $X$  and  $E$  is reduced by the lowest-order relativistic correction. The decrease is due to the presence of the  $H_\lambda$  parameter, which can take values between 0 and 1, where  $H_\lambda \rightarrow 1$  in the nonrelativistic limit. In  $X$  and  $E$  parameter space, the limit of the hyperpolarizability is located at  $X = 1/\sqrt[4]{3}$  and  $E = 0$ . The resulting limit corresponds to a two-level system, which is not surprising given the relationships in Eq. (14). Because the maximum is located when  $1/E_{20} \rightarrow 0$ , it seems counterintuitive that the maximum of the nonlinear-optical coefficients occur when  $X \neq 1$ ; however, we can no longer think in terms of simple linear optics. When calculating nonlinear-optical coefficients, the intermediate states and excited state sum rules are interwoven into Eqs. (1) and (3). The limit of the hyperpolarizability of nonrelativistic systems calculated using the three-level ansatz is given by

$$\beta_{\max} = \sqrt[4]{3} \frac{e^3 \hbar^3}{\sqrt{m^3 E_{10}^7}}. \quad (34)$$

The effects of linear, relativistic, kinetic energy on the fundamental limit of the hyperpolarizability may be studied by substituting  $X = 1/\sqrt[4]{3}$  and  $E = 0$  into Eq. (28). The lowest-order relativistic correction to the limit of the hyperpolarizability,  $\beta'(X, E)$ , is given by

$$\beta'(\sqrt[4]{3}, 0) = \frac{2}{\sqrt[4]{3}} \frac{e^3 \hbar^3}{\sqrt{m^3 E_{10}^7}} H_\lambda \times \left( \sqrt{6} H_\lambda \sqrt{\frac{3}{2}\lambda_{00} + \frac{3}{2}\lambda_{11} - 1 - 9\lambda_{10}} \right). \quad (35)$$

Note that when  $E = 0$ , the second excited state is infinitely large; however,  $E_{20}$  does not enter into the oscillator strength corrections as there is no  $\lambda_{22}$  term. The same is true for any number of truncated states, where there is no diagonal  $\lambda_{pp}$  term for a system truncated to  $p$  states. Thus, we may still assume that  $E_{20} \rightarrow \infty$  without any obvious negative consequences.

The limit to the hyperpolarizability for increasingly relativistic systems is shown in Fig. 1, where the intrinsic value,  $\beta'_{\text{int}} = \beta'/\beta_{\max}$ , is plotted as a function of  $\lambda_{00}$  and  $\lambda_{11}$ . We must place a lower bound on some parameters due to the low-order approximation. We observe that for real values of the off-resonant hyperpolarizability,  $\lambda_{00}$  and  $\lambda_{11}$  can have a minimum value of  $2/3$ . As shown in Fig. 1(a), the lowest-order relativistic correction to the limit of the hyperpolarizability is reduced, or even negative, when  $\lambda_{10} = 0$  while  $\lambda_{00}$  and  $\lambda_{11}$  increase. The hyperpolarizability is further reduced when the off-diagonal relativistic term  $\lambda_{10}$  is increased as illustrated in Fig. 1(b). If we further increase  $\lambda_{10}$  away from the nonrelativistic limit, there are values of  $\lambda_{00}$  and  $\lambda_{11}$  that correspond to a negative hyperpolarizability that is greater in magnitude than the fundamental limit. These occurrences where the limit is broken appear for values of  $\lambda_{11}$  that deviate from unity, but not for large deviations of  $\lambda_{00}$ , where the entire function of  $\beta'$  is multiplied by  $H_\lambda$ . Thus, large values of  $\lambda_{00}$  quickly decrease the effects of an increasing  $\lambda_{10}$ .

The red region shown in Fig. 1(c) corresponds to the region that is opposite in sign and greater in magnitude to the fundamental limit when  $\lambda_{10} = 0.2$ , which is still within the stability boundaries of the lowest-order approximation. There is the possibility that higher-order relativistic corrections may lessen the effects of the lowest-order correction; however, introducing higher-order corrections into an analytical framework is quite complicated and beyond the scope of the present study. The lowest-order correction to the  $(n, n)$  sum rules appears to damp the total strength of the transition probabilities by increasing the momentum at semirelativistic energies. Note that an exotic Hamiltonian with a small momentum correction of opposite sign to that of the lowest-order relativistic correction would instead produce a virtual increase in the total oscillator strength. Relativistic corrections to the  $(n, k)$  TRK sum rules, where  $n \neq k$ , appear to directly subtract from the total response as opposed to an apparent quadratic damping. These nonzero terms are what appear to allow the nonrelativistic fundamental limit to be broken when scaled to semirelativistic kinetic energies.

To get a general idea of how relativity affects the second hyperpolarizability, we first study the limits of the nonrelativistic three-level model, Eq. (33). The upper limit of




FIG. 1. (Color online) The intrinsic hyperpolarizability as a function of  $\lambda_{00}$  and  $\lambda_{11}$  when  $X = 1/\sqrt[4]{3}$  and  $E = 0$ . The relativistic behavior of the Kuzyk limit is illustrated for (a)  $\lambda_{10} = 0$ , (b)  $\lambda_{10} = 0.1$ , and (c)  $\lambda_{10} = 0.2$ . Values outside the magnitude for the nonrelativistic limit are given by the solid (red) region.

the nonrelativistic second hyperpolarizability, in the reduced parameter space, is located at  $E = 0$  and  $X = 0$ , which gives

$$\gamma_{\max} = \frac{4e^4\hbar^4}{m^2E_{10}^5}. \quad (36)$$

The lower limit is found when either  $E = 1$ , or when  $E = 0$  and  $X = 1$ . For the nonrelativistic case, the lower limit of the second hyperpolarizability is

$$\gamma_{\min} = -\frac{e^4\hbar^4}{m^2E_{10}^5}. \quad (37)$$

We may now substitute the corresponding three-level energy and first transition moment fractions,  $X$  and  $E$ , into the lowest-order corrected second hyperpolarizability expression to study the maximum value of semirelativistic systems. After substituting the parameters associated with the maximum for the nonrelativistic limit, Eq. (29) reduces to

$$\gamma'(0,0) = \frac{e^4\hbar^4}{m^2E_{10}^5} [16 + 9\lambda_{10} - 24\lambda_{11} + 3\lambda_{00}(12\lambda_{11} - 8) - 12\lambda_{10}\sqrt{(3\lambda_{00} - 2)(3\lambda_{11} - 2)}]. \quad (38)$$

The lowest-order relativistic corrections to the second hyperpolarizability are shown in Fig. 2. Note that the maximum intrinsic value,  $\gamma'_{\text{int}}$ , is 1 and the minimum is  $-1/4$ . As shown in Fig. 2(a), the maximum possible second hyperpolarizability is reduced for a potential with negligible off-diagonal  $\lambda$  parameters. The other two plots in Fig. 2 illustrate how a nonzero  $\lambda_{10}$  further reduces the second nonlinear response from the nonrelativistic maximum. Again, note that even though the intrinsic values are reduced, the net numerical values for the off-resonant response may be affected differently because of relativistic changes in  $E_{10}$ .

There are two regimes that are associated with the minimum value of the second hyperpolarizability. Focusing only on the minimum at the two-level limit, i.e.,  $E \rightarrow 0$ , there is an intrinsic value of  $-0.25$  when  $X = 1$ . The lower limit in this regime, with lowest-order relativistic corrections, is given by

$$\gamma'(1,0) = \frac{e^4\hbar^4}{m^2E_{10}^5} [12\lambda_{00} - 9\lambda_{00}^2 + 9\lambda_{10}^2 - 4]. \quad (39)$$

The minimum value in this regime is only affected by the lowest diagonal term  $\lambda_{00}$ , and the first off-diagonal term  $\lambda_{10}$ . Thus, it appears that, for well-behaved systems under these approximations, the first excited state does not contribute to the lowest-order relativistic correction at the (1,0) minimum.

The minimum at (1,0) is plotted in Fig. 3 as a function of  $\lambda_{00}$  and  $\lambda_{10}$ . The value of  $\lambda_{00}$  is “walked” away from the nonrelativistic value of 1, while  $\lambda_{10}$  is increased from the nonrelativistic limit of zero. Notice how the magnitude of the lower limit in this regime is also reduced which signifies response damping as the dominant mechanism as opposed to a subtraction of the net response. Similar to  $\beta'$ , under more extreme circumstances, it appears that a negative value of  $\gamma$  may also become zero or even positive. The positive is due to the off-diagonal  $\lambda_{kn}$  subtraction of the response which is less prominent for the second hyperpolarizability.

The second regime where there exists a minimum is found when  $E = 1$ , where the minimum also reaches the negative intrinsic limit of  $-1/4$ . In this regime, the lowest-order




FIG. 2. (Color online) The maximum intrinsic value of the second hyperpolarizability as a function of  $\lambda_{00}$  and  $\lambda_{11}$  from the relativistic corrections to the TRK sum rules. The corrected maximum is shown for (a)  $\lambda_{10} = 0$ , (b)  $\lambda_{10} = 0.1$ , and (c)  $\lambda_{10} = 0.2$ .

relativistic correction gives

$$\gamma'(X,1) = \frac{e^4 \hbar^4}{m^2 E_{10}^5} [12\lambda_{00} - 9\lambda_{00}^2 + 9\lambda_{10}^2 + 9\lambda_{20}^2 - 4]. \quad (40)$$

Thus, this degenerate minimum is more strongly affected by relativistic corrections with the inclusion of a positive  $\lambda_{20}$




FIG. 3. (Color online) The minimum intrinsic value of the second hyperpolarizability as a function of  $\lambda_{00}$  and  $\lambda_{10}$  from the relativistic corrections to the TRK sum rules.

parameter that increases the minimum value away from the negative limit.

#### Relativistic effects of H-like ions and the 3-level ansatz

It is well known that, unlike many organic molecules, the continuum states make a significant contribution to the total dipole response of a single hydrogen atom. Thus, these continuum states cause problems with the SOS method for the second hyperpolarizability. Other nonrelativistic methods have been developed such as a time-independent perturbation approach [58,59] to calculate the zero-frequency response and a method employing Sturmian functions used by Shelton [60] to calculate the frequency-dependent coefficients. Because the largest portion of the dipole strength is in the  $1s$ - $2p$  transition, a qualitatively study of the lowest-relativistic corrections to H-like ions can be performed with a simple three-level model. Here, problems with convergence and continuum states are washed away by placing the entire oscillator strength in the first two excited state transitions, which gives a reasonably approximate description for most systems.

The only nonzero angular contributions from the non-relativistic transitions are either  $1/\sqrt{3}$  for  $ns$ - $n^p$  or  $2/\sqrt{15}$  for  $np$ - $n^d$ . The similarity between the two nonzero angular contributions, the low frequency of  $np$ - $n^d$  transitions in the SOS expression, and the fact that we can limit our study to the  $\gamma_{zzzz}$  component allows us to make a one-dimensional approximation; thus, the total response is given by Eq. (11). Note that the spin-orbit term will still enter into the calculation, but it will be later introduced as a perturbation in the energy so that we can further simplify the example.

We can treat the lowest-order linear moment, spin-orbit, and Darwin terms as first-order perturbations in the energy [61]. This provides a simpler approach when solving Eq. (9), where we may write

$$P_{nk}^2 \approx \delta_{k,n} \frac{Z^4 \alpha^2}{(n+1)^2} \left[ \frac{2}{(1+2j)(n+1)} - \frac{3}{4(n+1)^2} \right] - \delta_{k,n} \frac{Z^2}{(n+1)^2} - 2V_{kn}, \quad (41)$$




FIG. 4. The second hyperpolarizability as a function of atomic number for an H-like ion using the three-level model with lowest-order relativistic corrections to the TRK sum rules.

given in atomic units ( $\hbar \rightarrow 1$ ,  $m \rightarrow 1$ ,  $e \rightarrow 1$ ), where  $Z$  is the number of protons,  $\alpha$  is the fine structure constant,  $n = 0, 1, 2, \dots$ , and  $V = -Z/r$ . We can simplify the example even further by assuming a single energy level from averaging the  $j = l \pm 1/2$  splitting for  $l \neq 0$ . To make this simplification, the transition probabilities for the  $1s_{1/2}$ - $1p_{1/2}$  and  $1s_{1/2}$ - $1p_{3/2}$  doublet as well as for the second transition's  $2p_{1/2}$ - $3p_{3/2}$ ,  $2p_{3/2}$ - $3p_{3/2}$ , and  $2p_{3/2}$ - $3p_{5/2}$  multiplet are evaluated from the Dirac equation [62–64], and used to perform the weighted averages for the excited state energies.

The  $\lambda_{kn}$  terms are then given by

$$\lambda_{nk}^{\text{H-like}} \approx \delta_{k,n} - \alpha^2 p_{kn}^2, \quad (42)$$

where the off-diagonal terms in Eq. (42) are taken to be zero under the current set of approximations with energy perturbations from a three-dimensional central potential.

The second hyperpolarizability can now be calculated by substituting the approximate  $Z$ -dependent coefficients  $E_{10}^{\text{H-like}}$ ,  $E^{\text{H-like}}$ ,  $X^{\text{H-like}}$ ,  $\lambda_{01}^{\text{H-like}}$ , and  $\lambda_{02}^{\text{H-like}}$ , into Eq. (29). The ratio of the second hyperpolarizability for H-like atoms for the  $z$ -diagonal tensor component,  $\lambda_{zzzz}^{\text{H-like}}$ , divided by the

approximate second hyperpolarizability of the hydrogen atom,  $\lambda_{zzzz}^{\text{hydrogen}}$ , is given in Fig. 4 as a function of the atomic number. The total strength of the transitions decrease causing a drop in the static nonlinear optical response. Note that the severity of damping to the second nonlinear response is lessened by a decrease in the first transition energy as the atomic number increases.

#### IV. CONCLUSION

The lowest-order relativistic correction to the TRK sum rules was shown to limit the oscillator strength below the value derived from the nonrelativistic Hamiltonian. This correction was applied to both the static linear and first two nonlinear optical responses; the magnitude of this correction is no longer a constant and depends on the potential energy function. This lowest-order relativistic correction has been applied to the three-level ansatz, where in the relativistic regime, the magnitudes of the fundamental limits of the polarizability, hyperpolarizability, and second hyperpolarizability are reduced. Thus, the nonrelativistic regime gives the largest values of the fundamental limit for closed quantum systems.

In the regime where the relativistic parameters pull the hyperpolarizability, at the positive fundamental limit, to below the negative bound, we find that it may be possible to break the Kuzyk limit (although with opposite sign). This is a disturbing result and further studies with higher degrees of accuracy must be performed for this consequence to be supported. Further studies with additional corrections may also help in understanding the peculiar influences of the off-diagonal sum rules on the linear and nonlinear responses. Originally, these off-diagonal terms were equal to zero in the nonrelativistic limit, where they are the primary reason that the Kuzyk limit of the hyperpolarizability could possibly be broken when referencing to the lowest-order correction.

#### ACKNOWLEDGMENTS

I would like to thank Professor Kenneth D. Singer and Professor Mark G. Kuzyk for useful discussions. I would also like to thank the National Science Foundation, Grant No. OISE-1243313, for partial support of this project.

- [1] M. G. Kuzyk, Physical limits on electronic nonlinear molecular susceptibilities, *Phys. Rev. Lett.* **85**, 1218 (2000).
- [2] W. Thomas, Über die zahl der dispersionselektronen, die einem stationären Zustände zugeordnet sind. (vorläufige Mitteilung), *Naturwissenschaften* **13**, 627 (1925).
- [3] F. Reiche and W. Thomas, Über die Zahl der Dispersionselektronen, die einem stationären Zustand zugeordnet sind, *Z. Phys.* **34**, 510 (1925).
- [4] W. Kuhn, Über die Gesamtstärke der von einem Zustände ausgehenden Absorptionslinien, *Z. Phys.* **33**, 408 (1925).
- [5] M. G. Kuzyk, Fundamental limits on third-order molecular susceptibilities, *Opt. Lett.* **25**, 1183 (2000).
- [6] M. G. Kuzyk, Quantum limits of the hyper-Rayleigh scattering susceptibility, *IEEE J. Sel. Top. Quantum Electron.* **7**, 774 (2001).
- [7] K. Clays, Theoretical upper limits and experimental overestimates for molecular hyperpolarizabilities: A symbiosis, *Opt. Lett.* **26**, 1699 (2001).
- [8] M. G. Kuzyk, Fundamental limits on two-photon absorption cross-sections, *J. Chem. Phys.* **119**, 8327 (2003).
- [9] K. Clays and B. J. Coe, Design strategies versus limiting theory for engineering large second-order nonlinear optical polarizabilities in charged organic molecules, *Chem. Mater.* **15**, 642 (2003).
- [10] K. Tripathi, P. Moreno, M. G. Kuzyk, B. J. Coe, K. Clays, and A. M. Kelley, Why hyperpolarizabilities Fall short of the fundamental quantum limits, *J. Chem. Phys.* **121**, 7932 (2004).
- [11] J. Pérez Moreno and M. G. Kuzyk, Fundamental limits of the dispersion of the two-photon absorption cross section, *J. Chem. Phys.* **123**, 194101 (2005).

- [12] M. G. Kuzyk, Compact sum-over-states expression without dipolar terms for calculating nonlinear susceptibilities, *Phys. Rev. A* **72**, 053819 (2005).
- [13] M. G. Kuzyk, Truncated sum rules and their use in calculating fundamental limits of nonlinear susceptibilities, *J. Nonl. Opt. Phys. & Mat.* **15**, 77 (2006).
- [14] M. G. Kuzyk, Fundamental limits of all nonlinear-optical phenomena that are representable by a second-order susceptibility, *J. Chem. Phys.* **125**, 154108 (2006).
- [15] J. Pérez Moreno, Y. Zhao, K. Clays, and M. G. Kuzyk, Modulated conjugation as a means for attaining a record high intrinsic hyperpolarizability, *Opt. Lett.* **32**, 59 (2007).
- [16] J. Pérez Moreno, I. Asselberghs, Y. Zhao, K. Song, H. Nakanishi, S. Okada, K. Nogi, O.-K. Kim, J. Je, J. Matrai, M. De Mayer, and M. G. Kuzyk, Combined molecular and supramolecular bottom-up nano-engineering for enhanced nonlinear optical response: Experiments, modelling and approaching the fundamental limit, *J. Chem. Phys.* **126**, 074705 (2007).
- [17] X. Pérez Moreno, K. Clays, and M. G. Kuzyk, A new dipole-free sum-over-states expression for the second hyperpolarizability, *J. Chem. Phys.* **128**, 084109 (2008).
- [18] J. Zhou and M. G. Kuzyk, Intrinsic Hyperpolarizabilities as a figure of merit for electro-optic molecules, *J. Phys. Chem. C* **112**, 7978 (2008).
- [19] M. G. Kuzyk, Using fundamental principles to understand and optimize nonlinear-optical materials, *J. Mater. Chem.* **19**, 7444 (2009).
- [20] N. J. Dawson, B. R. Anderson, J. L. Schei, and M. G. Kuzyk, Classical model of the upper bounds of the cascading contribution to the second hyperpolarizability, *Phys. Rev. A* **84**, 043406 (2011).
- [21] N. J. Dawson, B. R. Anderson, J. L. Schei, and M. G. Kuzyk, Quantum mechanical model of the upper bounds of the cascading contribution to the second hyperpolarizability, *Phys. Rev. A* **84**, 043407 (2011).
- [22] S. Shafei and M. G. Kuzyk, Paradox of the many-state catastrophe of fundamental limits and the three-state conjecture, *Phys. Rev. A* **88**, 023863 (2013).
- [23] M. G. Kuzyk, J. Pérez-Moreno, and S. Shafei, Sum rules and scaling in nonlinear optics, *Phys. Rep.* **529**, 297 (2013).
- [24] J. Zhou, M. G. Kuzyk, and D. S. Watkins, Pushing the hyperpolarizability to the limit, *Opt. Lett.* **31**, 2891 (2006).
- [25] J. Zhou, U. B. Szafruga, D. S. Watkins, and M. G. Kuzyk, Optimizing potential energy functions for maximal intrinsic hyperpolarizability, *Phys. Rev. A* **76**, 053831 (2007).
- [26] G. A. Wiggers and R. G. Petschek, Comment on "Pushing the hyperpolarizability to the limit," *Opt. Lett.* **32**, 942 (2007).
- [27] M. C. Kuzyk and M. G. Kuzyk, Monte Carlo studies of the fundamental limits of the intrinsic hyperpolarizability, *J. Opt. Soc. Am. B* **25**, 103 (2008).
- [28] D. S. Watkins and M. G. Kuzyk, Optimizing the hyperpolarizability tensor using external electromagnetic fields and nuclear placement, *J. Chem. Phys.* **131**, 064110 (2009).
- [29] S. Shafei, M. C. Kuzyk, and M. G. Kuzyk, Monte Carlo studies of the intrinsic second hyperpolarizability, *J. Opt. Soc. Am. B* **27**, 1849 (2010).
- [30] D. S. Watkins and M. G. Kuzyk, The effect of electron interactions on the universal properties of systems with optimized off-resonant intrinsic hyperpolarizability, *J. Chem. Phys.* **134**, 094109 (2011).
- [31] T. J. Atherton, J. Lesnifsky, G. A. Wiggers, and R. G. Petschek, Maximizing the hyperpolarizability poorly determines the potential, *J. Opt. Soc. Am. B* **29**, 513 (2012).
- [32] C. J. Burke, T. J. Atherton, J. Lesnifsky, and R. G. Petschek, Optimizing the second hyperpolarizability with minimally parametrized potentials, *J. Opt. Soc. Am. B* **30**, 1438 (2013).
- [33] J. M. Cole, R. C. B. Copley, G. J. McIntyre, J. A. K. Howard, M. Szablewski, and G. H. Cross, Charge-density study of the nonlinear optical precursor DED-TCNQ at 20 K, *Phys. Rev. B* **65**, 125107 (2002).
- [34] J. M. Cole, Organic materials for second-harmonic generation: Advances in relating structure to function, *Phil. Trans. R. Soc. Lond.* **361**, 2751 (2003).
- [35] A. P. Higginbotham, J. M. Cole, M. A. Blood-Forsythe, and D. D. Hickstein, Identifying and evaluating organic nonlinear optical materials via molecular moments, *J. Appl. Phys.* **111**, 033512 (1993).
- [36] S. Shafei, R. Lytel, and M. G. Kuzyk, Geometry-controlled nonlinear optical response of quantum graphs, *J. Opt. Soc. Am.* **29**, 3419 (2012).
- [37] R. Lytel and M. G. Kuzyk, Dressed quantum graphs with optical nonlinearities approaching the fundamental limit, *J. Nonlinear Optic. Phys. Mater.* **22**, 1350041 (2013).
- [38] R. Lytel, S. Shafei, J. H. Smith, and M. G. Kuzyk, Influence of geometry and topology of quantum graphs on their nonlinear optical properties, *Phys. Rev. A* **87**, 043824 (2013).
- [39] B. J. Orr and J. F. Ward, Perturbation theory of the non-linear optical polarization of an isolated system, *Mol. Phys.* **20**, 513 (1971).
- [40] J. S. Levinger and C. G. Bethea, Relativistic corrections to the dipole sum rule, *Phys. Rev.* **106**, 1191 (1957).
- [41] P. T. Leung, M. L. Rustgi, and S. A. T. Long, Relativistic corrections to the Bethe sum rule, *Phys. Rev. A* **33**, 2827 (1986).
- [42] S. M. Cohen and P. T. Leung, General formulation of the semirelativistic approach to atomic sum rules, *Phys. Rev. A* **57**, 4994 (1998).
- [43] H. Sinky and P. T. Leung, Relativistic corrections to a generalized sum rule, *Phys. Rev. A* **74**, 034703 (2006).
- [44] L. L. Foldy and S. A. Wouthuysen, On the Dirac theory of spin 1/2 particles and its non-relativistic limit, *J. Opt. Soc. Am. B* **78**, 29 (1950).
- [45] W. Greiner, *Relativistic Quantum Mechanics. Wave Equations*, 3rd ed. (Springer, New York, 2000).
- [46] M. Douglas and N. M. Kroll, Quantum electrodynamic corrections to the fine structure of helium, *Ann. Phys.* **82**, 89 (1974).
- [47] B. A. Hess, Relativistic electronic-structure calculations employing a two-component no-pair formalism with external-field projection operators, *Phys. Rev. A* **33**, 3742 (1986).
- [48] T. Nakajima and K. Hirao, The higher-order DouglasKroll transformation, *J. Chem. Phys.* **113**, 7786 (2000).
- [49] M. Reiher, Relativistic DouglasKrollHess theory, *WIREs Comput. Mol. Sci.* **2**, 139 (2012).
- [50] E. van Lenthe, E. J. Baerends, and J. G. Snijders, Relativistic regular two-component Hamiltonians, *J. Chem. Phys.* **99**, 4597 (1993).
- [51] E. van Lenthe, E. J. Baerends, and J. G. Snijders, Relativistic total energy using regular approximations, *J. Chem. Phys.* **101**, 9783 (1994).



- [52] E. van Lenthe, R. v. Leeuwen, E. J. Baerends, and J. G. Snijders, Relativistic regular two-component Hamiltonians, *Int. J. Quantum Chem.* **57**, 281 (1996).
- [53] M. Filatov and D. Cremer, Analytic energy derivatives for regular approximations of relativistic effects applicable to methods with and without correlation corrections, *J. Chem. Phys.* **118**, 6741 (2003).
- [54] M. Filatov and D. Cremer, Representation of the exact relativistic electronic Hamiltonian within the regular approximation, *J. Chem. Phys.* **119**, 11526 (2003).
- [55] W. Kutzelnigg and W. Liu, Quasirelativistic theory equivalent to fully relativistic theory, *J. Chem. Phys.* **123**, 241102 (2005).
- [56] W. Kutzelnigg and W. Liu, Quasirelativistic theory I. Theory in terms of a quasi-relativistic operator, *Mol. Phys.* **104**, 2225 (2006).
- [57] W. Liu and W. Kutzelnigg, Quasirelativistic theory. II. Theory at matrix level, *J. Chem. Phys.* **126**, 114107 (2007).
- [58] G. L. Sewell, Stark effect for a hydrogen atom in its ground state, *Math. Proc. Cambridge* **45**, 678 (1949).
- [59] L. L. Boyle, A. D. Buckingham, R. L. Disch, and D. A. Dunmur, Higher polarizability of the Helium atom, *J. Chem. Phys.* **45**, 1318 (1966).
- [60] D. P. Shelton, Hyperpolarizability of the hydrogen atom, *Phys. Rev. A* **36**, 3032 (1987).
- [61] D. J. Griffiths, *Introduction to Quantum Mechanics* (Prentice Hall, Upper Saddle River, NJ, 1995).
- [62] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One and Two Electron Atoms* (Plenum, New York, 1977).
- [63] R. H. Garstang, Relativistic radiative transitions, in *Topics in Modern Physics, A Tribute to Edward U. Condon*, edited by W. E. Brittin and H. Odabasi (Colorado Associated University Press, Boulder, 1971), pp. 153–167.
- [64] S. M. Younger and A. W. Weiss, Relativistic effects on line strengths in the hydrogenic isoelectronic for transitions sequence, *J. Res. Nat. Stand. Sec. A* **79A**, 629 (1975).