

Enhancing teleportation of quantum Fisher information by partial measurementsXing Xiao,¹ Yao Yao,² Wo-Jun Zhong,¹ Yan-Ling Li,³ and Ying-Mao Xie^{1,*}¹*College of Physics and Electronic Information, Gannan Normal University, Ganzhou Jiangxi 341000, China*²*Microsystems and Terahertz Research Center, China Academy of Engineering Physics, Chengdu, Sichuan 610200, China*³*School of Information Engineering, Jiangxi University of Science and Technology, Ganzhou, Jiangxi 341000, China*

(Received 14 October 2015; published 7 January 2016)

The purport of quantum teleportation is to completely transfer information from one party to another distant partner. However, from the perspective of parameter estimation, it is the information carried by a particular parameter, not the information of total quantum state that needs to be teleported. Due to the inevitable noise in environments, we propose two schemes to enhance quantum Fisher information (QFI) teleportation under amplitude damping noise with the technique of partial measurements. We find that post-partial measurement can greatly enhance the teleported QFI, while the combination of prior partial measurement and post-partial measurement reversal could completely eliminate the effect of decoherence. We show that, somewhat consequentially, enhancing QFI teleportation is more economic than that of improving fidelity teleportation. Our work extends the ability of partial measurements as a quantum technique to battle decoherence in quantum information processing.

DOI: [10.1103/PhysRevA.93.012307](https://doi.org/10.1103/PhysRevA.93.012307)**I. INTRODUCTION**

Quantum teleportation, one of the most fascinating protocols predicted by quantum mechanics [1], is a critical ingredient for quantum communication and quantum computation networks [2,3]. It is the faithful transfer of quantum states between two distant parties which have established prior entanglement and can communicate classically. In the past two decades, quantum teleportation has attracted considerable attention and has been studied both theoretically and experimentally [4–11]. However, in many scenarios, it is not the whole quantum state, but rather the information of a particular parameter physically encoded in it, that is needed to be transmitted. Therefore, there is no need to teleport the full information of the quantum states themselves, but only the relevant parameter information is our practical concern. In contrast to the quantum-state teleportation where the credibility of teleportation is measured by fidelity, the transmission of information that is carried by a physical parameter is usually quantified by quantum Fisher information (QFI) [12–16]. QFI plays a significant role in the fields of quantum geometry of state spaces [12,13,17], quantum information theory [18], and quantum metrology [19,20]. Particularly, the inverse of QFI characterizes the ultimate achievable precision in parameter estimation [21].

Unfortunately, any realistic quantum system inevitably couples to other uncontrollable environments which influence it in a non-negligible way [22]. Then the issue of robustness of QFI against various sources of decoherence has soon been raised. Numerous researchers have indeed demonstrated that QFI is fragile and easily broken by environmental noise [23–27]. This would be the most limiting factor for the applications of QFI in quantum teleportation, quantum metrology, and other quantum tasks. In this context, it is an extremely important issue to protect the QFI from decoherence during the

procedure of teleportation, especially for the atomic, trapped ions and other solid-state systems [6–9].

Partial measurements, which are generalizations of von Neumann measurements, are associated with positive-operator valued measures (POVM). For partial measurements [28–32], the information extracted from the quantum state is deliberately limited, thereby keeping the measured state alive (i.e., without completely collapsing towards an eigenstate). Thus, it would be possible to retrieve the initial information with some operations, even when the quantum state has suffered decoherence. Recently, many proposals that exploit partial measurements to protect the fidelity of a single qubit, the quantum entanglement of two qubits, and two qutrits from amplitude damping (AD) decoherence have been demonstrated both theoretically [33–35] and experimentally [36–38]. This motivates us to study the QFI teleportation under decoherence by utilizing the partial measurements.

In this paper we propose two schemes to show that partial measurements can greatly enhance the QFI teleportation under decoherence. In particular, we find that the combination of prior partial measurement and post partial measurement reversal is able to completely circumvent the influence of AD noise. Our schemes for enhancing QFI are based on the fact that partial measurements are nondestructive and can be reversed with a certain probability. Moreover, we analytically obtain the optimal parameters of the enhancement of QFI teleportation. We also demonstrate that the success probability of enhancing the QFI is higher than that of enhancing the fidelity of the quantum state, which indicates that enhancing the teleported QFI is a more reasonable and economic way in the scenario of parameter estimation.

This paper is organized as follows: we introduce QFI and partial measurements in Sec. II. In Sec. III, we then show how the QFI teleportation could be enhanced by partial measurements. We consider two different protocols, as illustrated in Fig. 1, and compare the results in Refs. [39,40], where the fidelity of teleportation is improved by partial measurements. We show that the cost of enhancing QFI is smaller than that of improving fidelity, which is represented

*xieyingmao@126.com




FIG. 1. Schematic illustrations of enhancing QFI teleportation under decoherence using partial measurements. (a) The circuit of scheme A is similar to the standard circuit of quantum teleportation, while a post partial measurement is added before the unitary operations. (b) In the circuit of scheme B, Alice performs a prior partial measurement (with strength p) on the second qubit before she sends it to Bob. During their classical communications, Alice has to send the results of Bell measurement and the prior partial measurement strength p to Bob. When Bob receives the results, he first performs a post partial measurement reversal (i.e., bit-flip, partial measurement, and bit-flip) and then sequentially performs the local unitary operations.

as the higher success probability. Finally, the conclusions are summarized in Sec. IV.

II. PRELIMINARIES

A. Quantum Fisher information

The QFI of a parameter characterizes the sensitivity of the state with respect to changes of the parameter. We consider that a parameter ϕ is encoded in quantum states ρ_ϕ (in general, mixed states). The QFI of ϕ is defined as [12,13,21]

$$\mathcal{F}_\phi = \text{Tr}(\rho_\phi \mathcal{L}_\phi^2) = \text{Tr}[(\partial_\phi \rho_\phi) \mathcal{L}_\phi], \quad (1)$$

where \mathcal{L}_ϕ is the so-called symmetric logarithmic derivative, which is defined by $\partial_\phi \rho_\phi = (\mathcal{L}_\phi \rho_\phi + \rho_\phi \mathcal{L}_\phi)/2$ with $\partial_\phi = \partial/\partial\phi$. Typically, there are three methods to calculate the QFI [41]. The most frequently used one is diagonalizing the matrix as $\rho_\phi = \sum_n \lambda_n |\psi_n\rangle\langle\psi_n|$. Then one can rewrite the QFI as [42,43]

$$\mathcal{F}_\phi = \sum_n \frac{(\partial_\phi \lambda_n)^2}{\lambda_n} + \sum_n \lambda_n \mathcal{F}_{\phi,n} - \sum_{n \neq m} \frac{8\lambda_n \lambda_m}{\lambda_n + \lambda_m} |\langle\psi_n|\partial_\phi\psi_m\rangle|^2, \quad (2)$$

where $\mathcal{F}_{\phi,n}$ is the QFI for the pure state $|\psi_n\rangle$ with the form $\mathcal{F}_{\phi,n} = 4[\langle\partial_\phi\psi_n|\partial_\phi\psi_n\rangle - |\langle\psi_n|\partial_\phi\psi_n\rangle|^2]$. According to Eq. (2), the last term stemming from the mixture of pure states suggests that the QFI of a mixed state is smaller than the pure-state case.

For the single-qubit state, a simple and explicit expression of QFI could be obtained. In the Bloch sphere representation,

any qubit state can be written as

$$\rho = \frac{1}{2}(1 + \vec{r} \cdot \hat{\sigma}), \quad (3)$$

where $\vec{r} = (r_x, r_y, r_z)^T$ is the real Bloch vector and $\hat{\sigma} = (\hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z)$ denotes the Pauli matrices. Therefore, for the single-qubit state, \mathcal{F}_ϕ can be represented as follows [44]:

$$\mathcal{F}_\phi = \begin{cases} |\partial_\phi \vec{r}|^2 + \frac{\vec{r} \cdot \partial_\phi \vec{r}}{1 - |\vec{r}|^2}, & \text{if } |\vec{r}| < 1, \\ |\partial_\phi \vec{r}|^2, & \text{if } |\vec{r}| = 1. \end{cases} \quad (4)$$

B. Partial measurements

In quantum physics, the standard von Neumann projective measurements are referred to as ‘‘sharp measurements,’’ which project the initial state to one of the eigenstates of the measurement operator. As generalizations of the standard von Neumann projective measurements, partial measurements don’t completely collapse the initial state (i.e., nonprojective measurements); hence they have the interesting property that they can be reversed in a probabilistic way. For a single qubit with computational basis $|0\rangle$ and $|1\rangle$, the so-called partial measurement is

$$\mathcal{M}_0 = |0\rangle\langle 0| + \sqrt{1-p}|1\rangle\langle 1|, \quad (5)$$

$$\mathcal{M}_1 = \sqrt{p}|1\rangle\langle 1|, \quad (6)$$

where parameter p ($0 \leq p \leq 1$) is usually named as the strength of partial measurement. Note that \mathcal{M}_0 and \mathcal{M}_1 are not necessarily projectors and are also nonorthogonal to each other, but $\mathcal{M}_0^\dagger \mathcal{M}_0 + \mathcal{M}_1^\dagger \mathcal{M}_1 = I$. Though the measurement operator \mathcal{M}_1 is the same as the von Neumann projective measurement and is associated with irrevocable collapse, \mathcal{M}_0 is a partial measurement, which we focus on in this paper.

C. Partial measurements reversal

The partial measurement \mathcal{M}_0 has some interesting properties: (i) the strength p is controllable and (ii) it could be reversed for the case $p \neq 1$. The reverse procedure can be noticed immediately to be

$$\begin{aligned} \mathcal{M}_0^{-1} &= |0\rangle\langle 0| + \frac{1}{\sqrt{1-p}}|1\rangle\langle 1| \\ &= \frac{1}{\sqrt{1-p}} X \mathcal{M}_0 X, \end{aligned} \quad (7)$$

where $X = |0\rangle\langle 1| + |1\rangle\langle 0|$ is the qubit bit-flip operation. The second line of Eq. (7) indicates that the reverse process \mathcal{M}_0^{-1} can be achieved physically by three sequential operations: bit-flip, partial measurement, and bit-flip.

III. ENHANCING QFI TELEPORTATION BY PARTIAL MEASUREMENTS

In realistic quantum teleportation, the maximally entangled state may lose its coherence and become a mixed state due to the interaction with its environment. Note that the quantum channel which is less entangled will reduce the teleported QFI. Here, we propose two schemes to enhance the QFI teleportation under AD noise, as shown in Fig. 1. For simplicity, we consider the scenario that Alice has a perfect

quantum memory while Bob doesn't. Alice prepares the Einstein-Podolsky-Rosen (EPR) state and then sends one particle to Bob. Therefore, the entangled part possessed by Bob will suffer AD noise and hence reduces the entanglement. The situation that both Alice and Bob are influenced by noise could be treated similarly.

A. Scheme A

We first examine the efficiency of scheme A, in which only one post partial measurement is performed by Bob before local operations (e.g., X or Z). We assume the shared EPR state is prepared in $|\Psi^+\rangle = (|00\rangle + |11\rangle)/\sqrt{2}$ by Alice and the input quantum state is an arbitrary superposition state,

$$|\psi_{\text{in}}\rangle = \cos\frac{\theta}{2}|0\rangle + \sin\frac{\theta}{2}e^{i\phi}|1\rangle, \quad (8)$$

which carries a phase parameter ϕ . Note that the QFI of ϕ is our concern during this teleportation procedure. The input state of the quantum teleportation circuit in Fig. 1(a) is the product state of $|\psi_{\text{in}}\rangle$ and $|\Psi^+\rangle$.

While the EPR state is prepared and stored perfectly by Alice, the qubit kept by Bob may be affected by AD noise. The dynamics of an entangled state subject to AD noise is described by the quantum operation Λ acting on the pure state [18]:

$$\begin{aligned} \rho^{AD} &= \Lambda(\rho_0) = \sum_{i=1,2} E_i |\Psi^+\rangle \langle \Psi^+| E_i^\dagger \\ &= \frac{1}{2} (|00\rangle \langle 00| + \gamma |10\rangle \langle 10| + \bar{\gamma} |11\rangle \langle 11| \\ &\quad + \sqrt{\bar{\gamma}} |00\rangle \langle 11| + \sqrt{\bar{\gamma}} |11\rangle \langle 00|), \end{aligned} \quad (9)$$

where $\rho_0 = |\Psi^+\rangle \langle \Psi^+|$ and the subscript AD denotes the pure AD noise case (i.e., without partial measurements). E_i are the Kraus operators of AD noise,

$$E_1 = I_1 \otimes \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1-\gamma} \end{pmatrix}, \quad E_2 = I_1 \otimes \begin{pmatrix} 0 & \sqrt{\bar{\gamma}} \\ 0 & 0 \end{pmatrix}, \quad (10)$$

where I_1 is the two-dimensional identity operator of the first qubit (Q1) since we have assumed Alice is not affected by AD noise. Note that following the standard circuit of the quantum teleportation process, we get the teleported QFI under AD noise:

$$\mathcal{F}_\phi^{AD} = \sin^2 \theta (1 - \gamma). \quad (11)$$

Then we consider the effect of a post partial measurement (with the strength p_r^A) performed by Bob before he makes the corresponding unitary operations. Here, the superscript A (B) indicates it belongs to the scheme A (B),

$$\begin{aligned} \rho^A &= M_0 [\Lambda(\rho_0)] M_0^\dagger \\ &= \frac{2}{1 + \bar{p}_r^A} (|00\rangle \langle 00| + \bar{p}_r^A \gamma |10\rangle \langle 10| + \bar{p}_r^A \bar{\gamma} |11\rangle \langle 11| \\ &\quad + \sqrt{\bar{p}_r^A \bar{\gamma}} |00\rangle \langle 11| + \sqrt{\bar{p}_r^A \bar{\gamma}} |11\rangle \langle 00|), \end{aligned} \quad (12)$$

where $M_0 = I_1 \otimes \mathcal{M}_0$. After the implementation of the quantum circuit of Fig. 1(a), Bob gets the teleported state ρ_{out}^A ,

whose three Bloch vector components are

$$r_x^A = \frac{2 \sin \theta \cos \phi}{N^A} \sqrt{\bar{p}_r^A \bar{\gamma}}, \quad (13)$$

$$r_y^A = \frac{2 \sin \theta \sin \phi}{N^A} \sqrt{\bar{p}_r^A \bar{\gamma}}, \quad (14)$$

$$r_z^A = \frac{\cos \theta}{N^A} (1 + \bar{p}_r^A) \bar{\gamma}, \quad (15)$$

where p_r^A is the strength of partial measurement, and $\bar{p}_r^A = 1 - p_r^A$, $\bar{\gamma} = 1 - \gamma$. $N^A = 2 - p_r^A - \gamma p_r^A$ is the normalized factor. According to Eq. (4), the QFI of the teleported state could be obtained,

$$\mathcal{F}_\phi^A = \frac{4 \sin^2 \theta \bar{p}_r^A \bar{\gamma}}{(N^A)^2}. \quad (16)$$

Since the strength of partial measurement is tunable, we can maximize the QFI by choosing the optimal partial measurement strength,

$$p_r^{A,\text{opt}} = 2\gamma/(1 + \gamma). \quad (17)$$

Then the maximal QFI is

$$\mathcal{F}_\phi^{A,\text{opt}} = \frac{\sin^2 \theta}{1 + \gamma}. \quad (18)$$

To quantify the efficiency of partial measurement on enhancing QFI, we introduce the teleported QFI improvement,

$$\mathcal{F}_{\text{imp}}^A \equiv \mathcal{F}_\phi^{A,\text{opt}} - \mathcal{F}_\phi^{AD}, \quad (19)$$

which is plotted in Fig. 2 as a function of decoherence strength γ and initial parameter θ . It is remarkable to find that the teleported QFI improvement $\mathcal{F}_{\text{imp}}^A$ is always non-negative. Particularly, when $\gamma = 1$, we find $\mathcal{F}_\phi^{AD} = 0$ while $\mathcal{F}_\phi^{A,\text{opt}} = \sin^2 \theta / 2$, which means that half of the QFI is still teleported under complete decoherence with the help of post partial measurement. Our result indicates that the application of quantum partial measurement indeed enhances the QFI




FIG. 2. The improved QFI $\mathcal{F}_{\text{imp}}^A$ by post partial measurement as a function of θ and dimensionless parameter γ .

teleportation, which is very important for quantum metrology and other quantum information tasks.

The underlying physical mechanism of the enhancement of QFI needs to be clarified further. One might deduce that the enhancement of teleported QFI should be attributed to the improvement of entanglement between Alice and Bob. We argue that this is not the case. According to Eqs. (9) and (12) and following the definition of concurrence [45], we can obtain the concurrence of ρ^{AD} and ρ^A , respectively, which are $C^{AD} = \sqrt{\bar{\gamma}}$ and $C^A = 2\sqrt{\bar{p}_r^A \bar{\gamma}/(1 + \bar{p}_r^A)}$. Employing the optimal partial measurement strength of QFI, i.e., Eq. (17), we have $C^{A,\text{opt}} = \bar{\gamma}\sqrt{1 + \bar{\gamma}}$. However, it is contrary to one's expectation that $C^{A,\text{opt}}/C^{AD} = \sqrt{1 - \bar{\gamma}^2} \leq 1$, which means that, on the optimal condition of teleported QFI, the entanglement between Alice of Bob is not improved with the help of post partial measurement. Therefore the enhancement of QFI in scheme A could not be attributed to the improvement of entanglement, but rather to the probabilistic nature of partial measurement.

In contrast to the results in Refs. [39,40], our ultimate aim is not to enhance the whole state fidelity but rather to improve the information of a particular parameter encoded in the teleported state (i.e., the QFI). Therefore, as one might expect, improving the QFI teleportation is easier than enhancing the fidelity. This intuition could be confirmed by comparing the success probability. In Refs. [39,40], the success probability of improving the fidelity is $P_{\text{fid}}^A = 1 - \frac{\gamma(3+\gamma)}{2(1+\gamma)}$, while in our scenario the success probability of enhancing QFI is

$$P_{\text{QFI}}^A = 1 - \gamma. \quad (20)$$

Similarly, we define the success probability improvement P_{imp}^A and it is easy to check that

$$P_{\text{imp}}^A \equiv P_{\text{QFI}}^A - P_{\text{fid}}^A = \frac{\gamma(1 - \gamma)}{2(1 + \gamma)} \geq 0, \quad (21)$$

as $\gamma \in [0,1]$. This means if we focus on teleporting the information of a particular parameter, we have no need to improve the teleportation fidelity of the whole state. Enhancing the QFI teleportation would be more reasonable and economic in this case.

B. Scheme B

Although a post partial measurement can enhance the QFI teleportation, part of QFI is still lost in the decoherence process. This is not our ultimate purpose. In this section, we show that an improved scheme can completely circumvent the decoherence and retrieve all the initial QFI. The key improvement is that Alice performs a prior partial measurement (with the measurement strength p) on the second qubit (Q2) before she sends it to Bob. During their classical communications, Alice has to send the results of Bell measurement and the information of p to Bob. When Bob receives the results, he firstly carries out a post partial measurement reversal (i.e., bit-flip, partial measurement and bit-flip) and then sequentially performs the local unitary operations. Remarkably, we show below that the prior partial measurement plays a significant role in enhancing the QFI teleportation.

According to the circuit depicted in Fig. 1(b), the second qubit (Q2) experiences three processes: prior partial measurement, AD noise, and post partial measurement reversal. The final entangled state could be represented as $\rho^B = M_0^{-1}[\Lambda(M_0\rho_0M_0^\dagger)](M_0^{-1})^\dagger$, where $M_0^{-1} = I_1 \otimes \mathcal{M}_0^{-1}$. The following QFI teleportation is based on the above entangled state and the final teleported state is ρ_{out}^B . The corresponding three Bloch vector components are as follows:

$$r_x^B = \frac{2 \sin \theta \cos \phi}{N^B} \sqrt{\bar{p} \bar{p}_r^B \bar{\gamma}}, \quad (22)$$

$$r_y^B = \frac{2 \sin \theta \sin \phi}{N^B} \sqrt{\bar{p} \bar{p}_r^B \bar{\gamma}}, \quad (23)$$

$$r_z^B = \frac{\cos \theta}{N^B} (\bar{p}_r^B \bar{\gamma} + \bar{p} \bar{\gamma} + p\gamma \bar{p}_r^B), \quad (24)$$

with $\bar{p} = 1 - p$ and $N^B = \bar{p}_r^B + \bar{p} \bar{\gamma} + \bar{p} \gamma \bar{p}_r^B$. The teleported QFI is now given by

$$\mathcal{F}_\phi^B = \frac{4 \sin^2 \theta \bar{p} \bar{p}_r^B \bar{\gamma}}{(N^B)^2}. \quad (25)$$

The optimal post measurement strength p_r^B could be obtained by calculating the following conditions: $\partial \mathcal{F}_\phi^B / \partial p_r^B = 0$ and $\partial^2 \mathcal{F}_\phi^B / (\partial p_r^B)^2 < 0$. The result turns out to be

$$p_r^{B,\text{opt}} = 1 - \frac{\bar{p} \bar{\gamma}}{1 + \bar{p} \gamma}, \quad (26)$$

and the maximally teleported QFI is

$$\mathcal{F}_\phi^B = \frac{\sin^2 \theta}{1 + \bar{p} \gamma}. \quad (27)$$

Hence it can be seen clearly that Eqs. (26) and (27) reduce to Eqs. (17) and (18) if we set $p = 0$ (i.e., without the prior partial measurement). However, in contrast to scheme A, the most intriguing result is that the existence of prior partial measurement provides the possibility of dramatically enhancing the teleported QFI. As shown in Fig. 3(a), with the combination of prior and post partial measurement, the QFI could be greatly recovered by adjusting the partial measurement strength p . Particularly, when $p \rightarrow 1$, $\mathcal{F}_\phi^B \rightarrow \sin^2 \theta$ without regard to the strength of AD noise.

By comparing these two schemes, it is easy to conclude that scheme B is much more efficient than scheme A on enhancing the QFI teleportation. However, why does scheme B work much better than scheme A with the assistance of a prior partial measurement? The underlying physics could be understood as follows: from Eq. (5), we know that the stronger the partial measurement strength p , the closer the initial EPR state is reversed towards the $|00\rangle$ state which is immune to AD noise. In scheme A, no prior partial measurement is carried out before the qubit goes through the AD noise; thus the amount of reversed QFI highly depends on the decoherence strength γ . While in the second scheme, a prior partial measurement is performed to move the state towards $|00\rangle$, which does not experience AD decoherence. Then an optimal partial measurement reversal is applied to revert the qubit back to the initial state. Therefore, the teleported QFI is not related to the decoherence strength γ but depends on the prior partial measurement strength p . Full QFI can entirely be recovered by




FIG. 3. (a) Teleported QFI $\mathcal{F}_\phi^{B,\text{opt}}$, $\mathcal{F}_\phi^{A,\text{opt}}$, and \mathcal{F}_ϕ^{AD} as a function of dimensionless parameters p and γ . The three surfaces from top to bottom correspond to scheme B , scheme A , and pure AD noise cases, respectively. Note that $\mathcal{F}_\phi^{A,\text{opt}}$ and \mathcal{F}_ϕ^{AD} are independent of p so they are planes. The parameter θ has been set to $\pi/2$. (b) The success probability improvement P_{imp}^B as a function of dimensionless parameters p and γ .

the combination of prior partial measurement and post partial measurement reversal when $p \rightarrow 1$.

The success probability of scheme B is given by

$$P_{\text{QFI}}^B = (1-p)(1-\gamma), \quad (28)$$

which obviously decays with the increase of p and γ . Note that when $p \rightarrow 1$, $P_{\text{QFI}}^B \rightarrow 0$. This means that the complete retrieval of QFI is attained at the expense of infinite low success probability. Nevertheless, the price of recovering QFI is still smaller than that of fidelity. As discussed in Ref. [39], the success probability of fidelity teleportation is

$$P_{\text{fid}}^B = \frac{\bar{\gamma} \bar{p}(2 + \gamma \bar{p})}{2(1 + \gamma \bar{p})}. \quad (29)$$

From Fig. 3(b), we note that $P_{\text{imp}}^B \equiv P_{\text{QFI}}^B - P_{\text{fid}}^B \geq 0$, which means the success probability of enhancing QFI is always higher than that of enhancing fidelity, except for boundary values of p and γ . This further confirms our conclusion that enhancing QFI is more economic than enhancing fidelity.

C. Generalizations

In the above analyses, we have assumed that Alice is not affected by noises. In fact, these two schemes are universal for the case that both Alice and Bob suffer AD noise. Since scheme A is a reduced version of scheme B , we only consider the latter situation as an example. Here, Alice has to make prior partial measurements separately on Q1 and Q2 and then sends Q2 to Bob. It should be noted that Alice must act on the post partial measurement before she does the Bell measurement, while Bob should perform the post partial measurement after he has received the results sent by Alice. The final teleported state can be characterized by the following Bloch vectors:

$$r_x = \frac{2 \sin \theta \cos \phi}{N} \sqrt{\bar{p}_1 \bar{p}_2 \bar{\gamma}_1 \bar{\gamma}_2 \bar{p}_{r_1} \bar{p}_{r_2}}, \quad (30)$$

$$r_y = \frac{2 \sin \theta \sin \phi}{N} \sqrt{\bar{p}_1 \bar{p}_2 \bar{\gamma}_1 \bar{\gamma}_2 \bar{p}_{r_1} \bar{p}_{r_2}}, \quad (31)$$

$$r_z = \frac{\cos \theta}{N} \left[\bar{p}_{r_1} \bar{p}_{r_2} + \bar{p}_1 \bar{p}_2 \bar{p}_{r_1} \bar{p}_{r_2} \gamma_1 \gamma_2 + \bar{p}_1 \bar{p}_2 \bar{\gamma}_1 \bar{\gamma}_2 - \bar{p}_1 \bar{p}_2 (\gamma_1 \bar{\gamma}_2 \bar{p}_{r_1} + \bar{\gamma}_1 \gamma_2 \bar{p}_{r_2}) \right], \quad (32)$$

with the normalized factor $N = \bar{p}_{r_1} \bar{p}_{r_2} + \bar{p}_1 \bar{p}_2 \bar{p}_{r_1} \bar{p}_{r_2} \gamma_1 \gamma_2 + \bar{p}_1 \bar{p}_2 \bar{\gamma}_1 \bar{\gamma}_2 + \bar{p}_1 \bar{p}_2 (\gamma_1 \bar{\gamma}_2 \bar{p}_{r_1} + \bar{\gamma}_1 \gamma_2 \bar{p}_{r_2})$. With these equations in hand, we can calculate the teleported QFI and optimize the variables p_{r_1} and p_{r_2} . In order to simplify the calculations, we assume that both qubit 1 and 2 interact with the same environments, i.e., $\gamma_1 = \gamma_2 = \gamma$. Consequently, we have $p_{r_1} = p_{r_2} = p$ and $p_{r_1} = p_{r_2} = p_r$. The final results reduce to $p_r^{\text{opt}} = 1 - \bar{p} \bar{\gamma} / \sqrt{1 + \bar{p}^2 \gamma^2}$ and

$$\mathcal{F}_\phi^{\text{opt}} = \frac{\sin^2 \theta}{(\sqrt{1 + \bar{p}^2 \gamma^2} + \bar{p} \gamma)^2}. \quad (33)$$

From the above equation, we note that given the strength of AD noise γ , $\mathcal{F}_\phi^{\text{opt}}$ achieves the minimum value with $p = 0$. Namely, no prior partial measurement is performed, which corresponds to the method of scheme A . Moreover, the minimum value of $\mathcal{F}_\phi^{\text{opt}}$ is still larger than the pure AD noise case, which indicates that partial measurement indeed can be used for enhancing QFI teleportation, even when both Alice and Bob are subject to AD noise. If $p \neq 0$, i.e., prior partial measurements are carried out before the qubits undergo AD noise, the teleported QFI is further enhanced. Particularly, when $p \rightarrow 1$, the initial QFI is almost entirely teleported to Bob.

IV. DISCUSSION AND CONCLUSIONS

Before concluding, we point out that our proposals are entirely feasible with the present experimental techniques. The standard quantum teleportation has already been realized from photonic systems to trapped ion and atomic systems. The techniques of partial measurement have also developed rapidly in recent years [36,38,46]. As demonstrated in Ref. [38], the partial measurements can be implemented with a Brewster-angle glass plate (BAGP) for photon-polarization qubits, because the BAGP probabilistically rejects vertical polarization

($|1\rangle$ state) and completely transmits horizontal polarization ($|0\rangle$ state), which functions exactly as the measurement depicted by Eqs. (5) and (6). On the other hand, the post weak measurement reversal could be realized by sequential bit-flip (a half wave plate for polarization qubit), partial measurement, and bit-flip. Alternatively, the partial measurements could be described with projective measurements in a larger Hilbert space that includes an “ancilla qubit” [31,32]. Hence, the performance of partial measurement on the target qubit is equivalent to the action of von Neumann projective measurement on the ancilla qubit which is previously coupled to it. Motivated by this guideline, partial measurements can be realized in any quantum system and are not restricted to photon and superconducting qubits.

In summary, we propose the enhancement of QFI teleportation under decoherence utilizing partial measurements.

Thanks to the probabilistic nature of partial measurements, the teleported QFI could be greatly enhanced with the assistance of post partial measurement. Remarkably, we further show that the combined action of prior and post partial measurement can even completely keep QFI from AD noise. In addition, we demonstrate that the price of enhancing teleported QFI is smaller than that of improving fidelity. Our work extends the ability of partial measurements as a technique in various quantum information processing tasks, particularly when the research objects are subject to AD noise.

ACKNOWLEDGMENTS

This work is supported by the Funds of the National Natural Science Foundation of China under Grants No. 11247006 and No. 11365011.

-
- [1] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, *Phys. Rev. Lett.* **70**, 1895 (1993).
- [2] D. Gottesman and I. Chuang, *Nature (London)* **402**, 390 (1999).
- [3] R. Ursin, T. Jennewein, M. Aspelmeyer, R. Kaltenbaek, M. Lindenthal, P. Walther, and A. Zeilinger, *Nature (London)* **430**, 849 (2004).
- [4] S. L. Braunstein and H. J. Kimble, *Phys. Rev. Lett.* **80**, 869 (1998).
- [5] D. Bouwmeester, J. W. Pan, K. Mattle, M. Eibl, H. Weinfurter, and A. Zeilinger, *Nature (London)* **390**, 575 (1997).
- [6] M. Riebe, H. Häffner, C. F. Roos, W. Hänsel, J. Benhelm, G. P. T. Lancaster, T. W. Körber, C. Becher, F. Schmidt-Kaler, D. F. V. James *et al.*, *Nature (London)* **429**, 734 (2004).
- [7] M. D. Barrett, J. Chiaverini, T. Schaetz, J. Britton, W. M. Itano, J. D. Jost, E. Knill, C. Langer, D. Leibfried, R. Ozeri *et al.*, *Nature (London)* **429**, 737 (2004).
- [8] S. Olmschenk, D. N. Matsukevich, P. Maunz, D. Hayes, L. M. Duan, and C. Monroe, *Science* **323**, 486 (2009).
- [9] W. Pfaff, B. J. Hensen, H. Bernien, S. B. van Dam, M. S. Blok, T. H. Taminiau, M. J. Tiggelman, R. N. Schouten, M. Markham, D. J. Twitchen *et al.*, *Science* **345**, 532 (2014).
- [10] X.-L. Wang, X.-D. Cai, Z.-E. Su, M.-C. Chen, D. Wu, L. Li, N.-L. Liu, C.-Y. Lu, and J.-W. Pan, *Nature (London)* **518**, 516 (2015).
- [11] H. Takesue, S. D. Dyer, M. J. Stevens, V. Verma, R. P. Mirin, and S. W. Nam, *Optica* **2**, 832 (2015).
- [12] C. W. Helstrom, *Quantum Detection and Estimation Theory* (Academic, New York, 1976).
- [13] A. S. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory* (North-Holland, Amsterdam, 1982).
- [14] S. M. Kay, *Fundamentals of Statistical Signal Processing: Estimation Theory* (Prentice Hall, Upper Saddle River, NJ, 1993).
- [15] M. G. Genoni, S. Olivares, and M. G. A. Paris, *Phys. Rev. Lett.* **106**, 153603 (2011).
- [16] X. M. Lu, Z. Sun, X. Wang, S. Luo, and C. H. Oh, *Phys. Rev. A* **87**, 050302(R) (2013).
- [17] W. K. Wootters, *Phys. Rev. D* **23**, 357 (1981).
- [18] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2000).
- [19] V. Giovannetti, S. Lloyd, and L. Maccone, *Phys. Rev. Lett.* **96**, 010401 (2006).
- [20] V. Giovannetti, S. Lloyd, and L. Maccone, *Nat. Photon.* **5**, 222 (2011).
- [21] S. L. Braunstein and C. M. Caves, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [22] H. P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
- [23] J. Kołodziej and R. Demkowicz-Dobrzański, *Phys. Rev. A* **82**, 053804 (2010).
- [24] J. Ma, Y. X. Huang, X. Wang, and C. P. Sun, *Phys. Rev. A* **84**, 022302 (2011).
- [25] K. Berrada, S. Abdel-Khalek, and A. S. Obada, *Phys. Lett. A* **376**, 1412 (2012).
- [26] Y. M. Zhang, X. W. Li, W. Yang, and G. R. Jin, *Phys. Rev. A* **88**, 043832 (2013).
- [27] Y. L. Li, X. Xiao, and Y. Yao, *Phys. Rev. A* **91**, 052105 (2015).
- [28] A. N. Korotkov, *Phys. Rev. B* **60**, 5737 (1999).
- [29] A. N. Korotkov and A. N. Jordan, *Phys. Rev. Lett.* **97**, 166805 (2006).
- [30] A. N. Korotkov and K. Keane, *Phys. Rev. A* **81**, 040103(R) (2010).
- [31] G. S. Paraoanu, *Found. Phys.* **41**, 1214 (2011).
- [32] G. S. Paraoanu, *Europhys. Lett.* **93**, 64002 (2011).
- [33] Q. Q. Sun, M. Al-Amri, and M. S. Zubairy, *Phys. Rev. A* **80**, 033838 (2009).
- [34] Z. X. Man, Y. J. Xia, and N. B. An, *Phys. Rev. A* **86**, 012325 (2012).
- [35] X. Xiao and Y. L. Li, *Eur. Phys. J. D* **67**, 204 (2013).
- [36] N. Katz, M. Neeley, M. Ansmann, R. C. Bialczak, M. Hofheinz, E. Lucero, A. O’Connell, H. Wang, A. N. Cleland, J. M. Martinis, and A. N. Korotkov, *Phys. Rev. Lett.* **101**, 200401 (2008).
- [37] Y. S. Kim, Y. W. Cho, Y. S. Ra, and Y. H. Kim, *Opt. Express* **17**, 11978 (2009).
- [38] Y. S. Kim, J. C. Lee, O. Kwon, and Y. H. Kim, *Nat. Phys.* **8**, 117 (2012).
- [39] T. Pramanik and A. S. Majumdar, *Phys. Lett. A* **377**, 3209 (2013).

- [40] L. Qiu, G. Tang, X. Yang, and A. Wang, *Ann. Phys. (New York)* **350**, 137 (2014).
- [41] Y. Yao, L. Ge, X. Xiao, X. Wang, and C. P. Sun, *Phys. Rev. A* **90**, 062113 (2014).
- [42] S. Knysh, V. N. Smelyanskiy, and G. A. Durkin, *Phys. Rev. A* **83**, 021804(R) (2011).
- [43] J. Liu, X. Jing, and X. Wang, *Phys. Rev. A* **88**, 042316 (2013).
- [44] W. Zhong, Z. Sun, J. Ma, X. Wang, and F. Nori, *Phys. Rev. A* **87**, 022337 (2013).
- [45] W. K. Wootters, *Phys. Rev. Lett.* **80**, 2245 (1998).
- [46] G. S. Paraoanu, *Phys. Rev. Lett.* **97**, 180406 (2006).